

Free Full Text from Publisher

Look Up Full Text

Full Text from Publisher

Find PDF

Export...

Add to Marked List

1 of 6

## Investigating the Effects of Primary Amine Linkers with Different Carbon Chain Lengths on the Acid Dissociation Constant (pKa) for Covalently Immobilized Anthraquinone at the Electrode Surface Using Linear and Non-Linear Fittings

By: [Hamzah, HH](#) (Hamzah, Hairul Hisham)<sup>[1]</sup>; [Chein, WC](#) (Chein, Wee Chew)<sup>[1]</sup>; [Rahiman, SSF](#) (Rahiman, Siti Sarah Fazalul)<sup>[2]</sup>; [Shafiee, SA](#) (Shafiee,

JOURNAL OF THE ELECTROCHEMICAL SOCIETY

Volume: 166 Issue: 16 Pages: H877-H887

DOI: 10.1149/2.0301916jes

Published: DEC 6 2019

Document Type: Article

[View Journal Impact](#)

### Abstract

Electrografting of primary amine linkers is a widely used technique to covalently immobilize redox molecules or biomolecules at the electrode surface as it can produce a uniform and highly stable monolayers. Herein, we discussed the effects of having primary amine linkers with different carbon chain lengths as a medium to attach anthraquinone (AQ) at the electrode surface to study its acid dissociation constant (pKa); anthraquinone was covalently attached to glassy carbon (GC) electrodes in aqueous buffer solutions at different pH values ranging from pH 1 to pH 13. In this study, ethylenediamine (EDA, C2) and hexanediamine (HDA, C6) were used as the primary amine linkers. The pKa values of the surface-bound AQ were determined from the linear and non-linear fittings performed on the graphs of the mid potential (E-mid) of the redox peaks for AQ against the pH. The pKa values obtained from the linear fittings were using the value at the intersection point of two linear slopes between two pH gradients, whilst the pKa values for the non-linear fittings were determined according to the Nernst pH reduction theory. Overall, the non-linear fittings can provide us with a reliable and accurate approach in determining the pKa values of AQ. (C) The Author(s) 2019. Published by ECS.

### Keywords

**KeyWords Plus:** SOLID-PHASE SYNTHESIS; MODIFIED GLASSY-CARBON; ELECTROCATALYTIC REDUCTION; AQUEOUS ELECTROCHEMISTRY; OXYGEN REDUCTION; GOLD ELECTRODES; QUINONE; DERIVATIVES; MONOLAYERS; OXIDATION

### Author Information

**Reprint Address:** Hamzah, HH (reprint author)

+ USM, Sch Chem Sci, George Town 11800, Malaysia.

#### Addresses:

+ [ 1 ] USM, Sch Chem Sci, George Town 11800, Malaysia

+ [ 2 ] USM, Sch Pharmaceut Sci, George Town 11800, Malaysia

+ [ 3 ] Univ Texas Austin, Sch Chem, Austin, TX 78712 USA

**E-mail Addresses:** [hishamhamzah@usm.my](mailto:hishamhamzah@usm.my)

### Funding

Funding Agency	Show details	Grant Number
Universiti Sains Malaysia		304/PKIMIA/6315208

[View funding text](#)

### Publisher

ELECTROCHEMICAL SOC INC, 65 SOUTH MAIN STREET, PENNINGTON, NJ 08534 USA

### Journal Information

**Impact Factor:** [Journal Citation Reports](#)

### Categories / Classification

**Research Areas:** Electrochemistry; Materials Science

**Web of Science Categories:** Electrochemistry; Materials Science, Coatings & Films

[See more data fields](#)

1 of 6

### Citation Network

In Web of Science Core Collection

0

Times Cited

[Create Citation Alert](#)

70

Cited References

[View Related Records](#)

### Use in Web of Science

Web of Science Usage Count

0

Last 180 Days

0

Since 2013

[Learn more](#)

This record is from:

Web of Science Core Collection

- Science Citation Index Expanded

[Suggest a correction](#)

If you would like to improve the quality of the data in this record, please [suggest a correction](#).

### Cited References: 70

Showing 30 of 70 [View All in Cited References page](#)

(from Web of Science Core Collection)