

Contents

| | |
|--|-----|
| Review Article Streptococcosis in Tilapia (<i>Oreochromis niloticus</i>): A Review <i>Amal, M.N.A. and Zamri-Saad, M.</i> | 195 |
| Regular Articles Effects of Deficit Irrigation on Water Productivity and Maize Yields in Arid Regions of Iran <i>Hanidreza Salemi, Mohd, Amin Mohd, Soom, Teang Shui Lee, Mohd Kamil Yusoff and Desa Ahmad</i> | 207 |
| Trans-scrotal Ultrasonography and Breeding Soundness Evaluation of Bulls in a Herd of Dairy and Beef Cattle with Poor Reproductive Performance <i>N. Yamer, Y. Rosnina, H. Wahid, A.A. Soharvee, K.C. Yap, P. Ganesamurthi and M.M. Fahmi</i> | 217 |
| Relationships of Distribution of Macroinvertebrates and the Physico-chemical Parameters from Semenyih River by Using Correlation and Multiple Linear Stepwise Regression Analyses <i>Yap, C.K. and Rahim Ismail, A.</i> | 229 |
| Carcass Compositions in Three Different Breeds of Chicken and Their Correlation with Growth Performance <i>Lokman, I.H., Zaki, A.B.Z., Goh, Y.M., Sazili, A.Q. and Noordin, M.M.</i> | 247 |
| Antioxidant Study of Garlic and Red Onion: A Comparative Study <i>Siti Fairuz Che Othman, Syed Zahir Idd, Mastapha Saleiman Koya, Aisyah Mohamed Rehan and Kamarul Rahim Kamarudin</i> | 253 |
| Phylogenetic Relationships among Different Breeds of Domestic Chickens in Selected Areas of Peninsular Malaysia Using Rapid Markers <i>Yap Fook Choy and Jayaraj Vijayaraj Kumaran</i> | 263 |
| Effect of Seminal Plasma Removal, Washing Solutions, and Centrifugation Regimes on Boer Goat Semen Cryopreservation <i>Sue Win Nating, Abd Wahid Haron, Mohd Azam Khan Goriman, Rosnina Yusoff, Md Zaki Abu Bakar, Kazhal Sarsaifi, Muhammad Mada Bakar, Myint Thein, Than Kyaw and Maung Maung San</i> | 271 |
| Phylogenetic Analysis of the Malaysian <i>Rhinolopus</i> and <i>Hipposideros</i> Inferred from Partial Mitochondrial DNA Cytochrome b Gene Sequences <i>Sazali, S.N., Besar, K. and Abdullah, M.T.</i> | 281 |



Research Management Centre (RMC)
18 Floor, IDE'A Tower II
UPM-MITDC Technology Centre
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia
<http://www.rmc.upm.edu.my>
E-mail : rlc@upm.edu.my
Tel : +603 8947 1622/1620

UPM Press
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia

<http://penerbit.upm.edu.my>
E-mail : ace@lib@putra.upm.edu.my
Tel : +603 8946 8855/8854
Fax : +603 8941 0172



Pertanika JTAS
Vol. 34 (2) Aug. 2011
UPM Press, Malaysia

Pertanika Journal of
TROPICAL
AGRICULTURAL SCIENCE
JTAS

VOL. 34 (2) AUG. 2011



A scientific journal published by Universiti Putra Malaysia Press

Journal of Tropical Agricultural Science

About the Journal

Pertanika is an international peer-reviewed journal devoted to the publication of original papers, and it serves as a forum for practical approaches to improving quality in issues pertaining to tropical agriculture and its related fields. *Pertanika* Journal of Tropical Agricultural Science which began publication in 1978 is a leading agricultural journal in Malaysia. After 29 years as a multidisciplinary journal, the revamped *Pertanika* Journal of Tropical Agricultural Science (JTAS) is now focusing on tropical agricultural research. Other *Pertanika* series include *Pertanika* Journal of Science and Technology (JST) and *Pertanika* Journal of Social Sciences and Humanities (JSSH).

JTAS is published in **English** and it is open to authors around the world regardless of the nationality. It is currently published two times a year, i.e. in **February** and **August**.

Goal of *Pertanika*

Our goal is to bring the highest quality research to the widest possible audience.

Quality

We aim for excellence, sustained by a responsible and professional approach to journal publishing. Submissions are guaranteed to receive a decision within 12 weeks. The elapsed time from submission to publication for the articles averages 5-6 months.

Indexing of *Pertanika*

Pertanika is now over 30 years old; this accumulated knowledge has resulted in *Pertanika* journals being indexed in SCOPUS (Elsevier), EBSCO, CABI and AGRICOLA, etc.

Future vision

We are continuously improving access to our journal archives, content, and research services. We have the drive to realise exciting new horizons that will benefit not only the academic community, but society itself.

We also have views on the future of our journals. The emergence of the online medium as the predominant vehicle for the 'consumption' and distribution of much academic research will be the ultimate instrument in the dissemination of research news to our scientists and readers.

Aims and scope

Pertanika Journal of Tropical Agricultural Science aims to provide a forum for high quality research related to tropical agricultural research. Areas relevant to the scope of the journal include: *agricultural biotechnology, biochemistry, biology, ecology, fisheries, forestry, food sciences, genetics, microbiology, pathology and management, physiology, plant and animal sciences, production of plants and animals of economic importance, and veterinary medicine.*

Editorial Statement

Pertanika is the official journal of Universiti Putra Malaysia. The abbreviation for *Pertanika* Journal of Tropical Agricultural Science is *Pertanika J. Trop. Agric. Sci.*

Editorial Board

2011-2013

Editor-in-Chief

Soon Guan TAN, Malaysia
Molecular population genetics

Executive Editor

Nayan D.S. KANWAL, Malaysia
Environmental issues- landscape plant modelling applications

Editorial Board Members

- Anuar Abd. Rahim** (Associate Professor Dr), *Soil Fertility and Management*, Universiti Putra Malaysia, Malaysia.
- Baharuddin Salleh** (Professor Dr), *Plant Pathologist/ Mycologist*, Universiti Sains Malaysia, Malaysia.
- Chee-Kong YAP** (Associate Professor Dr), *Biology, Ecotoxicology*, Universiti Putra Malaysia, Malaysia.
- David Edward BIGNELL** (Professor Dr), *Soil Biology and Termite Biology*, University of London, U.K.
- Eric STANBRIDGE** (Professor Dr), *Microbiology, Molecular Genetics*, University California, USA.
- Ghizan SALEH** (Professor Dr), *Plant Breeding and Genetics*, Universiti Putra Malaysia, Malaysia.
- Idris Abd. Ghani** (Professor Dr), *Entomology, Insect taxonomy and Biodiversity, Integrated Pest Management, Biological Control, Biopesticides*, Universiti Kebangsaan Malaysia, Malaysia.
- Jamilah BAKAR** (Professor Dr), *Food Science and Technology, Food Quality /Processing and Preservation*, Universiti Putra Malaysia, Malaysia.
- Kadambot H.M. SIDDIQUE, FTSE** (Winthrop Professor Dr), Chair in Agriculture and Director, UWA Institute of Agriculture, *Crop and Environment Physiology, Germplasm Enhancement*, The University of Western Australia, Australia.
- Leng-Guan SAW** (Dr), *Botany and Conservation, Plant Ecology*, Forest Research Institute Malaysia (FRIM), Kepong, Malaysia.
- Mahani Mansor CLYDE** (Professor Dr), *Genetics, Cytogenetics*, Universiti Kebangsaan Malaysia, Malaysia.
- Mohd. Azmi AMBAK** (Professor Dr), *Fisheries and Aquaculture*, Universiti Malaysia Terengganu, Malaysia.
- Mohd. Zamri-Saad** (Professor Dr), *Veterinary Pathology*, Universiti Putra Malaysia, Malaysia.
- Nor Aini AB-SHUKOR** (Professor Dr), *Tree Improvement, Forestry Genetics and Biotechnology*, Universiti Putra Malaysia, Malaysia.
- Richard T. CORLETT** (Professor Dr), *Biological Sciences, Terrestrial Ecology, Climate Change, Conservation Biology, Biogeography*, National University of Singapore, Singapore.
- Shamshuddin Jusop** (Professor Dr), *Soil Science, Soil Mineralogy*, Universiti Putra Malaysia, Malaysia.
- Son RADU** (Professor Dr), *Food Safety, Risk Assessment, Molecular Biology*, Universiti Putra Malaysia, Malaysia.
- Srini KAVERI** (Dr), *Veterinary, Immunology*, INSERM, Centre de Recherche Cordeliers, Paris, France.
- Suman KAPUR** (Professor Dr), *Biological Sciences, Agricultural and Animal Biotechnology, Biochemistry*, Birla Institute of Technology and Science (BITS-Pilani), Hyderabad, India.
- Wen-Siang TAN** (Professor Dr), *Molecular Biology, Virology, Protein Chemistry*, Universiti Putra Malaysia, Malaysia.
- Yusof IBRAHIM** (Professor Dr), *Agricultural Entomology*, Universiti Pendidikan Sultan Idris, Malaysia.

International Advisory Board

Alexander SALENIKOVICH (Associate Professor Dr), *Forestry, Wood and Forest Sciences*, Université Laval, Canada.

Banpot NAPOMPETH (Professor Dr), *Entomology*, Kasetsart University, Thailand.

Denis J. WRIGHT (Professor Dr), *Pest Management*, Imperial College London, U.K.

Graham MATTHEWS (Professor Emeritus Dr), *Pest Management*, Imperial College London, U.K.

Jane M. HUGHES (Professor Dr), *Genetics*, Griffith University, Australia.

Malcolm WALKINSHAW (Professor Dr), *Biochemistry*, University of Edinburgh, Scotland.

Manjit S. KANG (Professor Emeritus Dr), Vice Chancellor, PAU, *Plant Breeding and Genetics*, Vice Chancellor, Punjab Agricultural University, India and Emeritus Professor, Louisiana State University Agric. Center, Baton Rouge, USA.

Peter B. MATHER (Professor Dr), *Ecology and Genetics*, Queensland University of Technology, Australia.

Syed M. ILYAS (Professor Dr), Project Director, National Institute of Rural Development, *Post Harvest Engineering and Technology*, Indian Council of Agricultural Research, Hyderabad, India.

Tanveer N. KHAN (Professor Dr), *Plant Breeding and Genetics*, Department of Agriculture and Food, South Perth, Western Australia.

Pertanika Editorial Office

Office of the Deputy Vice Chancellor (R&I)
1st Floor, IDEA Tower II, UPM-MTDC Technology Centre
Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia
Tel: +603 8947 1622
E-mail: ndeeps@admin.upm.edu.my

Publisher

The UPM Press
Universiti Putra Malaysia
43400 UPM, Serdang, Selangor, Malaysia
Tel: +603 8946 8855, 8946 8854 • Fax: +603 8941 6172
penerbit@putra.upm.edu.my
URL: <http://penerbit.upm.edu.my>

The publisher of *Pertanika* will not be responsible for the statements made by the authors in any articles published in the journal. Under no circumstances will the publisher of this publication be liable for any loss or damage caused by your reliance on the advice, opinion or information obtained either explicitly or implied through the contents of this publication.

All rights of reproduction are reserved in respect of all papers, articles, illustrations, etc., published in *Pertanika*. *Pertanika* provides free access to the full text of research articles for anyone, web-wide. It does not charge either its authors or author-institution for refereeing/ publishing outgoing articles or user-institution for accessing incoming articles.

No material published in *Pertanika* may be reproduced or stored on microfilm or in electronic, optical or magnetic form without the written authorization of the Publisher.

Copyright © 2011 Universiti Putra Malaysia Press. All Rights Reserved.

Pertanika Journal of Tropical Agricultural Science
Vol. 34 (2) Aug. 2011

Contents

Review Article

- Streptococcosis in Tilapia (*Oreochromis niloticus*): A Review 195
Amal, M.N.A. and Zamri-Saad, M.

Regular Articles

- Effects of Deficit Irrigation on Water Productivity and Maize Yields in Arid Regions of Iran 207
Hamidreza Salemi, Mohd. Amin Mohd. Soom, Teang Shui Lee, Mohd Kamil Yusoff and Desa Ahmad
- Trans-scrotal Ultrasonography and Breeding Soundness Evaluation of Bulls in a Herd of Dairy and Beef Cattle with Poor Reproductive Performance 217
N. Yimer, Y. Rosnina, H. Wahid, A.A. Saharee, K.C. Yap, P. Ganesamurthi and M.M. Fahmi
- Relationships of Distribution of Macrobenthic Invertebrates and the Physico-chemical Parameters from Semenyih River by Using Correlation and Multiple Linear Stepwise Regression Analyses 229
Yap, C.K. and Rahim Ismail, A.
- Carcass Compositions in Three Different Breeds of Chicken and Their Correlation with Growth Performance 247
Lokman, I.H., Zuki, A.B.Z., Goh, Y.M., Sazili, A.Q. and Noordin, M.M.
- Antioxidant Study of Garlic and Red Onion: A Comparative Study 253
Siti Fairuz Che Othman, Syed Zahir Idid, Mustapha Suleiman Koya, 'Aisyah Mohamed Rehan and Kamarul Rahim Kamarudin
- Phylogenetic Relationships among Different Breeds of Domestic Chickens in Selected Areas of Peninsular Malaysia Using Rapd Markers 263
Yap Fook Choy and Jayaraj Vijaya Kumaran
- Effect of Seminal Plasma Removal, Washing Solutions, and Centrifugation Regimes on Boer Goat Semen Cryopreservation 271
Soe Win Naing, Abd Wahid Haron, Mohd Azam Khan Goriman, Rosnina Yusoff, Md Zuki Abu Bakar, Kazhal Sarsaifi, Muhammad Modu Bukar, Myint Thein, Than Kyaw and Maung Maung San
- Phylogenetic Analysis of the Malaysian *Rhinolopus* and *Hipposideros* Inferred from Partial Mitochondrial DNA Cytochrome *b* Gene Sequences 281
Sazali, S.N., Besar, K. and Abdullah, M.T.

| | |
|---|-----|
| Management Practices Affecting Helminthiasis in Goats <i>Nor-Azlina, A.A., Sani, R.A. and Ariff, O.M.</i> | 295 |
| The Prevalence of Antimicrobial Resistant <i>Salmonella</i> spp. and the Risk Factors Associated with Their Occurrence in Finisher Pigs in Seberang Perai, Malaysia <i>D.W. Choe, L. Hassan and T.C. Loh</i> | 303 |
| Selected Articles from the Malaysian Biological Symposium 2009 | |
| Guest Editor: Tan Soon Guan | |
| Guest Editorial Board: Nur Ain Izzati Mohd. Zainudin and Omar Md. Yusoh | |
| Short Communications | |
| Microsatellite Markers for the Identification of Commercially Important Groupers <i>Epinephelus lanceolatus</i> , <i>Cromileptes altivelis</i> and <i>Epinephelus fuscoguttatus</i> <i>Rodrigues, K.F., Shigeharu, S. and Ch'ng, C.L.</i> | 311 |
| Regular Articles | |
| Sequence Variation in the Cellulose Synthase (<i>SpCesA1</i>) Gene from <i>Shorea parvifolia</i> ssp. <i>parvifolia</i> Mother Trees <i>Ho Wei Seng, Pang Shek Ling, Pauline Lau and Ismail Jusoh</i> | 317 |
| Isolation and Identification of <i>Fusarium</i> Species Associated with Fusarium Ear Rot Disease of Corn <i>Nur Ain Izzati Mohd Zainudin, Siti Nordahliawate Mohamed Sidique, Nor Azlina Johari, Darnetty, Azmi Abd. Razak and Baharuddin Salleh</i> | 325 |
| α -Tocopherol, Ascorbic Acid and Carotenoid Content in <i>Centella asiatica</i> Leaf Tissues and Callus Cultures <i>Norhayati, Y., Nor `Aini, M.F., Misri, K., Marziah, M. and Azman, J.</i> | 331 |
| Selected Articles from the 2nd Malaysian Association of Veterinary Pathology Conference 2010 | |
| Guest Editorial Board: Noordin Mohamed Mustapha, Mohd. Zamri Saad, Sabri Mohd. Yusof, Hazilawati Hamzah and Jasni Sabri | |
| Subchronic Oral Toxicity Study of <i>Morinda citrifolia</i> (Mengkudu) in Sprague Dawley Rats <i>Rosly, S.M., Shanmugavelu, S., Murugaiyah, M., Hadijah, H., Ahmad Tarmizi, S., Noridayusni, Y. and Subramaniam, K.</i> | 341 |
| Histopathological Features of Peripheral T-cell Lymphoma in Sprague Dawley Rats Induced with <i>N</i> -methyl- <i>N</i> -nitrosourea <i>A.H. Hutheyfa, H. Hazilawati, S.M. Rosly, S. Jasni, M.M. Noordin and S. Shanmugavelu</i> | 351 |
| Urolithiasis in Boer Bucks <i>M.S. Shahrom and M. Zamri-Saad</i> | 363 |

| | |
|--|-----|
| Pulmonary Lesions Associated with Intratracheal Benzo(a)pyrene Instillation in Sprague Dawley Rats <i>M. Mazlina, A.J. Karim, M. Zamri-Saad and M.M. Noordin</i> | 367 |
| Detection of Bcl-2 Gene in Leukaemic Rats Using an EvaGreen Real-time RT-PCR Assay <i>H. Nursyuhada, H. Hazilawati, A.H. Hutheyfa, S.M. Rosly, S. Shanmugavellu, M.M. Noordin and S. Jasni</i> | 373 |
| Expression of Circulating CD146 Associated with Endovascular Dysfunction in Adenine-induced Chronic Renal Damage in Rats Using an EvaGreen Real-time RT-PCR Assay <i>P. Farah Dina, H. Hazilawati, S.M. Rosly, S. Shanmugavelu and M.M. Noordin</i> | 381 |

Isolation and Identification of *Fusarium* Species Associated with Fusarium Ear Rot Disease of Corn

Nur Ain Izzati Mohd Zainudin^{1*}, Siti Nordahliawate Mohamed Sidique², Nor Azlina Johari¹, Darnetty³, Azmi Abd. Razak⁴ and Baharuddin Salleh³

¹Department of Biology, Faculty of Science, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

²Department of Agrotechnology, Faculty of Agrotechnology and Food Science, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia

³School of Biological Sciences, Universiti Sains Malaysia, 11800 Minden, Pulau Pinang, Malaysia

⁴Malaysian Agricultural Research and Development Institute, Bukit Raya, 06707 Pendang, Kedah, Malaysia

*E-mail: izzati@science.upm.edu.my

ABSTRACT

Fusarium species associated with corn (*Zea mays*) are very diverse and widely distributed throughout Malaysia. Out of 657 samples cultured, a total of 220 *Fusarium* isolates were obtained from corn plants showing the typical symptoms of Fusarium ear rot in 12 locations throughout Malaysia. All the isolates were identified into 10 species based on the morphological characteristics that emphasized on growth rates, colony features and microscopic characteristics. A total of 117 *Fusarium* isolates were classified into four species in the section *Liseola* and their allied, tentatively identified as *F. proliferatum* (58), *F. subglutinans* (34), *F. verticillioides* (24), and *F. nygamai* (1). Meanwhile, *F. proliferatum* was the most prevalent species in all the sampling areas. 103 isolates, which were classified into six other *Fusarium* species belonging to different sections, were also isolated and identified, and these included *F. semitectum* (47), *F. oxysporum* (20), *F. pseudograminearum* (19), *F. solani* (15), *F. equiseti* (1), and *F. longipes* (1). *F. semitectum* was the highest among other common saprophytic fungi in corn. *F. pseudograminearum* was only isolated from the samples obtained from Cameron Highlands, Pahang. In term of species diversity, *Fusarium* species was the highest obtained in Semenyih, Selangor, with $H' = 1.72$.

Keywords: *Fusarium* species, Fusarium ear rot, diversity, corn, morphology

INTRODUCTION

Genus *Fusarium* is classified under phylum Ascomycota and ubiquitous fungi that are extensively distributed worldwide, from the temperate to the tropical regions (Leslie & Summerell, 2006). *Fusarium* species are

commonly reported as endophytes, saprophytes, and pathogens of various plants, especially economically important crops, including corn (Nelson *et al.*, 1983; Burgess *et al.*, 1994). Corn (*Zea mays*) belongs to dicotyledonous angiosperm plant and it is a member of the grass family Poaceae.

Received: 5 March 2010

Accepted: 11 November 2010

*Corresponding Author

Fusarium species can cause plant diseases, such as wilt, rot, abnormal growth and decay, on vegetables, wood, herbs, and ornamental plants. Besides being plant pathogens, *Fusarium* species may also produce secondary metabolites, such as mycotoxins (i.e. beauvericin, fumonisins, and moniliformin), as well as phytotoxins (fusaric acid and gibberellic acid) problems (Booth, 1971; Summerell *et al.*, 2003). Fumonisin and moniliformin can cause serious mycotoxicoses on humans and animals (Summerell *et al.*, 2003), whereas, fusaric acid and gibberellic acids can respectively lead to stunting and abnormal elongation on the growth of plants.

Meanwhile, the *Fusarium* species in the section *Liseola* and their allies have been reported as potential pathogens on corn and caused Fusarium ear rot disease. However, the identity and aetiology of this pathogen is still highly debated. Therefore, this study was conducted to isolate and identify the *Fusarium* from the samples showing typical symptoms of the disease. Recently, *Fusarium* species such as *F. proliferatum*, *F. oxysporum*, *F. nygamai*, *F. semitectum*, *F. solani*, and *F. verticillioides* were successfully isolated from corn showing typical symptoms of ear rot disease in four states of Malaysia, namely, Perlis, Pulau Pinang, Sabah and Sarawak (Darnetty *et al.*, 2008). However, no report is available on the distribution and diversity of *Fusarium* isolates obtained from the west coast (Selangor), east coast (Pahang) and Southeast areas (Johor) of Peninsular Malaysia.

MATERIALS AND METHODS

Corn Samples

A total of 657 corn samples were obtained from 12 main corn growing areas throughout Malaysia. All the samples were surface sterilized with 10% Clorox® and rinsed in several changes of sterile water.

The Isolation of Fusarium Isolates and Monospore Culture

The samples were placed on semi-selective medium for *Fusarium*, peptone pentachloronitrobenzene agar (PPA) as described by Nash & Snyder (1962), and incubated for 7 days under standard growth conditions (Salleh & Sulaiman, 1984). The cultures were single-spored following a standard protocol by Burgess *et al.* (1994). After 7 days of incubation, the cultures were transferred onto potato dextrose agar (PDA) and carnation leaf agar (CLA; Fisher *et al.* 1982) for species identification.

Morphological Characteristics and Species Identification

The cultures on PDA were used for observing the macroscopic characteristics such as colony features, growth rate and pigmentation. For microscopic characterization, pure cultures were transferred onto CLA and soil extract agar (SEA; Klotz *et al.*, 1988). The microscopic characteristics such as conidia ontogeny, as well as the presence of chlamydo-spores and types of conidiophores were examined following the procedure by Burgess *et al.* (1994) and Leslie & Summerell (2006). The observations were done using a light microscope (Olympus model BX-50F4) and photographed using a JVC camera model KY-F55BE, with an image analyzer-SIS programme. The *Fusarium* isolates were identified based on the morphological characteristics into species level following Nelson *et al.* (1983), Burgess *et al.* (1994) and the *Fusarium* Laboratory Manual (Leslie & Summerell, 2006). The pure cultures were maintained on water agar (WA) as short-term working cultures (Burgess *et al.*, 1994).

Diversity of the Fusarium Species

The diversity of the *Fusarium* species isolated within Peninsular Malaysia was calculated based

on Shannon-Weiner Index (Spellerberg, 2008), as follows:

$$H' = -\sum_{i=1}^s p_i \ln p_i$$

where: H' = value of Shanon-Weiner Index
 \sum refers to “the sum of”
 there are s species in the community
 p_i = is the relative abundance (proportion) of the i species in the community
 \ln = natural log

RESULTS AND DISCUSSION

A total of 220 isolates of *Fusarium* were obtained from the corn plant samples showing typical symptoms of *Fusarium* ear rot disease in 12 locations throughout the main corn growing areas in Malaysia. The typical symptoms of *Fusarium* ear rots are a white to pink- or salmon-coloured mold (fungal mass), beginning anywhere on the ear or scattered throughout. Some infected ears show brown necrotic lesions at the end of the cob (Figs. 1A-C). Often, the decay begins with insect-damaged kernels, by corn borer or bird feeding as a first infection and is later infected by fungi as a secondary infection. Normally, the disease does not involve the whole ear or kernels, but a portion of the

corn ear. The infected kernels are frequently appeared as tan or brown, or have white streaks of the fungal mycelia.

The *Fusarium* isolates were single-spored and tentatively identified into 10 species (namely, *F. equiseti*, *F. longipes*, *F. nygamai*, *F. oxysporum*, *F. pseudograminearum*, *F. proliferatum*, *F. semitectum*, *F. solani*, *F. subglutinans*, and *F. verticillioides*) as shown in Table 1. Among the 10 species, *F. proliferatum* (Section Liseola) was the most prevalent species and widely distributed (it was found to be present in seven locations examined with 58 isolates). On the contrary, three species, namely, *F. equiseti*, *F. longipes* and *F. nygamai*, were noted as the least frequent species with a single isolate each. The *Fusarium* species in the Section Gibbosum (*F. equiseti* and *F. longipes*) and Section Arthrosporiella (*F. semitectum*) were also isolated and have previously been reported to be frequently present as saprophytes on various plants including maize (Nelson *et al.*, 1983, Summerell *et al.*, 2003; Leslie & Summerell, 2006). However, *F. proliferatum*, *F. verticillioides*, *F. solani* and *F. oxysporum* commonly cause plant diseases on a variety of crops (Summerell *et al.*, 2003; Leslie & Summerell, 2006).

The *Fusarium* species associated with the corn samples, showing the typical symptoms of Fusarium ear rot disease in Malaysia, are very




Fig. 1: Samples of the corn obtained from Cameron Highland, Pahang; A) Healthy corn; and B-C) infected cobs showing brown necrotic lesion on the ears (arrows). Scale bar = 0.7cm

diverse, based on Shanon-Weiner Index. The *Fusarium* species isolated from the samples obtained from Semenyih was the highest in term of its diversity, with $H' = 1.72$, and this was followed by Banting (1.46), TPU UPM and MARDI Pulau Pinang (1.41), Cameron Highland (1.25), Jerantut (0.69), Lanchang (0.51), and Seri Medan (0.46). The lowest index (0) was obtained by the samples from Jengka, Kuala Berang and Kota Kinabalu (Table 1).

The cultures were identified as the *Fusarium* species based on several diagnostic characteristics, such as the presence of chlamydoconidia, microconidia, macroconidia and the type of conidiophores. The number of macroconidia septation is variable, depending on the species. Meanwhile, *F. equiseti*, *F. longipes*, and *F. pseudograminearum* only produced macroconidia and without microconidia. Most of the species in Section Liseola and their allied (which were tentatively identified as

F. nygamai, *F. proliferatum*, and *F. verticillioides*) produced microconidia in chains and/or false heads. Another archetypal characteristic of this particular section is that the majority of the species were unable to produce chlamydoconidia.

The most dominant species was morphologically identified as *F. proliferatum* that belongs to the section Liseola and their allied. This species has a worldwide distribution and is frequently isolated on an economically important plant including maize (Nelson *et al.*, 1983; Marasas *et al.*, 1984). Oláh *et al.* (2006) state that *F. proliferatum* is a weak pathogen of maize and it enters into host tissues during germination.

The second highest isolate is *F. semitectum* which is regularly found as a secondary invader in diseased tissues (Summerell *et al.*, 2003), soils (Burgess *et al.*, 1994), as well as from diverse aerial parts of several plants, including maize (Andrés Ares *et al.*, 2004), asparagus

TABLE 1
The occurrence and frequency of *Fusarium* species isolated from corns in Malaysia

| <i>Fusarium</i> species | Total (number of isolates) | Main corn growing areas | | | | | | | | | | | |
|-----------------------------|----------------------------------|-------------------------|----------------|------------------|--------------------------|-------------------|--------------------|--------------------|-------------------|-------------------|--------------------------|---------------------|----------------------|
| | | Jerantut, Pahang | Jengka, Pahang | Lanchang, Pahang | Cameron Highland, Pahang | Banting, Selangor | Semenyih, Selangor | TPU, UPM, Selangor | Seri Medan, Johor | Senggarang, Johor | Kuala Berang, Terengganu | MARDI, Pulau Pinang | Kota Kinabalu, Sabah |
| <i>F. equiseti</i> | 1 | - | - | - | 1 | - | - | - | - | - | - | - | - |
| <i>F. longipes</i> | 1 | - | 1 | - | - | - | - | - | - | - | - | - | - |
| <i>F. nygamai</i> | 1 | - | - | - | - | - | 1 | - | - | - | - | - | - |
| <i>F. oxysporum</i> | 20 | - | - | - | 10 | 3 | 3 | 2 | - | - | - | 2 | - |
| <i>F. pseudograminearum</i> | 19 | - | - | - | 19 | - | - | - | - | - | - | - | - |
| <i>F. proliferatum</i> | 58 | - | - | 3 | - | 2 | 6 | 21 | 5 | 3 | - | 18 | - |
| <i>F. semitectum</i> | 47 | - | - | - | 4 | 1 | 9 | 18 | - | 1 | - | 6 | 8 |
| <i>F. solani</i> | 15 | 1 | - | - | - | 3 | 9 | - | - | - | - | 2 | - |
| <i>F. subglutinans</i> | 34 | 1 | - | 11 | 1 | 6 | 5 | 8 | - | - | 1 | 1 | - |
| <i>F. verticillioides</i> | 24 | - | - | - | 1 | - | 1 | 11 | 1 | 3 | - | 7 | - |
| Percentage (%) | - | 0.9 | 0.4 | 6.4 | 16.4 | 6.8 | 15.5 | 27.3 | 2.7 | 3.2 | 0.4 | 16.4 | 3.6 |
| Total (n) | 220 | 2 | 1 | 14 | 36 | 15 | 34 | 60 | 6 | 7 | 1 | 36 | 8 |
| Shannon-Weiner Index | - | 0.69 | 0.0 | 0.51 | 1.25 | 1.46 | 1.72 | 1.41 | 0.46 | 1.0 | 0.0 | 1.41 | 0.0 |

(Al-Amodi, 2006), kangaroo paw (Satou *et al.*, 2001), beans (Dhingra & Muchovej, 1979), sorghum (Gopinath *et al.*, 1985), millet (Mathur *et al.*, 1973; 1975) and potatoes (Kim *et al.*, 1995). Meanwhile, *F. verticillioides* has been reported as a pathogen on maize which causes epidemics of maize ear rot. The species can be found worldwide, or wherever maize is cultivated (Leslie & Summerell, 2006).

Three species (namely, *F. equiseti*, *F. nygamai*, and *F. longipes*) were also isolated from the maize samples showing typical symptoms of Fusarium ear rot disease which were previously recovered, particularly from diverse hosts. These species are cosmopolitan soil inhabitants that have been recovered from many parts of the world, primarily as saprophytes or endophytes (Nelson *et al.*, 1983; Summerell *et al.*, 2003; Leslie & Summerell, 2006) and have also been isolated from maize (Logrieco *et al.*, 1998). Similarly, Logrieco *et al.* (1998) have reported that those species produce beauvericin and may be one of the contaminants of maize.

A total of 14 isolates of *F. pseudograminearum* were obtained from Cameron Highlands in Pahang, whereby the day and night temperature range is 18-25°C. Based on the colony growth requirement, this species is categorized as a low temperature tolerates fungus, and therefore, the species is usually found in the temperate region. Moreover, *F. pseudograminearum* is morphologically and culturally indistinguishable from *F. graminearum*, whereby both species can only be differentiated by observing the formation of perithecia on media such as CLA. *F. graminearum* is homothallic and able to abundantly produce perithecia on the media, although *F. pseudograminearum* is not capable of producing perithecia (Leslie & Summerell, 2006). *F. pseudograminearum* and *F. graminearum* are important plant pathogens worldwide, including the maize disease. Diseases of cereals, including maize, caused by these species are responsible for large economic losses due to the reduction in seed quality and contamination of grain with their secondary metabolites that are known as mycotoxin (Russell *et al.*, 2005).

In conclusion, ten species of *Fusarium* were isolated from 12 locations in Malaysia and they were morphologically identified as *F. equiseti*, *F. longipes*, *F. nygamai*, *F. oxysporum*, *F. proliferatum*, *F. pseudograminearum*, *F. semitectum*, *F. solani*, *F. subglutinans*, and *F. verticillioides*. This is a rather comprehensive report on the diversity of the *Fusarium* species associated with corn in Malaysia and the role of these fungi in causing plant diseases; nonetheless, their biological species and toxigenicity still require further investigation.

ACKNOWLEDGEMENT

The authors wish to thank the Ministry of Higher Education for the financial support under the Fundamental Research Grant Scheme (FRGS), project no. 01-11-08-661FR.

REFERENCES

- Al-Amodi, M.O.S. (2006). Morphological, pathogenic, genetic and molecular variabilities of *Fusarium* spp., the pathogens of asparagus crown and root rot in Malaysia and Brunei Darussalam. PhD thesis. Universiti Sains Malaysia.
- Andrés Ares, J.L., Alonso Ferro, R.C., Campo Ramírez, L., & Moreno González, J. (2004). *Fusarium graminearum* Schwabe, a maize root and stalk rot pathogen isolated from lodged plants in northwest Spain. *Spanish Journal of Agricultural Research*, 2, 249-252.
- Booth, C. (1971). *The genus fusariumm*. Commonwealth Mycological Institute, Kew, Surrey.
- Burgess, L.W., Summerell, B.A., Bullock, S., Gott, K.P., & Backhouse, D. (1994). *Laboratory manual for fusarium research*. University of Sydney.
- Darnetty, Dian Mariana, M.N., Nur Ain Izzati, M.Z., Nor Azliza, I., Nik Mohd Izham, M.N., & Salleh, B. (2008). Diversity of *Fusarium* species associated with ear rot of corn in Indonesia and Malaysia. In A.R. Adb Wahab, L.K. Chan, S. Shaida Fariza, M.Y. Ahmad Ramli, B. Amirul Al-Ashraf, J. Zairi *et al.* 6th Regional IMT-GT UNINET Conference 2008 (pp. 422-425).

- Dhingra, O.D., & Muchovej, J.J. (1979). Pod rot, seed rot and root rot of snap bean caused by *Fusarium semitectum*. *Plant Disease Reporter*, 63, 84-87.
- Fisher, N.L., Burgess, L.W., Toussoun, T.A., & Nelson, P.E. (1982). Carnation leaves as a substrate and for preserving cultures of *Fusarium* species. *Phytopathology*, 72, 151-153.
- Gopinath, A., Prakash, H.S., & Shetty, H.S. (1985). Grain mold of sorghum: Role associated on the health and variability of seeds. *International Journal of Tropical Plant disease*, 3, 177-182.
- Kim, J.C., Lee, Y.W., & Yu, S.H. (1995). Sambutoxin producing isolates of *Fusarium* species and occurrence of sambutoxin in rotten potato tubers. *Applied and Environmental Microbiology*, 61, 3750-3751.
- Klotz, L.V., Nelson, P.E., & Toussoun, T.A. (1988). A medium for enhancement of chlamydospore formation in *Fusarium* species. *Mycologia*, 80, 108-109.
- Leslie, J.F., & Summerell, B.A. (2006). *The fusarium laboratory manual*. UK: Blackwell Publishing Ltd.
- Logrieco, A., Moretti, A., Castella, G., Kostecki, M., Golinski, P., Ritieni, A., & Chelkowski, J. (1998). Beauvericin production by *Fusarium* species. *Applied and Environmental Microbiology*, 64, 3084-3088.
- Marasas, W.F.O., Nelson, P.E., & Toussoun, T.A. (1984). *Toxigenic Fusarium species, identity and mycotoxicology*. The Pennsylvania State University Press.
- Mathur, S.K., Mathur, S.B., & Neergaard, P. (1975). Detection of seed-born fungi in sorghum and location of *Fusarium moniliforme* in the seed. *Seed Science and Technology*, 3, 683-690.
- Mathur, S.K., Nath, R., & Mathur, S.B. (1973). Seedborn fungi of pearl millet (*Pennisetum typhoides*) and their significance. *Seed Science and Technology*, 1, 811-820.
- Nash, S.M., & Snyder, W.C. (1962). Quantitative estimations by plate counts of propagules of the bean root rot *Fusarium* in field soils. *Phytopathology*, 73, 458-462.
- Nelson, P.E., Toussoun, T.A., & Marasas, W.F.O. (1983). *Fusarium species: An illustrated manual for identification*. The Pennsylvania State University Press.
- Oláh, B., Jeney, A., & Hornok, L. (2006). Monitoring the endophytic colonization of *Fusarium proliferatum* in maize tissues. *Acta Phytopathologica et Entomologica Hungarica*, 41, 185-191.
- Russell, D.M., Matthew, G.C., Denis, R.L., Margaret, D.M., Lincoln, & John, M. (2005). *Fusarium graminearum*, *F. cortaderiae* and *F. pseudograminearum* in New Zealand: Molecular phylogenetic analysis, mycotoxin chemotypes and co-existence of species. *Mycological Research*, 109, 410-420.
- Salleh, B., & Sulaiman, B. (1984). Fusaria associated with naturally diseased plants in Penang. *Journal of Plant Protection in the Tropics*, 1, 47-53.
- Satou, M., Ichinoe, F., Fukumoto, N., Tezuka, & Horiuchi, S. (2001). Fusarium blight of kangaroo paw (*Anigozanthos* spp.) caused by *Fusarium chlamydosporum* and *F. semitectum*. *Journal of Phytopathology*, 149, 203-206.
- Spellerberg, I.F. (2008). *Encyclopaedia of ecology*. Lincoln, New Zealand: Lincoln University.
- Summerell, B.A., Salleh, B., & Leslie, J.F. (2003). A utilitarian approach to *Fusarium* identification. *Plant Disease*, 87, 117-128.