


Giant knee “ganglion”—a case report

**Nuno Vieira Ferreira, Luis Filipe
Carriço, Bruno Pereira, Rui Duarte,
Ricardo Maia, Nuno Sevivas, Ramiro
Fidalgo & Manuel Vieira da Silva**

**European Orthopaedics and
Traumatology**
Official Journal of the European
Federation of National Associations
of Orthopaedics and Traumatology
(EFORT)

ISSN 1867-4569

Eur Orthop Traumatol
DOI 10.1007/s12570-013-0223-1



Your article is protected by copyright and all rights are held exclusively by EFORT. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Giant knee “ganglion”—a case report

Nuno Vieira Ferreira · Luis Filipe Carriço ·
Bruno Pereira · Rui Duarte · Ricardo Maia ·
Nuno Sevivas · Ramiro Fidalgo · Manuel Vieira da Silva

Received: 6 August 2013 / Accepted: 24 September 2013
© EFORT 2013

Introduction

“Ganglion” is a cystic lesion that originates in the joint capsules or tendon sheaths usually containing clear liquid and jelly. The lesions are composed of cystic space with a wall containing dense fibrous and adipose tissue without epithelial lining. Rare occurrence [1–5] in the general population (about 1 %) [2] uncommonly produces specific symptoms or shows classic signs. These lesions frequently occur on the back of the wrist, palm, and dorsum/lateral foot [2]; however, they are rare in locations such as the shoulder, peri-acetabular region, or knee [1, 6]. When they do occur, ganglions are usually small, whereas very few cases of giant ganglion are reported [3, 7, 8].

Although the etiology is still unclear, it is thought to result from myxoid degeneration [1, 2]. The diagnosis is usually clinical and by imaging [5, 6], and biopsy is rarely necessary [1, 2]. The treatment of choice is complete excision with free margin [2, 8].

Case presentation

The authors present the case of a male patient, 47 years old, who had a left infrapatellar mass with progressive growth of 5 years of evolution, which is spherical, measuring approximately 10 cm in diameter (Fig. 1). No other signs or symptoms were observed, nor association with other changes to the clinical examination of the knee.

It was investigated initially by ultrasound and computed tomography (CT). The lesion was best characterized by

magnetic resonance imaging (MRI), describing the presence of a large nodule measuring 8 cm in longitudinal diameter in the anterior left knee, well delineated by thick surrounding regular wall. Imaging revealed hyperintensity in T2 and slight hyperintensity in T1, with internal debris that revealed hyposignal in different sequences, not enhanced after contrast administration.

This lesion was located within the subcutaneous fat, independent of patellar tendon or bone structures (Fig. 2).

The patient was proposed for surgical excision under spinal anesthesia. Cross-cutting approach was carried out (Fig. 3a), followed by excision with skin flap. Cleavage plane was found throughout the extent of injury.

The postoperative period was good with the patient discharged at the third day. Four weeks after, the patient was completely asymptomatic and returned to his usual active life.

The specimen (Fig. 3b) was sent for histological examination; it has smooth outer surface and partially surrounded by adipose tissue, with a cavity containing blood clots and wall composing of dense fibrous tissue, without epithelial lining (Fig. 3c).



Fig. 1 Macroscopic view of the ganglion

N. V. Ferreira (✉) · L. F. Carriço · B. Pereira · R. Duarte · R. Maia ·
N. Sevivas · R. Fidalgo · M. V. da Silva
Hospital de Braga, Braga, Portugal
e-mail: nvferreira@gmail.com

Fig. 2 MRI showing well-defined lesion: sagittal and axial




Fig. 3 **a** Cross-cutting approach. **b** Lesion after resection. **c** Histological view of the specimen



Discussion

The case presented is uncommon either by location or by the size of the lesion. As a slow-growing lesion, it is justified by elapsed time to treatment.

The MRI examination was more accurate to describe the injury and its relations with the neighboring structures. However, it was not capable to make an accurate preoperative diagnosis of the lesion. This is due to known limitations of the technique, but it did not interfere with the effectiveness of the treatment.

According to the literature, it was considered unnecessary to carry out a diagnostic biopsy [1, 2]. Our work confirms this option for treating these lesions, given the irrelevance of its results for the preoperative planning.

Conclusion

The giant ganglion of the knee is an extremely rare injury. The preoperative diagnosis is not always easy; however, it is a benign lesion, for which surgical excision is an effective treatment.

Acknowledgments The authors thank the Pathology Department of Hospital de Braga.

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Canale ST, Campbell WC (2003) Campbell's operative orthopaedics, 10th edn. Mosby, St. Louis
2. Cohen R et al (1999) "Ganglion" intra-articular do joelho: comportamento clinico-patológico. *Revista Brasileira de Otopedia* 34(2):159–64
3. Mine T et al (2003) A giant ganglion cyst that developed in the infrapatellar fat and partly extended into the knee joint. *Arthroscopy* 19(5):E40
4. David KS, Korula RJ (2004) Intra-articular ganglion cyst of the knee. *Knee Surg Sports Traumatol Arthrosc* 12(4):335–337
5. Yilmaz T et al (2004) Ganglion cysts of the knee originating from tendons and ligaments. *Tani Girisim Radyol* 10(3):246–251
6. Yilmaz E et al (2004) A ganglion cyst that developed from the infrapatellar fat pad of the knee. *Arthroscopy* 20(7):e65–e68
7. Vayvada H et al (2003) Giant ganglion cyst of the quadriceps femoris tendon. *Knee Surg Sports Traumatol Arthrosc* 11(4):260–262
8. O'Rourke PJ, Byrne JJ (1995) Giant ganglion of the proximal tibiofibular joint: a case report. *Ir J Med Sci* 164(4):295–296