

Odynophagia and neck pain after exercise

Catarina Faria, Inês Medeiros, Fábía Carvalho, Ana Antunes

Pediatric Department, Hospital de Braga, Braga, Portugal

Correspondence to
Dr Catarina Faria,
catmagalhaesfaria@gmail.com

Accepted 18 March 2018

DESCRIPTION

A previously healthy, non-asthmatic, non-smoker, 16-year-old male came for medical attention because of odynophagia and pain located at anterior cervical area and upper retrosternal area, increasing with swallowing and deep breathing, that begun after physical exercise at gymnastic class. The adolescent also mentioned a felling of air bubbles running up and down that area. There was no history of local trauma. He referred stuffed nose and mild cough but no other respiratory complaints or fever. Physical examination was unremarkable, with normal auscultation and no alterations on cervical and thoracic examination. Lateral soft tissue neck X-ray revealed free air in retropharyngeal space (figure 1, arrows). Chest X-ray showed a small amount of air in the upper mediastinum. Blood count was normal, and C reactive protein was negative. Mycoplasma pneumoniae IgM was negative. The patient was treated with high concentration oxygen and oral

analgesia. The next day, he only referred mild pain when coughing. Control neck and chest X-rays showed residual air on retropharyngeal space and no air on mediastinum, respectively. He was discharged with recommendation to avoid exercise and physical efforts for the next week as well to avoid tobacco smoke. He was re-evaluated 1 week later and was symptom free. Neck and chest X-rays were normal. After 1 month, he remained with no symptoms and with no abnormal findings on neck and chest X-rays, including pleural effusion.

Spontaneous pneumomediastinum (SPM) is a rare, usually benign and self-limited entity.¹ Valsalva manoeuvre (eg, induced by physical exercise) can be a trigger. Typical symptoms include dyspnoea and retrosternal pleuritic pain.² The authors want to emphasise the fact that SPM can present with head and neck complaints, with less obvious respiratory symptoms. SPM with retropharyngeal emphysema is rare but should not be forgotten in differential diagnosis of odynophagia and neck pain, as it can be complicated with upper airway obstruction in punctual cases.²



Figure 1 Free air in the retropharyngeal space (arrows).

Learning points

- ▶ Spontaneous pneumomediastinum presenting as retropharyngeal emphysema is a rare entity that requires a high degree of suspicion.
- ▶ Lateral soft tissue neck X-ray is a fundamental tool on making this diagnosis.
- ▶ Prognosis is usually good with conservative treatment.

Contributors CF, IM and FC: conception and design, drafting the article and final approval of the version to be published. AA: conception and design, and final approval of the version to be published.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Smith JL, Hsu JM. Spontaneous pneumomediastinum presenting with retropharyngeal emphysema. *Am J Otolaryngol* 2004;25:290–4.
- 2 Cho DY, Aaron GP, Shepard KG. Spontaneous retropharyngeal and mediastinal emphysema. *Clin Exp Otorhinolaryngol* 2016;9:178–81.



To cite: Faria C, Medeiros I, Carvalho F, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224581

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow