

New records of *Astyanax pelegrini* Eigenmann, 1907 and *Triportheus pantanensis* Malabarba, 2004 (Actinopterygii: Characiformes: Characidae) for the Río Pilcomayo National Park and Ramsar Site in the province of Formosa, Argentina

Florencia Brancolini^{1,3*}, Lucila Protogino², Priscilla Minotti^{1,3}, Claudio Baigún⁴, Hugo López⁵ and Darío Colautti⁴

¹ Centro de Altos Estudios en Ciencias Exactas (CAECE). Junín 516, 1026. Ciudad Autónoma de Buenos Aires, Argentina.

² Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Instituto de Limnología "Dr. Raúl A. Ringuelet". C.C. 712, 1900 La Plata, BA, Argentina.

³ Universidad Nacional de San Martín (UNSAM). Belgrano 3563. 1650. San Martín, BA, Argentina.

⁴ Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Instituto Nacional de Biotecnología e Instituto Nacional de Tecnología Chascomús (IBB-INTECH). Camino Circunvalación Laguna Km 6. B7130IWA. Chascomús, BA, Argentina.

⁵ Museo de La Plata, División Zoología Vertebrados. Paseo del Bosque s/n. 1900. La Plata, BA, Argentina.

* Corresponding author. E-mail: florencia.brancolini@gmail.com

ABSTRACT: This note presents the first records of *Astyanax pelegrini* Eigenmann, 1907 and *Triportheus pantanensis* Malabarba, 2004 for the province of Formosa, Argentina from fish surveys conducted in the Río Pilcomayo National Park and Ramsar Site. In the case of *T. pantanensis*, this occurrence represents the first record of this species for Argentina.

The objective of this note is to provide information on fish occurrences recorded in the Río Pilcomayo National Park and Ramsar Site (RPNPRS) in the province of Formosa, which represent new records for this province and for Argentina. The Park is located in the eastern corner of Formosa province, northern Argentina (Figure 1A). The landscape is typical of the Great Chaco region, with a remarkable number of seasonal swamps, with poorly known fish fauna. General characteristics of Formosa's ichthyofauna can be found in López and Castello (1966), López (1972), Bayley (1973), Azpelicueta and Braga (1980), Pignalberi de Hassan and Cordiviola de Yuan (1988), Menni *et al.* (1992), Azpelicueta and Yanosky (1992) and Azpelicueta (2005). Menni (2004) mentioned the presence of 143 fish species in Formosa from wetlands related to the Paraguay, Pilcomayo and Bermejo river systems. From an ichthyogeographic perspective, the National Park belongs to the Great Rivers Province (López *et al.* 2008) and is located within the Subtropical Potamic Axis Ecoregion of López *et al.* (2002).

Between January 2007 and January 2008, we conducted bimonthly surveys to assess fish biodiversity in the RPNPRS larger wetlands, using different fishing gears. For species identification we followed Ringuelet *et al.* (1967), Malabarba (2004) and Mirande *et al.* (2006), whereas osteological preparations were performed according to the technique of Taylor and Van Dyke (1985). Counts of gill-rakers on the first arch were taken from cleared and stained specimens (c&s). The repository of the material is maintained in the collection of the Instituto de Limnología "Dr. Raúl A. Ringuelet" (Institutional Code: ILPLA). Details on the material examined can be found in Appendix 1.

CHARACIFORMES

CHARACIDAE

Astyanax pelegrini Eigenmann, 1907. Figure 2.

Diagnostic features were considered consistent with the work of Mirande *et al.* (2006) on the re-description of *Astyanax correntinus* (Holmberg, 1891) and its comparison with *A. pelegrini*.

Scale rows between dorsal fin and lateral line: 12; scale rows between lateral line and anal-fin: 8; lateral line complete, perforated scales: 48; scales around caudal peduncle: 19; rows of scales along the anal fin base: 3; anal-fin rays: v, 41 (46); teeth on outer premaxillary series: 5; the anal fin origin, is located below the last third of dorsal fin; partially naked area between the supraoccipital process and dorsal fin origin; gill rakes on the first gill arch: 26; base anal fin: 40.2% in standard length (SL); distance pectoral-ventral origin: 16.3% SL; pectoral-fin length: 21.8% SL; maxilar length: 21.5% in head length (HL); upper jaw length: 34.2% HL.

Triportheus pantanensis Malabarba, 2004. Table 1, Figure 3.

Diagnostic features: depth at dorsal fin origin: 33.4-41.4 (38.0)% SL, number (n) = 23; depth at pectoral fin insertion: 33.0-44.2 (37.8)% SL, n = 23; presence of two longitudinal scales rows between the pectoral fin insertion and the ventral keel; few gill rakes on the lower limb of first gill arch: 27-31 (28), n = 11 c&s specimens; the lateral-line scales counts: 28-31 (30); the bony head length: 22.5-27.1 (24.6) % SL, n = 23; the snout to anal-fin origin distance:




FIGURE 1. Distribution maps of species recorded: A. Study area, Río Pilcomayo National Park and Ramsar Site in Formosa province, Argentina. B. red triangles: *Astyanax pelegrini*; C. yellow circle: *Triportheus pantanensis*.

TABLE 1. Morphometric data for *Triportheus pantanensis* from Argentina compared to holotype and paratype data (taken from Malabarba 2004).

| CHARACTER | LOW | HIGH | MEAN | SD | N | HOLOTYPE | PARATYPES | N |
|---------------------------------|--------|--------|--------|-----|----|----------|-----------|----|
| Standard length (mm) (SL) | 65.2 | 126.4 | 98.6 | | | 99.5 | 67.7-158 | 31 |
| As a percentage of SL | | | | | | | | |
| Bony head length | 22.5 | 27.1 | 24.6 | 1.2 | 23 | 26.0 | 22.5-27.3 | 31 |
| Snout to anal-fin origin | 65.4 | 73.8 | 67.9 | 2.2 | 23 | 70.0 | 66.4-74.1 | 31 |
| Snout to dorsal-fin origin | 59.3 | 64.2 | 61.5 | 1.2 | 23 | 60.6 | 59.3-64.6 | 31 |
| Snout to pelvic-fin origin | 45.4 | 55.3 | 49.4 | 2.2 | 23 | 51.9 | 49.9-66.9 | 30 |
| Anal-fin base length | 27.3 | 32.1 | 30.0 | 1.3 | 23 | 28.7 | 25.5-31.0 | 30 |
| Caudal peduncle length | 6.2 | 9.8 | 7.6 | 1.0 | 23 | 6.7 | 5.5-11.3 | 31 |
| Caudal peduncle depth | 10.1 | 11.6 | 10.7 | 0.4 | 23 | 11.0 | 9.2-11.1 | 31 |
| Depth at dorsal-fin origin | 33.4 | 41.4 | 38.0 | 2.0 | 23 | 36.4 | 34.6-42.5 | 31 |
| Depth at pectoral fin | 33.0 | 44.2 | 37.8 | 2.7 | 23 | 36.8 | 30.7-42.8 | 31 |
| Dorsal-fin height | 17.9 | 21.9 | 19.9 | 1.0 | 23 | 19.8 | 18.0-22.8 | 31 |
| Pelvic-fin length | 12.1 | 17.2 | 15.1 | 1.5 | 22 | 13.9 | 12.8-17.0 | 30 |
| Pectoral-fin length | 33.6 | 44.2 | 38.6 | 2.7 | 23 | 38.8 | 32.9-42.7 | 31 |
| As a percentage of head length | | | | | | | | |
| Snout length | 18.8 | 25.9 | 21.5 | 2.0 | 23 | 21.6 | 18.2-25.2 | 31 |
| Horizontal eye diameter | 29.1 | 36.6 | 32.2 | 1.7 | 23 | 30.1 | 25.0-32.5 | 31 |
| Least interorbital width | 33.9 | 41.4 | 38.1 | 1.4 | 23 | 35.1 | 31.6-38.9 | 31 |
| Meristic | | | | | | | | |
| Gill rakers | 27 | 31 | 28 | 1.1 | 11 | 30 | 26-33 | 29 |
| Scales in lateral line | 28 | 31 | 30 | 1.0 | 23 | 30 | 28-33 | 32 |
| Scale series above lateral line | 6 | 6 | 6 | 0.0 | 23 | 6 | 6 | - |
| Scale series below lateral line | 2 | 2 | 2 | 0.3 | 23 | 2 | 2 | - |
| Scales along mid-dorsal line | 10 | 13 | 12 | 0.8 | 23 | - | 6-13 | 29 |
| Dorsal-fin rays | ii,8 | ii,9 | ii,9 | 0.4 | 23 | ii,9 | ii,9 | 31 |
| Anal-fin rays | iii,24 | iii,29 | iii,27 | 1.8 | 23 | iii,29 | iii,25-30 | 31 |
| Pectoral-fin rays | i,10 | i,11 | i,11 | 0.6 | 23 | i,12 | i,10-12 | - |
| Pelvic-fin rays | i,5 | i,6 | i,6 | 0.4 | 23 | i,6 | i,6 | - |

65.4-73.8 (67.9)% SL, n = 23; the length of anal fin base: 27.3-32.1 (30.0)% SL, n= 23, and possession of median caudal-fin rays that extend well beyond the caudal fin margin.

As a result of these surveys, two new species records are added to the ichthyofauna of the Formosa: *A. pelegrini* (Figure 1B) and *T. pantanensis* (Figure 1C). The first one was previously reported in Chaco, Buenos Aires, and Santa Fe provinces. In turn, the occurrence of *T. pantanensis* in the RPNPRS represents the first record for Argentina and the southernmost for this species, being until now only mentioned for the Pantanal region in Mato Grosso, Brazil. These results indicate the need to gather more ichthyological information from these dynamic environments, where their hydrological regime supported by regional floods from the Pilcomayo River favors a constant rate of recolonization and exchange of species.



FIGURE 2. *Astyanax pelegrini*, ILPLA 1964, 59.9 mm SL, lower Río Pilcomayo, Pilcomayo Departament, Pilcomayo river basin, Formosa province, Argentina.



FIGURE 3. *Triportheus pantanensis*, ILPLA 1969, 107.46 mm SL, Laguna Blanca, Pilcomayo Departament, Pilcomayo river basin, Formosa province, Argentina.

ACKNOWLEDGMENTS: We are grateful to Carlos A. S. Lucena (MCP) for the loan of comparative material and to Jeff Williams and Sandra Raredon (USNM) for sending photographs and radiographs of type material. This paper is part of the project "Assessment of fish diversity patterns and development of guidelines for sustainable fishery resources in the Ramsar Site and Río Pilcomayo National Park" corresponding to Wetlands for the Future Program. We acknowledge RAMSAR for providing a grant to the National Park Administration from Argentina for overall logistical support. We also want thank anonymous reviewers who improved the manuscript with their comments and suggestions.

LITERATURE CITED

- Azpelicueta, M.M. 2005. Peces de la Reserva El Bagual; p. 99-116. In A.G. Di Giacomo and S.F. Krapovickas (ed.). *Historia natural y paisaje de la Reserva El Bagual, Formosa, Argentina. inventario de la fauna de vertebrados y de la flora vascular de un área del Chaco Húmedo*. Temas de Naturaleza y Conservación 4. Buenos Aires: Aves Argentinas / Asociación Ornitológica del Plata.
- Azpelicueta, M.M. and L. Braga. 1980. Una nueva cita y ampliación de la distribución de dos especies para la ictiofauna argentina. *Neotropica* 26: 163-169.
- Azpelicueta, M.M. and A.A. Yanosky. 1992. A check-list of fishes from El Bagual Ecological Reserve, Argentina. *Ichthyological Exploration of Freshwaters* 3: 73-76.
- Bayley, P.B. 1973. Studies on the migratory characin, *Prochilodus platensis* Berg, 1889 (Pisces Characoidei), in the River Pilcomayo, South America. *Journal of Fish Biology* 5: 25-40.
- López, H.L., C.C. Morgan and M.J. Montenegro. 2002. *Ichthyological ecoregions of Argentina*. La Plata: Probiota, FCNyM, UNLP - Serie Documentos 1. 68 p.
- López, H.L., R.C. Menni, M. Donato and A.M. Miquelarena. 2008. Biogeographical revision of Argentina (Andean and Neotropical Regions): an analysis using freshwater fishes. *Journal of Biogeography* 35: 1564-1579.
- López, R.B. 1972. Migración de peces en el río Bermejo. *Gaea, Anales de la Sociedad Argentina de Estudios Geográficos* 15:138-143.
- López, R.B and H.P. Castello. 1966. *Sternopygus macrurus* (Bloch and Schneider) (Teleostomi, Sternopyginae). Primera cita para la Argentina. *Comunicaciones del Museo Argentino de Ciencias Naturales, Zoológica* 4: 13-16.
- Malabarba, M.C.S.L. 2004. Revision of the Neotropical genus *Triportheus* Cope, 1872 (Characiformes: Characidae). *Neotropical Ichthyology* 2(4): 167-204.
- Menni, R., A.M. Miquelarena, H. López, J. Casciotta, A. Almirón and L. Protogino. 1992. Fish fauna and environments of the Pilcomayo-Paraguay basins in Formosa, Argentina. *Hydrobiología* 245: 129-146.
- Menni, R.C. 2004. Peces y ambientes en la Argentina continental. *Monografías del Museo Argentino de Ciencias Naturales* 5: 1-316.
- Mirande, J.M., M.M. Azpelicueta and G. Aguilera. 2006. Redescription of *Astyanax correntinus* (Holmberg, 1891) (Teleostei: Characiformes: Characidae), more than one hundred years after original description. *Zoologische Abhandlungen* 55: 9-15.
- Pignalberi de Hassan, C. and E. Cordiviola de Yuan. 1988. Fish population in the Paraguay River of the Formosa area, Argentina. *Studies on Neotropical Fauna and Environment* 23: 165-175.
- Ringuelet, R. A., R. H. Arámburu and A. Alonso de Arámburu. 1967. *Los peces argentinos de agua dulce*. La Plata: Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CIC). 602 p.
- Taylor, W.R. and Van Dyke, G.C. 1985. Revised procedures for staining and clearing small fishes and other vertebrates for bone and cartilage study. *Cybium* 9: 107-119.

RECEIVED: February 2011

LAST REVISED: July 2011

ACCEPTED: July 2011

PUBLISHED ONLINE: October 2011

EDITORIAL RESPONSIBILITY: Sergio Maia Queiroz Lima

APPENDIX 1. Museum records of species analyzed. ILPLA: Instituto de Limnología "Dr. Raúl A. Ringuelet" (CONICET-UNLP), La Plata, Buenos Aires, Argentina; MCP: Museu da Católica Pontifícia de Rio Grande do Sul, Rio Grande do Sul, Porto Alegre, Brazil, and MLP: Museo de La Plata, La Plata, Buenos Aires, Argentina and USNM: Smithsonian National Museum of Natural History, Washington, D. C, USA.

Astyanax pelegrini: Argentina: ILPLA 1964, 59.9 mm SL, Formosa province, Pilcomayo Department, Pilcomayo river basin, lower Río Pilcomayo, 25°00'53" S, 58°07'46" W, Col. C. Baigún, P. Minotti y F. Brancolini, 11/10/07; ***Triportheus pantanensis: Argentina***: ILPLA 1966, 6, 3 c&s, 89.8-118.7 mm SL, Formosa province, Pilcomayo Department, Pilcomayo river basin, Laguna Blanca, 25°10'16" S, 58°07'53" W, Col. C. Baigún *et al.*, 19/02/07, ILPLA 1967, 12, 4 c&s, 50.1-75.6 mm SL, Formosa province, Pilcomayo Department, Pilcomayo river basin, Laguna Blanca in outlet of Estero Poi, 25°08'57" S, 58°10'58" W, Col. C. Baigún *et al.*, 19/02/07, ILPLA 1968, 6, 3 c&s, 49.3-75.1 mm SL, Formosa province, Pilcomayo Department, Pilcomayo river basin, South Pilcomayo River, 24°58'46" S, 58°18'11" W, Col. C. Baigún *et al.*, 05/07/07, ILPLA 1969, 3, 1 c&s, 107.5-122 mm SL, Formosa province, Pilcomayo Department, Pilcomayo river basin, Laguna Blanca, 25°10'17.33" S, 58°07'51.24" W, Col. C. Baigún *et al.*, 14/08/07, ILPLA 1970, 6, 1 c&s, 108.6-126.4 mm SL, Formosa province, Pilcomayo Department, Pilcomayo river basin, Laguna Blanca, 25°10'02.5" S, 58°07'50.7" W, Col. C. Baigún *et al.*, 10/10/07; ***Paraguay***: USNM 181682, 4 paratypes radiographs, 100.3-124.0 mm SL, Asunción Bay, Paraguay River, near Asunción, 25°15' S, 57°40' W, Col. C.J.D. Brown, 10/01/57, USNM 181689, 2 paratypes photographs and radiographs, 133.9-154.0 mm SL, Lago Ypacaray near San Bernardino, 25°28' S, 57°33' W, Col. C.J.D. Brown, 11/10/56; ***Brasil***: MCP 35824, 2, 73.8-78.9 mm SL, from América do Sul, Brasil, Mato Grosso, Paraguay, Paraguay River in Cáceres and surrounding areas, 16°03'00" S, 57°42'00" W, Col. R.E. Reis *et al.*, 11/08/91; ***Triportheus nematurus* (Kner, 1858)**: ***Argentina***: ILPLA 1971, 2 c&s, 93.8-97.6 mm SL, Buenos Aires, second section of San Fernando delta, Cangrejo Stream, 34°17'35.3" S, 58°32'03.7" W, Col. F. Brancolini, A. Brancolini and M.C. Nouzaret, 5/05/07.