

DIFFERENCES IN LARVAL MORPHOLOGY OF ALLOPATRIC ISOLATED POPULATIONS OF THE *ODONTOPHRYNUS OCCIDENTALIS* GROUP FROM WESTERN ARGENTINA

J. M. CEI & E. G. CRESPO



ARQUIVOS
DO
MUSEU
BOCAGE

SÉRIE A

Publicação do Museu e Laboratório Zoológico e Antropológico

FACULDADE DE CIÊNCIAS DE LISBOA

Vol. I n.º 14
1982

Arq. Mus. Boc. (Série A)
vol. I n.º 14 págs. 335-340

DIFFERENCES IN LARVAL MORPHOLOGY
OF ALLOPATRIC ISOLATED POPULATIONS
OF THE *ODONTOPHRYNUS OCCIDENTALIS*
GROUP FROM WESTERN ARGENTINA

J. M. CEI * & E. G. CRESPO **

ABSTRACT: A morphological comparative study of two larval populations — one from Mendoza and the other an isolated one from Sierra Famatina (La Rioja) — of the Argentine toad, *Odontophrynus occidentalis*, is reported.

As a result of field studies and comparative karyological research on *Odontophrynus occidentalis* from western Argentina, some remarkable morphological and biological differences between most of its populations and an isolated mountain population from La Rioja became evident.

Toads of the La Rioja population were found at about 2000-2400 m high in the creek of Aguadita, at the geographically isolated Sierra Famatina, East of Andean mountains. Tadpoles at different stages were observed all over the year, in the running waters and ponds of the creek; adults were only exceptionally found, except in the short breeding season. Significant karyological differences between specimens of *Odontophrynus occidentalis* from Mendoza Province and from Famatina have been reported (RUIZ *in press*).

Our attention is drawn to the probable taxonomic value of their morphological larval characters. Tadpoles of the Famatina population are easily recognizable from those of the Mendoza populations by its larger

* Universidad Nacional de Rio Cuarto, Province of Cordoba, Argentina.

** Centro de Fauna da Universidade de Lisboa (INIC) — Faculdade de Ciências de Lisboa.

size, darker coloration and a less acuminate tip of tail. Moreover, further examinations showed significant differences in shape and measurements of the head and/or other somatic characters. These differences in larval morphological parameters are often comparable to those observed in the adults of both forms.

The purpose of the present paper is therefore a more detailed analysis of tadpole morphology, at different stages of development, both in Famatina and in Mendoza samples. Larval character states may be of interest for a further definitive taxonomic recognition of the isolated *Odontophrynus* form from Famatina mountains.

MATERIAL AND METHODS

Odontophrynus occidentalis tadpoles fixed in 5% formalin from the Mendoza region and Sierra Famatina (La Rioja), were available in stages of development between 26 and 38 (cf. GOSNER 1960). All the measurements were made through a stereoscopic microscope using compasses and accurated to 0.1 mm. For the study of the chondrocranial structures of tadpoles, some specimens of both populations were cleared and stained, in accordance with the methodology preconized by WASSERSUG (1976).

After a previous selection, the following exosomatic parameters under a ratio form, were used: total length, body length, spiracle-snout (dorsal and ventral) distance, internasal distance, interocular distance and eye-snout distance.

We mean by:

- total length — the distance between the anteriormost tip of the snout and the tip of the tail;
- body length — the distance between the snout end and the middle point of the anterior margin of the vent tube;
- spiracle-snout length (dorsal and ventral) — the distances, taken along the midline, dorsal and ventral, between the level of the posterior edge of the spiracular opening and the snout end;
- internasal distance — the distance between the margins of the external nares;
- interocular distance — the distance between the internal sides of the eyeballs;
- eye-snout distance — the distance taken in the dorsal midline, between the level of the posterior margins of the eyeballs and the snout tip.

RESULTS AND DISCUSSION

From the several ratios considered as biometric parameters of the tadpoles, only the following differ significantly (by application of the t — STUDENT test):

A) Total length / body length

	<i>Mean</i>	<i>Min.</i>	<i>Max.</i>	<i>Standard deviation</i>	<i>n</i> ^o
<i>Mendoza</i>	2.45	(2.33 — 2.60)		<i>s</i> = 0.077	<i>n</i> = 25
<i>Famatina</i>	2.52	(2.32 — 2.86)		<i>s</i> ₁ = 0.143	<i>n</i> ₁ = 25
		<i>t</i> = 2.14	0.05 < P < 0.02		

B) Body length / spiracle-snout distance (ventral)

<i>Mendoza</i>	1.80	(1.60 — 2.00)	<i>s</i> = 0.100	<i>n</i> = 25
<i>Famatina</i>	1.62	(1.42 — 1.76)	<i>s</i> ₁ = 0.082	<i>n</i> ₁ = 25
		<i>t</i> = 6.75	P < 0.001	

C) Body length / spiracle-snout distance (dorsal)

<i>Mendoza</i>	1.71	(1.52 — 1.93)	<i>s</i> = 0.094	<i>n</i> = 25
<i>Famatina</i>	1.57	(1.38 — 1.70)	<i>s</i> ₁ = 0.080	<i>n</i> ₁ = 25
		<i>t</i> = 5.62	P < 0.001	

D) Internasal distance / body length

<i>Mendoza</i>	0.10	(0.05 — 0.13)	<i>s</i> = 0.025	<i>n</i> = 25
<i>Famatina</i>	0.17	(0.10 — 0.20)	<i>s</i> ₁ = 0.017	<i>n</i> ₁ = 25
		<i>t</i> = 11.53	P < 0.001	

E) Interocular distance / body length

<i>Mendoza</i>	0.15	(0.13 — 0.16)	<i>s</i> = 0.008	<i>n</i> = 25
<i>Famatina</i>	0.17	(0.12 — 0.20)	<i>s</i> ₁ = 0.017	<i>n</i> ₁ = 25
		<i>t</i> = 5.73	P < 0.001	

F) Eyes-snout / body length } × 10

<i>Mendoza</i>	1.00	(0.89 — 1.14)	<i>s</i> = 0.065	<i>n</i> = 25
<i>Famatina</i>	1.15	(0.87 — 1.41)	<i>s</i> ₁ = 0.153	<i>n</i> ₁ = 25
		<i>t</i> = 4.46	P < 0.001	

Therefore, from the comparison of these two populations, the Famatina one shows the following:

1) a smaller body length in relation to the total length; correlatively the tail length is proportionally greater (figs. 1-2); 2) the spiracular opening located in a posterior position; 3) in relation to the body length, greater inter-nasal and interocular distances (figs. 2-4, 9); 4) in relation to the body length, a greater distance between the posterior level of the eyeballs and the snout tip (figs. 2-4, 9).

Besides these differences, there are others, which if hardly in quantification, are notwithstanding well conspicuous.

The specimens of Sierra Famatina are also distinguished from those of Mendoza, because they have a very much pronounced corporal pigmentation, the snout end and tail tips more roundish (this difference is less evident in the younger samples) (figs. 2-6, 9) the spiracular opening directed more dorsally (figs. 7-8), the labial *papilla* more developed, the second superior row of labial teeth less extensively interrupted in the midline (figs. 12-13), the horny beaks more pigmented and with the margin much more serrated (figs. 14-15) and the eyeballs generally smaller and more lateral (figs. 2-4, 9).





The architectural plan of the chondrocrania in individuals of both populations under study does not reveal any differences which are worth of mention (figs. 16-18). There exist only differences in the proportions that are however correlated with those detected at the level of the external morphology.





Although the larval characters are relatively labile, and so, a taxonomic criterion which meaning must be considered with prudence, we believe that we are facing such a kind of differences, which are based in a considerable degree of genetic differentiation.




The morphological differences between the samples of both populations, which we have the chance to observe are, for example, greater than those we have recorded between the Iberian species of *Alytes* (*A. obstetricans* and *A. cisternasi*) studied by one of us in some detail (CRESPO 1979). These differences also does not seem lesser than those existing between another pair of amphibian species of the European fauna, *Discoglossus pictus* and *D. sardus* (KNOEPFFLER 1962).

SUMÁRIO

Observações de campo e a análise cariológica de várias populações de *O. occidentalis* da região ocidental da Argentina, revelaram a existência de diferenças bem marcadas numa população isolada da Serra de Famatina, em relação às demais populações conhecidas da espécie.







Relativamente à população de Mendoza, tomada como padrão das restantes, os espécimes da Serra de Famatina, têm:

1) menor comprimento do corpo em relação ao comprimento total (o mesmo é dizer que têm a cauda proporcionalmente maior); 2) a abertura espiracular em posição mais posterior; 3) em relação ao comprimento do corpo, maiores distâncias interoculares e internasais; 4) em relação ao comprimento do corpo, maior distância entre o nível posterior dos globos oculares e a extremidade do focinho.

Além destas diferenças, os indivíduos de Famatina têm ainda: pigmentação geral muito mais pronunciada; as extremidades do focinho e da cauda mais arredondadas (esta diferença é menos acentuada nos exemplares mais jovens); as aberturas espiraculares dirigidas mais dorsalmente; as papilas labiais mais desenvolvidas; a 2.^a fiada superior de dentículos labiais menos extensivamente interrompida na parte média; o bico cárneo mais pigmentado e com a margem muito mais serreada; os globos oculares geralmente menores e mais laterais.

Não existem diferenças significativas nos planos arquitecturais dos condrocrânios dos exemplares das duas populações.

Embora com a prudência com que os caracteres larvares, devido à sua relativa labilidade, têm de ser considerados, cremos que neste caso as diferenças observadas reflectem uma acentuada diferenciação genética. Tendo por termo de comparação outros Anfíbios, nomeadamente da fauna europeia, pode atribuir-se a estas diferenças um nível específico.

LITERATURE CITED

- CRESPO, E. G. (1979) — Contribuição para o conhecimento da biologia dos *Alytes* ibéricos, *Alytes obstetricans boscai* Lataste, 1879 e *Alytes cisternasi* Boscá, 1879 (Amphibia, Salientia): — a problemática da especiação de *Alytes cisternasi*. Tese. Faculdade de Ciências. Universidade de Lisboa.
- GOSNER, K. L. (1960) — A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica*, **16**: 183-190.
- KNOEPFFLER, L. P. (1962) — Contribution à l'étude du genre *Discoglossus* (Amphibians, Anoures), *Vie et Milieu* **13** (1): 1-94.
- WASSERSUG, R. J. (1976) — A procedure for differential staining of cartilage and bone in whole formalin — fixed Vertebrates. *Stain Technology* **51** (2): 131-134.

EXPLANATION OF PLATES

- 1 — Lateral views of the tadpoles of *O. occidentalis*. Above: population of La Rioja (stage 28); below: population of Mendoza (stage 29).
- 2 — Dorsal views of *O. occidentalis* tadpoles. Above: population of Mendoza (stage 30); below: population of La Rioja (stage 30).
- 3 — Dorsal view of the snout of the tadpole of *O. occidentalis*, population of Mendoza (stage 33).
- 4 — Dorsal view of the snout of the tadpole of *O. occidentalis*, population of L. Rioja (stage 32).
- 5 — Tail of the tadpole of *O. occidentalis*, population of Mendoza (stage 30).
- 6 — Tail of the tadpole of *O. occidentalis*, population of La Rioja (stage 28).
- 7 — Lateral view of the tadpole of *O. occidentalis*, population of Mendoza — stage 30 (position of the spiracle, *arrow*).
- 8 — Lateral view of the tadpole of *O. occidentalis*: population of La Rioja — stage 28 (position of the spiracle, *arrow*).
- 9 — Lateral views of the tadpoles of *O. occidentalis*. Above: population of Mendoza (stage 28); below: population of La Rioja (stage 27).
- 10 — Oral region of the tadpole of *O. occidentalis*, population of Mendoza (stage 37): development of the labial *papillae*.
- 11 — Oral region of the tadpole of *O. occidentalis*, population of La Rioja (stage 37): different development of the labial *papillae*.
- 12 — Oral region of the tadpole of *O. occidentalis*, population of La Rioja (stage 37): interruption of the second superior row of labial teeth.
- 13 — Oral region of the tadpole of *O. occidentalis*, population of Mendoza (stage 37): second superior row of labial teeth interrupted.
- 14 — Horny beak of the tadpole of *O. occidentalis*, population of Mendoza (stage 37).
- 15 — Horny beack of the tadpole of *O. occidentalis*, population of La Rioja (stage 37).
- 16 — Ventral view of cleared and stained chondrocranium of the tadpole of *O. occidentalis*, population of La Rioja (stage 40).
- 17 — Dorsal view of cleared and stained chondrocranium of the tadpole of *O. occidentalis*, population of La Rioja (stage 40).
- 18 — Dorsal view of cleared and stained chondrocranium of the tadpole of *O. occidentalis*, population of Mendoza (stage 40).

COMPOSTO E IMPRESSO NAS OFICINAS
DA «IMPRENSA DE COIMBRA, LIMITADA»
LARGO DE S. SALVADOR, 1-3 — COIMBRA

Os ARQUIVOS DO MUSEU BOCAGE são constituídos por quatro séries, SÉRIE A; SÉRIE B — NOTAS; SÉRIE C — SUPLEMENTOS e SÉRIE D — EXTENSÃO CULTURAL e ENSINO, publicando trabalhos originais de Zoologia, Antropologia e Biologia Geral, realizados ou não no Museu e Laboratório Zoológico e Antropológico da Faculdade de Ciências da Universidade de Lisboa. Os originais devem ser enviados ao Editor e cada trabalho constituirá um fascículo, que será distribuído separadamente. Cada volume será constituído, em princípio, por quatrocentas a quinhentas páginas, podendo corresponder a um ou mais anos civis.

O *título* dos originais, dactilografados a dois espaços, deverá ser conciso e bem definido e o(s) nome(s) do(s) autor(es) virão seguidos pelo da Instituição onde trabalha(m).

Recomenda-se que ao título se siga um «*abstract*» que não excederá as 50 palavras.

O texto do artigo deverá ser completado por um *resumo* em Francês, Inglês ou Alemão.

Recomenda-se que a *bibliografia* seja apresentada como abaixo se indica.

Aos autores serão concedidas 100 separatas grátis. Poderão ser encomendadas cópias extra. O contexto dos artigos é da inteira responsabilidade dos respectivos autores.

Les ARQUIVOS DO MUSEU BOCAGE sont constitués par quatre séries, SÉRIE A; SÉRIE B — NOTAS; SÉRIE C — SUPLEMENTOS e SÉRIE D — EXTENSÃO CULTURAL e ENSINO, publiant des travaux sur Zoologie, Anthropologie et Biologie Générale, réalisés soit par des auteurs portugais, soit par des auteurs étrangers. Les travaux doivent être envoyés à l'Editeur et chacun d'eux composera un fascicule qui sera distribué tout de suite. Les volumes seront constitués par un nombre variable de fascicules, qui auront quatre cents à cinq cents pages, sans périodicité annuelle.

Le *titre* des travaux devra être concis et bien défini. Le nom(s) de l'auteur(s) sera suivi par celui de l'Institution où il travaille. Il est souhaitable qu'en tête de l'article l'auteur fasse figurer un «*abstract*» qui ne devra pas excéder 50 mots.

Le texte de l'article devra être suivi d'un *résumé* en français, anglais ou allemand.

La bibliographie devra être présentée selon l'exemple donné plus bas.

L'auteur aura droit à 100 tirés à part gratuits. Des copies supplémentaires peuvent être commandées. Le texte des articles sera de la responsabilité exclusive des auteurs respectifs.

The ARQUIVOS DO MUSEU BOCAGE are constituted by four series, SÉRIE A; SÉRIE B — NOTAS; SÉRIE C — SUPLEMENTOS and SÉRIE D — EXTENSÃO CULTURAL e ENSINO, containing original papers, made by Portuguese or foreign authors, within the general field of Zoology, Anthropology and General Biology. Manuscripts should be sent to the Editor. Each paper will form a separate number which appears at irregular intervals, as it becomes ready. Volumes will have about four or five hundred pages, and will not necessarily be completed within the calendar year.

The *title* of the papers should be concise and specific and the name(s) of the author(s) should be accompanied by that of the institution to which the author belongs.

An *abstract* will be printed at the head of all papers, that should generally not exceed 50 words. A *summary*, in English, French or German, should give a succinct account of the subject, results and conclusions.

Authors are requested to present the *literature* arranged alphabetically according the surnames of the authors. References will be indicated as below.

One hundred reprints of each paper will be supplied free, additional copies may be purchased. The text is of Author's exclusive responsibility.

SACARRÃO G. F. (1965) — On the origin and development of the epistellar body of the Octopus (*Octopus vulgaris* Lam.). *Arq. Mus. Boc.*, (2.^a série), 1, n.^o 1, 1-8.

Conselho de Redacção:

Prof. Dr. G. F. SACARRÃO

Prof. Dr. C. ALMAÇA

Lic.^a ANA M. NEVES

Editor:

Museu e Laboratório Zoológico

e Antropológico (Museu Bocage)

Faculdade de Ciências

R. Escola Politécnica, 58

1200 LISBOA. PORTUGAL

**I M P R E N S A
D E C O I M B R A ,
L I M I T A D A**