

## The University of Southern Mississippi The Aquila Digital Community

---

### Faculty Publications

---

10-1-2012

# Nutrient Depletion as a Proxy for Microbial Growth in Deepwater Horizon Subsurface Oil/Gas Plumes

Alan M. Shiller

*University of Southern Mississippi*, alan.shiller@usm.edu

Dong Joo Joung

*University of Southern Mississippi*, DongJoo.Joung@usm.edu

Follow this and additional works at: [http://aquila.usm.edu/fac\\_pubs](http://aquila.usm.edu/fac_pubs)



Part of the [Marine Biology Commons](#)

---

### Recommended Citation

Shiller, A. M., Joung, D. (2012). Nutrient Depletion as a Proxy for Microbial Growth in Deepwater Horizon Subsurface Oil/Gas Plumes. *Environmental Research Letters*, 7(4).

Available at: [http://aquila.usm.edu/fac\\_pubs/7653](http://aquila.usm.edu/fac_pubs/7653)

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Faculty Publications by an authorized administrator of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

## Nutrient depletion as a proxy for microbial growth in Deepwater Horizon subsurface oil/gas plumes

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2012 Environ. Res. Lett. 7 045301

(<http://iopscience.iop.org/1748-9326/7/4/045301>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 131.95.218.41

This content was downloaded on 17/07/2017 at 16:49

Please note that [terms and conditions apply](#).

You may also be interested in:

[The weathering of oil after the Deepwater Horizon oil spill: insights from the chemical composition of the oil from the sea surface, salt marshes and sediments](#)

Zhanfei Liu, Jiqing Liu, Qingzhi Zhu et al.

[Rapid microbial respiration of oil from the Deepwater Horizon spill in offshore surfacewaters of the Gulf of Mexico](#)

Bethanie R Edwards, Christopher M Reddy, Richard Camilli et al.

[Radiocarbon evidence that carbon from the Deepwater Horizon spill entered the planktonic food web of the Gulf of Mexico](#)

J P Chanton, J Cherrier, R M Wilson et al.

[Marine snow formation in the aftermath of the Deepwater Horizon oil spill in the Gulf of Mexico](#)

U Passow, K Ziervogel, V Asper et al.

[The influence of droplet size and biodegradation on the transport of subsurface oil droplets during the Deepwater Horizon spill: a model sensitivity study](#)

Elizabeth W North, E Eric Adams, Anne E Thessen et al.

[Evolution of the optical properties of seawater influenced by the Deepwater Horizon oil spill in the Gulf of Mexico](#)

Zhengzhen Zhou and Laodong Guo

[On the possible long-term fate of oil released in the Deepwater Horizon incident, estimated using ensembles of dye release simulations](#)

Mathew Maltrud, Synte Peacock and Martin Visbeck

# Nutrient depletion as a proxy for microbial growth in Deepwater Horizon subsurface oil/gas plumes

Alan M Shiller and DongJoo Joung

Department of Marine Science, The University of Southern Mississippi, 1020 Balch Boulevard, Stennis Space Center, MS 39529, USA

E-mail: [alan.shiller@usm.edu](mailto:alan.shiller@usm.edu)

Received 5 July 2012

Accepted for publication 25 September 2012


Published 18 October 2012

Online at [stacks.iop.org/ERL/7/045301](http://stacks.iop.org/ERL/7/045301)

## Abstract

The Deepwater Horizon accident resulted in a substantial uncontrolled hydrocarbon release to the northern Gulf of Mexico, much of which was entrained in deep submerged plumes. While bio-degradation of the hydrocarbons has been inferred from microbial biomass and genetics, the amount of conversion of oil and gas carbon to biomass remains uncertain having only been estimated in modeling studies. Here we examine correlated depletions of nitrate, phosphate and oxygen in the submerged plumes and conclude that a substantial portion of hydrocarbons in these plumes was converted to biomass ( $0.8\text{--}2 \times 10^{10}$  mol C). This contrasts with nutrient-limited surface waters where other work has suggested hydrocarbon-induced microbial growth to have been minimal. Our results suggest the need for better monitoring of changes in nutrients as well as study of nutrient recycling in similar future hydrocarbon releases.


**Keywords:** Deepwater Horizon, hydrocarbon degradation, Gulf of Mexico, oil spill, nutrients, oxygen


 Online supplementary data available from [stacks.iop.org/ERL/7/045301/mmedia](http://stacks.iop.org/ERL/7/045301/mmedia)

## 1. Introduction

The Deepwater Horizon blowout resulted in a release of  $\sim 5 \times 10^6$  bbl of crude oil (McNutt *et al* 2011) and  $\sim 10^{10}$  moles of methane (Kessler *et al* 2011, Reddy *et al* 2011) in the northern Gulf of Mexico. Essentially all of the methane and much of the oil entered deep water plumes (McNutt *et al* 2011). These hydrocarbons constituted a pool of potentially oxygen-depleting organic matter (Joye *et al* 2011), substrate for microbial growth (Hazen *et al* 2010, Redmond and Valentine 2011), and an injection of toxic materials such as polycyclic aromatic hydrocarbons (Diercks *et al* 2010). Based

on dissolved oxygen and methane distributions, Kessler *et al* (2011) found that the plume hydrocarbons were primarily consumed by the microbial community. Implicit is that the hydrocarbon release stimulated a substantial deep water microbial bloom (Hazen *et al* 2010, Redmond and Valentine 2011, Kessler *et al* 2011). A physical-metabolic model of the release (Valentine *et al* 2012) yielded an estimate of the microbial bloom of  $10^{11}$  g C, equivalent to about half of the carbon injected into the Gulf during the blowout. And, an oxygen-based calculation yielded a similar estimate (Du and Kessler 2012). Such estimates are important because they help provide an understanding of the fate of the hydrocarbons, including both the rapidity with which the methane disappeared (Kessler *et al* 2011) as well as in what forms and to which geochemical reservoirs the released hydrocarbons were delivered. Herein, we examine nutrient distributions in deep waters in the vicinity of the wellhead

 Content from this work may be used under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 licence](http://creativecommons.org/licenses/by-nc-sa/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



**Figure 1.** Dissolved nutrient and oxygen profiles in four cruises to the vicinity of the Deepwater Horizon blowout site. For the May and October 2010 cruises, profiles were constructed from samples collected at different locations around the wellhead.

as a means of providing another estimate of how much the hydrocarbon consumption supported growth rather than just respiration to carbon dioxide.

## 2. Methods and materials


Water sampling in the vicinity of the Deepwater Horizon blowout was conducted aboard the R/V *Pelican* (10–14 May 2010), R/V *F.G. Walton Smith* (26 May–1 June 2010), and R/V *Cape Hatteras* (11–20 October 2010 and 20–29 October 2011) using each ship's rosette-mounted Niskin bottles. During the first two cruises, samples were collected during multiple hydrocasts in a southwesterly trend within 27 km of the wellhead. During October 2010, samples were also collected during multiple hydrocasts, this time in all directions and ranging to over 100 km away from the wellhead. For October 2011, the data are from one hydrocast at a station within 5 km of the wellhead. Station locations and hydrocast data are listed in the supplementary data tables (available at [stacks.iop.org/ERL/7/045301/mmedia](http://stacks.iop.org/ERL/7/045301/mmedia)).

Samples were filtered ( $0.45 \mu\text{m}$ ) and frozen ( $-20^\circ\text{C}$ ) until analysis. For the first three cruises, nutrient analyses were performed at USM using an Astoria-Pacific A2+2 nutrient auto-analyzer (Astoria-Pacific International, OR, USA). For samples from October 2011, nutrients were analyzed by auto-analyzer at the Geochemical and Environmental Research Group (GERG; Texas A&M University). A subset of frozen samples from the first three cruises was also analyzed at the same time at GERG to make sure that results from all four cruises were properly intercalibrated. Dissolved oxygen data were taken from the ship's *in situ* oxygen sensor mounted on the rosette sampling system. Oxygen calibrations were performed by standard Winkler titrations. Failure of the titration system during the late May 2010 cruise probably accounts for the slight ( $<10 \mu\text{mol kg}^{-1}$ ) overall negative shift of those oxygen data relative to October 2011, but did not affect the slope of the oxygen–nutrient relationship, which was nearly identical in the 2 May 2010 cruises despite the data being obtained at different times and aboard different ships.


## 3. Results and discussion

During the May 2010 cruises, profiles of *in situ* fluorescence and beam attenuation showed peaks in the depth range of 600–1300 m, which were subsequently identified as subsurface plumes of oil and gas (e.g., Diercks *et al* 2010, Camilli *et al* 2010). We observed slight decreases in dissolved nitrate and phosphate as well as oxygen in this depth range of the subsurface plumes during the 2 May 2010 cruises (figure 1). In contrast, the two post-blowout cruises (October 2010 and 2011) showed no evidence of submerged oil/gas plumes and likewise did not show the nutrient or oxygen decreases (figure 1). Nutrient and oxygen values for samples outside of the submerged plumes were comparable among all cruises. Also, for dissolved silica, no depletions were observed. Dissolved nitrite and ammonium concentrations were less than 1% of the nitrate in the May 2010 nutrient-depleted samples, indicating that nitrate decreases were not reflected in proportionate increases in other forms of dissolved inorganic nitrogen. Interestingly, although elevated levels of polycyclic aromatic hydrocarbons (PAHs) were found at plume depths, no correlation was found between PAH concentrations and nutrient depletions. All of these observations lend confidence to the interpretation of the nutrient depletions as not resulting from analytical or sampling artifacts, but rather from biological uptake. Our results can be contrasted with those of Hazen *et al* (2010), who found a significant decrease in nitrate concentrations inside of submerged plume waters (mean decrease  $6 \mu\text{M}$ ) in late May 2010, but found no significant decrease in plume phosphate concentrations (mean decrease  $0.03 \mu\text{M}$ ).

We also examined other chemical parameters in the water samples (e.g., trace elements) to verify that the nutrient depletions were not caused by sample bottles accidentally tripping at a shallow depth. Pre/post-tripped bottles would not have affected the oxygen data since these were obtained from the CTD oxygen sensor. A few samples were eliminated from our dataset because all examined water parameters suggested that the Niskin bottle closed at a shallower depth than expected. In order to quantify the extent of the oxygen and nutrient depletions, profiles of oxygen and nutrients in the 600–1600 m depth range from October 2011 were fitted



**Figure 2.** Dissolved nutrient and oxygen anomalies in the subsurface near the Deepwater Horizon blowout site during two cruises in May 2010. (A) Oxygen anomaly versus depth. (B) Oxygen anomaly versus nitrate anomaly. (C) Phosphate anomaly versus nitrate anomaly.



**Figure 3.** Dissolved nitrite versus the nitrate anomaly in the subsurface near the Deepwater Horizon blowout site during May 2010.

**Table 1.** Coefficients for third order polynomial fits to oxygen and nutrient data, Oct. 2011, 600–1600 m. Depth ( $z$ ) in meters; concentrations in  $\mu\text{M}$ .

	Oxygen	Nitrate	Phosphate
$z^3$	$-8.6567 \times 10^{-8}$	$1.6612 \times 10^{-8}$	$-1.7679 \times 10^{-10}$
$z^2$	$2.1681 \times 10^{-4}$	$-5.3416 \times 10^{-5}$	$6.6621 \times 10^{-7}$
$z$	$-5.5765 \times 10^{-2}$	$4.6401 \times 10^{-2}$	$-1.2431 \times 10^{-3}$
Const.	$1.0662 \times 10^2$	$1.7482 \times 10^1$	$2.5665 \times 10^0$

to third order polynomials (table 1) in order to estimate concentrations at depths that were not sampled in 2011. Thus, the May 2010 oxygen and nutrient anomalies are differences relative to the concentrations determined in the October 2011 profile.

Phosphate and nitrate anomalies had a positive correlation with the oxygen anomaly, and the phosphate and nitrate anomalies were also well correlated (figure 2). Again, these observations suggest that the nutrients were removed together with the oxygen in the subsurface plume. The molar oxygen/nitrate anomaly ratio was approximately 8, while the N/P depletion ratio was 12–13 (table 2). Scatter in the graphs of  $\text{O}_2$ , N and P anomalies makes it difficult to

**Table 2.** Submerged plume oxygen and nutrient molar anomaly ratios determined during two cruises in the vicinity of the Deepwater Horizon wellhead during May 2010.

	$\text{O}_2/\text{NO}_3^-$	$\text{NO}_3^-/\text{PO}_4^{3-}$	$n$
10–14 May 2010	$7.7 \pm 1.9$	$12.0 \pm 1.6$	7
26 May–1 June 2010	$8.3 \pm 1.4$	$12.8 \pm 0.7$	29

predict which could have theoretically been depleted to zero first (i.e., which was most limiting); however, our empirical relationships suggest that all three would have run out at nearly the same time. Nonetheless, given that reported oxygen depletions throughout the blowout period did not exceed ~25% of ambient concentrations (e.g., Kessler *et al* 2011, Du and Kessler 2012), it seems unlikely that anything close to N or P limitation would have occurred during or after the blowout. We also observed that the highest deep water values of nitrite were associated with the highest nitrate depletions (figure 3), suggesting the possibility of nutrient recycling associated with the nutrient depletions.

Assuming that nutrient depletion dominantly reflects microbial growth, the correlated nutrient and oxygen depletions reported here provide a way to estimate the overall microbial production in the submerged oil/gas plumes. We do this by starting with the total integrated subsurface oxygen depletion from Du and Kessler (2012), who estimated  $1.9 (\pm 0.4) \times 10^{10}$  mol of  $\text{O}_2$  were respired in the plume during the blowout. Assuming our oxygen/nitrate depletion ratio (table 2; figure 2) held for the entire blowout, then dividing that ratio into the integrated oxygen depletion yields a total blowout nitrate uptake of  $1.6\text{--}4.0 \times 10^9$  mol N. (The range in this estimate is determined by both the uncertainty in the oxygen depletion and the uncertainty in the  $\text{O}_2/\text{NO}_3^-$  anomaly ratio.) We then multiply this result by a bacterial C/N molar ratio of 5 (Goldman *et al* 1987), yielding  $0.77\text{--}2.0 \times 10^{10}$  mol microbial C produced. This is similar in magnitude to the  $10^{11}$  g C of bacterial productivity (i.e.,  $0.8 \times 10^{10}$  mol C) estimated in the modeling study of Valentine *et al* (2012). To put these numbers into a local context, the estimated bacterial carbon production in the submerged oil/gas plume is about one tenth of the annual primary production associated



with the outflow of the Mississippi River (e.g., Lohrenz *et al* 1997, Chen *et al* 2000). While the time, specific location, and areal scales of these two events are clearly different, the comparison nonetheless indicates that the bloom associated with the blowout was not outside of the range of organic matter fixation occurrences in the northern Gulf of Mexico. Toxic effects associated with the crude oil, however, are another matter.

While all of the numbers in our calculation have significant uncertainties, the result is similar to the total moles of C released in the blowout ( $2.0\text{--}2.4 \times 10^{10}$  mol C; Kessler *et al* 2011), supporting the notion of a massive bloom response (i.e.,  $66 \pm 33\%$  conversion of hydrocarbon-C to biomass C). Because we did not obtain samples during later stages of the blowout nor have nutrient data been reported as part of the government response ([www.nodc.noaa.gov/General/DeepwaterHorizon/support.html](http://www.nodc.noaa.gov/General/DeepwaterHorizon/support.html)), we cannot be certain that microbial growth occurred throughout the period of uncontrolled hydrocarbon release. This would suggest our estimate is an upper bound. We also have no information on the extent of nutrient recycling, which would result in our oxygen/nitrate depletion ratio underestimating the extent of nitrate uptake. And, we cannot state whether the sole C source for the bloom was hydrocarbons. However, methanotrophs are known to use methane as their sole carbon source, though not all species may be obligate in that regard (Dedysh *et al* 2005). We do note that isotopic evidence from shallower nearshore regions affected by the blowout supports the hypothesis of transfer of hydrocarbon-C to biomass, at least in that environment (Graham *et al* 2010). We conclude, therefore, that hydrocarbons in the subsurface plume were likely substantially converted to biomass. In contrast, in surface waters contaminated with Deepwater Horizon crude oil, Edwards *et al* (2011) found evidence of hydrocarbon-supported respiration but with only limited microbial growth due to low nutrient concentrations.

The results presented here suggest the need for and utility of nutrient measurements in studies of similar future hydrocarbon releases. Such studies, including concentration measurements throughout the course of the event, as well as estimates of the extent of nutrient recycling, have the potential to shed light on the biological response to and the initial fate of the hydrocarbons.

## Acknowledgments

Funding for this work was provided by the National Science Foundation (OCE-1042934) and the Northern Gulf Institute/BP (10-BP-GRI-USM-01, Task 2). The comments of three reviewers substantially improved the manuscript. We thank R C Highsmith (U Miss), S B Joye (U Ga), and K M Yeager (U Ky)/C A Brunner (USM) for the opportunity to participate in the R/V *Pelican*, R/V *Walton Smith* and R/V *Cape Hatteras* cruises, respectively. Funding for those cruises was provided by NOAA (to RCH), NSF

(OCE-1043225 to SBJ), and the Northern Gulf Institute/BP (10-BP-GRI-USM-01 & 11-BP-GRI-24 to KMY & CAB); NSF also provided some supplemental funding for the 2010 R/V *Cape Hatteras* cruise.

## References

- Camilli R, Reddy C M, Yoerger D R, Van Mooy B A S, Jakuba M V, Kinsey J C, McIntyre C P, Sylva S P and Maloney J V 2010 Tracking hydrocarbon plume transport and biodegradation at Deepwater Horizon *Science* **330** 201–4
- Chen X, Lohrenz S E and Wiesenburg D A 2000 Distribution and controlling mechanisms of primary production on the Louisiana–Texas continental shelf *J. Mar. Syst.* **25** 179–207
- Dedysh S N, Knief C and Dunfield P F 2005 Methylocella species are facultatively methanotrophic *J. Bacteriol.* **187** 4665–70
- Diercks A R *et al* 2010 Characterization of subsurface polycyclic aromatic hydrocarbons at the Deepwater Horizon site *Geophys. Res. Lett.* **37** L20602
- Du M and Kessler J D 2012 Assessment of the spatial and temporal variability of bulk hydrocarbon respiration following the Deepwater Horizon oil spill *Environ. Sci. Technol.* **46** 10499–507
- Edwards B R, Reddy C M, Camilli R, Carmichael C A, Longnecker K and Van Mooy B A S 2011 Rapid microbial respiration of oil from the *Deepwater Horizon* spill in offshore surface waters of the Gulf of Mexico *Environ. Res. Lett.* **6** 035301
- Goldman J C, Caron D A and Dennett M R 1987 Regulation of gross growth efficiency and ammonium regeneration in bacteria by substrate C:N ratio *Limnol. Oceanogr.* **32** 1239–52
- Graham W M, Condon R H, Carmichael R H, D’Ambr I, Patterson H K, Linn L J and Hernandez F J Jr 2010 Oil carbon entered the coastal planktonic food web during the Deepwater Horizon oil spill *Environ. Res. Lett.* **5** 045301
- Hazen T C *et al* 2010 Deep-sea oil plume enriches indigenous oil-degrading bacteria *Science* **330** 204–8
- Joye S B, MacDonald I R, Leifer I and Asper V 2011 Magnitude and oxidation potential of hydrocarbon gases released from the BP oil well blowout *Nature Geosci.* **4** 160–4
- Kessler J D *et al* 2011 A persistent oxygen anomaly reveals the fate of spilled methane in the deep Gulf of Mexico *Science* **331** 312–5
- Lohrenz S E, Fahnenstiel G L, Redalje D G, Lang G A, Chen X and Dagg M J 1997 Variations in primary production of northern Gulf of Mexico continental shelf waters linked to nutrient inputs from the Mississippi river *Mar. Ecol. Prog. Ser.* **155** 45–54
- McNutt M K, Camilli R, Crone T J, Guthrie G D, Hsieh P A, Ryerson T B, Savas O and Shaffer F 2011 Review of flow rate estimates of the *Deepwater Horizon* oil spill *Proc. Natl Acad. Sci. USA* at press (doi:10.1073/pnas.1112139108)
- Reddy C M *et al* 2011 Composition and fate of gas and oil released to the water column during the Deepwater Horizon oil spill *Proc. Natl Acad. Sci. USA* at press (doi:10.1073/pnas.1101242108)
- Redmond M C and Valentine D L 2011 Natural gas and temperature structured a microbial community response to the Deepwater Horizon oil spill *Proc. Natl Acad. Sci. USA* at press (doi:10.1073/pnas.1108756108)
- Valentine D L, Mezić I, Mačević S, Črnjarić-Žic N, Ivić S, Hogan P J, Fonoberov V A and Loire S 2012 Dynamic autoinoculation and the microbial ecology of a deep water hydrocarbon irruption *Proc. Natl Acad. Sci. USA* at press (doi:10.1073/pnas.1108820109)