

## Northeast Gulf Science

---

Volume 2  
Number 2 Number 2

Article 11

---

12-1978

# Spawning of the Striped Mullet, *Mugil cephalus*, in the Northwestern Gulf of Mexico

John H. Finucane  
*National Marine Fisheries Service*

L. Alan Collins  
*National Marine Fisheries Service*

Lyman E. Barger  
*National Marine Fisheries Service*

DOI: 10.18785/negs.0202.11

Follow this and additional works at: <https://aquila.usm.edu/goms>

---

### Recommended Citation

Finucane, J. H., L. Collins and L. E. Barger. 1978. Spawning of the Striped Mullet, *Mugil cephalus*, in the Northwestern Gulf of Mexico. *Northeast Gulf Science* 2 (2). Retrieved from <https://aquila.usm.edu/goms/vol2/iss2/11>

This Article is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Gulf of Mexico Science by an authorized editor of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

*Southeast Fisheries Center, Panama City Laboratory, 3500 Delwood Beach Road, Panama City, FL 32407.*

## SPAWNING OF THE STRIPED MULLET, *Mugil cephalus*, IN THE NORTHWESTERN GULF OF MEXICO<sup>1</sup>

Pelagic eggs of striped mullet, *Mugil cephalus*, were identified from ichthyoplankton samples taken on December 2, 1977, from 89 to 98 km off the Texas coast in the northwestern Gulf of Mexico. This is the first reported spawning of striped mullet in this area. Diameters of eggs ranged from 0.91 to 0.99 mm and averaged 0.95 mm, and diameters of oil globules ranged from 0.30 to 0.36 mm and averaged 0.33 mm. Newly hatched yolk sac larvae were 2.1 mm long.

The striped mullet, *Mugil cephalus* Linneaus, has a cosmopolitan distribution. It occurs in the western Atlantic from Nova Scotia to Santos, Brazil (Briggs 1958). During 1977, 9,926 metric tons were caught by commercial fishermen in the United States (U. S. Department of Commerce 1978). It is an important species in Florida waters where it is also called black mullet and liza.

Although much has been published on the biology of the striped mullet, little is known about the exact locations of their spawning grounds in the Gulf of Mexico. Many authors reported that striped mullet spawn either inshore or within a few miles of the beach (Higgins 1927, Hildebrand and Schroeder 1928, Breder 1940, Gunter 1945, and Taylor 1951). Broadhead (1953) inferred that striped mullet probably spawn 9-37 kilometers off the northwest Florida coast.

Anderson (1958) also inferred that striped mullet spawned offshore from the 37 meter line to the Gulf Stream along the Atlantic coast from North Carolina to lower Florida. Prior to our ichthyoplankton survey the only known spawning area in the Gulf of Mexico was reported by Arnold and Thompson (1958), who found striped mullet eggs, larvae and juveniles 74-92 kilometers southeast of the Mississippi River Delta over the continental shelf in water 914 to 1645 m deep.

Our mullet eggs were collected in 1977 during one of nine ichthyoplankton cruises made in the northwestern Gulf of Mexico. The area covered about 8,760 sq km extending from Port Isabel, Texas near the Mexican border to Matagorda Bay (Figure 1).

All ichthyoplankton collections were made with paired bongo net samplers, using standard sampling techniques and equipment (Jossi *et al.* 1975, Smith and Richardson 1977). Tows were double




Figure 1. Ichthyoplankton sampling area showing station locations and locations of striped mullet egg collections during 1977 off the south Texas coast.

<sup>1</sup>Contribution No. 79-01PC, NMFS, SEFC, Panama City Laboratory, 3500 Delwood Beach Road, Panama City, FL 32407.

oblique and were made during both day and night from the stern of the ship. Salinity and temperature measurements were taken with STD casts at each station.

Plankton samples were preserved in about 7% buffered Formalin in seawater. Ichthyoplankton was sorted from the entire 0.333 mm net sample. All measurements were made with a stereoscopic microscope using an ocular micrometer. Identifications were made using published descriptions of striped mullet (Sanzo 1936, Vodianitskii and Kazanova 1954, Tang 1964, Yashouv and Berner-Samsonov 1970, and Shehadeh and Norris 1972).

Striped mullet eggs were found in samples collected on December 2, 1977, at Stations II-3 and II-7 located 89 to 98 km offshore from Port Aransas, Texas, at depths of 131 and 183 m (Figure 1). Mean water temperatures and salinities through the water column were 22.2°C and 36.4‰ at Station II-3 and 22.4°C and 36.5‰ at Station II-7. The total number of eggs collected was 1,112 at Station II-3 and 4,128 at Station II-7. Egg abundances were estimated to be 3,763 and 12,684 under 10m<sup>2</sup> of sea surface, respectively. Diameters of eggs ranged from 0.91 to 0.99 mm and averaged 0.95 mm, while the oil globules ranged from 0.30 to 0.36 mm and averaged 0.33 mm in diameter. Three yolk sac larvae with mean notochord length of 2.1 mm were also recorded from the samples.

The spawning times of the striped mullet are well documented. Anderson (1958) stated that striped mullet spawned along the South Atlantic coast of the United States from late November to early December or into February. Observations by Gunter (1945) indicated that spawning of striped mullet off the Texas coast occurs from late October

to early January with the peak in late November and early December. More recent work by Moore (1974) in the estuarine and coastal waters off Port Aransas showed that post-spawning striped mullet of both sexes appeared from December through May, indicating that some fish had completed spawning by December and that all fish had done so by May. He also suggested that individual mullet spawn more than once during the same spawning season. Our findings indicate that the striped mullet spawns in early winter in the northwestern Gulf of Mexico and support those of Arnold and Thompson (1958).

Based on water temperatures during the time of collection, the mullet eggs probably were about two days old. Hatching of *M. cephalus* eggs in the laboratory at 22°C required 48-50 hrs (Shehadeh and Norris 1972). Drift bottles released in the sampling area in December generally moved in a southwestern direction at a velocity of 10 km or more per day (U. S. Department of Commerce 1976). Thus the approximate spawning site was over the edge of the continental shelf, about 20 km northeast of our sampling site.

#### ACKNOWLEDGEMENTS

We thank Anne H. Naplin and Elbert H. Ahlstrom for their verification of our mullet egg identification. They and Edward D. Houde reviewed the paper.

#### LITERATURE CITED

- Anderson, W. W. 1958. Larval development, growth and spawning of striped mullet (*Mugil cephalus*) along the south Atlantic coast of the United States. Fish. Bull., U. S. 58:501-519.

- Arnold, E. L., Jr., and J. R. Thompson. 1958. Offshore spawning of the striped mullet, *Mugil cephalus*, in the Gulf of Mexico. *Copeia* 1958:130-132.
- Breder, C. M., Jr. 1940. The spawning of *Mugil cephalus* on the Florida coast. *Copeia* 1940:138-139.
- Briggs, J. C. 1958. A list of Florida fishes and their distribution. *Bull. Fla. St. Mus., Bio. Sci.* 2(8):318 p.
- Broadhead, G. C. 1953. Investigation of the black mullet, *Mugil cephalus* L., in northwest Florida. *Fla. St. Bd. Conserv., Tech. Ser.* 7, 21 p.
- Gunter, G. 1945. Studies on marine fishes of Texas. *Pub. Inst. Mar. Sci.* 1(1):190 p.
- Higgins, E. 1927. Progress in biological inquiries, 1926. *Rept. U. S. Comm. Fish., Bur. Fish. Doc.* 1029:624-627.
- Hildebrand, S. F., and W. C. Schroeder. 1928. Fishes of Chesapeake Bay. *Fish. Bull., U. S.* 43(pt. 1):366 p.
- Jossi, J. W., R. R. Marak, and H. Peterson, Jr. 1975. At-sea data collection and laboratory procedures. *In: MAR-MAP Survey I Manual, NMFS, Wash., D. C.* 115 p.
- Moore, R. H. 1974. General ecology, distribution and relative abundance of *Mugil cephalus* and *Mugil curema* on the south Texas coast. *Contr. Mar. Sci.* 18:241-245.
- Sanzo, L. 1936. Contributi ala conoscenza dello sviluppo embrionario e postembrionario nei Mugilidi. I. Uova e larve di *Mugil cephalus* Cuv. ottenute per fecondazione artificiale. II. Uova e larve di *Mugil chelo* Cuv. *R. Com. Talassogr. Ital. Mem.* 230, 11 p.
- Shehadeh, Z. H., and K. S. Norris. 1972. The grey mullet (*Mugil cephalus*) induced breeding and larval rearing research 1970-1972. *Nat. Sea Grant Program, NOAA, U. S. Dept. Comm. Rockville, Md.*, 202 p.
- Smith, P. S., and S. R. Richardson. 1977. Standard techniques for pelagic fish egg and larva surveys. *F. A. O. Fish. Tech. Paper No. 175*, 100 p.
- Tang, Y. 1964. Induced spawning of striped mullet by hormone injection *Jap. J. Ichthyol.* 12(1/2):23-28.
- Taylor, H. F. 1951. Survey of marine fisheries of North Carolina. *Chapel Hill Univ. N. C. Press*, 555 p.
- Vodianitskii, V. A., and I. I. Kazanova. 1954. Diagnostic descriptions of the eggs and larvae of the Black Sea fishes. (In Russian). *Trudy VNIRO* 28: 240-323.
- U. S. Department of Commerce. 1976. *In: NOAA Final Report to BLM: Environmental studies of the south Texas outer continental shelf 1975. Vol II Physical oceanography*, 290 p.
- \_\_\_\_\_. 1978. *Fisheries of the United States, 1977. U. S. Dept. Comm., NOAA, NMFS*, 112 p.
- Yashouv, A., and E. Berner-Samsonov. 1970. Contributions to the knowledge of eggs and early larval stages of mullets (MUGILIDAE) along the Israel coast. *Bamidgeh* 22(3):22-89.
- John H. Finucane, L. Alan Collins and Lyman E. Barger. *National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southeast Fisheries Center, Panama City Laboratory, 3500 Delwood Beach Road, Panama City, FL 32407.*

## REVIEWERS

Below are listed those scientists who donated their time during 1978 as referees for manuscripts submitted to Northeast Gulf Science. To them we extend our sincere thanks.

- Stephen Bortone  
University of West Florida
- Mark Chittenden  
Texas A & M University
- Bruce Coull  
University of South Carolina
- John Dean  
University of South Carolina
- John Dindo  
Dauphin Island Sea Lab
- Joseph Dorn  
Spring Hill College
- Charles Duncan  
University of Alabama at Birmingham
- William Fink  
MCZ, Harvard University
- Carter Gilbert  
University of Florida
- Gordon Gunter  
Gulf Coast Research Laboratory
- Robert Hastings  
Rutgers University
- Richard Heard  
Gulf Coast Research Laboratory
- William Kruczynski  
EPA, Atlanta, Georgia
- Michel Lelong  
University of South Alabama
- Thomas Lewis  
Florida State University
- Robert Livingston  
Florida State University
- Don Reish  
California State University, Long Beach
- Sally Richardson  
Gulf Coast Research Laboratory
- Joseph Simon  
Florida Dept. of Natural Resources
- Franklin Snelson  
Florida Technological University
- Marlin Tagatz  
EPA, Gulf Breeze, FL
- James C. Tyler  
National Marine Fisheries Service