

Spring 5-1-2015

## Impacts of the Deepwater Horizon Oil Spill on Microbial-Mediated Cellulose Decomposition in Mississippi Gulf Coast Salt Marshes

Jerrid Shawn Boyette  
*University of Southern Mississippi*

Follow this and additional works at: [https://aquila.usm.edu/masters\\_theses](https://aquila.usm.edu/masters_theses)



Part of the [Biology Commons](#), and the [Environmental Microbiology and Microbial Ecology Commons](#)

---

### Recommended Citation

Boyette, Jerrid Shawn, "Impacts of the Deepwater Horizon Oil Spill on Microbial-Mediated Cellulose Decomposition in Mississippi Gulf Coast Salt Marshes" (2015). *Master's Theses*. 103.  
[https://aquila.usm.edu/masters\\_theses/103](https://aquila.usm.edu/masters_theses/103)

This Masters Thesis is brought to you for free and open access by The Aquila Digital Community. It has been accepted for inclusion in Master's Theses by an authorized administrator of The Aquila Digital Community. For more information, please contact [Joshua.Cromwell@usm.edu](mailto:Joshua.Cromwell@usm.edu).

The University of Southern Mississippi

IMPACTS OF THE DEEPWATER HORIZON OIL SPILL ON  
MICROBIAL-MEDIATED CELLULOSE DECOMPOSITION IN  
MISSISSIPPI GULF COAST SALT MARSHES

by

Jerrid Shawn Boyette

A Thesis  
Submitted to the Graduate School  
of The University of Southern Mississippi  
in Partial Fulfillment of the Requirements  
for the Degree of Master of Science

Approved:

Dr. Kevin A. Kuehn  
Committee Chair

Dr. Micheal A. Davis

Dr. Carl P. Qualls

Dr. Karen S. Coats  
Dean of the Graduate School

May 2015

ABSTRACT

IMPACTS OF THE DEEPWATER HORIZON OIL SPILL ON  
MICROBIAL-MEDIATED CELLULOSE DECOMPOSITION IN  
MISSISSIPPI GULF COAST SALT MARSHES

by Jerrid Shawn Boyette

May 2015

Field studies were conducted to examine the effects of the Deepwater Horizon oil spill on rates of marsh organic matter decomposition. Decomposition in surface and subsurface marsh sediments was assessed in stands of *Spartina alterniflora* and *Juncus roemerianus* in 9 Mississippi Gulf Coast marshes exposed to differing oiling intensities. The cotton strip bioassay technique was used as a proxy for cellulose decomposition. In addition, rates of microbial respiration, fungal biomass (ergosterol), and nutrients (C:N, C:P) of surface sediment cotton strips were also quantified. Subsurface cotton strip decay, as determined by losses in tensile strength, were significantly different among marsh sites, with higher overall rates being observed in oiled versus unoiled *S. alterniflora* plant zones ( $p < 0.05$ ). No differences were observed in subsurface sediments between oiled and unoiled *J. roemerianus* plant zones ( $p > 0.05$ ). In contrast to subsurface sediments, cotton strip decay in surface sediments displayed an opposite pattern, with significantly ( $p > 0.05$ ) higher rates of decay in unoiled versus oiled *S. alterniflora* and *J. roemerianus* plant zones. Cotton strip C:N and C:P ratios were negatively correlated with losses in cotton strip tensile strength. In addition, both fungal ergosterol concentrations and microbial respiration rates were positively correlated with cotton strip decay and negatively correlated with C:N and C:P ratios, providing evidence that N and P availabilities in

marsh sediments may have limited the activity of microbial communities. Although conducted ~1.5 years after the Deep Water Horizon oil spill, this study suggests that both subsurface and surface microbial processes may still be affected by oil.

## ACKNOWLEDGMENTS

I would like to thank my committee chair, Dr. Kevin A. Kuehn, and my committee members, Dr. Micheal A. Davis and Dr. Carl P. Qualls, for their assistance and input throughout this project. I would like to thank Dr. Scott Teigs for assisting in data collection and analysis. I would like to thank Dr. Jake Schaefer and his lab for providing us with transportation to our sites. I would like to extend a big thank you to Rong Su, Stephanie Koury, and Rachel Smilo, for all their support and help, Knox Flowers for assisting me in navigating between sites, and Jessica Eves for all the assistance she provided in the lab. I would like to thank Denise and Jamey Boyette for all their support, and also I would like to thank my loving partner, Eric Tribunella, for all his patience and support throughout this project. Last, I would like to thank the Gulf of Mexico Research Initiative (GoMRI) for funding this project and making this research possible.

TABLE OF CONTENTS

ABSTRACT..... ii

ACKNOWLEDGMENTS.....iv

LIST OF TABLES.....vi

LIST OF ILLUSTRATIONS.....vii

CHAPTER

I. INTRODUCTION..... 1

II. MATERIALS AND METHODS.....4

    Study Sites

    Field Procedures

    Cellulose Decomposition Rates

    Rates of Microbial Respiration

    Fungal Ergosterol and Nutrient Concentrations

    Data Analyses

III. RESULTS..... 11

    Cementation Effects

    Environmental Parameters

    Cotton Strip Decay - Subsurface Sediments

    Cotton Strip Decay - Surface Sediments

    Cotton Strip Nutrient Dynamics

    Cotton Strip Fungal Colonization and Rates of Microbial Respiration

IV. DISCUSSION.....34

REFERENCES.....42

LIST OF TABLES

Table

1. Marsh study sites examine and their varying degrees of oil exposure as determined from Shoreline Cleanup Assessment Team (SCAT) surveys.....6

2. Salinity, pH and reduction potential (ORP) of sediment pore water within *S. alterniflora* and *J. roemerianus* plant zones. Values are the mean±1SD (n=12, except site 4 for *J. roemerianus* where n=1).....13

3. Nested ANOVA summary table indicating the effect of site, plant species zone and oiling on sediment pore water salinity, pH and reduction potential (ORP) at each marsh site.....15

4. Nested ANOVA summary table indicating the effect of site, plant species zone and oiling on subsurface and surface percent cotton strip tensile strength loss.....18

5. Overall tensile strength loss (%CTSL/d) of surface and subsurface cotton strips in oiled and unoled *S. alterniflora* and *J. roemerianus* plant zones. Values are the mean±1SD (n=15-30).....18

6. Nested ANOVA summary table indicating the effect of site, plant species zone and oiling on the carbon:nitrogen and carbon:phosphorus ratios of cotton strips incubated at the marsh sediment surface.....24

7. Overall patterns of carbon:nitrogen and carbon:phosphorus ratios for cotton strips incubated at the marsh surface sediments in *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoled marsh sites. Values are the mean±1SD (n=15-30).....24

8. Pearson product-moment intercorrelation matrix showing the relationships between rates of surface sediment cotton strip decay (%CTSL/d) and cotton strip-associated fungal ergosterol concentrations, rates of microbial respiration and carbon:nitrogen and carbon:phosphorus ratios within oiled and unoled marsh sites. Values are the mean±1SD (n=15-30).....27

9. Nested ANOVA summary table indicating the effect of site, plant species zone and oiling on cotton strip-associated fungal ergosterol concentrations and microbial respiration rates in cotton strips incubated at the marsh sediment surface.....30

10. Overall patterns of fungal ergosterol concentrations and rates of microbial respiration associated with cotton strips incubated at the marsh surface sediments in *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoled marsh sites. Values are the mean±1SD (n=15-30).....31

LIST OF ILLUSTRATIONS

Figure

1. Map of the Mississippi Gulf Coast illustrating the sample study sites and the known oiling impacts based on the Shoreline Cleanup Assessment Team (SCAT) surveys conducted immediately after the oil spill.....5

2. Changes in surface (A) and subsurface (B) sediment temperatures during the study period. Values are the overall mean±SD for all study sites combined.....12

3. Losses in cotton strip tensile strength observed in subsurface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean±SE (n=5).....17

4. Overall patterns of cotton strip tensile strength loss observed in subsurface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).....19

5. Losses in cotton strip tensile strength observed in surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean±SE (n=5).....20

6. Overall patterns of cotton strip tensile strength loss observed in surface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).....21

7. Carbon:nitrogen ratio patterns for cotton strips incubated at the marsh surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean±SE (n=5).....23

8. Overall patterns of carbon:nitrogen (A) and carbon:phosphorus (B) ratios for cotton strip incubated at the marsh surface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).....25

9. Carbon:phosphorus ratio patterns for cotton strips incubated at the marsh surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean±SE (n=5).....26

10. Fungal ergosterol concentrations associated with cotton strips incubated at the marsh surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean±SE (n=5).....29



11. Overall patterns of fungal ergosterol concentrations associated with cotton strips incubated at the marsh surface sediments in the <i>S. alterniflora</i> and <i>J. roemerianus</i> plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).....	30
12. Rates of microbial respiration (CO <sub>2</sub> evolution) associated with cotton strips incubated at the marsh surface sediments in the <i>S. alterniflora</i> (A) and <i>J. roemerianus</i> (B) plant zones at each marsh site. Values are the mean±SE (n=5).....	32
13. Overall patterns of microbial respiration rates (CO <sub>2</sub> evolution) associated with cotton strips incubated at the marsh surface sediments in the <i>S. alterniflora</i> and <i>J. roemerianus</i> plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).....	33

## CHAPTER I

### INTRODUCTION

In the United States alone, approximately 62,000 km<sup>2</sup> of wetland area occur along the Gulf Coast (Stedman and Dahl 2008) and assessing the health and stability of these ecosystems is of vital importance. These coastal wetlands provide a variety of ecological and economic services to the region, which include water quality enhancement, flood protection, defense of coastal shorelines from erosion, and the provision of habitat and food resources for wildlife. Furthermore, these wetlands provide an important source and sink for carbon (C) and nutrients (nitrogen and phosphorus), and their processing and storage is widely recognized as a critical ecosystem service (McLeod et al. 2011).

Globally, salt marshes are recognized as among the most productive ecosystems on earth, with rates of annual net primary productivity ranging from 800-4,000 g/m<sup>2</sup>/y (Mitsch and Gosselink 2007). In the Gulf Coast region, the emergent vascular plants *Spartina alterniflora* and *Juncus roemerianus* are the two most common marsh species and frequently form the largest fraction of the annual marsh plant biomass produced (White et al. 1978; Christian et al. 1990; Dai and Wiegert 1996; Ewe et al. 2006). Most of this plant biomass enters the detrital pool following senescence and death of the plant shoot, where the accumulation and subsequent decomposition of plant litter is important for marsh soil development and accretion, storage of C and nutrients, and supply of detrital-based food resources that fuels both estuarine and near-shore food webs (Cebrian and Lartigue 2004; Moore et al. 2004; Craft 2007; Mitsch and Gosselink 2007; Neubauer 2008).

There are justifiable concerns over the effects of the Deepwater Horizon Oil Spill and its unintended consequences on salt marsh ecological processes (DeLaune and Wright 2011, Mendelssohn et al. 2012). Prior studies have examined the effects of oil-spills on marsh vegetation (DeLaune et al. 1984; Lin and Mendelssohn 1996; Hester and Mendelssohn 2000; Pezeshki et al. 2000; DeLaune et al. 2003) and marsh sediment microbial community structure (Wright et al. 1997; Paisse et al. 2008; Bordenave et al. 2009). However, very few have addressed sub-lethal effects of oil exposure on the physiology of marsh vegetation (Smith et al. 1984; Pezeshki and DeLaune 1993; DeLaune et al. 2003), or its potential impacts on salt marsh ecosystem functional processes. Recently, efforts have been made to assess the specific impacts of the Deepwater Horizon oil spill on various components of salt marsh plant productivity (e.g. Lin and Mendelssohn 2012; Mishra et al. 2012; Khanna et al. 2013; Judy et al. 2014). Mishra et al. (2012) observed reduced marsh canopy chlorophyll contents in response to oiling in Louisiana marshes, but did not assess the exact mechanisms behind this response. Additional studies have also reported alterations in marsh sediment microbial communities in response to oil intrusion (Bik et al. 2012; Cravo-Laureau and Duran 2014; Mason et al. 2014; Newell et al. 2014), but very little is known about how these shifts in microbial communities may have impacted marsh biogeochemical processes and the detrital-based ecosystem services they provide (DeLaune and Wright 2011; Mendelssohn et al. 2012).

Currently, no published study has examined effects of the Deepwater Horizon oil spill on decomposition and nutrient cycling processes within Mississippi Gulf Coast salt marshes. The present study was conducted to examine organic matter decomposition

patterns in Mississippi Gulf Coast marshes exposed to different oiling intensities as a result of the Deepwater Horizon oil spill. In this study, we assessed both surface and subsurface marsh sediment cellulose decomposition in nine coastal salt marshes (6 oiled, 3 unoiled) using a standardized cotton strip bioassay technique as a proxy for organic matter decomposition (French 1988b; Slocum et al. 2009, Tiegs et al. 2013). In addition, rates of microbial respiration, fungal biomass (ergosterol) and nutrient ratios (CNP) associated with surface sediment cotton strips were also quantified in order to examine their relationship with cotton strip decomposition (i.e., tensile strength loss) and their potential importance to nutrient cycling functions. Overall, this research hypothesized that oiling would have a significant impact on organic matter decomposition and predicted that sites impacted by oiling would exhibit altered cotton strip decomposition rates in comparison to unoiled sites.

## CHAPTER II

### MATERIALS AND METHODS

#### Study Sites

This study was conducted at 9 saltmarsh sites along the Mississippi Gulf Coast, which differed in their degree of oiling exposure (Figure 1). Study sites were initially screened based on the degree of oiling exposure, as determined by cumulative maximum observed oiling data gathered from Shoreline Cleanup Assessment Team (SCAT) maps (Environmental Response Management Application [ERMA], 2014). Sites were also initially ground-truthed based on observations of oiling and from on-site interviews with personnel from the MS Department of Marine Resources and Emergency Management Agencies of Jackson and Hancock County. A total of six oiled and three unoiled sites were examined (Table 1), with the final site selections chosen for similarity in plant physiognomy and tidal influences.

#### Field Procedures

The effects of oiling on cellulose decomposition in 2 differing salt marsh vegetative zones, *J. roemerianus* and *S. alterniflora*, was evaluated by the standardized cotton strip bioassay using Fredrix-brand #548 style art canvas fabric (see Slocum et al. 2009, Tiegs et al. 2013). Cotton strips (9 x 2.5 cm) were enclosed in fiberglass mesh (200  $\mu\text{m}$ ) litter bags, transported to the field sites (September 2011), and placed in three randomly selected plots within monotypic stands of *S. alterniflora* or *J. roemerianus*. At each plot and within each vegetation zone, five replicate litter bags were placed at and within the marsh sediments to assess the influence of oiling on both surface (i.e. aerobic)




Figure 1. Map of the Mississippi Gulf Coast illustrating the sample study sites and the known oiling impacts based on the Shoreline Cleanup Assessment Team (SCAT) surveys conducted immediately after the oil spill.

Table 1

*Marsh study sites examine and their varying degrees of oil and contaminant exposure as determined from Shoreline Cleanup Assessment Team (SCAT) surveys.*

Site	Maximum oil observed*	Latitude	Longitude
Site 1, Herron Bay	No oil observed	30.193646°	-89.481996°
Site 2, Lower Point Clear	Light	30.176753°	-89.462508°
Site 3, St. Joseph Point	Very light	30.189712°	-89.446815°
Site 4, Grand Bayou	No oil observed	30.263125°	-89.355899°
Site 5, Cat Island, Little Bay	Moderate	30.223953°	-88.756339°
Site 6, Cat Island	Moderate	30.224328°	-84.113933°
Site 7, Graveline Bay	No oil observed	30.364200°	-88.668912°
Site 8, Chevron Refinery	Moderate	30.320676°	-88.492312°
Site 9, Point Aux Chenes Bay	Very light	30.207264°	-88.418591°

and subsurface (anaerobic) cellulose decay, respectively (9 marsh sites x 2 vegetation zones x 3 plots x 2 sediment zones x 5 replicates = 540 total litter bags).

Subsurface litter bags containing 1 cotton strip were buried at ~5 cm beneath the marsh sediment surface by using a standard dibble bar to ensure their placement at a consistent uniform depth. Corresponding surface litter bags contained two cotton strips were placed in direct contact with surface sediment. These litter bags included one cotton strip for determination of cellulose decomposition rates and a second strip to examine

cotton strip associated fungal biomass (ergosterol), rates of microbial respiration (CO<sub>2</sub> evolution) and associated nutrients (carbon, nitrogen and phosphorus). Five litter bags from each vegetative zone were immediately collected returned to the laboratory for determination of initial cotton strip tensile strength. Thereafter, litter bags were retrieved at 2, 3 and 4 weeks, placed in clean zip lock bags on ice in a cooler, and immediately returned to laboratory for processing.

On each sampling date, interstitial pore water within each vegetative zone was monitored in established peziometers (n=3, ~60 cm depth) for pH, oxidation-reduction potential (mV), salinity (ppt) using a YSI professional plus multiparameter meter. In addition, surface and subsurface temperatures at each site were continuously monitored at 30 min intervals throughout the entire study period using two Onset HOBO water temp Pro V2 series data loggers.

#### Cellulose Decomposition Rates

Rates of cotton strip cellulose decomposition were inferred from losses in tensile strength of cotton strip cellulose fibers. Upon return to the laboratory, collected strips were gently cleaned with deionized water to remove adhering sediment, dried at 60°C, and stored in clean ziplock bags within a desiccator until analyzed. Cotton strip tensile strengths were subsequently determined using a Mark-10 Tensiometer following protocols described by Tiegs et al. (2013). Losses in tensile strength are expressed as the percent of initial tensile strength lost per day during the incubation period using the equation:



$$\%CTSL = 1 - [(TS_{treatment\ strips} / TS_{reference\ strips}) \times 100] / \text{incubation time}$$

where  $TS_{treatment\ strips}$  is the maximum tensile strength recorded for each cotton strip incubated in the field, and  $TS_{reference\ strips}$  is the mean tensile strength recorded for initial cotton strips that were transported to the field during litterbag deployment and immediately returned to the laboratory for processing.

#### Rates of Microbial Respiration

Rates of microbial respiration (CO<sub>2</sub> evolution) from collected surface cotton strips were determined using a LiCor LI-6400 Portable Infrared Gas Analyzer. Upon return to the laboratory, collected surface cotton strips were removed from litter bags and placed into sterile Petri dishes lined with sterile (autoclaved) filter paper. Cotton strips and filter paper were wetted with ~10 mL of 0.22 μm membrane filtered marsh surface water collected from each respective site and incubated in darkness for 2 h at 15°C. After 2 hours of incubation, rates of CO<sub>2</sub> evolution were measured by enclosing cotton strips into a LI 6400-89 insect respiration chamber connected to the LiCor LI-6400 Infrared Gas Analyzer. Following respiration measurements, cotton strips were stored frozen (-20°C) and later lyophilized and weighed to determine hourly rates of microbial respiration. Subsamples of lyophilized cotton strips were also used to determine cotton strip associated fungal biomass and nutrient concentrations (see below).

#### Fungal Ergosterol and Nutrient Concentrations

The extent of fungal colonization of collected cotton strips was estimated from ergosterol concentrations (Gessner 2005). Following lyophilization, subsamples were immediately cut from cotton strips, weighed, placed into clean 20 ml glass scintillation vial containing 5 ml of HPLC grade methanol, and stored at -20°C until analyzed. Later,

ergosterol in stored samples were extracted in alcoholic KOH (0.8% KOH in HPLC grade methanol, total extraction volume 10 ml) for 30 min at 80°C in tightly capped thick-walled Pyrex digestion tubes with constant stirring. Sample extracts were partially cleaned by solid-phase extraction (Gessner and Schmitt 1996), and ergosterol quantified by a Shimadzu high-pressure liquid chromatography (HPLC) system using a LichroSpher 100 RP-18 column (0.46 × 25 cm, mobile phase HPLC-grade methanol, flow rate of 1.5 mL min<sup>-1</sup>). Ergosterol was detected at 282 nm and identified and quantified on the basis of comparison with known ergosterol standards (Fluka Chemicals). Ergosterol concentrations were determined as the average of two HPLC injections per sample extract.

Subsamples of cotton strips were also analyzed for carbon, nitrogen and phosphorus concentrations. The carbon and nitrogen concentrations of cotton strips were determined using a Costech 4010 elemental combustion analyzer. Phosphorus concentrations were determined using a SEAL AA3 Flow Injection Nutrient Analyzer (molybdate-ascorbic acid method) following combustion (500°C) of cotton strip subsamples and hot 1N HCl solubilization of the resulting ash.

#### Data Analyses

Statistical analyses of the data was conducted using JMP software (SAS Institute Inc., Cary, NC, USA, version 11), with differences at the  $p < 0.05$  level being considered significant. If necessary, data were transformed prior to analysis to ensure normality and, heteroscedasticity. All data were analyzed using a nested ANOVA, with sites nested within oiling as the random factor and species zones and oiling as fixed factors. A

Tukey's HSD multiple comparison test was used to test for differences among treatments when significant main or interactive effects of ANOVAs were found.

## CHAPTER III

### RESULTS

#### Cementation Effects

Unfortunately, all sites examined in this research project experienced some degree of cotton strip cementation during the study period (French 1988a). Cementation is a phenomenon whereby the tensile strength of cotton fibers tends to increase during incubation in the field. This phenomenon is caused by a variety of biotic and abiotic factors, such as the production of polysaccharides and resins by inhabitant microorganisms or the physical binding of fine sediments to the cotton strip. Correction for cementation effects is often difficult, and any cementation correction when comparing multiple experimental sites must ensure that both physical and chemical conditions at the sites are similar (French 1988c). However, even if correction is possible, the resulting data should be interpreted with caution.

In the present study, both surface and subsurface cotton strips were deployed at the marsh study sites and retrieved at 2, 3 and 4 week intervals. The initial objective of this study was to calculate a rate of cotton strip tensile strength loss through time. However, due to cementation issues it was not possible to use all of the cotton strip tensile strength data collected. To deal with the effects of cementation the single sampling date that displayed the highest mean tensile strength loss (i.e. less cementation) at each site was chosen for analysis and inclusion in this thesis.

#### Environmental Parameters

Both surface and subsurface temperatures remained relatively consistent during the study period (Figure 2), and did not significantly vary among sites (surface sediments:




Figure 2. Changes in surface (A) and subsurface (B) sediment temperatures during the study period. Values are the overall mean $\pm$ SD for all study sites combined

$F_{(7,230)}=0.276$ ,  $p=0.96$ , subsurface sediments:  $F_{(7,230)}=0.341$ ,  $p=0.934$ ). In contrast, marsh pore water salinity, pH and reduction potential (ORP) varied significantly among sites and plant vegetative zones (Table 2). However, oiling showed no significant effects on any of the measured parameters (Table 3). In addition, no significant interactions between

Table 2

*Salinity, pH and reduction potential (ORP) of sediment pore water within S. alterniflora and J. roemerianus plant zones. Values are the mean±1SD (n=12, except site 4 for J. roemerianus where n=1).*

Site	Species	Salinity (ppm)	pH	ORP (mV)
1	Spartina	14.81±2.32	6.79±0.15	-254±13
	Juncus	14.82±1.33	6.59±0.07	-233±39
2	Spartina	13.75±2.19	6.87±0.17	-290±23
	Juncus	17.83±1.30	6.63±0.10	-242±47
3	Spartina	18.47±1.15	7.08±0.15	-178±57
	Juncus	19.68±0.55	7.16±0.15	-180±41
4	Spartina	15.93±5.65	6.91±0.18	-154±37
	Juncus	22.04	6.67±0.03	-69±31
5	Spartina	27.26±2.29	6.66±0.18	-180±97
	Juncus	28.58±1.81	6.89±0.22	-90±34
6	Spartina	25.51±2.92	6.94±0.44	-147±73
	Juncus	24.17±1.43	6.68±0.10	-174±12
7	Spartina	18.98±4.27	6.90±0.15	-298±33
	Juncus	26.31±1.07	6.61±0.06	-262±24
8	Spartina	24.96±0.83	6.95±0.09	-268±51
	Juncus	27.47±0.05	6.87±0.07	-263±11

Table 2 (continued).

Site	Species	Salinity (ppm)	pH	ORP (mV)
9	Spartina	28.28±1.97	6.85±0.28	-7±102
	Juncus	30.10±2.64	6.84±0.27	-139±95

Table 3

*Nested ANOVA summary table indicating the effect of site, plant species zone, and oiling on marsh sediment pore water salinity, pH and reduction potentials (ORP) at each marsh site.*

Effect	Salinity			pH			Reduction Potentials (ORP)		
	df	F-Ratio	p-value	df	F-Ratio	p-value	df	F-Ratio	p-value
Site [Oiling]	7,84	50.13	<0.001	7,94	3.85	<0.001	7,90	35.42	<0.001
Plant species zone	1,84	28.26	<0.001	1,94	11.12	0.001	1,90	8.32	0.005
Oiling	1,84	2.01	0.199	1,94	2.21	0.178	1,90	0.47	0.514
Plant species zone *Oiling	1,84	3.08	0.083	1,94	5.42	0.022	1,90	0.15	0.703



oiling and plant species zones were observed, except for pH where *J. roemerianus* zones had significantly lower pore water pH at oiled sites (Tukey's HSD,  $p > 0.05$ ).

#### Cotton Strip Decay - Subsurface Sediments

In the present study, tensile strength loss of cotton strips (CTSL %/d) in subsurface marsh sediment varied significantly among study sites (ANOVA,  $F_{(7,76)} = 10.46$ ,  $p < 0.001$ ) and between plant species zones (ANOVA,  $F_{(1,76)} = 7.62$ ,  $p < 0.01$ ), with a slightly higher rate of tensile strength loss being observed within sediments at the *J. roemerianus* versus *S. alterniflora* plant zone (Figure 3, Tables 4 and 5). Although oiling showed no significant effect (ANOVA,  $F_{(1,76)} = 1.37$ ,  $p = 0.280$ ) on tensile strength loss, there was a significant interaction between oiling and plant species zone (ANOVA,  $F_{(1,76)} = 5.33$ ,  $p = 0.024$ ), suggesting that tensile strength losses varied between vegetation zones (*S. alterniflora* vs. *J. roemerianus*) in response to oiling (Table 4). When analyzed for oiling effects across study sites and by overall plant species zone, tensile strength losses within oiled *S. alterniflora* plant zones were significantly higher (~26%, Tukey's HSD,  $p < 0.05$ ) than corresponding losses observed in unoiled *S. alterniflora* zones (Figure 4). In contrast, only minor differences in tensile strength losses were observed between oiled and unoiled *J. roemerianus* plant zones, which were not significantly different (Tukey's HSD,  $p > 0.05$ ) (Figure 4, Table 5).

#### Cotton Strip Decay - Surface Sediments

Tensile strength losses of cotton strips at the marsh surface sediments (CTSL %/d) were markedly lower than observed cotton decay rates in subsurface sediments, and did not vary significantly (ANOVA,  $F_{(7,79)} = 7.80$ ,  $p = 0.271$ ) among the study sites examined (Figure 5, Table 4). However, similar to marsh subsurface sediments,




Figure 3. Losses in cotton strip tensile strength observed in subsurface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean  $\pm$  SE (n=5).

Table 4

Nested ANOVA summary indicating the effect of site, plant species zone, and oiling on subsurface and surface percent cotton strip tensile strength loss.

Effect	Subsurface (%CTSL/d)			Surface (%CTSL/d)		
	df	F-Ratio	p-value	df	F-Ratio	p-value
Site [Oiling]	7,76	10.46	<0.001	7,79	7.80	0.271
Plant species zone	1,76	7.62	0.007	1,79	9.43	0.003
Oiling	1,76	1.37	0.280	1,79	3.99	0.086
Plant species zone *Oiling	1,76	5.33	0.024	1,79	5.98	0.071

Table 5

Overall tensile strength loss (%CTSL/d) of surface and subsurface cotton strips in oiled and unoiled *S. alterniflora* and *J. roemerianus* plant zones. Values are the mean $\pm$ 1SD ( $n=15-30$ ).

Species Zone	Site	Subsurface Sediments	Surface Sediments
<i>S. alterniflora</i>	Oiled	2.51 $\pm$ 0.82	0.35 $\pm$ 0.34
	Unoiled	1.73 $\pm$ 0.60	0.93 $\pm$ 0.36
<i>J. roemerianus</i>	Oiled	2.60 $\pm$ 0.84	0.21 $\pm$ 0.12
	Unoiled	2.34 $\pm$ 0.18	0.27 $\pm$ 0.20




Figure 4. Overall patterns of cotton strip tensile strength loss observed in subsurface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean $\pm$ SE (n=15-30).

significant differences (ANOVA,  $F_{(1,79)}=9.43$ ,  $p=0.003$ ) in tensile strength losses were noted between vegetative zones (*S. alterniflora* vs. *J. roemerianus*) (Table 4). These differences displayed an opposite pattern to those observed in marsh subsurface sediments, with on average higher tensile strength losses observed in the *S. alterniflora* versus the *J. roemerianus* plant zone. Similar to subsurface sediment cotton strips, oiling displayed no significant effect on tensile strength loss between sites; however, note that the p-value was low (ANOVA,  $F_{(1,79)}=3.99$ ,  $p=0.086$ ) (Table 4). Furthermore, the p-value for the interaction between oiling and plant species zone was also low (ANOVA,  $F_{(1,79)}=5.98$ ,  $p=0.071$ ), suggesting that tensile strength loss rates in surface sediments may have also varied in vegetative zones according to oiling exposure (Table 4). When analyzed for




Figure 5. Losses in cotton strip tensile strength observed in surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean $\pm$ SE (n=5).

oiling effects across study sites and by overall plant zone, tensile strength losses within surface marsh sediments displayed an opposite pattern when compared to corresponding cotton strip data from marsh subsurface sediments, with cotton strips in unoiled *S. alterniflora* plant zone having the significantly higher rates of decay (Tukey's HSD,  $p > 0.05$ ). Surface sediment tensile strength losses of cotton strips were on average ~63% and ~24% lower in the oiled versus unoiled *S. alterniflora* and *J. roemerianus* plant zones, respectively (Figure 6, Table 5), which implies that oiling might have differential effects on microbial decay dynamics between subsurface (anaerobic) and surface (aerobic) sediment environments.




Figure 6. Overall patterns of cotton strip tensile strength loss observed in surface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean  $\pm$  SE (n=15-30).

### Cotton Strip Nutrient Dynamics

Significant differences in the carbon:nitrogen (C:N) ratios of surface sediment cotton strips were observed among study sites (ANOVA,  $F_{(7,79)}=7.80$ ,  $p=0.041$ ) and between plant vegetative zones (ANOVA,  $F_{(1,79)}=4.85$ ,  $p=0.031$ ) (Figure 7, Table 6). Site oiling also had a significant effect on cotton strip C:N ratios (ANOVA,  $F_{(1,79)}=10.80$ ,  $p=0.013$ ), and there was a significant interaction between oiling and plant species zone (ANOVA,  $F_{(1,79)}=7.11$ ,  $p=0.009$ ) (Table 6). When analyzed across study sites and by plant species zone, C:N ratio's of cotton strips were on average significantly higher (~38%, Tukey's HSD,  $p<0.05$ ) in oiled versus unoiled *S. alterniflora* plant zones (Figure 8, Table 7). Similarly, the C:N ratio's of cotton strips in *J. roemerianus* plant zones were also higher in oiled versus unoiled sites, with cotton strips in unoiled *J. roemerianus* zones having on average lower C:N ratios (Figure 8, Table 7), however, this difference was not significant (Tukey's HSD,  $p<0.05$ ). Across study sites, cotton strip C:N ratios were negatively correlated with losses in cotton strip tensile strength (Table 8).

Significant differences (ANOVA,  $F_{(7,79)}=6.57$ ,  $p<0.001$ ) in the carbon:phosphorus (C:P) ratios of cotton strips were also observed among study sites (Figure 9, Table 6). In contrast to C:N ratios, no significant differences were observed between plant species zones (ANOVA,  $F_{(1,79)}=0.56$ ,  $p=0.457$ ), nor was there a significant oiling effect (ANOVA,  $F_{(1,79)}=4.105$ ,  $p=0.082$ ). However, a significant interaction (ANOVA,  $F_{(1,79)}=4.93$ ,  $p=0.029$ ) was observed between plant species zone and oiling, suggesting that cotton strip C:P ratios may have also varied with vegetative zones according to oiling exposure (Table 6). Overall, the C:P ratios of cotton strips followed a similar pattern as C:N ratios, with cotton strips in oiled *S. alterniflora* plant zones having much higher




Figure 7. Carbon:nitrogen ratio patterns for cotton strips incubated at the marsh surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean $\pm$ SE (n=5).



Table 6

*Nested ANOVA summary table indicating the effect of site, plant species zone and oiling on the carbon:nitrogen and carbon:phosphorus ratios of cotton strips incubated at the marsh sediment surface.*

Effect	Carbon:Nitrogen			Carbon:Phosphorus	
	df	F-Ratio	p-value	F-Ratio	p-value
Site [Oiling]	7,79	7.80	0.041	6.57	<0.001
Plant species zone	1,79	4.85	0.031	0.56	0.457
Oiling	1,79	10.80	0.013	4.11	0.082
Plant species zone *Oiling	1,79	7.11	0.009	4.93	0.029

Table 7

*Overall patterns of carbon:nitrogen and carbon:phosphorus ratios for cotton strips incubated at the marsh surface sediments in *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean $\pm$ 1SD (n=15-30).*

Species Zone	Site	Carbon:Nitrogen	Carbon:Phosphorus
<i>S. alterniflora</i>	Oiled	316 $\pm$ 42	22,991 $\pm$ 9,653
	Unoiled	230 $\pm$ 46	12,450 $\pm$ 5,428
<i>J. roemerianus</i>	Oiled	310 $\pm$ 55	20,377 $\pm$ 6,577
	Unoiled	281 $\pm$ 21	16,143 $\pm$ 4,703




Figure 8. Overall patterns of carbon:nitrogen (A) and carbon:phosphorus (B) ratios for cotton strip incubated at the marsh surface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).




Figure 9. Carbon:phosphorus ratio patterns for cotton strips incubated at the marsh surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean $\pm$ SE (n=5).

Table 8

*Pearson product-moment intercorrelation matrix showing the relationships between rates of surface sediment cotton strip decay (%CTSL/d) and cotton strip-associated fungal ergosterol concentrations, rates of microbial respiration and carbon:nitrogen and carbon:phosphorus ratios.*

	Cotton strip (%CTSL/d)	Ergosterol	Microbial Respiration	Carbon:Nitrogen	Carbon:Phosphorus
Cotton strip (%CTSL/d)	1.000				
Ergosterol	0.514***	1.000			
Microbial Respiration	0.672***	0.604***	1.000		
Carbon:Nitrogen	-0.628***	-0.629***	-0.616***	1.000	
Carbon:Phosphorus	-0.380**	-0.332*	-0.349*	0.637***	1.000

Note: \*\*\*p<0.001, \*\*p<0.01, and \*p<0.05

ratios than corresponding unoiled sites (Figure 8, Table 6). When analyzed for oiling across study sites and by vegetative zone, average C:P ratios of cotton strips were significantly higher in oiled *S. alterniflora* (~85%) and *J. roemerianus* (~26%) plant zones versus cotton strips in unoiled zones, respectively (Tukey's HSD  $p < 0.05$ ) (Figure 8, Table 7). In addition, similar to cotton strip C:N ratios, C:P ratios were also negatively correlated with observed losses in cotton strip tensile strength, however, the resulting correlation coefficient was low (Table 8).

#### Cotton Strip Fungal Colonization and Microbial Respiration Rates

Fungal colonization of cotton strips, as determined by ergosterol concentrations, were not significantly different among study sites (ANOVA,  $F_{(7,79)} = 7.80$ ,  $p = 0.171$ ) or between plant zones (ANOVA,  $F_{(1,79)} = 0.085$ ,  $p = 0.77$ ) (Figure 10, Table 9). However, marsh oiling did have a significant effect on ergosterol concentrations (ANOVA,  $F_{(1,79)} = 8.12$ ,  $p = 0.025$ ), and there was a significant interaction between plant species zone and oiling (ANOVA,  $F_{(1,79)} = 5.98$ ,  $p = 0.015$ ). When analyzed across study sites and by vegetative zone, cotton strip ergosterol concentrations were on average significantly lower (~76%, Tukey's HSD,  $p < 0.05$ ) in oiled *S. alterniflora* zones versus unoiled zones (Figure 11, Table 10). In contrast, cotton strip ergosterol concentrations in the *J. roemerianus* plant zones were only 7% lower in oiled versus unoiled sites and were not significantly different (Tukey's HSD,  $p > 0.05$ ).

Rates of microbial respiration associated with cotton strips varied significantly among marsh sites (ANOVA,  $F_{(7,79)} = 8.41$ ,  $p < 0.0001$ ) (Figure 12, Table 9). Consistent with patterns of fungal colonization (ergosterol), no significant difference was observed between plant zones (ANOVA,  $F_{(1,79)} = 1.85$ ,  $p = 0.177$ ). Unlike ergosterol concentrations,





Figure 10. Fungal ergosterol concentrations associated with cotton strips incubated at the marsh surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean $\pm$ SE (n=5)

Table 9

*Nested ANOVA summary table indicating the effect of site, plant species zone and oiling on cotton strip-associated fungal ergosterol concentrations and microbial respiration rates in cotton strips incubated at the marsh sediment surface.*

Effect	Ergosterol			Microbial Respiration	
	df	F-Ratio	p-value	F-Ratio	p-value
Site [Oiling]	7,79	7.80	0.171	8.41	<0.001
Plant species zone	1,79	0.09	0.772	1.85	0.177
Oiling	1,79	8.12	0.025	0.17	0.692
Plant species zone *Oiling	1,79	5.98	0.015	5.32	0.024



*Figure 11. Overall patterns of fungal ergosterol concentrations associated with cotton strips incubated at the marsh surface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).*

Table 10

*Overall patterns of fungal ergosterol concentrations and rates of microbial respiration associated with cotton strips incubated at the marsh surface sediments in *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean $\pm$ ISD (n=15-30).*

Species Zone	Site	Ergosterol ( $\mu\text{g/gC}$ )	Microbial Respiration ( $\mu\text{g CO}_2\text{-C/gC/h}$ )
<i>S. alterniflora</i>	Oiled	15.06 $\pm$ 4.73	53.26 $\pm$ 27.69
	Unoiled	26.45 $\pm$ 7.08	124.26 $\pm$ 157.71
<i>J. roemerianus</i>	Oiled	19.80 $\pm$ 8.33	60.38 $\pm$ 26.93
	Unoiled	21.26 $\pm$ 7.03	51.72 $\pm$ 32.37

marsh oiling had no significant effect on respiration rates (ANOVA,  $F_{(1,79)}=0.1705$ ,  $p=0.692$ ); however, there was a significant interaction between oiling and plant species zone (ANOVA,  $F_{(1,79)}=5.32$ ,  $p=0.024$ ) (Table 9). When analyzed across study sites by oiling and vegetative zone, rates of microbial respiration associated with cotton strips in oiled *S. alterniflora* plant zones were significantly lower (~133%, Tukey's HSD,  $p<0.05$ ) versus corresponding cotton strips in unoiled zones (Figure 13, Table 9). Rates of microbial respiration associated with cotton strips in *J. roemerianus* plant zones were on average slightly higher in oiled versus unoiled plant zones (Figure 13, Table 10), however, this difference was not significant (Tukey's HSD,  $p>0.05$ ). Cotton strip ergosterol concentrations and rates of microbial respiration were positively correlated with one another and positively correlated with observed losses in cotton strip tensile



strength (Table 8). Furthermore, both ergosterol concentrations and respiration rates were




Figure 12. Rates of microbial respiration (CO<sub>2</sub> evolution) associated with cotton strips incubated at the marsh surface sediments in the *S. alterniflora* (A) and *J. roemerianus* (B) plant zones at each marsh site. Values are the mean ± SE (n=5).




Figure 13. Overall patterns of microbial respiration rates (CO<sub>2</sub> evolution) associated with cotton strips incubated at the marsh surface sediments in the *S. alterniflora* and *J. roemerianus* plant zones within oiled and unoiled marsh sites. Values are the mean±SE (n=15-30).

negatively correlated with cotton strip C:N and C:P ratios, providing evidence that prevailing N and P availabilities within the marsh surface sediments may have limited the decay activity of fungal communities associated with cotton strips and hence the corresponding observed losses in cotton strip tensile strengths.

## CHAPTER VI

## DISCUSSION

Coastal salt marshes along the Northern Gulf of Mexico are widely regarded as among the most productive wetland-estuarine ecosystems within the United States (Mitsch and Gosselink 2007), and their conservation and management is critical due to the multitude of important economic and ecosystem services that they provide to the region. The emergent vascular plants, *Spartina alterniflora* and *Juncus roemerianus*, are the dominant vegetation within these coastal marshes and provide the base for many important ecosystem services (Maltby 2009; DeLaune and Wright 2011; Mendelsohn et al. 2012). Currently, published estimates indicate that ~4.9 million barrels of oil were released into the Gulf of Mexico from the Deepwater Horizon Mississippi Canyon Block 252 oil spill (Crone and Tolstoy 2010). As a result of the oil spill, approximately 430 miles of marsh shoreline was exposed to extensively weathered MC252 crude oil (Reddy et al. 2011), of which ~41% or 176 miles were either moderately or heavily oiled (Mendelsohn et al. 2012). Although recent studies from the Deepwater Horizon spill have attempted to assess oiling impacts on salt marsh vegetation (Anderson and Hess 2012; Lin and Mendelsohn 2012; Mishra et al. 2012; Silliman et al. 2012; Khanna et al. 2013; Judy et al. 2014), sediment microbial communities (Joye et al. 2014; Horel et al. 2012; Horel et al. 2014b), marsh fish communities (Fodrie et al. 2014; Echols et al. 2015), and benthic insect and invertebrate populations (McCall and Pennings 2012), very few studies have examined the effect of the oil spill on critical ecological processes, such as organic matter decomposition.

In the present study, we used a standardized cotton strip bioassay to examine rates of cellulose decomposition in 9 Mississippi coastal salt marsh sites that differed in the degree of oiling intensity. All of the study sites were chosen based on their similar plant physiognomy and degree of tidal influence, in order to control for other environmental parameters (e.g., pH, salinity, moisture and temperature), which can also significantly influence rates of organic matter decomposition. Although this study was conducted ~1.5 years after the oil spill, we observed significant differences in both surface and subsurface cotton strip decay among the marsh study sites, with oiling impacts being significantly noted within *S. alterniflora* plant zones. In contrast, no significant oiling effects were observed in the *J. roemerianus* plant zone, which was consistent with initial field observations that most of the visible oil inundation in these marshes were restricted to the lower *S. alterniflora* tidal zone at the outer periphery of the marsh. Although, note that recent studies in Louisiana coastal marshes have provided evidence that oil was distributed much deeper (100 m) into oiled marshes (i.e. *J. roemerianus* plant zones), which was not noted during the rapid Shoreline Cleanup Assessment Team (SCAT) surveys immediately following the oil spill (Turner et al. 2014b; see also Ramsey et al. 2014).

A notable and unexpected finding in the present study is the observation of opposing cotton strip decay dynamics between surface (aerobic) and subsurface (anaerobic) sediment environments, particularly within oiled and unoled *S. alterniflora* plant zones. Despite the presence of aerobic conditions, tensile strength losses of cotton strips placed at the marsh surface sediments were markedly lower than corresponding tensile strength losses for cotton strips incubated within the subsurface sediments. In

addition, the degree of oiling impact on cotton strip decay was also quite different between surface and subsurface sediments. Tensile strength losses of cotton strips in the surface sediment were markedly higher in unoiled versus oiled *S. alterniflora* plant zones, suggesting that oiling may have a negative impact on cellulose decomposition. In contrast, corresponding tensile strength losses of cotton strips in subsurface sediments were significantly higher in oiled versus unoiled *S. alterniflora* marsh sediments, suggesting that oiling inundation within sediments may have stimulated cellulose degradation. Collectively, these observations imply that oiling may have differential effects on microbial communities and hence organic matter decomposition between surface (aerobic) and subsurface (anaerobic) sediment environments.

Observations of oiling effects in the present study differ with earlier findings of Mendelsohn and Slocum (2004), which found no significance influence of oil composition and concentration on salt marsh sediment cellulose decomposition. Using the same cotton strip bioassay approach, Mendelsohn and Slocum (2004) assessed rates of subsurface cellulose decomposition at 8 brackish intertidal marshes that had been exposed to different oiling intensities as a result of the Chalk Point, Maryland fuel oil spill. Similar to the present study, Mendelsohn and Slocum (2004) conducted cotton strip decay studies ~2 years after the oil spill in order to assess potential long-term effects from the oil spill. Although no apparent oiling effect was observed, results of their study found that abiotic environmental characteristics, particularly pH and salinity, had a strong effect on cellulose decomposition rates among the study sites. In their study, strong negative effects of salinity and positive effects of pH on cellulose decomposition were observed among the sites examined. These findings led the authors to conclude that sediment decomposition in

these marshes was controlled more by natural environmental conditions than by its prior history of oiling.

As mentioned earlier, marsh sites chosen for this study all had a similar plant physiognomy and tidal influence in order to control for possible between-site differences in environmental conditions. Despite our best efforts in site selection, significant differences in some environmental parameters were observed between study sites. Although surface and subsurface temperatures did not significantly vary among sites during the study period, we did observe significant differences in both the pH and salinity of interstitial pore water. Mendelsohn and Slocum (2004) observed the slowest rates of cellulose decomposition at estuarine marsh sites with high salinities and low pH. In the present study, we observed no significant relationships between tensile strength losses of cotton strip and salinity or pH. Unlike Mendelsohn and Slocum (2004), we did not measure soil redox potentials, sediment nutrient pools, or other abiotic factors (e.g., bulk density) during the study period. As a consequence, we cannot rule out that some environmental conditions may have contributed, at least in part, to the cotton strip decay patterns observed.

Degradation of oil hydrocarbons by microorganisms is well documented, and deposition of crude oil along coastal shorelines (beaches and marshes) is often expected to alter wetland microbial processes and their important biogeochemical functions (e.g. Leahy and Colwell 1990; Atlas and Bartha 1998; Shin et al. 2000; Widdel and Rabus 2001; Horel et al. 2014a). Following contamination, crude oil typically leads to a shift in the microbial community, resulting in a decrease in microbial diversity and an increased production and abundance in microbial taxa that possess the enzymatic capacity to utilize

oil substrates as a carbon source (Shiaris 1989; Leahy and Colwell 1990; Li et al. 1990; So and Young 2001; Paissé et al. 2008; Haritash and Kaushik 2009; Horel et al. 2012).

However, there is no clear consensus in the literature regarding the impacts of oil contamination on microbial communities and hence the exact spatial and temporal outcomes on wetland biogeochemical cycles (DeLaune and Wright 2011; Mendelsohn et al. 2012). The overall effect of oil contamination on microbial communities appears to be variable and dependent on the composition (i.e., toxicity) and concentration of hydrocarbons in the oil and the environmental conditions present, such as temperature, oxygen, nutrient availability, pH and salinity (Mendelsohn et al. 2012; Horel et al. 2012; 2014a; 2014b). If large enough amounts of oil contaminants are introduced microbial communities may be negatively impacted due to increased toxicity (Walker et al. 1975; Siddiqui and Adams 2002; Suarez-Suarez et al. 2011), which may inhibit rates of organic matter decomposition. In contrast, if small to moderate amounts of oil contaminants are introduced it may actually stimulate or possibly prime (Bianchi 2011) microbial communities by providing a labile carbon source (Shiaris 1989; Leahy and Colwell 1990; Li et al. 1990; Nyman 1999; So and Young 2001; Haritash and Kaushik 2009), which may accelerate rates of organic matter decomposition.

Hydrocarbon degradation in salt marshes is primarily considered an aerobic bacterial process (Joye et al. 2014; McGenity 2014) where bacteria initially metabolize labile fractions of crude oil (e.g., linear, branched and cyclic alkanes) followed by limited degradation of more recalcitrant oil fractions (e.g., polycyclic aromatic hydrocarbons, PAH's), which are considered the most toxic and may potentially inhibit some microbial populations. Like bacteria, degradation of petroleum hydrocarbons by fungal organisms

(i.e. filamentous and yeast) is also well documented (e.g. Yanto and Tachibana 2014; Covino et al. 2015), however, the effects of specific crude oil fractions (e.g., PAH's) on fungal community growth and their contribution to crude oil degradation within salt marshes has not been extensively investigated. The pervasive hyphal growth of filamentous fungi suggests that they may play a role in metabolizing oil within marsh sediments (Mendelssohn et al. 2012), and recent studies have pointed to fungal-bacterial interactions as important in facilitating hydrocarbon degradation (McGenity et al. 2012; Banitz et al. 2013).

Under anaerobic conditions, microbial-mediated degradation of oil occurs much more slowly, which supports recent observations that petroleum hydrocarbons (e.g., PAH's) continue to persist in Gulf Coast salt marsh sediments after nearly 5 years since the Deep Water Horizon oil spill (Turner et al. 2014a; 2014b). Although higher under aerobic conditions, hydrocarbon degradation does occur under anaerobic conditions (Milhelcic and Luthy 1988; Widdel and Rabus 2001), where it can stimulate the activities of anaerobic bacterial populations. For example, recently Horel et al. (2014a) found that nitrogen fixation rates among sulfur reducing bacteria increased 142% in response to oiling, which demonstrated that these anaerobic microbial communities had the potential to both fix nitrogen and degrade hydrocarbons.

Prior studies have demonstrated that the addition of hydrocarbons can accelerate organic matter decomposition rates at the salt marsh sediment surface (Hershner and Lake 1980; Li et. al. 1990). For example, Hershner and Lake (1980) observed that oiling significantly increased the decomposition rate of *S. alterniflora* plant litter. This additional C source can fuel or possibly prime (Bianchi 2011) the microbial decomposer community



and accelerate rates of decomposition and the immobilization of nutrients (Jingguo and Bakken 1997). However, because oil provides little to no nitrogen or phosphorus to the system, increased decomposition and mineralization of organic matter can ultimately lead to long-term increases in C:N and C:P ratios, which can eventually limit microbial growth and production (Leah and Colwell 1990) and slow decomposition. In the present study, measured nutrient and microbial parameters associated with surface incubated cotton strips were significantly correlated with corresponding rates of cotton strip decay, with cotton strips from oiled sites collectively having much higher C:N and C:P ratios, lower ergosterol content (fungal biomass), lower respiration rates, and ultimately lower decomposition rates. This study was conducted approximately 1.5 years after the spill, and it is possible that shortly after the spill decomposition rates were possibly elevated, but as time progressed nitrogen and phosphorus may have become limited slowing microbial growth and leading to decreased rates of decomposition. Research by Hines et al. (2006) demonstrated that the addition of a labile carbon source to salt marsh ecosystems increased microbial communities' demand for nitrogen, and that this increased microbial immobilization of nitrogen negatively impacted both aboveground plant biomass (growth) and the resulting C:N ratios of plant litter. Their findings in conjunction with this study imply that oil contamination could possibly lead to progressive nitrogen limitation (and perhaps phosphorus) in aerobic marsh surface sediments, thereby impacting important biogeochemical functions.

In the present study, we did not measure nutrient and microbial parameters associated with cotton strips placed within the more anaerobic marsh sediments. Findings of faster decay rates of cotton strip decay in these zones, particularly *S. alterniflora* plant

zones exposed to oiling, suggest that anaerobic microbial communities may not be nutrient limited and that oiling enhanced their overall decay activities. If this trend is consistent and widespread, then increased rates of organic matter (i.e., cellulose) decomposition could alter the rate of organic matter accumulation and accretion within oiled Gulf Coast marsh sediments, which could further weaken the marsh soil profile and facilitate increased rates of erosion. Increased rates of marsh erosion have been documented in marsh sites exposed to a wide range of oiling intensity, despite an apparent recovery of the plant community (McClenachan et al. 2013).

In salt marshes, decomposition of organic matter is a complex process that includes an array of biotic and abiotic processes that result in the production of decomposer biomass (e.g., microbial and invertebrate), release of CO<sub>2</sub> and nutrients (N & P) through organic matter mineralization, as well as the release of dissolved and fine particulate organic matter (Kuehn 2008). From a microbial perspective, the rates of these decay processes are strongly influenced by the response of microbial communities to the prevailing environmental conditions, the intrinsic quality of the plant detrital resources they metabolize, and the myriad of potential interactions that may occur within detrital food web. Even though this study was conducted ~1.5 years after the Deep Water Horizon oil spill, it is evident that both the subsurface and surface microbial processes were still being affected by oil intrusion. Due to the continued persistence of oil contaminants in Gulf coast marshes, particularly PAH's in subsurface sediments (Turner et al. 2014a; 2014b), more long-term monitoring of marsh biogeochemical functions are warranted in order to understand and predict the spatial and temporal recovery of these vital ecosystems and possible restoration activities.

## REFERENCES

- Anderson CJ, Hess TA. 2012. The effects of oil exposure and weathering on black-needle rush (*Juncus roemerianus*) marshes along the Gulf of Mexico. *Marine Pollution Bulletin* 64:2749-2755.
- Atlas RM, Barth R. 1998. *Microbial ecology: fundamentals and applications*, 3rd edition. Benjamin Cummings, California, USA.
- Banitz T, Johst K, Wick Y, Schamfuß S, Harms H, Frank K. 2013. Highways versus pipelines: contributions of two fungal transport mechanisms to efficient bioremediation. *Environmental Microbiology Reports* 5:211-218.
- Bianchi TS. 2011. The role of terrestrially derived organic carbon in the coastal ocean: a changing paradigm and the priming effect. *Proceedings of the National Academy of Sciences of the United States of America* 108:19473-19481.
- Bik HM, Halanych KM, Sharma J, Thomas WK. 2012. Dramatic shifts in benthic microbial eukaryote communities following the Deepwater Horizon oil spill. *PLoSone* 6: doi:10.1371/journal.pone.0038550.
- Bordenave S, Goni-Urriza M, Caumette P, Duran R. 2009. Differential display analysis of cDNA involved in microbial mat response after heavy fuel oil contamination. *Journal of Microbial and Biochemical Technology* 1: doi:10.4172/1948-5948.1000001.
- Cebrian J, Lartigue J. 2004. Patterns of herbivory and decomposition in aquatic and terrestrial ecosystems. *Ecological Monographs* 74:237–259.

- Covino S, D'Annibale A, Stazi SR, Cajthaml T, Cvancarova M, Stella T, Petucioli M. 2015. Assessment of degradation potential of aliphatic hydrocarbons by autochthonous filamentous fungi from a historically polluted clay soil. *Science of the Total Environment* 505:545-554.
- Christian RR, Bryant WL, Brinson MM. 1990. *Juncus roemerianus* production and decomposition along gradients of salinity and hydroperiod. *Marine Ecology Progress Series* 68:137-145.
- Craft C. 2007. Freshwater input structures soil properties, vertical accretion, and nutrient accumulation of Georgia and U.S. tidal marshes. *Limnology and Oceanography* 52:1220-1230.
- Crone TJ, Tolstoy M. 2010. Magnitude of the 2010 Gulf of Mexico oil leak. *Science* 330:634.
- Cravo-Laureau C, Duran R. 2014. Marine coastal sediments microbial hydrocarbon degradation processes: contribution of experimental ecology in the omics' era. *Frontiers in Microbiology* 5: doi:10.3389/fmicb.2014.00039.
- Dai T, Wiegert RG. 1996. Estimation of the primary productivity of *Spartina alterniflora* using a canopy model. *Ecography* 19:410-423.
- DeLaune RD, Smith CJ, Patrick Jr. WH, Fleeger JW, Tolley MD. 1984. Effect of oil on salt marsh biota, methods for restoration. *Environmental Pollution* 36:207-227.
- DeLaune RD, Pezeshki SR, Jugsujinda A, Lindau CW. 2003. Sensitivity of US Gulf of Mexico coastal marsh vegetation to crude oil: Comparison of greenhouse and field responses. *Aquatic Ecology* 37:351-360.

- DeLaune RD, Wright AL. 2011. Projected impact of Deepwater Horizon oil spill on U.S. Gulf Coast wetlands. *Soil Science Society of America Journal* 75:1602-1612.
- Echols BS, Smith AJ, Rand GM, Seda BC. 2015. Factors affecting toxicity test endpoints in sensitive life stages of native Gulf of Mexico species. *Archives of Environmental Contamination and Toxicology* (in press): doi:10.1007/s00244-014-0122-7.
- Environmental Response Management Application. 2014. Web application. Gulf of Mexico. National Oceanic and Atmospheric Administration, Web. 10 Nov. 2014. Retrieved from <http://response.restoration.noaa.gov/erma/>.
- Ewe SL, Gaiser E, Childers D, Iwaniec D, Rivera-Monroy V, Twilley R. 2006. Spatial and temporal patterns of aboveground net primary productivity (ANPP) along two freshwater-estuarine transects in the Florida Coastal Everglades. *Hydrobiologia* 569:459-474.
- Fodrie FJ, Able KW, Galvez F, Heck KL Jr, Jensen OP, LÓpez-Duarte PC, Martin CW, Turner RE, Whitehead A. 2014. Integrating organismal and population responses of estuarine fishes in macondo spill research. *Bioscience* 64:778-788.
- French DD. 1988a. Problems with cementation. Pages 32-34. In: Harrison AF, Latter PM, Walton DWH, (eds), Cotton strip assay: an index of decomposition in soils. Paper presented at The Institute of Terrestrial Ecology symposium number 24, Grange-over-sands, Cumbria. Great Britain: Titus Wilson & Son Ltd.

- French DD. 1988b. Seasonal patterns in cotton strip decomposition in soils. Pages 46-49. In: Harrison AF, Latter PM, Walton DWH, (eds), Cotton strip assay: an index of decomposition in soils. Paper presented at The Institute of Terrestrial Ecology symposium number 24, Grange-over-sands, Cumbria. Great Britain: Titus Wilson & Son Ltd.
- French DD. 1988c. Some effects of changing soil chemistry on decomposition of plant litters and cellulose on a Scottish moor. *Oecologia* 75:608-618.
- Gessner M. 2005. Ergosterol as a measure of fungal biomass. Pages 189-196. In: Graça, MS, Bärlocher F, Gessner M, (eds.), *Methods to Study Litter Decomposition*. Springer, Dordrecht, The Netherlands.
- Gessner MO, Schmitt AL. 1996. Use of solid-phase extraction to determine ergosterol concentrations in plant tissue colonized by fungi. *Applied Environmental Microbiology* 62:415-419.
- Haritash AK, Kaushik CP. 2009. Biodegradation aspects of Polycyclic Aromatic Hydrocarbons (PAHs): A review. *Journal of Hazardous Materials* 169:1-15.
- Hines J, Megonial JP, Denno RF. 2006. Nutrient subsidies to below ground microbes impact aboveground food web interactions. *Ecology* 87:1542-1555.
- Hershner C, Lake J. 1980. Effects of chronic oil pollution on a salt marsh grass community. *Marine Biology* 56:163-173.
- Hester MW, Mendelsohn IA. 2000. Long-term recovery of a Louisiana brackish marsh plant community from oil-spill impact: Vegetation response and mitigating effects of marsh surface elevation. *Marine Environmental Research* 49:233-254.

- Horel A, Bernard RJ, Mortazavi B. 2014a. Impact of crude oil exposure on nitrogen cycling in a previous impacted *Juncus roemerianus* salt marsh in the northern Gulf of Mexico. *Environmental Science Pollution Research* 21:6982-6993.
- Horel A, Mortazavi B, Sobecky PA. 2014b. Biostimulation of weathered MC252 crude oil in northern Gulf of Mexico sandy sediments. *International Biodeterioration and Biodegradation* 93:1-9.
- Horel A, Mortazavi B, Sobecky PA. 2012. Responses of microbial community from Northern Gulf of Mexico sandy sediments following exposure to Deepwater Horizon crude oil. *Environmental Toxicology and Chemistry* 31:1004-1011.
- Howard PJA. 1988. A critical evaluation of the cotton strip assay. Pages 34-42. In: Harrison AF, Latter PM, Walton DWH, (eds), Cotton strip assay: an index of decomposition in soils. Paper presented at The Institute of Terrestrial Ecology symposium number 24, Grange-over-sands, Cumbria. Great Britain: Titus Wilson & Son Ltd.
- Jingguo W, Bakken LR. 1997. Competition for nitrogen during mineralization of plant residues in soil: microbial response to C and N availability. *Soil Biology and Biochemistry* 29:163-170.
- Joye SB, Teske AP, Kostka JE. 2014. Microbial dynamics following the Macondo oil well blowout across Gulf of Mexico environments. *BioScience* 64:766-777.
- Judy CR, Graham SA, Lin Q, Hou A, Mendelssohn IA. 2014. Impacts of Macondo oil from *Deepwater Horizon* spill on the growth response of the common reed *Phragmites australis*: a mesocosm study. *Marine Pollution Bulletin* 79:69-76.

- Khanna S, Santos, MJ, Ustin SL, Koltunov A, Kokaly RF. 2013. Detection of salt marsh vegetation stress and recovery after the Deepwater Horizon oil spill in Barataria Bay, Gulf of Mexico using AVIRIS data. *PLoSone*, 8: doi: 10.1371/journal.pone.0078989.
- Kuehn KA. 2008. The role of fungi in the decomposition of emergent wetland plants. Pages 19-41. In; Sridhar KR, Bärlocher F, Hyde HD (eds), *Novel Techniques and Ideas in Mycology*. Fungal Diversity Research Series 20, Fungal Diversity Press, Thailand.
- Leahy JG, Colwell RR. 1990. Microbial Degradation of Hydrocarbons in the Environment. *Microbiological Reviews* 54:305-315.
- Li Y, Morris JT, Yoch DC. 1990. Chronic low level hydrocarbon amendments stimulate plant growth and microbial activity in salt marsh microcosms. *Journal of Applied Ecology* 27:159-171.
- Lin Q, Mendelsohn IA. 1996. A comparative investigation of the effects of Louisiana crude oil on the vegetation of fresh, brackish, and saltmarsh. *Marine Pollution Bulletin* 32:202-209.
- Lin Q, Mendelsohn IA. 2012. Impacts and Recovery of the Deepwater Horizon Oil Spill on Vegetation Structure and Function of Coastal Salt Marshes in the Northern Gulf of Mexico. *Environmental Science and Technology* 46:3737–3743.
- Maltby E. 2009. *Functional assessment of wetlands: towards evaluation of ecosystem services*, 9th edition. Woodhead Publishing Limited, Cambridge, UK.



- Mason OU, Scot NM, Gonzolez A, Robbins-Pianka A, Bælum J, Kimbrel J, Bouskill NJ, Prestat E, Borglin S, Joyner DC, Fortney JL, Jurelevicius D, Stringfellow WT, Alvarez-Cohen L, Hazen TC, Knight R, Gilbert JA, Jansson JK. 2014. Metagenomics reveals sediment microbial community response to Deepwater Horizon oil spill. *International Society for Microbial Ecology Journal* 8:1464-1475.
- McCall BD, Pennings SC. 2012. Disturbance and recovery of salt marsh arthropod communities following BP Deepwater Horizon oil spill. *PLoS ONE* 7:e32735.
- McClenachan G, Turner RE, Tweel AW. 2013. Effects of oil on the rate and trajectory of Louisiana marsh shoreline erosion. *Environmental Research Letters* 8: doi:10.1088/1748-9326/8/4/044030.
- McLeod E, Chmura GL, Bouillon S, Salm R, Björk M, Duarte CM, Lovelock CE, Schlesinger WH, Silliman BR. 2011. A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO<sub>2</sub>. *Frontiers in Ecology and the Environment* 9:552–560.
- McGenity TJ. 2014. Hydrocarbon biodegradation in intertidal wetland sediments. *Current Opinion in Biotechnology* 27:46-54.
- McGenty TJ. 2012. Marine crude-oil biodegradation: a central role for interspecies interactions. *Aquatic Biosystems* 8: doi:10.1186/2046-9063-8-10.
- Mendelssohn IA, Andersen GL, Baltz DM, Caffey RH, Carman KR, Fleeger JW, Joye SB, Lin Q, Maltby E, Overton EB, Rozas LP. 2012. Oil impacts on coastal wetlands: Implications for the Mississippi river delta ecosystem after the Deepwater Horizon oil spill. *BioScience* 62:562-574.

- Mendelssohn IA, Slocum MG. 2004. Relationship between soil cellulose decomposition and oil contamination after an oil spill at Swanson Creek, Maryland. *Marine Pollution Bulletin* 48:359-370.
- Milhelcic JR, Luthy RG. 1988. Degradation of polycyclic aromatic hydrocarbons under various redox conditions in soil-water systems. *Applied Environmental Microbiology* 54:1182-1187.
- Mishra DR, Cho HJ, Ghosh S, Fox A, Downs C, Merani PBT, Kirui P, Jackson N, Mishra S. (2012). Post-spill state of the marsh: Remote estimation of the ecological impact of the Gulf of Mexico oil spill on Louisiana salt marshes. *Remote Sensing of Environment* 118:176-185.
- Mitsch WJ, Gosselink JG. 2007. *Wetlands*, 4th edition. Wiley and Sons, New York.
- Moore JC, Berlow EL, Coleman DC, de Ruiter PC, Dong Q, Hastings A, Collins-Johnson N, McCann KS, Melville K, Morin PJ, Nadelhoffer K, Rosemond AD, Post DM, Sabo JL, Scow KM, Vanni MJ, Wall DH. 2004. Detritus, trophic dynamics and biodiversity. *Ecology Letters* 7:584-600.
- Newell SE, Eveillard D, McCarthy MJ, Gardner WS, Liu Z, Ward BB. 2014. A shift in the archaeal nitrifier community in response to natural and anthropogenic disturbances in the northern Gulf of Mexico. *Environmental Microbiology Reports* 6:106-112.
- Neubauer SC. 2008. Contributions of mineral and organic components to tidal freshwater marsh accretion. *Estuarine, Coastal and Shelf Science* 78:78-88.
- Nyman A. 1999. Effect of crude oil and chemical additives on metabolic activity of mixed microbial populations in fresh marsh soils. *Microbial Ecology* 37:152-162.

- Pezeshki SR, De Laune RD. 1993. Effect of crude oil on gas exchange functions of *Juncus roemerianus* and *Spartina alterniflora*. *Water, Air and Soil Pollution* 68:461–468.
- Pezeshki SR, Hester MW, Lin Q, Nyman JA. 2000. The effects of oil spill and clean-up on dominant US gulf coast marsh macrophytes: a review. *Environmental Pollution* 108:129-139.
- Paissé S, Coulon F, Goñi-Urriza M, Peperzak L, McGenity TJ, Duran R. 2008. Structure of bacterial communities along a hydrocarbon contamination gradient in a coastal sediment. *Federation of European Microbiological Societies Microbiology Ecology* 66:295-305.
- Ramsey E, Meyer MB, Rangoonwala A, Overton E, Jones CE, Bannister T. 2014. Oil source-fingerprinting in support of polarimetric radar mapping of Macondo-252 oil in gulf Coast marshes. *Marine Pollution Bulletin* 89:85-89.
- Reddy CM, Arey JS, Seewald JS, Sylva SP, Lemkau KL, Nelson RK, Carmichael CA, McIntyre CP, Fenwick J, Ventura GT, Van Mooy BAS, Camilli R. 2011. Science applications in the Deepwater Horizon oil spill special feature: composition and fate of gas and oil released to the water column during the Deepwater Horizon oil spill. *Proceedings of the National Academy of Sciences of the United States of America*, 109:20292-20297.
- Siddiqui S, Adams WA. 2002. The fate of diesel hydrocarbons in soil and their effect on the germination of perennial ryegrass. *Environmental Toxicology* 17(1), 49-62.

- Silliman BR, van de Koppel J, McCoy MW, Diller J, Kasozi GN, Earl K, Adams PN, Zimmerman AR. 2012. Degradation and resilience in Louisiana salt marshes after the BP-Deepwater Horizon oil spill. *Proceedings of the National Academy of Sciences of the United States of America* 109:11234-11239.
- Shiaris MP. 1989. Seasonal Biotransformation of Naphthalene, Phenanthrene, and Benzo[ $\alpha$ ]pyrene in Surficial Estuarine Sediments. *Applied and Environmental Microbiology* 55:1391-1399.
- Shin WS, Pardue JH, Jackson WA. 2000. Oxygen demand and sulfate reduction in petroleum hydrocarbon contaminated salt marsh soils. *Water Research* 34:1345-1353.
- Slocum MG, Roberts J, Mendelsohn IA. 2009. Art canvas as a new standard for the cotton strip assay. *Journal of Plant Nutrition and Soil Science* 172:71-74.
- Smith CJ, DeLaune RD, Patrick WH, Fleeger JW. 1984. Impact of dispersed and undispersed oil entering a gulf coast salt marsh. *Environmental Toxicology and Chemistry* 3:609-616.
- So CM, Young LY. 2001. Anaerobic biodegradation of alkanes by enriched consortia under four different reducing conditions. *Environmental Toxicology and Chemistry* 20:473-478.
- Stedman S, Dahl TE. 2008. Status and trends of wetlands in the coastal watersheds of the Eastern United States 1998-2004. National Oceanic and Atmospheric Administration, National Marine Fisheries Service and U.S. Department of the Interior, Fish and Wildlife Service, 1-32.

- Suárez-Suárez A, López-López A, Tovar-Sánchez A, Yarza P, Orfila A, Terrados J, Arnds J, Marqués S, Niemann H, Schmitt-Kopplin P, Amann R, Rosselló-Móra R. 2011. Response of sulfate-reducing bacteria to an artificial oil-spill in a coastal marine sediment. *Environmental Microbiology* 13:1488-1499.
- Tiegs SD, Clapcott JE, Griffiths NA, Boulton AJ. 2013. A standardized cotton-strip assay for measuring organic-matter decomposition in streams. *Ecological Indicators* 32: 131-139.
- Turner RE, Overton EB, Meyer BM, Miles MS, Hooper-Bui L. 2014a. Changes in the concentration and relative abundance of alkanes and PAHs from the Deepwater Horizon oiling of coastal marshes. *Marine Pollution Bulletin* 86:291-297.
- Turner RE, Overton EB, Meyer BM, Miles MS, McClenchan G, Hooper-Bui L, Engel AS, Swenson EM, Lee JM, Milan CS. 2014b. Distribution and recovery trajectory of Macondo (Mississippi Canyon 252) oil in Louisiana coastal wetlands. *Marine Pollution Bulletin* 87:57-67.
- Walker JD, Seesman PA, Colwell RR. 1975. Effect of South Louisiana crude oil and no. 2 fuel oil on growth of heterotrophic microorganisms, including proteolytic lipolytic, chitinolytic and cellulolytic bacteria. *Environmental Pollution* 9:13-33.
- White DA, Weiss TE, Trapant JM, Thein LB. 1978. Productivity and decomposition of the dominant salt marsh plants in Louisiana. *Ecology* 59:751-759.
- Widdel F, Rabus R. 2001. Anaerobic biodegradation of saturated and aromatic hydrocarbons. *Current Opinion in Biotechnology* 12:259-276.

Wright AL, Weaver RW, Webb JW. 1997. Oil bioremediation in salt march mesocosms as influenced by N and P fertilization, flooding, and season. *Water, Air, and Soil Pollution* 95:179–191.

Yanto DHY, Tachibana S. 2014. Potential of fungal co-culturing for accelerated biodegradation of petroleum hydrocarbons in soil. *Journal of Hazardous Materials* 278:454-463.