



Technological University Dublin  
ARROW@TU Dublin

---

Articles

School of Biological Sciences

---

2018

## Predictors of Vitamin D Supplementation Amongst Infants in Ireland Throughout The First Year of Life

Annemarie Bennett  
*Trinity College Dublin, Ireland*

John M. Kearney  
*Technological University Dublin, [john.kearney@tudublin.ie](mailto:john.kearney@tudublin.ie)*

Follow this and additional works at: <https://arrow.tudublin.ie/scschbioart>

---

### Recommended Citation

Bennett, A.E. & Kearney, J.M. (2018). Predictors of Vitamin D Supplementation Amongst Infants in Ireland Throughout The First Year of Life. *Journal of Public Health*, vol. 26, no. 5, pg. 577-583. doi.org/10.1007/s10389-017-0891-3

This Article is brought to you for free and open access by the School of Biological Sciences at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact [yvonne.desmond@tudublin.ie](mailto:yvonne.desmond@tudublin.ie), [arrow.admin@tudublin.ie](mailto:arrow.admin@tudublin.ie), [brian.widdis@tudublin.ie](mailto:brian.widdis@tudublin.ie).



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/)



# 1 Predictors of vitamin D supplementation amongst infants in Ireland throughout the first year of life

2 AE Bennett<sup>1\*</sup> and JM Kearney<sup>2</sup>

3 <sup>1</sup> Department of Clinical Medicine, Trinity Centre for Health Sciences, St. James' Hospital Campus, Dublin 8, Ireland

4 <sup>2</sup> School of Biological Sciences, Dublin Institute of Technology, Kevin Street, Dublin 8, Ireland

## 5 Abstract

6 **Aim:** To investigate predictors of compliance with the recommendation that all infants in Ireland are supplemented  
7 daily from birth to 12 months of age with 5 micrograms ( $\mu\text{g}$ ) of vitamin D.

8 **Subjects and methods:** A prospective observational study was completed. Self-complete questionnaires recorded  
9 socio-demographic characteristics, health behaviours and supplementation practices for 158 mother-infant dyads at 4, 9  
10 and 12 months post-partum. A 2-day food diary was also obtained on 12-month-old infants to examine the contribution  
11 of diet to vitamin D intakes.

12 **Results:** At 4, 9 and 12 months of age, 57.6% ( $n=91$ ), 34.2% ( $n=54$ ) and 23.4% ( $n=37$ ) of infants, respectively, were  
13 supplemented as recommended. In multivariate analyses, receiving supplementation advice from health professionals in  
14 the early post-partum period was the most significant predictor of correctly supplementing 4-month-old ( $p<0.01$ , odds  
15 ratio (OR): 61.94 [95% confidence interval (CI): 11.53-332.83]), 9-month-old ( $p<0.01$ , OR: 10.30 [95% CI: 2.29-  
16 46.27]) and 12-month-old ( $p=0.04$ , OR: 3.85 [95% CI: 1.05-14.08]) infants. Amongst 12-month-olds, even intakes from  
17 diet and supplementation combined ( $7.6\pm 4.7\mu\text{g/day}$ ) were suboptimal.

18 **Conclusion:** Suboptimal vitamin D supplementation practices were evident throughout infancy. Dietary intakes of  
19 vitamin D did not compensate for suboptimal supplementation practices. Supplementation practices may improve if  
20 health professionals advocate safe supplementation during routine infant health checks.

## 21 Keywords

22 Vitamin D supplementation; Infancy; Supplementation policy; Infant bone health; Non-compliance

## 23 Introduction

24 While an infant's food intake should ideally meet all of their nutritional needs, it has been consistently observed that the  
25 diets of many infants in Ireland lack sufficient vitamin D (Food Safety Authority of Ireland [FSAI] 2007). The chief  
26 functions of vitamin D are the regulation of calcium homeostasis and bone mineral metabolism (Institute of Medicine  
27 [IOM] 2010). Across the lifespan, the most accelerated rates of growth and bone mineral accretion occur in infancy  
28 (Gallo *et al.* 2012), with evidence indicating that maximising bone accretion during this time benefits bone health at  
29 later stages of the life cycle (Cooper *et al.* 2002; Gallo *et al.* 2012).

30 Therefore, vitamin D deficiency adversely affects bone health, with chronic deficiency resulting in bone  
31 demineralisation. If the deficiency is particularly severe, the resulting demineralisation can reduce bone rigidity, causing  
32 rickets, a condition which manifests in infancy as deformed arms, legs and rib cage. Rickets is the most severe  
33 manifestation of vitamin D deficiency in infancy, and although a small number of cases have re-emerged in Ireland in  
34 recent years, it remains an uncommon condition (FSAI 2007). However, chronic mild vitamin D deficiency may be  
35 more widespread due to inadequate dietary vitamin D intakes and compliance with the recommended practice of not  
36 exposing infants to sunlight (FSAI 2007; Irish Universities Nutrition Alliance [IUNA] 2012).

37 In light of the attention called to vitamin D deficiency in infancy, it was recommended that all infants in  
38 Ireland are supplemented with 5 micrograms ( $\mu\text{g}$ ) of vitamin D every day from birth to their first birthday (FSAI 2007;

1 Health Service Executive [HSE] 2010). Although this recommendation was devised in 2007 (FSAI 2007) and  
2 implemented as national policy in 2010 (HSE 2010), compliance with the policy has not yet been assessed.

3 In addition, given the history of non-compliance with supplementation policies amongst Irish women  
4 (McKeating *et al.* 2015), it is important to identify factors which increase the likelihood of Irish mothers correctly  
5 supplementing their infants with vitamin D.

6 Therefore, to assess the effectiveness of strategies currently in place to safeguard infant bone health in Ireland,  
7 this study aimed to assess compliance with recommendations for vitamin D supplementation in infancy and to  
8 investigate the predictors of a mother supplementing her infant as recommended.

## 9 **Methods**

10 Ethical approval for this prospective observational study was obtained from the Coombe Women and Infants University  
11 Hospital, and Our Lady's Children's Hospital Crumlin and Dublin Institute of Technology.

12 Women were recruited to this study whilst waiting in antenatal clinics in the Coombe Women and Infants  
13 University Hospital between October 2013 and August 2014. Women were eligible to participate in the study if they:  
14 had a healthy singleton pregnancy; were at least 24 weeks pregnant; and were willing to be contacted in the post-partum  
15 period for follow-up. Written informed consent was obtained from all women. A questionnaire was administered in  
16 clinic by the lead author for participants to self-complete. This questionnaire collected socio-demographic and health  
17 behaviour data, to include: age; nationality; parity; education level; marital status; risk of deprivation; antenatal folic  
18 acid supplementation practices; and smoking status.

19 To build rapport with participants and reduce drop-out rates, study participants were followed-up in hospital  
20 after giving birth to ensure that they were still amenable to being contacted by the lead author for a home visit at 4, 9  
21 and 12 months post-partum. Therefore, each morning during data collection, the lead author checked hospital records to  
22 identify those women who had given birth to a healthy term infant in the previous 24 hours and who had consented to  
23 participate in this study. Upon identifying such women, the lead author visited them on the ward, re-introduced herself  
24 and reminded mothers of the study, asking them if they were still happy to be contacted by the researcher in the post-  
25 partum period. Home visits were arranged by phone approximately one week before the infant turned 4, 9 or 12 months  
26 of age. During each home visit, a quantitative questionnaire on infant feeding and supplementation practices was self-  
27 completed by mothers. Closed-ended questions on vitamin D supplementation practices at 4, 9 and 12 months post-  
28 partum included: whether a mother was supplementing her infant with vitamin D; the frequency of vitamin D  
29 supplementation; the brand name of the supplement; and whether advice had been received from a recognised health  
30 professional (*e.g.* medical doctor or public health nurse) on vitamin D supplementation.

31 In addition to completing a questionnaire at 12 months post-partum, mothers also completed a 2-day food diary  
32 for their 12-month-old infant using standard household measures. Food diaries were not completed at any other point of  
33 data collection. There is no national policy in Ireland on vitamin D supplementation for individuals aged over one year.  
34 Therefore, the food diaries were obtained to examine the separate contributions of food and supplements to vitamin D  
35 intakes and to compare vitamin D intakes from food alone with the recommended intake of 10µg/d (IOM 2010); this  
36 was important to assess the adequacy of vitamin D intakes from food alone as the first year of life draws to a close and  
37 guidance on supplementation is no longer provided. In the diary, mothers recorded the: time of eating occasion; source  
38 of food or drink; brand of food eaten (if applicable); volume of food or drink consumed using standard household  
39 measures; cooking methods used; and condiments added (if any). Data were entered into *Nutritics* Nutrition Analysis  
40 Software (*Nutritics*, Dublin, Ireland). The *Nutritics* software calculated mean energy and nutrient intakes for each infant  
41 and the resulting figures were entered into IBM SPSS for Windows, version 22.0 (IBM, New York, United States).

1 IBM SPSS for Windows, version 22.0 (IBM, New York, United States) was used for analysis. Normally  
2 distributed data were summarised numerically using the mean and standard deviation (SD). Variables associated with  
3 vitamin D supplementation practices at 4, 9 and 12 months of age were examined in univariate analyses. Such variables  
4 included: maternal age (years); education (college-educated/not college-educated); social class (high/middle/low);  
5 parity (primiparous/multiparous); first milk (breast/formula); breastfeeding duration (days); age of weaning (weeks);  
6 smoking status at conception (yes/no); smoked all throughout pregnancy (yes/no); smoking at time of questionnaire  
7 completion (yes/no); consumed alcohol in pregnancy (yes/no); correctly supplemented with folic acid (yes/no); and  
8 recalled receiving vitamin D supplementation advice from a qualified health professional (yes/no).

9 Associations with normally distributed continuous variables were assessed by an Independent Samples *t*-test.  
10 Associations between categorical variables were assessed using cross-tabulations and the Chi-squared statistics test  
11 assessed statistical significance. Variables which were significantly associated with recommended supplementation  
12 practices in univariate analyses were included in multivariate analyses. Binary logistic regression was used to predict  
13 recommended supplementation practices at 4, 9 and 12 months of age. The Forced Entry Method was used, whereby all  
14 predictor variables were tested in one block to assess their predictive ability whilst controlling for the effects of other  
15 predictors in the model. Statistical significance was taken at  $p < 0.05$ .

## 16 Results

### 17 *Baseline characteristics of participants*

18 One-hundred-and-seventy-two pregnant women were recruited for this study. Of these, 158 mothers (91.9% follow-up  
19 rate) allowed the lead researcher to conduct a home visit at 4, 9 and 12 months post-partum. There were no significant  
20 differences in the socio-demographic characteristics and health behaviours of mothers who did and did not withdraw  
21 from the study. The mean age of the 158 participating women upon giving birth was 32.0 (SD  $\pm$  4.9) years and the mean  
22 gestational age of infants was 40.3 (SD  $\pm$  1.2) weeks.

23 Participating mothers were either of Irish (98.1%,  $n$ 155) or British (1.9%,  $n$ 3) nationality. Almost three-  
24 quarters (73.4%,  $n$ 116) had planned their pregnancy and 43.0% ( $n$ 68) were primiparous. Two-thirds (65.2%,  $n$ 103) were  
25 college-educated and 86.7% ( $n$ 137) were married or cohabiting. Almost a third (32.3%,  $n$ 51) smoked around the time of  
26 conception, but only 8.2% ( $n$ 13) smoked all throughout pregnancy. Almost all (99.4%,  $n$ 157) took folic acid in  
27 pregnancy to some degree, but less than a third (32.3%,  $n$ 51) supplemented with folic acid in line with national  
28 recommendations.

### 29 *Vitamin D supplementation practices*

30 The proportions of infants being supplemented with vitamin D in line with recommendations at 4, 9 and 12 months of  
31 age were 57.6% ( $n$ 91), 34.2% ( $n$ 54) and 23.4% ( $n$ 37), respectively (**Figure I**).

### 32 *Predictors of recommended vitamin D supplementation practices*

33 As shown in the statistically significant adjusted model ( $\chi^2$  (5,  $n$ 158) = 55.42,  $p < 0.01$ ) in **Table I**, the strongest predictor  
34 of correctly supplementing an infant with vitamin D at 4 months post-partum was receiving advice from a health  
35 professional on doing so. Mothers who recalled receiving advice from a health professional on supplementation were  
36 almost 62 times (OR: 61.94 [95% CI: 11.53-332.83]) more likely to correctly supplement their infant when compared  
37 with mothers who did not receive supplementation advice (**Table I**). Maternal education also significantly predicted  
38 supplementation at this time; mothers who were college-educated were significantly ( $p = 0.03$ ) less likely to supplement  
39 their 4-month-old infant as recommended (OR: 0.37 [95% CI: 0.15-0.92]).

1 At 9 months post-partum, the strongest predictor of correctly supplementing an infant with vitamin D was  
2 receiving advice from a health professional on doing so. Mothers who recalled receiving supplementation advice in the  
3 early post-partum period were ten times (OR: 10.30 [95% CI: 2.29-46.27]) more likely to correctly supplement their 9-  
4 month-old infant when compared with mothers who did not receive this advice (**Table II**). As shown in the statistically  
5 significant adjusted model ( $\chi^2$  (5,  $n=158$ ) = 28.48,  $p < 0.01$ ) in **Table II**, smoking status in pregnancy also significantly  
6 ( $p = 0.02$ ) predicted vitamin D supplementation; mothers who did not smoke antenatally were 2.5 times more likely to  
7 correctly supplement their 9-month-old infant (OR: 2.57 [95% CI: 1.13-5.81]).

8 At 12 months post-partum, the strongest predictor of correctly supplementing an infant with vitamin D was  
9 receiving advice from a health professional on doing so. As shown in the statistically significant adjusted model ( $\chi^2$  (5,  
10  $n=158$ ) = 12.34,  $p = 0.03$ ) in **Table III**, mothers who recalled receiving advice on supplementation in the early post-  
11 partum period were almost four times (OR: 3.85 [95% CI: 1.05-14.08]) more likely to correctly supplement their 12-  
12 month-old infant when compared with mothers who did not receive advice at this time (**Table III**). Maternal education  
13 also significantly ( $p = 0.04$ ) predicted supplementation practices, where mothers who were college-educated were  
14 significantly less likely to supplement their 12-month-old infant as recommended (OR: 0.42 [95% CI: 0.19-0.95]).

#### 15 *Dietary vitamin D intakes at 12 months of age*

16 The mean age at which complementary feeding commenced in this study was 20.7 (SD  $\pm$  4.6) weeks. The mean intake  
17 of vitamin D from food and beverages amongst 12-month-old infants ( $n=153$ ) was 6.3 (SD  $\pm$  4.4)  $\mu\text{g}$  per day. When  
18 vitamin D from supplementation was included, the mean daily intake of vitamin D from all sources increased to 7.6 (SD  
19  $\pm$  4.7)  $\mu\text{g}$ .

#### 20 **Discussion**

21 The current recommended dietary allowance (RDA) for vitamin D amongst infants is 10 $\mu\text{g}$  per day (IOM 2010). To  
22 help achieve this intake, it is recommended that all infants in Ireland consume 5 $\mu\text{g}$  of vitamin D by supplementation  
23 each day from birth to their first birthday (HSE 2010).

24 However, despite the development of a widely disseminated policy to promote daily vitamin D  
25 supplementation for all infants in Ireland (HSE 2010), supplementation practices in this study were suboptimal  
26 throughout the first year of life. At age one year, less than a quarter of infants were being correctly supplemented and  
27 their vitamin D intakes from diet alone were inadequate to meet the RDA. Therefore, it is clear that this is an aspect of  
28 paediatric health which should continue to receive attention.

29 It is notable that the advice of health professionals was the most consistent and significant predictor of  
30 recommended vitamin D supplementation practices in this study, surpassing many maternal sociodemographic  
31 characteristics and health behaviours. Mothers who recalled receiving supplementation advice from a health  
32 professional in the weeks following their infant's birth were significantly more likely to correctly supplement their  
33 infant for the remainder of the first year of life. The strength of this association waned as the first year of life  
34 progressed; however, better adherence to recommendations may result if health professionals are reminded to advocate  
35 supplementation and appropriate food sources of vitamin D during routine infant health checks. During infant health  
36 checks, it is expected that a public health nurse or medical officer will discuss child and family health issues and assess  
37 an infant's general physical health, to include developmental progress and growth (HSE 2017). A brief discussion on  
38 the importance of vitamin D supplementation would fall within the remit of such health checks and would be a cost-  
39 effective approach to promoting adherence to recommended supplementation practices.

40 Although varying degrees of effectiveness have been reported (Arditi *et al.* 2012), electronic clinical reminders  
41 have generally been shown to achieve small to modest improvements in healthcare quality (Gandhi *et al.* 2003; Sequist

1 *et al.* 2005; Vashitz *et al.* 2009). It has been observed that if health professionals receive specific reminders to do so,  
2 they are more likely to promote adherence to particular health guidelines amongst their patients (Sequist *et al.* 2005).  
3 However, while the research to date has examined the use of reminders with health issues such as vaccinations  
4 (Kersting and Weltermann 2016) and health screenings (Kenealy *et al.* 2005), the use of reminders to encourage health  
5 professionals to promote adherence to supplementation policies has not been investigated. Given the positive impact of  
6 advice from health professionals in this study on supplementation practices, future research should investigate whether  
7 a reminder system (Shojania *et al.* 2010) for health professionals in primary care elicits improved adherence to  
8 recommended supplementation practices in infancy.

9 Interestingly, mothers with a college education were significantly less likely to supplement their infant as  
10 recommended. Using the Health Belief Model (Abraham and Sheeran 2005), it could be argued that these mothers felt  
11 confident in the nutritional adequacy of the diet which they provided to their infant, and therefore were less likely to  
12 believe that supplementation was necessary to avoid poor health outcomes (Fulford *et al.* 2014; Malek *et al.* 2016;  
13 Touskova *et al.* 2015). However, given the small number of foods which are both a source of vitamin D and appropriate  
14 for infants (FSAI 2011), educational campaigns should state that even a high quality varied diet does not circumvent the  
15 need for vitamin D supplementation (Fulford *et al.* 2014; FSAI 2007).

16 Mothers who smoked during pregnancy were also less likely to supplement their infant in line with  
17 recommendations. It has been suggested (Fulford *et al.* 2014) that women who engage in adverse health behaviours but  
18 have seemingly healthy births may perceive themselves and their offspring to be invulnerable to poor health outcomes  
19 and therefore may not see a need to invest in certain protective health behaviours (Barbour *et al.* 2012). However, given  
20 the insidious and often silent progression of bone demineralisation over the life course (Touskova *et al.* 2015), public  
21 health campaigns must emphasise to all parents that no infant is exempt from the need for vitamin D supplementation.

22 As expected, vitamin D intakes from diet alone were insufficient to meet the RDA, and this further underpins  
23 the need to emphasise appropriate supplementation practices as soon as is practicable. Since approximately half of the  
24 maternal stores of vitamin D are transferred to the foetus during pregnancy (Cooper *et al.* 2002), health professionals  
25 could advise pregnant women on appropriate vitamin D-rich foods and safe vitamin D supplementation (FSAI 2011).  
26 Educating women during pregnancy on the value of vitamin D for mother and infant could help parents to establish  
27 positive supplementation practices which optimise infant bone health from the earliest days of development (Bener *et*  
28 *al.* 2013; Christesen *et al.* 2012).

### 29 *Limitations of this study*

30 The data presented here were collected as part of a longitudinal observational study conducted by one researcher in  
31 Dublin and its surrounding counties. Although inter-observer variation is not a concern, the results are not nationally  
32 representative and causal inferences cannot be made due to the observational study design (Grimes and Schulz 2002).  
33 The study population was limited to participants who were Caucasian and of Irish or British nationality, and therefore  
34 future research should investigate vitamin D supplementation practices amongst a nationally representative sample.  
35 Future research should also triangulate supplementation practices amongst parents with the perspectives of health  
36 professionals and the directives issued by health authorities. Once the views of all stakeholders have been considered, a  
37 better understanding of the most effective means of promoting adherence to recommendations (to include the timing,  
38 frequency, content and mode of delivery of reminders) should become clearer.

39 Dietary intakes amongst 12-month-old infants were measured using estimated household measures. Weighed  
40 measures would be considered preferable to assess food intake in infants (Burrows *et al.* 2010; Smith 2011), but this  
41 was not feasible when study resources and participant burden were considered. Nevertheless, the food diaries used did  
42 obtain detailed data on the foods consumed and measuring spoons were provided to help carers to complete the diary.

1 The carer who made an entry in the diary (*e.g.* mother, father, childminder) was recorded for each eating occasion. To  
2 account for the possibility of different carers completing the diary, clear instructions were included to reduce variation  
3 in the method of recording the food consumed. Finally, although estimated household measures were used, the vitamin  
4 D intakes in this study were comparable to those reported by the nationally representative National Preschool Nutrition  
5 Survey (IUNA 2012) for 12-month-olds, in which weighed measures were used to assess dietary intakes.

## 6 **Conclusion**

7 Dietary practices throughout the first year of life have received considerable attention in Ireland (Bennett *et al.* 2012;  
8 Tarrant *et al.* 2010). However, to the authors' knowledge, this is the first study to examine compliance with  
9 recommended vitamin D supplementation practices amongst Irish infants since the implementation of a national policy  
10 in 2010 (HSE 2010).

11 This study highlights the degree to which compliance with recommendations is suboptimal, and provides  
12 insights into potentially modifiable factors which affect compliance, in addition to identifying sub-groups at which  
13 educational campaigns could be specifically targeted. The identification of strategies which promote adherence to  
14 supplementation policy is challenging (McKeating *et al.* 2015). However, the development of such strategies is made  
15 possible by a better understanding of the factors which help or hinder adherence to supplementation policy (Patience  
16 2015). This study has contributed to our understanding of factors which potentially impact bone health during a time of  
17 immeasurable investment into life-long health.

## 18 **Acknowledgements**

19 The authors wish to acknowledge the mothers who participated in this study; their time was greatly appreciated.

## 20 **Compliance with Ethical Standards**

21 Funding: The lead author was funded by a Dublin Institute of Technology *Fiosraigh* Scholarship. This scholarship was  
22 50% funded by Dublin Institute of Technology and 50% funded by Danone Nutricia Early Life Nutrition.

23 Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical  
24 standards of the institutional research committees and with the 1964 Helsinki declaration and its later amendments.

25 Informed consent: Informed consent was obtained from all individual participants included in the study.

26 Conflict of interest: The authors have no conflict of interest to declare.

## 27 **References**

- 28 Abraham C, Sheeran P (2005) *The health belief model*. In: Conner M, Norman P (ed) *Predicting health behaviour*,  
29 Open University Press. Berkshire, England, pp 28-80.
- 30 Arditi C, Rège-Walther M, Wyatt JC, Durieux P, Burnand B (2012) Computer-generated reminders delivered on paper  
31 to healthcare professionals; effects on professional practice and healthcare outcomes. *Cochrane Database Syst*  
32 *Review*; doi: 10.1002/14651858.CD001175.pub3.
- 33 Barbour RS, Macleod M, Mires G, Anderson AS (2012) Uptake of folic acid supplements before and during pregnancy:  
34 focus group analysis of women's views and experiences. *J Hum Nutr Diet*; **25**:140-147.
- 35 Bener A, Al-Hamaq AOAA, Saleh NM (2013) Association between vitamin D insufficiency and adverse pregnancy  
36 outcome: global comparisons. *Int J Women Health*; **5**:523-531.
- 37 Bennett AE, O'Connor AL, Canning N, Kenny A, Keaveney E, Younger K, Flynn MA (2012) Weaning onto solid  
38 foods: some of the challenges. *Ir Med J*; **105**:266-268.


- 1 Burrows TL, Martin RJ, Collins CE (2010) A systematic review of the validity of dietary assessment methods in  
2 children when compared with the method of doubly labeled water. *J Am Diet Assoc*; **110**:1501-1510.
- 3 Christesen HT, Elvander C, Lamont RF, Jørgensen JS (2012) The impact of vitamin D in pregnancy on extraskeletal  
4 health in children: a systematic review. *Acta Obstet Gynecol Scand*; **91**:1368-1380.
- 5 Cooper C, Javaid MK, Taylor P, Walker-Bone K, Dennison E, Arden N (2002) The fetal origins of osteoporotic  
6 fracture. *Calcif Tissue Int*; **70**:391-394.
- 7 Food Safety Authority of Ireland (2011) *Scientific recommendations for a national infant feeding policy* (2<sup>nd</sup> ed.)  
8 Dublin, Ireland: Author.
- 9 Food Safety Authority of Ireland (2007) *Recommendations for a national policy on vitamin D supplementation for*  
10 *infants in Ireland*. Dublin, Ireland: Author.
- 11 Fulford B, Macklon N, Boivin J (2014) Mental models of pregnancy may explain low adherence to folic acid  
12 supplementation guidelines: a cross-sectional international survey. *Eur J Obstet Gynaecol Reprod Biol*; **176**:99-  
13 103.
- 14 Gallo S, Vanstone CA, Weiler HA (2012) Normative data for bone mass in healthy term infants from birth to 1 year of  
15 age. *J Osteoporos*; ID 672403.
- 16 Gandhi TK, Sequist TD, Poon EG, Karson AS, Murff H, Fairchild DG, Kuperman GJ, Bates DW (2003) Primary care  
17 clinician attitudes towards electronic clinical reminders and clinical practice guidelines. *AMIA Annu Symp*  
18 *Proc*;848.
- 19 Grimes DA, Schulz KF (2002) Bias and causal associations in observational research. *Lancet* **359**:248-252.
- 20 Health Service Executive (2017) *A guide to your child's growth and development*. Available from:  
21 <http://www.hse.ie/eng/health/child/cfyb/2-5yrs/growthdevelopment/>. Accessed 10 December 2017.
- 22 Health Service Executive (2010) *Policy on vitamin D supplementation for infants in Ireland, 2010*. Available from:  
23 <https://www.hse.ie/eng/health/child/vitaminD/Vitamin%20D%20Policy.pdf>. Accessed 17 July 2015.
- 24 Institute of Medicine (2010) *Dietary reference intakes for calcium and vitamin D* (1st ed.). Washington DC, United  
25 States of America: The National Academies Press.
- 26 Irish Universities Nutrition Alliance (2012) *National Pre-school Nutrition Survey: summary report*. Cork, Ireland:  
27 Author.
- 28 Kenealy T, Arroll B, Petrie KJ (2005) Patients and computers as reminders to screen for diabetes in family practice. *J*  
29 *Gen Intern Med*; **20**:916-921.
- 30 Kersting C, Weltermann B (2016) Electronic reminders to facilitate longitudinal care: a mixed-methods study in general  
31 practices. *BMC Med Inform Decis Mak*; **16**:148.
- 32 Malek L, Umberger W, Makrides M, Zhou SJ (2016) Poor adherence to folic acid and iodine supplement  
33 recommendations in preconception and pregnancy: a cross-sectional analysis. *Aust N Z J Public Health*; **40**:424-  
34 429.
- 35 McKeating A, Farren M, Cawley S, Daly N, McCartney D, Turner M (2015) Maternal folic acid supplementation trends  
36 2009-2013. *Acta Obstet Gynecol Scand*; **94**:727-733.
- 37 Patience S (2015) Promoting vitamin D uptake in infants and children. *BJM*; **23**:10S-13S.
- 38 Sequist TD, Gandhi TK, Karson AS, Fiskio JM, Bugbee D, Sperling M, Cook EF, Orav EJ, Fairchild DG, Bates DW  
39 (2005) A randomized trial of electronic clinical reminders to improve quality of care for diabetes and coronary  
40 artery disease. *J Am Med Inform Assoc*; **12**:431-437.
- 41 Shojanian KG, Jennings A, Mayhew A, Ramsay C, Eccles M, Grimshaw J (2010) Effect of point-of-care computer  
42 reminders on physician behaviour: a systematic review. *CMAJ*; **182**:216-25.



- 1 Smith AE (2011) Validation studies of diets of children and adolescents. *J Am Diet Assoc*; **111**:1124-1125.
- 2 Tarrant RC, Younger KM, Sheridan Pereira M, White MJ, Kearney JM (2010) Factors associated with weaning  
3 practices in term infants: a prospective observational study in Ireland. *Br J Nutr*; **104**:1544-1554.
- 4 Touskova T, Vytrisalova M, Palicka V, Hendrychova T, Fуска L, Holcova R, Konopacova J, Kubena AA (2015) Drug  
5 holidays: the most frequent type of non-compliance with calcium plus vitamin D supplementation in persistent  
6 patients with osteoporosis. *Patient Prefer Adherence*; **9**:1771-1779.
- 7 Vashitz G, Meyer J, Parmet Y, Peleg R, Goldfarb D, Porath A, Gilutz H (2009) Defining and measuring physicians'  
8 responses to clinical reminders. *J Biomed Inform*; **42**:317-326.

9

Corrected proof



1

2 **Figure I. Proportion of infants receiving a daily supplement of 5 micrograms (µg) of vitamin D**  
3 **during the first year of life**

4

5

6

Corrected proof

**Table I.** Binary logistic regression model examining factors associated with recommended vitamin D supplementation amongst 158 4-month-old infants

Characteristic	$\beta$	<i>n</i>	OR	95% CI	<i>p</i> -value*
Maternal third level education					
Yes	-	103	0.37	0.15 – 0.92	<b>0.03</b>
No		55	1.0	Ref.	
Recommended folic acid supplementation					
Yes	+	51	1.56	0.67 – 3.64	0.30
No		107	1.00	Ref.	
Smoking in pregnancy					
Yes	+	51	1.00	Ref.	0.32
No		107	1.51	0.67 – 3.41	
Vitamin D advice from health professional					
Yes	+	127	61.94	11.53 – 332.83	<b>&lt;0.01</b>
No		31	1.0	Ref.	
Maternal age	-	158	0.92	0.84 – 0.99	0.06
Model summary:					
R <sup>2</sup> = 0.29, Cox & Snell R Square = 29.6, Nagelkerke R Square = 39.8, 74.7% predictive of variance					
* <i>p</i> -value <0.05 was significant		<b>OR:</b> Odds ratio		<b>CI:</b> Confidence interval	

2

3

**Table II.** Binary logistic regression model examining factors associated with recommended vitamin D supplementation amongst 158 9-month-old infants

Characteristic	$\beta$	<i>n</i>	OR	95% CI	<i>p</i> -value*
Maternal third level education					
Yes	-	103	0.67	0.31 – 1.46	0.32
No		55	1.0	Ref.	
Recommended folic acid supplementation					
Yes	+	51	2.06	0.95 – 4.45	0.07
No		107	1.00	Ref.	
Smoking in pregnancy					
Yes	+	51	1.00	Ref.	<b>0.02</b>
No		107	2.57	1.13 – 5.81	
Vitamin D advice from health professional					
Yes	+	127	10.30	2.29 – 46.27	<b>&lt;0.01</b>
No		31	1.0	Ref.	
Maternal age	+	158	1.04	0.96 – 1.12	0.40
Model summary:					
R <sup>2</sup> = 0.23, Cox & Snell R Square = 16.5, Nagelkerke R Square = 22.8, 71.5% predictive of variance					
* <i>p</i> -value <0.05 was significant		<b>OR:</b> Odds ratio		<b>CI:</b> Confidence interval	

4

1

**Table III.** Binary logistic regression model examining factors associated with recommended vitamin D supplementation amongst 158 12-month-old infants

Characteristic	$\beta$	<i>n</i>	OR	95% CI	<i>p</i> -value*
Maternal third level education					
Yes	-	103	0.42	0.19 – 0.95	<b>0.04</b>
No		55	1.0	Ref.	
Recommended folic acid supplementation					
Yes	+	51	1.21	0.53 – 2.80	0.65
No		107	1.00	Ref.	
Smoking in pregnancy					
Yes	+	51	1.00	Ref.	0.09
No		107	2.16	0.88 – 5.33	
Vitamin D advice from health professional					
Yes	+	127	3.85	1.05 – 14.08	<b>0.04</b>
No		31	1.0	Ref.	
Maternal age	+	158	1.03	0.94 – 1.12	0.56

Model summary:

 $R^2 = 0.16$ , Cox & Snell R Square = 7.5, Nagelkerke R Square = 11.3, 77.8% predictive of variance\* *p*-value <0.05 was significant**OR:** Odds ratio**CI:** Confidence interval

2

3

Corrected proof