



Technological University Dublin  
ARROW@TU Dublin

Articles

School of Chemical and Pharmaceutical  
Sciences

2012

## Analogues of the epoxy resin monomer diglycidyl ether of bisphenol F: Effects on contact allergenic potency and cytotoxicity

Niamh O'Boyle

Technological University Dublin, [niamh.oboyle@tudublin.ie](mailto:niamh.oboyle@tudublin.ie)

Tamara Delaine

University of Goteborg

Kristina Luthman

University of Goteborg, [luthman@chem.gu.se](mailto:luthman@chem.gu.se)

Andreas Natsch

Givaudin Schweiz AG, Switzerland, [andreas.natsch@givaudan.com](mailto:andreas.natsch@givaudan.com)

Follow this and additional works at: <https://arrow.tudublin.ie/scschcpsart>

Therese Karlberg  
Pan European Commons Actions and Uses Commons, Environmental Health Commons, Organic  
University of Goteborg, [karlberg@chem.gu.se](mailto:karlberg@chem.gu.se)  
Chemicals Commons, and the Toxicology Commons

### Recommended Citation

O'Boyle, N. et al (2013). Analogues of the epoxy resin monomer diglycidyl ether of bisphenol F: Effects on cont. act allergenic potency and cytotoxicity. *Chemical Research in Toxicology*, 2012, 25(11) pp. 2469-2478.

This Article is brought to you for free and open access by the School of Chemical and Pharmaceutical Sciences at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact [yvonne.desmond@tudublin.ie](mailto:yvonne.desmond@tudublin.ie), [arrow.admin@tudublin.ie](mailto:arrow.admin@tudublin.ie), [brian.widdis@tudublin.ie](mailto:brian.widdis@tudublin.ie).



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/)



**Final Draft of:**

**Analogue of the epoxy resin monomer diglycidyl ether of bisphenol F: Effects on contact allergenic potency and cytotoxicity**

Niamh M. O'Boyle\*, Tamara Delaine, Kristina Luthman, Andreas Natsch, Ann-Therese Karlberg\*

Chemical Research in Toxicology, 2012, Vol. 25, Issue 11, Pages 2469-2478

DOI: <http://dx.doi.org/10.1021/tx300305k>

**Analogue of the Epoxy Resin Monomer Diglycidyl Ether of Bisphenol F: Effects on Contact Allergenic Potency and Cytotoxicity**

Niamh O'Boyle<sup>†\*</sup>, Tamara Delaine<sup>†</sup>, Kristina Luthman<sup>‡</sup>, Andreas Natsch<sup>§</sup>, Ann-Therese Karlberg<sup>†\*</sup>


<sup>†</sup>Department of Chemistry and Molecular Biology, Dermatochemistry and Skin Allergy, University of Gothenburg, SE-412 96 Gothenburg, Sweden

<sup>‡</sup>Department of Chemistry and Molecular Biology, Medicinal Chemistry, University of Gothenburg, SE-412 96 Gothenburg, Sweden

<sup>§</sup>Givaudan Schweiz AG, Ueberlandstrasse 138, CH-8600 Duebendorf, Switzerland

\*Corresponding authors. Email: oboyleni@tcd.ie; karlberg@chem.gu.se

**Table of Contents Graphic**



## Abstract

Diglycidyl ethers of bisphenol A (DGEBA) and bisphenol F (DGEBF) are widely used as components in epoxy resin thermosetting products. They are known to cause occupational and non-occupational allergic contact dermatitis. The aim of this study is to investigate analogues of DGEBF with regard to contact allergy and cytotoxicity. A comprehensive knowledge of the structural features that contribute to the allergenic and cytotoxic effects of DGEBF will guide the development of future novel epoxy resin systems with reduced health hazards for those coming into contact with them. It was found that the allergenic effects of DGEBF were dependent on its terminal epoxide groups. In contrast, it was found that the cytotoxicity in monolayer cell culture was not only dependent on the presence of epoxide groups, but also on other structural features.

## Abbreviations

DGEBA	Diglycidyl ether of bisphenol A
DGEBF	Diglycidyl ether of bisphenol F
ERS	Epoxy resin systems
IARC	International Agency for Research on Cancer
LLNA	Local lymph node assay
MTT	3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
N-ACME	N-Acetyl-L-cysteine methyl ester
PGE	Phenyl glycidyl ether
PPAR $\gamma$	Peroxisome proliferator-activated receptor-gamma

## Introduction

Epoxy resin systems (ERS) are commercial thermosetting products that are combinations of epoxy resins, curing agents, modifiers, and reactive diluents used in applications where strong, flexible, and light-weight construction materials are required. The global epoxy resin market is projected to reach over 3 million tons in annual sales by 2017.<sup>1</sup> Examples of their uses are in paints, adhesives, coatings and electrical laminates. Due to their technical advantages, ERS continue to find new applications. Epoxy resin monomers are polymer precursor units which are reacted with hardeners to give the polymeric material. The most commonly used epoxy resin monomers are diglycidyl ethers based on bisphenol A (DGEBA) (also known as BADGE) and bisphenol F (DGEBF or BFDGE) (Figure 1). Alternatives to DGEBA and DGEBF are receiving commercial interest, as epoxy resins with a variety of properties are desired. ERS are often modified by the addition of reactive diluents, which are used mainly to reduce the viscosity and improve polymerization. Phenylglycidyl ether (PGE, Figure 1) is an example of a commonly used reactive diluent.

The main concerns with use of ERS at present are their environmental impact, their cytotoxicity and their ability to cause contact allergy. Epoxy resin monomers are among the most common causative agents of occupational contact dermatitis.<sup>2</sup> Skin sensitization is known to be particularly common amongst construction workers and is caused by ERS present in cement and other building materials,<sup>3,4</sup> but is also frequent in newer settings such as the production of wind turbine rotor blades<sup>5</sup> and epoxy pipe relining.<sup>6</sup> Epoxy resin monomers are also implicated in non-occupational contact allergy.<sup>7,8</sup> DGEBA is included in the European baseline series for diagnosis of contact allergy.<sup>9</sup> ERS components other than the resin monomers have also been shown to be potent sensitizers.<sup>10</sup> Allergic contact dermatitis from various epoxy resin system components<sup>6,11-</sup>

<sup>18</sup> including DGEBA and DGEBF<sup>5, 19</sup> has been reported. The skin sensitization potential has been investigated experimentally *in vivo* in mice and guinea pigs for DGEBA,<sup>20, 21</sup> DGEBF,<sup>21, 22</sup> PGE<sup>23, 24</sup> and others.<sup>25</sup> DGEBA, DGEBF and PGE are strong sensitizers in both species according to regulatory classifications.<sup>26</sup>

A number of ERS components have been demonstrated to be cytotoxic. PGE and resorcinol diglycidyl ether are classified by IARC as possibly carcinogenic to humans (class 2B).<sup>27</sup> PGE is known to bind to and damage DNA *in vitro*.<sup>28</sup> DNA damage, with sufficient dose and length of exposure, will trigger apoptosis and cytotoxicity.<sup>29</sup> The genotoxicity of epoxides including PGE is known<sup>30</sup> and PGE was suggested to be a direct-acting mutagen as long ago as 1979.<sup>31</sup> A previous study of a series of six epoxides noted that diepoxide functionality and aromaticity increased the induction of apoptosis and cytotoxicity.<sup>29</sup> It has been shown that DGEBA alkylates the isolated nucleophiles 4-(p-nitrobenzyl)-pyridine and deoxyguanosine.<sup>32</sup> DGEBF is cytotoxic in the hepatoma cell line HepG2<sup>33</sup> and in intestinal Caco-2 cells.<sup>34</sup> DGEBA was shown to be an antagonist of the nuclear transcription factor peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) in 3T3-L1 and 3T3-F442A preadipocytes.<sup>35</sup> PPAR $\gamma$  inhibitors cause apoptosis and cell cycle arrest in colorectal carcinoma cells.<sup>36</sup> In contrast, DGEBA acted as a PPAR $\gamma$  agonist in an ECV403 cell line.<sup>37</sup> Further research demonstrated that DGEBA can also induce apoptosis independently of PPAR $\gamma$  in both caspase-dependent and independent manners.<sup>38, 39</sup>

In this work we sought to investigate which structural features of DGEBF are important for its skin sensitizing potency. The *in vivo* murine local lymph node assay (LLNA)<sup>40</sup> and the *in vitro* KeratinoSens™ assay<sup>41</sup> were used to determine the sensitizing effects of the compounds. We also assessed the *in vitro* cytotoxicity of DGEBF analogues in the KeratinoSens™ cell line. The

chemical reactivity was investigated using a model peptide and a model amino acid to estimate the possibility of binding to reactive residues in skin proteins.

## **Experimental Procedures**

**Caution:** This study involves skin sensitizing compounds which must be handled with care.

**Instrumentation and Mode of Analysis.**  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy was performed on a Jeol Eclipse 400 spectrometer at 400 and 100 MHz, respectively, using  $\text{CDCl}_3$  solutions (residual  $\text{CHCl}_3$   $\delta$  7.26 and  $\text{CHCl}_3$   $\delta$  77.0 as internal standards). Electron-ionization mass spectral analysis (70 eV) was performed on a Hewlett-Packard 5973 mass spectrometer connected to a gas chromatograph (Hewlett-Packard 6890). The GC was equipped with a cool on-column capillary inlet and an HP-5MSi fused silica capillary column (30 m  $\times$  0.25 mm, 0.25  $\mu\text{m}$ , Agilent Technologies, Palo Alto, CA). Helium was used as carrier gas, and the flow rate was 1.2 mL/min. The temperature program started at 70  $^\circ\text{C}$  for 1 min, increased by 10  $^\circ\text{C}/\text{min}$ , and ended at 270  $^\circ\text{C}$  for 5 min. For mass spectral analysis, the mass spectrometer was used in the scan mode detecting ions with  $m/z$  values ranging from 50 to 1500.

High performance liquid chromatography/mass spectrometry (LC/MS) analyses were performed using electrospray ionization (ESI) on a Hewlett-Packard 1100 HPLC/MS. The system included a vacuum degasser, a binary pump, an autoinjector, a column thermostat, a diode array detector, and a single quadrupole mass spectrometer. The HPLC was equipped with a HyPURITY C18 column (150  $\times$  3 mm i. d., particle size 3  $\mu\text{m}$ , Thermo Hypersil-Keystone, Thermo Electron Corp., Bellafonte, PA). The mobile phase consisted of 0.005% pentafluoropropanoic acid, 0.1% acetic acid, and 5% acetonitrile in water (solvent A) and 0.005% pentafluoropropanoic acid,

0.1% acetic acid, and 50% water in acetonitrile (solvent B). A linear gradient from 0% to 100% B in 20 min, followed by 10 min of isocratic elution was used. The flow rate was 0.40 mL/min and the column temperature was set to 40 °C. The electrospray interface was used with the following spray chamber settings: nebulizer pressure, 40 psig; capillary voltage, 3500 V; drying gas temperature, 350 °C; and drying gas flow rate, 10 L/min. Fragmentor voltage was set to 120 V. The mass spectrometer was used in scan mode detecting molecular ions with  $m/z$  values ranging from 50 to 2000.

DGEBA, DGEBF, PGE and **3** (Figure 1) were purchased from Sigma-Aldrich; DGEBF was obtained as a mixture of three isomers. Resorcinol diglycidyl ether (**3**) was 91% pure when purchased and was further purified by column chromatography (hexane: ethyl acetate 7:3) prior to evaluation. Acetone was purchased from Merck (Darmstadt, Germany) and olive oil from Apoteket AB (Goteborg, Sweden). The peptide AcPHCKRM was purchased from Peptide 2.0 Inc. (Chantilly, USA). Unless otherwise indicated, reagents were obtained from commercial suppliers and used without further purification. TLC was performed using silica gel coated aluminium plates. The purity of both synthesized and purchased test compounds was >98% (GC/MS) before evaluation in biochemical and biological assays.

## Chemical Synthesis

**2-((4-(4-(2-Methoxyethoxy)benzyl)phenoxy)methyl)oxirane (1) (Scheme 1A).**

**4-(2-Methoxyethoxy)benzoic acid (7):** A solution of KOH (9.40 g, 166 mmol) in anhydrous ethanol (85 mL) was added to a solution of 4-hydroxybenzoic acid (10.0 g, 72.4 mmol) in

anhydrous ethanol (115 mL) under nitrogen atmosphere. The white suspension was stirred with dropwise addition of 2-bromo-1-methoxyethane (8.84 mL, 13.0 g, 94.1 mmol). The reaction mixture was refluxed for 24 h under nitrogen atmosphere. A solution of KOH (8.25 g, 63.5 mmol) in anhydrous ethanol (75 mL) was added and the reaction mixture was refluxed for a further 2 h. After evaporation of the solvent, the residue was dissolved in water (500 mL) and acidified to pH=1 with diluted HCl (1 N). The precipitate was isolated by filtration, dried and purified by recrystallization in ethanol, to give **7** (12.14 g, 86%) as white crystals. <sup>1</sup>H NMR δ 3.46 (3H, s), 3.78 (2H, t, J=9.26 Hz), 4.19 (2H, t, J=9.52 Hz), 6.97 (2H, d, J=8.79 Hz) 8.05 (2H, d, J=8.79 Hz). <sup>13</sup>C NMR δ 59.3, 67.4, 70.7, 114.3, 121.8, 132.3, 163.2, 171.3. ESI-MS (70 eV), *m/z* (%) 196 (66) (M<sup>+</sup>), 179 (2), 164 (1), 151 (6), 138 (13), 121 (33), 105 (6), 93 (5), 76 (6), 59 (100).

***N*-Methoxy-4-(2-methoxyethoxy)-*N*-methylbenzamide (8):** *N*-Methylmorpholine (2.8 mL, 2.49 g, 25.4 mmol) was added dropwise to a solution of **7** (4.64 g, 23.7 mmol) in freshly distilled THF (35 mL) under nitrogen atmosphere. Isopropyl chloroformate (27.8 mL, 26 mmol) was added dropwise at -20 °C and the reaction mixture was stirred for 30 min at -20 °C. Freshly distilled triethylamine (5.2 mL, 37.3 mmol) was added to a suspension of *N,O*-dimethylhydroxylamine hydrochloride (3.63 g, 37.6 mmol), in anhydrous DMF (20 mL). The suspension was added to the reaction mixture at 0 °C. After 3.5 hours at 0 °C, the reaction mixture was quenched with water (300 mL). The aqueous phase was extracted with ethyl acetate (3 × 100 mL). The combined organic layers were washed with HCl (1 N) (125 mL), saturated aqueous NaHCO<sub>3</sub> (125 mL) and brine (125 mL) and were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo*. The yellow oil was purified by column chromatography (CH<sub>3</sub>OH:



CH<sub>2</sub>Cl<sub>2</sub>, 0.5: 99.5) (*R<sub>f</sub>* = 0.32), to give **8** (3.22 g, 57%) as a colorless oil. <sup>1</sup>H NMR δ 3.35 (3H, s), 3.46 (3H, s), 3.55 (3H, s), 3.74-3.80 (2H, m), 4.13-4.18 (2H, m), 6.93 (2H, d, *J*=8.79 Hz), 7.71 (2H, d, *J*=8.79 Hz). <sup>13</sup>C NMR δ 33.9, 59.2, 60.8, 67.3, 70.8, 113.8, 126.2, 130.5, 160.7, 169.3. ESI-MS (70 eV), *m/z* (%) 239 (2) (M<sup>+</sup>), 209 (1), 179 (100), 135 (4), 121 (82), 104 (7), 93 (9), 76 (6), 59 (20).

**(4-*tert*-Butyldimethylsilyloxyphenyl)(4-(2-methoxyethoxy)phenyl)methanone (9):** A solution of (4-bromophenoxy)-*tert*-butyldimethylsilane (97%) (5.02 mL, 5.92 g, 20 mmol) in freshly distilled THF (20.0 mL) under nitrogen atmosphere was stirred for 5 min and then cooled to -78 °C. After stirring for a further 10 min, *N*-butyllithium (2.5 M in hexanes) (8.4 mL, 22 mmol, 2 equiv) was added dropwise to the solution. The temperature was kept at -78 °C for 20 min, and then kept at -40 °C for 40 min. A solution of **8** (2.39 g, 10 mmol) in freshly distilled THF (15 mL) was added dropwise to the *in situ* generated aryllithium under nitrogen atmosphere at -78 °C. The temperature was kept at -78 °C for 1 h, then kept at -30 °C for 40 min and then kept at room temperature for 1.5 h. Aqueous HCl (1 N) (13 mL) was added and after 30 min, the biphasic mixture was partitioned between chloroform (30 mL) and water (50 mL). The organic phase was washed with saturated, aqueous NaHCO<sub>3</sub> (3 × 100 mL). The combined organic layers were dried with anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The yellow oil was purified by column chromatography (hexane: ethyl acetate, 5:1) (*R<sub>f</sub>* = 0.22), to give **9** (3.66 g, 95%) as a colorless oil. <sup>1</sup>H NMR δ 0.25 (6H, s), 1.00 (9H, s), 3.47 (3H, s), 3.77-3.81 (2H, m), 4.18-4.22 (2H, m), 6.89 (2H, d, *J*=8.79 Hz), 6.98 (2H, d, *J*=8.79 Hz), 7.72 (2H, d, *J*=8.79 Hz), 7.78 (2H, d, *J*=8.79 Hz). <sup>13</sup>C NMR δ -4.4, 18.2, 25.6, 59.3, 67.4, 70.8, 114.0, 119.6, 130.9, 131.3, 132.1,

132.2, 159.5, 162.0, 194.6. ESI-MS (70 eV),  $m/z$  (%) 386 (100) ( $M^+$ ), 329 (99), 270 (21), 209 (91), 179 (68), 121 (69).

**1-tert-Butyldimethylsilyloxy-4-(4-(2-methoxyethoxy)benzyl)benzene (10):** Solid zinc iodide (3.89 g, 12.2 mmol) and sodium cyanoborohydride (3.82 g, 60.9 mmol) were added to a solution of **9** (3.14 g, 8.12 mmol) in 1,2-dichloroethane (40 mL) at room temperature. After 20 h, the reaction mixture was filtered through Celite and the Celite was washed with  $\text{CH}_2\text{Cl}_2$ . The filtrate was concentrated *in vacuo*, to give **10** (3.01 g, 98%) as a colorless oil.  $^1\text{H}$  NMR  $\delta$  0.17 (6H, s), 0.97 (9H, s), 3.45 (3H, s), 3.72-3.77 (2H, m), 3.85 (2H, s), 4.08-4.12 (2H, m), 6.74 (2H, d,  $J=8.42$  Hz), 6.85 (2H, d,  $J=8.42$  Hz), 7.00 (2H, d,  $J=8.42$  Hz), 7.07 (2H, d,  $J=8.42$  Hz).  $^{13}\text{C}$  NMR  $\delta$  -4.5, 18.2, 25.7, 40.2, 59.2, 67.2, 71.0, 114.5, 119.9, 129.6, 129.7, 133.9, 134.2, 153.8, 157.0. ESI-MS (70 eV),  $m/z$  (%) 372 (87) ( $M^+$ ), 316 (19), 287 (6), 256 (13), 195 (100), 165 (18), 150 (8), 133 (29), 107 (19), 89 (25), 73 (6), 59 (14).

**4-(4-(2-Methoxyethoxy)benzyl)phenol (11):** A solution of tetra-*n*-butylammonium fluoride in THF (1 M) (8.5 mL, 8.5 mmol) was added to a solution of **10** (3.0 g, 8.08 mmol) in freshly distilled THF (33 mL) under nitrogen atmosphere. After 3 h at room temperature, the reaction mixture was diluted with ethyl acetate (75 mL) and washed with a saturated solution of  $\text{NH}_4\text{Cl}$  (140 mL). The aqueous layer was extracted with ethyl acetate ( $3 \times 75$  mL). The combined organic layers were washed with water ( $2 \times 175$  mL), brine (175 mL), dried over anhydrous  $\text{Na}_2\text{SO}_4$  and concentrated *in vacuo*. The white crystals were purified by column chromatography ( $\text{CH}_3\text{OH}$ : toluene, 6:94) ( $R_f = 0.55$ ), to give **11** (1.87 g, 90%) as white crystals.  $^1\text{H}$  NMR  $\delta$  3.45 (3H, s), 3.75 (2H, t,  $J=9.52$  Hz), 3.84 (2H, s), 4.10 (2H, t,  $J=9.52$  Hz), 6.74 (2H, d,  $J=8.06$  Hz),

6.84 (2H, d, J=8.42 Hz), 7.02 (2H, d, J=8.42 Hz), 7.06 (2H, d, J=8.42 Hz).  $^{13}\text{C}$  NMR  $\delta$  40.1, 59.2, 67.2, 71.1, 114.6, 115.2, 129.7, 129.9, 133.7, 133.9, 153.8, 157.0. ESI-MS (70 eV),  $m/z$  (%) 258 (100) (M<sup>+</sup>), 199 (40), 183 (16), 165 (9), 152 (11), 107 (51), 94 (6), 77 (6), 59 (47).

**2-((4-(4-(2-Methoxyethoxy)benzyl)phenoxy)methyl)oxirane (1):** Epichlorohydrin (1.14 mL, 14.4 mmol) was added dropwise to a reaction mixture of **11** (0.92 g, 3.74 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (2.34 g, 7.12 mmol) in freshly distilled acetonitrile (60 mL). The reaction mixture was refluxed for 3 h under nitrogen atmosphere. Water (100 mL) and ethyl acetate (50 mL) were added at room temperature and the separated water phase was extracted with ethyl acetate (3 × 60 mL). The combined organic layers were washed with brine (200 mL), water (200 mL) and brine (200 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The mixture was purified by column chromatography (CH<sub>3</sub>OH: CH<sub>2</sub>Cl<sub>2</sub>, 1:99) (R<sub>f</sub> = 0.47) to give **1** (0.91 g, 78%) as white crystals.  $^1\text{H}$  NMR  $\delta$  2.4 (1H, dd, J=4.76, 2.56 Hz), 2.89 (1H, t, J=8.79 Hz), 3.31-3.37 (1H, m), 3.44 (3H, s), 3.71-3.76 (2H, m), 3.86 (2H, s), 3.90-3.97 (1H, m), 4.06-4.12 (3H, m), 4.15-4.21 (1H, m), 6.81-6.88 (4H, m), 7.04-7.11 (4H, m).  $^{13}\text{C}$  NMR  $\delta$  40.1, 44.7, 50.1, 59.2, 67.2, 68.8, 71.0, 114.6, 129.7, 129.8, 133.8, 134.3, 156.8, 157.1. LC/MS (API-ES, 120 V)  $m/z$  (%): 337.1 [M+Na] (40), 332.1 [M+H<sub>2</sub>O] (100), 315.1 [M+H] (3), 165.1 (58), 163.1 (51.5), 107.1 (9), 59.1 (8.3).

**Bis(4-(2-methoxyethoxy)phenyl)methane (2) (Scheme 1B).** A solution of potassium hydroxide (0.74 g, 13.2 mmol) in anhydrous ethanol (ca 10 mL) was added to a solution of 4,4'-methylenediphenol (1.10 g, 5.5 mmol) in anhydrous ethanol. 2-Bromo-1-methoxyethane (1.34 mL, 14.3 mmol) was added and the reaction was refluxed for 4 h. The solvent was evaporated

and the residue was diluted with ethyl acetate (70 mL) and washed successively with HCl (1 M, 50 mL), NaOH (2 M, 50 mL) and water (50 mL), and then dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated *in vacuo*. The product was purified by column chromatography (CH<sub>3</sub>OH: CH<sub>2</sub>Cl<sub>2</sub>, 0:100, then 0.25:99.75 then 0.5:99.5) to give pure **2** (0.50 g, 29%). <sup>1</sup>H NMR δ 3.43 (6H, s), 3.71-3.74 (4H, m), 3.85 (2H, s), 4.07-4.09 (4H, m), 6.83 (4H, d, J=6.6 Hz), 7.96 (4H, d, J=8.8 Hz). <sup>13</sup>C NMR δ 40.2, 59.3, 67.3, 71.2, 114.6, 123.8, 134.0, 157.2. ESI-MS (70 eV) *m/z* (%) 316 (100) (M<sup>+</sup>), 257 (18), 207 (34), 199 (19), 107 (19), 59 (66).

**(2-Methoxyethoxy)benzene (4) (Scheme 1C)**. A solution of KOH (12 mmol) in dry ethanol (5 mL) was added to phenol (0.49 g, 5.3 mmol) in dry ethanol (5 mL). 2-Bromo-1-methoxyethane (0.65 mL, 6.9 mmol) was added and the reaction was refluxed for 4 h. The solvent was evaporated and the residue was dissolved in ethyl acetate (70 mL). The organic layer was washed successively with HCl (1M, 50 mL), NaOH (2 M, 50 mL) and water (50 mL) and was dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated *in vacuo*. The product was purified by column chromatography (hexane: ethyl acetate 9:1) to yield **4** (0.22 g, 27%). <sup>1</sup>H NMR δ 3.45 (3H, s), 3.73 - 3.76 (2H, m), 4.11-4.13 (2H, m), 6.93-6.95 (3H, m), 7.26-7.30 (2H, m). <sup>13</sup>C NMR δ 59.3, 67.2, 71.1, 114.7, 120.9, 129.5, 158.8. ESI-MS (70eV) *m/z* (%) 152 (61) (M<sup>+</sup>), 107 (11), 94 (55), 77 (51), 65 (17), 59 (100), 51 (17).

**1,5-Bis(2,3-epoxypropoxy)pentane (5) (Scheme 2)**.

**1,5-Bis(allyloxy)pentane (12)**: Sodium hydride (0.62 g, 26 mmol, 5.2 equiv) was washed with hexane (20 mL × 2), suspended in anhydrous THF (40 mL) and cooled to 0 °C. Pentane-1,5-diol (0.52 g, 5.0 mmol) was dissolved in anhydrous THF (20 mL) and added to the suspension. The

mixture was stirred at 0 °C for 10 min. Allyl bromide (1.12 mL, 13 mmol, 2.6 equiv) was added and the mixture was stirred at room temperature for 90 min before refluxing overnight. The reaction was continued until TLC indicated disappearance of the starting material. The mixture was cooled to 0 °C and saturated aqueous NH<sub>4</sub>Cl (70 mL) was added slowly to quench the reaction. The aqueous layer was extracted with ethyl acetate (70 mL × 3). The combined organic fractions were washed with brine (150 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and reduced *in vacuo*. Compound **12** was obtained as a colorless oil (0.70 g, 76%). <sup>1</sup>H NMR δ 1.35-1.41 (m, 2H), 1.53-1.59 (m, 4H), 3.36 (t, 4H, J=6.6 Hz), 3.90-3.91 (m, 4H), 5.09-5.12 (m, 2H), 5.18-5.25 (m, 2H), 5.81-5.90 (m, 2H). <sup>13</sup>C NMR δ 22.9 (CH<sub>2</sub>), 29.6 (CH<sub>2</sub>), 70.3 (CH<sub>2</sub>), 71.8 (CH<sub>2</sub>), 116.7 (CH<sub>2</sub>), 135.1 (CH). EI-MS (70 eV), *m/z* (%) 143 (12), 127 (13), 97 (7), 85 (100), 69 (47), 57 (26).

**1,5-Bis(2,3-epoxypropoxy)pentane (5):** A solution of **12** (0.7 g, 3.8 mmol) in chloroform (20 mL) was cooled to 0 °C. 3-Chloroperbenzoic acid (≤77%, 1.96 g, 11.4 mmol) was added and the mixture was stirred at 0 °C for 2 h. The mixture was then stirred at room temperature with further additions of 3-chloroperbenzoic acid until the reaction was complete according to TLC. Aqueous NaOH (40 mL) (10% w/v) was added and the aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (40 mL). The organic phase was washed with brine (40 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and reduced *in vacuo*. Compound **5** was isolated as a colorless oil (0.50 g, 61%) after column chromatography (hexane: ethyl acetate, 8:2). <sup>1</sup>H NMR δ 1.35-1.39 (m, 2H), 1.52-1.60 (m, 4H), 2.54-2.56 (m, 2H), 2.73-2.75 (m, 2H), 3.07-3.11 (m, 2H), 3.29-3.33 (m, 2H), 3.39-3.49 (m, 4H), 3.67 (dd, 2H, J=2.96, 11.72 Hz). <sup>13</sup>C NMR δ 22.7, 29.5, 44.3, 50.9, 71.5. EI-MS (70 eV), *m/z* (%) 143 (4), 113 (24), 100 (5), 85 (82), 69 (78), 57 (100).

**1,7-Bis(2,3-Epoxypropoxy)heptane (6) (Scheme 2).**

**1,7-Bis(allyloxy)heptane (13):** Sodium hydride (0.62 g, 26 mmol, 5.2 equiv) was washed with hexane (2 × 20 mL), suspended in anhydrous THF (40 mL) and cooled to 0 °C. Heptane-1,7-diol (0.66 g, 5.0 mmol) was dissolved in anhydrous THF (20 mL) and added to the suspension. The mixture was stirred at 0 °C for 10 min. Allyl bromide (1.56 g, 13 mmol, 2.6 equiv) was added and the mixture was stirred at room temperature for 90 min before refluxing overnight. The reaction was continued until TLC indicated disappearance of the starting material. The mixture was cooled to 0 °C and saturated aqueous NH<sub>4</sub>Cl (70 mL) was added slowly to quench the reaction. The aqueous layer was extracted with ethyl acetate (3 × 70 mL). The combined organic fractions were washed with brine (150 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and reduced *in vacuo*. Compound **13** was obtained as a colorless oil (0.81 g, 76%). <sup>1</sup>H NMR δ 1.24-1.36 (m, 6H), 1.51-1.57 (m, 4H), 3.38 (t, 4H, J=6.6 Hz), 3.91-3.93 (m, 4H), 5.11-5.14 (m, 2H), 5.21-5.26 (m, 2H), 5.83-5.92 (m, 2H). <sup>13</sup>C NMR δ 26.2, 29.4, 29.8, 70.5, 71.9, 116.8, 135.1. EI-MS (70 eV), *m/z* (%) 171 (19), 125 (5), 113 (21), 97 (65), 81 (17), 71 (42), 55 (100).

**1,7-Bis(2,3-epoxypropoxy)heptane (6):** A solution of **13** (0.8 g, 3.8 mmol) in chloroform (20 mL) was cooled to 0 °C. 3-Chloroperbenzoic acid (≤77%, 1.96 g, 11.4 mmol) was added and the mixture was stirred at 0 °C for 2 h. The mixture was then stirred at room temperature with further additions of 3-chloroperbenzoic acid until the reaction was complete according to TLC. Aqueous NaOH (40 mL) (10% w/v) was added and the aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (40 mL). The organic phase was washed with brine (40 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and reduced *in vacuo*. Purification by column chromatography (hexane: ethyl acetate, 8:2) gave **6** as a colorless oil (0.54 g, 58%). <sup>1</sup>H NMR δ 1.29-1.32 (m, 6H), 1.51-1.55 (m, 4H),

2.55-2.57 (m, 2H), 2.74-2.76 (m, 2H), 3.07-3.11 (m, 2H), 3.30-3.35 (m, 2H), 3.38-3.48 (m, 4H), 3.64-3.68 (m, 2H).  $^{13}\text{C}$  NMR  $\delta$  26.1, 29.3, 29.7, 44.4, 50.9, 71.5, 71.7. EI-MS (70 eV),  $m/z$  (%) 141 (1), 127 (3), 113 (12), 95 (74), 87 (10), 81 (16), 75 (12), 69 (48), 55 (100).

**Experimental Animals.** Female CBA/Ca mice, 8 or 9 weeks of age, were purchased from NOVA SCB Charles River, Germany. The mice were housed in “hepa” filtered air flow cages and kept on standard laboratory diet and water ad lib. The local ethics committee in Gothenburg approved the study.

**Skin Sensitization Potency of Epoxy Resins in Mice.** The local lymph node assay (LLNA)<sup>40</sup> was used to assess the sensitization potency. Mice in six groups of three animals in each were treated by topical application on the dorsum of both ears with the test compound (25  $\mu\text{L}$ ) dissolved in acetone:olive oil (AOO) (4:1 v/v) or with the vehicle control. All solutions were freshly prepared for each application. Each compound was tested at five different concentrations. The test concentrations used were as follows: **1** and **2**: 1, 5, 10, 20 and 30% (w/v); Treatments were performed daily for three consecutive days (days 0, 1, and 2). Sham treated control animals received vehicle alone. On day 5, all mice were injected intravenously via the tail vein with [methyl- $^3\text{H}$ ]thymidine (2.0 Ci/mmol, Amersham Biosciences, UK) (20  $\mu\text{Ci}$ ) in phosphate-buffered saline (PBS, containing 137 mM NaCl, 2.7 mM KCl and 10 mM phosphate buffer, pH 7.4) (250  $\mu\text{L}$ ). After 5 h the mice were sacrificed, the draining lymph nodes were excised and pooled for each group, and single cell suspensions of lymph-node cells in PBS were prepared using cell strainers (Falcon, BD labware, 70  $\mu\text{m}$  pore size). Cell suspensions were washed twice with PBS, precipitated with TCA (5%) and left in the refrigerator overnight. The samples were

then centrifuged, resuspended in TCA (5%) (1 mL) and transferred to a scintillation cocktail (10 mL) (EcoLume, INC Radiochemicals, USA). The [methyl-<sup>3</sup>H]thymidine incorporation into DNA was measured by  $\beta$ -scintillation counting on Beckman LS 6000TA Instruments. Results are expressed as mean dpm/lymph node for each experimental group and as stimulation index (SI), i.e., test group/control group ratio. Test materials that at one or more concentrations caused an SI greater than 3 were considered to be positive in the LLNA. EC3 values (the estimated concentration required to induce an SI of 3) were calculated by linear interpolation. The sensitization potency was classified to the following:  $\leq 0.2\%$  w/v, extreme;  $>0.2$  to  $\leq 2\%$  w/v, strong;  $>2\%$  w/v moderate.<sup>26</sup>

**KeratinoSens™ Assay for Sensitization and Cellular Viability.** The KeratinoSens™ assay was performed as previously described in detail.<sup>41, 42</sup> Briefly, cells were grown for 24 h in 96-well plates. The medium was then replaced with medium containing the test chemical and a final level of 1% DMSO. Each compound was tested at 12 binary dilutions in the range from 0.98 to 2000  $\mu$ M. Each test plate contained seven test chemicals, six wells with the solvent control, one well with no cells for background value and five wells with the positive control cinnamic aldehyde in five different concentrations. In each repetition, three parallel replicate plates were run with this same set-up and a fourth parallel plate was prepared for cytotoxicity determination. Cells were incubated for 48 h with the test agents, after which luciferase activity and cytotoxicity (with the MTT assay<sup>43</sup>) were determined. This full procedure was repeated three times for each chemical.

**Thiol Reactivity with *N*-Acetyl Cysteine Methyl Ester.** Compound DGEBF or **2** (1 mmol) was stirred with *N*-ACME (1 mmol) at 37 °C overnight in a 2:1 solution of DMSO: ammonium



acetate buffer (100 mM pH 7.4). The reaction was analyzed using LC/MS (see parameters above).

**Reactivity with the Model Peptide Ac-Pro-His-Cys-Lys-Arg-Met-OH (AcPHCKRM).** All solvents were degassed with argon prior to use. The hexapeptide AcPHCKRM (Peptide 2.0) was >99% pure as determined by HPLC (UV<sub>220nm</sub>). Bz-His-OMe (Bachem) was used as an internal standard. Solutions of PGE, DGEBF or compound **1** in DMSO (40 mM, 100  $\mu$ L) together with potassium phosphate buffer (100 mM, pH 7.4), (200  $\mu$ L) were added to a vial purged with argon containing AcPHCKRM in DMSO (4 mM, 100  $\mu$ L). Accordingly, final concentrations of compound and model peptide in the reaction mixture were 10 mM and 1 mM, respectively. The reaction mixture was kept under argon at room temperature and was monitored with UV<sub>210nm</sub>/ESI-MS. As the HPLC run time was 40 minutes, data was combined from two experiments. In the first experiment, samples were collected at t=0, 40, 80, 120 mins and in the second experiment at t=20, 60, 100 mins. In a modified experiment using two haptens simultaneously, 50  $\mu$ L of both PGE and DGEBF (50 mM) were added to the solution containing the peptide, with the other experimental details remaining as above.

## Results and Discussion

### Chemical Synthesis

Efficient synthetic strategies were employed to produce the biaromatic compounds **1** and **2** illustrated in Figure 1. A six-step synthesis was used to obtain the non-symmetrical compound **1** (Scheme 1, A). 2-Bromoethyl methyl ether and 4-hydroxybenzoic acid were reacted (*step i*) and the isolated product **7** was subsequently reacted with *N,O*-dimethylhydroxylamine hydrochloride to produce the Weinreb amide **8** (*step ii*). The amide **8** was then coupled to 1-bromo-4-*tert*-butyldimethylsilyloxybenzene to yield the biaromatic product **9** (*step iii*). After reduction (*step iv*) and deprotection (*step v*), the final product **1** was obtained by alkylation of the free phenolic group with epichlorohydrin in 32% overall yield (*step vi*).

Synthesis of **2** was achieved in a one-step reaction in a moderate yield of 29% (Scheme 1, B). The monoalkylated compound was isolated as a by-product. Synthesis of **4** was achieved from phenol in 27% yield (Scheme 1, C). Both **1** and **2** were synthesized as tools to investigate the effects of the epoxide functionalities of DGEBF and cannot be used as epoxy resin monomer replacements for polymer synthesis. Compound **1**, with one terminal epoxide group and one terminal methoxy group, can only form dimers and compound **2**, with two terminal methoxy groups, cannot react with itself or other epoxide-containing polymeric precursors.

Linear compounds **5** and **6** were synthesized to investigate the influence of sterical shape on the cytotoxicity of **2**. A 5-carbon chain was chosen as it most closely imitates the inter-atomic distance between the two epoxide groups of DGEBF (14.47 Å for **5** compared to 14.75 Å for DGEBF) (Table S1, Supporting Information). The compound with a 7-carbon chain, **6**, was also synthesized to obtain supplemental information. These linear epoxides have greater flexibility

than DGEBF and hence there are a large number of conformations possible. Compounds **5** and **6** were obtained in two steps from 1,5-pentanediol and 1,7-heptanediol, respectively (Scheme 2). The diallylic intermediates **12** and **13** were each obtained in 76% yield by refluxing the starting material with allyl bromide and sodium hydride in an inert atmosphere (*step i*). The final compounds are formed by epoxidation using *meta*-chloroperbenzoic acid (*mCPBA*) (*step ii*). The corresponding monoepoxides were observed as by-products.

### **Skin Sensitization Potency Studies: LLNA and KeratinoSens™**

Biaromatic compounds **1** and **2** were assessed *in vivo* for skin sensitization potency using the murine local lymph node assay (LLNA).<sup>44</sup> LLNA results are expressed as EC3 values, which is the estimated concentration of a substance required to induce a 3-fold increase in sensitization compared to a control.

The EC3 value obtained for compound **1** was 74 mM (2.3% w/v) (Figure 2 and Table S2, Supporting Information). The stimulation index decreases for the 20% and 30% concentrations compared to that of 10%. This effect was seen in repeated LLNA experiments with **1**. Previously reported LLNA EC3 values for DGEBA, DGEBF and PGE indicate that they are equipotent at 36 mM, 36 mM and 31 mM (1.2, 1.1 and 0.46% w/v), respectively.<sup>21, 23</sup> According to suggested regulatory classifications, all three are sensitizers of strong potency.<sup>26</sup> The monoepoxide **1** has reduced sensitization potency in the LLNA compared to DGEBF and is classified as a moderate sensitizer according to its EC3 value of 2.3% w/v. Compound **2** does not contain any reactive epoxide groups and was non-sensitizing in concentrations up to 30% w/v (950 mM) (Figure 2). This compound did not show any dose-response relationship at the concentrations tested. The only structural difference between compound **2** and DGEBF is lack of epoxide groups in

compound **2** (Figure 1). Thus, our results show that epoxide groups are directly responsible for the skin sensitization effects of DGEBF in the LLNA.

To obtain further information, all compounds were evaluated in the *in vitro* KeratinoSens™ assay.<sup>42</sup> In this assay, compounds are rated as skin sensitizers if they induce the luciferase gene by greater than 1.5-fold at non-cytotoxic concentrations. Previously published values predict DGEBA, DGEBF and PGE to be sensitizers, giving a 1.5-fold induction at low micromolar concentrations.<sup>21</sup> The KeratinoSens™ assay predicts **1** as a sensitizer (Table 1). Partial cytotoxicity of a chemical may lead to a luciferase induction in dying cells and therefore chemicals are rated as sensitizers only when luciferase induction starts at non-cytotoxic concentrations.<sup>42</sup> For compound **2**, >1.5-fold induction can only be noted at cytotoxic concentrations, and thus **2** is classified as a non-sensitizer by this method. The KeratinoSens™ results for both biaromatic compounds are in agreement with the LLNA data.

Linear diepoxides **5** and **6** were predicted to be sensitizers by the KeratinoSens™, with maximum induction values ( $I_{\max}$ ) of 209 and 43, respectively. These compounds were not evaluated in the LLNA due to ethical considerations, but it is not surprising that they are predicted to be sensitizers as both contain two terminal, reactive epoxide groups. A structurally related compound, 1,6-hexanediol diglycidyl ether, has a reported EC3 value of 1.9% w/v, classifying it as a strong sensitizer.<sup>25</sup>

Two monoaromatic compounds were also assessed in the KeratinoSens™. Of these, epoxide containing compound **3** was predicted to be a skin sensitizer ( $I_{\max} = 120$ ), while non-epoxide **4** was predicted as a non-sensitizer ( $I_{\max} = 1.2$ ). Neither **3** nor **4** were assessed in the LLNA, again due to ethical considerations. On the basis of the results above, **3** would be expected to be a

strong sensitizer in the LLNA due to the presence of the epoxide groups, analogous to PGE.<sup>21</sup> Conversely, **4** is structurally related to **2** and would not be expected to act as a sensitizer due to the lack of epoxide groups.

### **Reactivity Towards the Thiol Residue in Cysteine and in a Model Peptide**

Small molecules react with reactive amino acid residues of proteins in the skin to form immunogenic complexes, which are then recognized by the immune systems as ‘non-self’ and provoke an immune response. Reactive amino acid residues that play a role in skin sensitization include cysteines (thiol), lysines (amino) and to a lesser extent arginines, histidines, methionines and tyrosines.<sup>45</sup> It has been previously demonstrated that the epoxy reactive diluent PGE reacts selectively with the thiol residue of a model peptide.<sup>23</sup> We firstly investigated the reactivity of DGEBF and compound **2** with a cysteine derivative (*N*-acetyl-L-cysteine methyl ester, *N*-ACME). LC/MS analysis of the DGEBF/*N*-ACME reaction after 24 hours showed the formation of both monoadducts and diadducts, indicating that DGEBF is thiol-reactive. There was no reaction between **2** and *N*-ACME, indicating that **2** is non-reactive towards free thiols. Further reactions of compound **2** were not investigated, but we would predict that the methoxy group is also unreactive towards other amino acids implicated in skin sensitization. The results of the reactivity experiment with *N*-ACME indicate that the epoxide groups of DGEBF are necessary for thiol binding.

Having confirmed that the epoxide group was required to react with cysteine, the reactivities of DGEBF, compound **1** and PGE towards a model peptide were investigated. The three compounds were chosen to investigate the relative reactivity of a biaromatic diepoxide, a biaromatic monoepoxide and a monoaromatic monoepoxide. The hexapeptide chosen,

AcPHCKRM, contains both cysteine and lysine amino acid residues. All compounds were stable in the conditions of the assay.

The percentage peptide depletion was calculated for the three compounds (Figure 3 and Table S3, Supporting Information). DGEBF and compound **1** are structurally similar with the only difference being one terminal epoxide group in compound **1** compared to two in DGEBF. The depletion is slightly faster for DGEBF compared to compound **1**. The reactivity of DGEBF is similar to that of PGE at 120 min (7.1% of free peptide remaining). To confirm the latter result, a second experiment was performed in which both DGEBF and PGE were allowed to react simultaneously with the peptide. 51.5% of DGEBF-peptide adduct was formed compared to 48.5% of PGE-peptide adduct, confirming that DGEBF and PGE react with this peptide in essentially equimolar amounts.

Interestingly, the diepoxide DGEBF was observed to only form mono-adducts and did not cross-link peptides. This result is unexpected, as the ability to cross-link proteins could be a potential reason for the enhanced response of DGEBF in the LLNA over compound **1**. A potential reason for the lack of cross-linking observed is that an excess of DGEBF compared to peptide is used in this assay (10:1 ratio).

A cysteine adduct was observed for all three compounds, indicated by haptenated y5 and y4 fragments in the MS, but non-haptenated y3 and y2 fragments (Table 2 and Figure S1, Supporting Information). No adducts with lysine or other residues were observed. This is in agreement with previous results showing that the epoxides bind to cysteine residues in peptides.<sup>23, 46</sup>

In summary, the reactivity experiments indicate that both the biaromatic and monoaromatic epoxide-containing compounds DGEBF, compound **1** and PGE can bind to thiol residues, while compound **2** cannot. This explains the lack of sensitizing effects shown by **2** in the LLNA and KeratinoSens™.

### Cytotoxicity

All compounds were screened for *in vitro* cytotoxicity in the KeratinoSens™ cell line using the MTT assay (Table 1).<sup>43</sup> The IC<sub>50</sub> values for the four biaromatic compounds DGEBA, DGEBF, **1** and **2** fall in the low micromolar range between 22 and 69 μM. Linear diepoxide **6** (IC<sub>50</sub>=169 μM) was over 7-fold less cytotoxic compared to both DGEBA and DGEBF. Compound **5** was the least cytotoxic of the epoxide-containing compounds (IC<sub>50</sub>=284 μM). Cytotoxicity of monoaromatic diepoxide **3** (IC<sub>50</sub>=38 μM) was in the same range as that of DGEBF, **1** and **2**. The monoepoxide PGE has an IC<sub>50</sub> value of 182 μM while compound **4** did not reduce viability by 50% or more at concentrations up to 2,000 μM.

DGEBA is known both as an alkylating agent and as a PPARγ antagonist.<sup>32, 35</sup> It is unknown at present if DGEBA covalently modifies the PPARγ receptor.<sup>47</sup> It is anticipated that DGEBF causes cell death by similar mechanisms to DGEBA due to their structural similarity. The relative loss in cytotoxic potency of the two linear epoxides **5** and **6** compared to DGEBF may indicate that terminal epoxides are not the only structural feature required for potent cytotoxicity in this cell line. DGEBF and compound **2** have equipotent cytotoxicity indicating that a structurally similar compound without reactive epoxides may have a similar cytotoxicity in monolayer cell culture. This is in contrast to the results observed for skin sensitization, where the presence of epoxide groups was essential for sensitizing effects.

The nature of the alkyl chain has a large impact on *in vitro* cytotoxicity when comparing monoaromatic compounds PGE and **4**. PGE is known to cause cell death *via* reactive mechanisms, including DNA-damage,<sup>48</sup> activation of heat shock response and antioxidant response.<sup>49</sup> PGE forms DNA-adducts via its epoxide group, causing cytotoxicity and apoptosis. Compound **4** lacks the epoxide group, and therefore cannot bind to cellular thiols or cause cell death by reactive mechanisms. Our results indicate that a reactive group is necessary for the cytotoxic effect of the monoaromatic epoxides PGE and **3** in the cell line used in our study.

## **Conclusion**

The aim of our work was to investigate the skin sensitizing potency and *in vitro* cytotoxicity of analogues of the epoxy resin monomer DGEBF. These results have implications for the development of new ERS components. We have shown that the skin sensitization potency of the epoxy resin monomer DGEBF is due to its terminal epoxide groups. Based on our findings, newly reported epoxy resin monomers<sup>50, 51</sup> would be expected to cause skin sensitization and allergic contact dermatitis as they contain terminal epoxide groups. Our results also imply that there is a need for further research to try to decrease the reactivity of the epoxide groups to a level where the contact sensitization is reduced but the polymerization capacity is still intact. We have shown that the cytotoxicity of DGEBF in monolayer cell culture may not only be dependent on the epoxide groups. Depending on the structure of alternative epoxy resins,<sup>50-52</sup> reduced cytotoxicity might be obtained. It is important to keep the potential adverse health effects of epoxy resins in mind when developing a new system and to assess these effects at an early stage in the development process.



**Supporting Information:** Inter-atomic distances for linear epoxides and DGEBF, complete LLNA information, results of peptide depletion assays and the structure and fragments of peptide AcPHCKRM are available free of charge via the internet at <http://pubs.acs.org>.

**Funding Support:** Financial support for this project was obtained from Swedish Council for Working Life and Social Research and from AFA Försäkring. The work was performed within the Centre for Skin Research at the University of Gothenburg.

**Acknowledgements:** We would like to acknowledge Daniel Westerberg and Karla Bolbaran for assistance with synthesis, Ida B. Niklasson, Suzanne Exing and Anders Eliasson for assistance with the LLNA experiments and Tina Haupt for KeratinoSens<sup>™</sup> testing.

## References

- (1) Global Industry Analysts, Inc. (2011) Epoxy Resins. A Global Strategic Business Report, pp 1-605, San Jose.
- (2) Dickel, H., Kuss, O., Schmidt, A. and Diepgen, T. L. (2002) Occupational relevance of positive standard patch-test results in employed persons with an initial report of an occupational skin disease. *Int. Arch. Occup. Environ. Health* 75, 423-434.
- (3) Condé-Salazar, L., Guimaraens, D., Villegas, C., Rumero, A. and Gonzalez, M. A. (1995) Occupational allergic contact dermatitis in construction workers. *Contact Dermatitis* 33, 226-230.
- (4) Canelas, M. M., Gonçalo, M. and Figueiredo, A. (2010) Contact allergy to epoxy resins— a 10-year study. *Contact Dermatitis* 62, 55-55.
- (5) Pontén, A., Carstensen, O., Rasmussen, K., Gruvberger, B., Isaksson, M. and Bruze, M. (2004) Epoxy-based production of wind turbine rotor blades: occupational dermatoses. *Contact Dermatitis* 50, 329-338.
- (6) Berglind, I. A., Lind, M.-L. and Lidén, C. (2012) Epoxy pipe relining—an emerging contact allergy risk for workers. *Contact Dermatitis* 67, 59-65.
- (7) Majasuo, S., Liippo, J. and Lammintausta, K. (2012) Non-occupational contact sensitization to epoxy resin of bisphenol A among general dermatology patients. *Contact Dermatitis* 66, 148-153.
- (8) Amado, A. and Taylor, J. S. (2008) Contact allergy to epoxy resins. *Contact Dermatitis* 58, 186-187.

- (9) Andersen, K., White, I. R. and Goossens, A. (2001) Allergens from the Standard Series, In *Textbook of Contact Dermatitis* (Rycroft, R. J. G., Menne, T. and Frosch, P. J., Eds.), Springer-Verlag, Berlin.
- (10) Gamer, A. O., Nies, E. and Vohr, H. W. (2008) Local lymph node assay (LLNA): Comparison of different protocols by testing skin-sensitizing epoxy resin system components. *Regul. Toxicol. Pharmacol.* 52, 290-298.
- (11) Kanerva, L., Jolanki, R. and Estlander, T. (1991) Allergic contact dermatitis from non-diglycidyl-ether-of-bisphenol-A epoxy resins. *Contact Dermatitis* 24, 293-300.
- (12) Kanerva, L., Jolanki, R., Estlander, T., Henriks-Eckerman, M.-L., Tuomi, M.-L. and Tarvainen, K. (2000) Airborne occupational allergic contact dermatitis from triglycidyl-*p*-aminophenol and tetraglycidyl-4,4'-methylene dianiline in preimpregnated epoxy products in the aircraft industry. *Dermatology* 201, 29-33.
- (13) Burrows, D., Fregert, S., Campbell, H. and Trulsson, L. (1984) Contact dermatitis from the epoxy resins tetraglycidyl-4,4'-methylene dianiline and *o*-diglycidyl phthalate in composite material. *Contact Dermatitis* 11, 80-82.
- (14) Geier, J., Oestmann, E., Lessmann, H. and Fuchs, T. (2001) Contact allergy to terephthalic acid diglycidylester in a powder coating. *Contact Dermatitis* 44, 35-36.
- (15) Nishioka, K., Ogasawara, M. and Asagami, C. (1988) Occupational contact allergy to triglycidyl isocyanurate (TGIC, Tepic®). *Contact Dermatitis* 19, 379-380.
- (16) Jensen, C. and Andersen, K. (2003) Two cases of occupational allergic contact dermatitis from a cycloaliphatic epoxy resin in a neat oil: Case Report. *Environ. Health* 2, 3.
- (17) Maibach, H. I. and Mathias, C. T. (2001) Allergic contact dermatitis from cycloaliphatic epoxide in jet aviation hydraulic fluid. *Contact Dermatitis* 45, 56-56.

- (18) Jolanki, R., Kanerva, L., Estlander, T. and Tarvainen, K. (1994) Concomitant sensitization to triglycidyl isocyanurate, diaminodiphenylmethane and 2-hydroxyethyl methacrylate from silk-screen printing coatings in the manufacture of circuit boards. *Contact Dermatitis* 30, 12-15.
- (19) Pontén, A. and Bruze, M. (2001) Contact allergy to epoxy resin based on diglycidyl ether of bisphenol F. *Contact Dermatitis* 44, 98-100.
- (20) Thorgeirsson, A., Fregert, S. and Ramnäs, O. (1978) Sensitization capacity of epoxy resin oligomers in the guinea pig. *Acta Derm.-Venereol.* 58, 17 - 21.
- (21) Delaine, T., Niklasson, I. B., Emter, R., Luthman, K., Karlberg, A.-T. and Natsch, A. (2011) Structure-activity relationship between the *in vivo* skin sensitizing potency of analogues of phenyl glycidyl ether and the induction of Nrf2-dependent luciferase activity in the KeratinoSens *in vitro* assay. *Chem. Res. Toxicol.* 24, 1312-1318.
- (22) Pontén, A., Zimerson, E., Sörensen, Ö. and Bruze, M. (2002) Sensitizing capacity and cross-reaction pattern of the isomers of diglycidyl ether of bisphenol F in the guinea pig. *Contact Dermatitis* 47, 293-298.
- (23) Niklasson, I. B., Broo, K., Jonsson, C., Luthman, K. and Karlberg, A.-T. (2009) Reduced sensitizing capacity of epoxy resin systems: a structure-activity relationship study. *Chem. Res. Toxicol.* 22, 1787-1794.
- (24) Pontén, A., Zimerson, E. and Bruze, M. (2009) Sensitizing capacity and cross-reactivity of phenyl glycidyl ether studied in the guinea-pig maximization test. *Contact Dermatitis* 60, 79-84.

- (25) Gamer, A. O., Nies, E. and Vohr, H.-W. (2008) Local lymph node assay (LLNA): Comparison of different protocols by testing skin-sensitizing epoxy resin system components. *Regul. Toxicol. Pharmacol.* 52, 290-298.
- (26) Basketter, D. A., Andersen, K. E., Liden, C., Loveren, H. V., Boman, A., Kimber, I., Alanko, K. and Berggren, E. (2005) Evaluation of the skin sensitizing potency of chemicals by using the existing methods and considerations of relevance for elicitation. *Contact Dermatitis* 52, 39-43.
- (27) IARC Monographs Database on Carcinogenic Risks to Humans. <http://monographs.iarc.fr/ENG/Classification/index.php> Accessed: 07 May 2012.
- (28) Deforce, D. L. D., Ryniers, F. P. K., Van den Eeckhout, E. G., Lemiere, F. and Esmans, E. L. (1996) Analysis of DNA adducts in DNA hydrolysates by capillary zone electrophoresis and capillary zone electrophoresis-electrospray mass spectrometry. *Anal. Chem.* 68, 3575-3584.
- (29) Brockmann, W. G., Kostoryz, E. L. and Eick, J. D. (2006) Correlation of apoptotic potential of simple oxiranes with cytotoxicity. *Toxicol. In Vitro* 20, 729-735.
- (30) von der Hude, W., Seelbach, A. and Basler, A. (1990) Epoxides: comparison of the induction of SOS repair in *Escherichia coli* PQ37 and the bacterial mutagenicity in the Ames test. *Mutat. Res., Fundam. Mol. Mech. Mutagen.* 231, 205-218.
- (31) Greene, E. J., Friedman, M. A., Sherrod, J. A. and Salerno, A. J. (1979) *In vitro* mutagenicity and cell transformation screening of phenylglycidyl ether. *Mutat. Res., Genet. Toxicol.* 67, 9-19.

- (32) Hemminki, K., Falck, K. and Vainio, H. (1980) Comparison of alkylation rates and mutagenicity of directly acting industrial and laboratory chemicals. *Arch. Toxicol.* 46, 277-285.
- (33) Cabaton, N., Dumont, C., Severin, I., Perdu, E., Zalko, D., Cherkaoui-Malki, M. and Chagnon, M.-C. (2009) Genotoxic and endocrine activities of bis(hydroxyphenyl)methane (bisphenol F) and its derivatives in the HepG2 cell line. *Toxicology* 255, 15-24.
- (34) Ramilo, G., Valverde, I., Lago, J., Vieites, J. and Cabado, A. (2006) Cytotoxic effects of BADGE (bisphenol A diglycidyl ether) and BFDGE (bisphenol F diglycidyl ether) on Caco-2 cells *in vitro*. *Arch. Toxicol.* 80, 748-755.
- (35) Wright, H. M., Clish, C. B., Mikami, T., Hauser, S., Yanagi, K., Hiramatsu, R., Serhan, C. N. and Spiegelman, B. M. (2000) A synthetic antagonist for the peroxisome proliferator-activated receptor gamma inhibits adipocyte differentiation. *J. Biol. Chem.* 275, 1873-1877.
- (36) Schaefer, K. L., Takahashi, H., Morales, V. M., Harris, G., Barton, S., Osawa, E., Nakajima, A. and Saubermann, L. J. (2007) PPAR $\gamma$  inhibitors reduce tubulin protein levels by a PPAR $\gamma$ , PPAR $\delta$  and proteasome-independent mechanism, resulting in cell cycle arrest, apoptosis and reduced metastasis of colorectal carcinoma cells. *Int. J. Cancer* 120, 702-713.
- (37) Bishop-Bailey, D., Hla, T. and Warner, T. D. (2000) Bisphenol A diglycidyl ether (BADGE) is a PPAR $\gamma$  agonist in an ECV304 cell line. *Br. J. Pharmacol.* 131, 651-654.

- (38) Fehlbeg, S., Trautwein, S., Göke, A. and Göke, R. (2002) Bisphenol A diglycidyl ether induces apoptosis in tumour cells independently of peroxisome proliferator-activated receptor-g, in caspase-dependent and -independent manners. *Biochem. J.* 362, 573-578.
- (39) Fehlbeg, S., Gregel, C. M., Göke, A. and Göke, R. (2003) Bisphenol A diglycidyl ether-induced apoptosis involves Bax/Bid-dependent mitochondrial release of apoptosis-inducing factor (AIF), cytochrome c and Smac/DIABLO. *Br. J. Pharmacol.* 139, 495-500.
- (40) Gerberick, G. F., Ryan, C. A., Dearman, R. J. and Kimber, I. (2007) Local lymph node assay (LLNA) for detection of sensitization capacity of chemicals. *Methods* 41, 54-60.
- (41) Natsch, A. and Emter, R. (2008) Skin sensitizers induce antioxidant response element dependent genes: application to the in vitro testing of the sensitization potential of chemicals. *Toxicol. Sci.* 102, 110-119.
- (42) Emter, R., Ellis, G. and Natsch, A. (2010) Performance of a novel keratinocyte-based reporter cell line to screen skin sensitizers *in vitro*. *Toxicol. Appl. Pharmacol.* 245, 281-290.
- (43) Mosmann, T. (1983) Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J. Immunol. Methods* 65, 55-63.
- (44) Basketter, D. A., Evans, P., Fielder, R. J., Gerberick, G. F., Dearman, R. J. and Kimber, I. (2002) Local lymph node assay: validation, conduct and use in practice. *Food Chem. Toxicol.* 40, 593-598.
- (45) Divkovic, M., Pease, C. K., Gerberick, G. F. and Basketter, D. A. (2005) Hapten-protein binding: from theory to practical application in the *in vitro* prediction of skin sensitization. *Contact Dermatitis* 53, 189-200.

- (46) Niklasson, I. B., Delaine, T., Luthman, K. and Karlberg, A.-T. (2011) Impact of a heteroatom in a structure-activity relationship study on analogues of phenyl glycidyl ether (PGE) from epoxy resin systems. *Chem. Res. Toxicol.* 24, 542-548.
- (47) Elbrecht, A., Chen, Y., Adams, A., Berger, J., Griffin, P., Klatt, T., Zhang, B., Menke, J., Zhou, G., Smith, R. G. and Moller, D. E. (1999) L-764406 is a partial agonist of human peroxisome proliferator-activated receptor-gamma. *J. Biol. Chem.* 274, 7913-7922.
- (48) Fekry, M. I., Price, N. E., Zang, H., Huang, C., Harmata, M., Brown, P., Daniels, J. S. and Gates, K. S. (2011) Thiol-activated DNA damage by alpha-bromo-2-cyclopentenone. *Chem. Res. Toxicol.* 24, 217-228.
- (49) Trott, A., West, J. D., Klaic, L., Westerheide, S. D., Silverman, R. B., Morimoto, R. I. and Morano, K. A. (2008) Activation of heat shock and antioxidant responses by the natural product celastrol: transcriptional signatures of a thiol-targeted molecule. *Mol. Biol. Cell* 19, 1104-1112.
- (50) Cheng, J., Chen, J. and Yang, W. T. (2007) Synthesis and characterization of novel multifunctional epoxy resin. *Chin. Chem. Lett.* 18, 469-472.
- (51) Nouailhas, H., Aouf, C., Le Guerneve, C., Caillol, S., Boutevin, B. and Fulcrand, H. (2011) Synthesis and properties of biobased epoxy resins. part 1. Glycidylation of flavonoids by epichlorohydrin. *J. Polym. Sci., Part A: Polym. Chem.* 49, 2261-2270.
- (52) Kishi, H., Fujita, A., Miyazaki, H., Matsuda, S. and Murakami, A. (2006) Synthesis of wood-based epoxy resins and their mechanical and adhesive properties. *J. Appl. Polym. Sci.* 102, 2285-2292.



## Tables

**Table 1. Results from the KeratinoSens™ assay<sup>a</sup>**

Compound	Luciferase Induction				Cytotoxicity
	I <sub>max</sub>	Classification	EC <sub>KS1.5</sub> (μM)	EC <sub>KS4.5</sub> (μM)	IC <sub>50</sub> (μM)
<b>DGEBA</b> <sup>21</sup>	13	Sensitizer	5.2	10	22
<b>DGEBF</b> <sup>21</sup>	5	Sensitizer	6.5	12	23
<b>PGE</b> <sup>21</sup>	56	Sensitizer	16	63	182
<b>1</b>	76	Sensitizer	5.7	19	69
<b>2</b>	2.2 <sup>b</sup>	Non-sensitizer	9.9 <sup>b</sup>	<i>No value</i>	31
<b>3</b>	120	Sensitizer	13	16	38
<b>4</b>	1.2	Non-sensitizer	<i>No value</i>	<i>No value</i>	> 2000
<b>5</b>	209	Sensitizer	60	105	284
<b>6</b>	43	Sensitizer	37	73	169

<sup>a</sup> KeratinoSens™ assay performed using KeratinoSens™ reporter keratinocytes, which contain a stable insertion of a luciferase gene under control of the ARE element of the AKR1C2 gene. I<sub>max</sub> is defined as the average maximal induction of gene activity. A luciferase induction of greater than 1.5-fold at non-cytotoxic concentrations was considered to be significant for skin sensitization. All assays were performed in triplicate on at least 3 separate occasions.

<sup>b</sup> Fold-induction of > 1.5 was only noted at cytotoxic concentrations; therefore this compound is classified as a non-sensitizer according to the KeratinoSens™

**Table 2. Ions and fragments observed in the reactivity experiment with peptide AcPHCKRM<sup>a</sup>**

<b>Hapten</b>	<b>Molecular weight</b>	<b>[M* + H]<sup>+</sup></b> <i>m/z</i>	<b>[M* + 2H]<sup>2+</sup></b> <i>m/z</i>	<b>y<sub>5</sub><sup>*</sup></b> <i>m/z</i>	<b>y<sub>4</sub><sup>*</sup></b> <i>m/z</i>	<b>y<sub>3</sub></b> <i>m/z</i>	<b>y<sub>2</sub></b> <i>m/z</i>
<b>DGEBF</b>	312.4	1125.4	563.3	986.4	849.3	434.2	306.2
<b>1</b>	314.4	1127.4	564.2	988.3	851.4	434.2	306.2
<b>PGE</b>	150.2	963.3	482.2	824.2	687.2	434.2	Nd <sup>b</sup>

<sup>a</sup>M\* or y\* indicates haptened peptide or peptide fragment, respectively; y indicates unhaptened peptide fragment. The mass of the unhaptened peptide (M) is 813.2. The masses of the unhaptened fragments are as follows: **y<sub>2</sub>**: 306.2; **y<sub>3</sub>**: 434.2; **y<sub>4</sub>**: 537.3; **y<sub>5</sub>**: 674.3

<sup>b</sup>Not detected.

## Figure Legends

**Figure 1. Structures of DGEBA (diglycidyl ether of bisphenol A), DGEBF (diglycidyl ether of bisphenol F), PGE (phenyl glycidyl ether) and analogues**

**Figure 2. Local Lymph Node Assay (LLNA) results for DGEBF, 1 and 2.** The EC<sub>3</sub> value (estimated concentration required to induce a stimulation index (SI) of 3) was calculated using linear interpolation. EC<sub>3</sub> values are as follows: DGEBF: 0.036 M (1.1 % w/v);<sup>21</sup> **1**: 0.074 M (2.3 % w/v); **2** > 0.95 M (> 30 % w/v).

**Figure 3. Peptide depletion by DGEBF, 1 and PGE.** The model peptide AcPHCKRM was incubated with the relevant compound (1: 10) in a mixture of 1:1 DMSO: potassium phosphate buffer (100 mM, pH 7.4). The reaction mixture was kept under argon at room temperature and was monitored by UV/ESI-MS.

Figure 1.




Figure 2.





Figure 3.



## Scheme Legends.

### Scheme 1. Synthesis of compounds 1, 2 and 4<sup>a</sup>

\*TBDMS=*tert*-butyldimethylsilyl

<sup>a</sup>Reagents and conditions: A. (i) KOH, EtOH, N<sub>2</sub>, 24 h, reflux; (ii) 4-Methylmorpholine, THF, -20 °C (30 min), N<sub>2</sub>, triethylamine, DMF, 0 °C (3.5 h); (iii) C<sub>4</sub>H<sub>9</sub>Li, THF, N<sub>2</sub>, -78 °C; (iv) ZnI<sub>2</sub>, NaBH<sub>3</sub>CN, ClCH<sub>2</sub>CH<sub>2</sub>Cl, 20 h, rt; (v) TBAF, THF, N<sub>2</sub>, 3 h, rt; (vi) Cs<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>CN, N<sub>2</sub>, 3 h, reflux. B & C. KOH, EtOH, N<sub>2</sub>, 4 h, reflux.

### Scheme 2. Synthesis of compounds 5 and 6<sup>a</sup>

<sup>a</sup>Reagents and conditions: (i) NaH, THF, N<sub>2</sub>, reflux, overnight; (ii) CHCl<sub>3</sub>, 0 °C then RT until complete on TLC

**Scheme 1.**

