

Cedarville University DigitalCommons@Cedarville

Pharmacy and Nursing Student Research and Evidence-Based Medicine Poster Session

12-5-2014

Azithromycin Use in Upper Respiratory Infection

Charles D. Snyder Cedarville University, charlessnyder@cedarville.edu

Sarah Winey Cedarville University, sarahwiney@cedarville.edu

Scarlet Lau Cedarville University, scarletlau@cedarville.edu

Ryley Uber Cedarville University, ryleyuber@cedarville.edu

Mouhannad Saad Cedarville University, msaad@cedarville.edu

See next page for additional authors

Follow this and additional works at: http://digitalcommons.cedarville.edu/ pharmacy_nursing_poster_session



Part of the Nursing Commons, and the Pharmacy and Pharmaceutical Sciences Commons

Recommended Citation

Snyder, Charles D.; Winey, Sarah; Lau, Scarlet; Uber, Ryley; Saad, Mouhannad; Widder, Rebecca; and Anderson, Douglas, "Azithromycin Use in Upper Respiratory Infection" (2014). Pharmacy and Nursing Student Research and Evidence-Based Medicine Poster

http://digitalcommons.cedarville.edu/pharmacy_nursing_poster_session/57

This Poster Session is brought to you for free and open access by DigitalCommons@Cedarville, a service of the Centennial Library. It has been accepted for inclusion in Pharmacy and Nursing Student Research and Evidence-Based Medicine Poster Session by an authorized administrator of DigitalCommons@Cedarville. For more information, please contact digitalcommons@cedarville.edu.



Authors Charles D. Snyder, Sarah Winey, Scarlet Lau, Ryley Uber, Mouhannad Saad, Rebecca Widder, and Douglas Anderson



Azithromycin use in Upper Respiratory Infection Patients

Scarlet Lau, B.S., Mouhannad Saad, B.S., Charles Snyder, Ryley Uber, Sarah Winey, Rebecca Widder, B.A., B.S.P.S., Douglas Anderson, Pharm.D., Cedarville University School of Pharmacy



STATEMENT OF THE PROBLEM

Background:

- Definition: Upper respiratory tract infection (URI)- an infectious disease affecting the portion of the respiratory tract spanning from the nasal cavity to the larynx; can range in severity from acute to life threating ¹
- URIs are generally caused by a viral infection²
 - 44% viral infection
 - 18% bacterial
- 38% no detectable pathogen
- Current guidelines for care:3
 - Not recommended: Initial regimen of antibiotics, except in immunosuppressed or chronically-infected patients
- Recommended: Delayed antibiotic prescribing (not prescribing antibiotics until 48-72 hours after initial symptom presentation)
- Definition: Antibiotic resistance- the process by which microorganisms exposed to antibiotic treatment survive and multiply as a resistant population 4

Significance of the Problem:

- Prescribing guidelines for antibiotics are not being followed⁵
- It is estimated that in the United States doctors overprescribe antibiotics at a rate of at least 50%.4
- Antibiotic overprescribing has been linked to the increasing rate of antibiotic resistance.4
- In the past several years, research and educational promotion concerning antibiotic resistance and stewardship has increased. However, the effectiveness of these materials on prescribing rates has not been determined.4

OBJECTIVES

- 1. To determine adherence to antibiotic prescribing guidelines for upper respiratory infections, specifically regarding azithromycin usage.
- 2. To determine if published literature about antibiotic resistance is correlated with a change in rates of azithromycin prescriptions for acute bronchitis.
- 3. To determine if patients diagnosed with an upper respiratory infection and prescribed a course of azithromycin are being prescribed a second round of antibiotics.
- 4. To determine if education are correlated with a change in rates of second antibiotic after an initial course of azithromycin

HYPOTHESES

Alternate Hypotheses

specifically azithromycin.

azithromycin prescribing.

Null Hypotheses

H01: The majority of prescribers adhere to HA1: The majority of prescribers do not antibiotic prescribing guidelines for upper adhere to antibiotic prescribing guidelines respiratory infections, specifically regarding azithromycin usage.

H02: No correlation exists between published literature about antibiotic resistance and azithromycin prescribing.

H03: The majority of patients prescribed an initial course of azithromycin were not prescribed a second round of antibiotics.

H04: No correlation exists between national antibiotic resistance education and the rate of an additional antibiotic course following azithromycin.

HA2: A correlation exists between national antibiotic resistance education and

for Upper Respiratory Infections,

HA3: The majority of patients prescribed an initial course of azithromycin were prescribed a second round of antibiotics

HA4: A correlation exists between national antibiotic resistance education and the rate of an additional antibiotic course following azithromycin.

PROPOSED METHODS

Study Design

A two-pronged study design

- 1. Retrospective observational study to determine the rates of antibiotic prescribing.
- 2. Literature Review to determine if the increasing body of antibiotic resistance literature is correlated with the prescribing rates.

Data Collection

- 1. Federally Qualified Health Centers chart review
- Date of appointment
- Diagnosis code (ICD-9)
 - Will not include participants who are under age 21 or pregnant
 - Will only include acute respiratory infections, not exacerbations of chronic
- Antibiotic prescription
- 2. PubMed
- Number of articles published per year
 - · Based on search criteria (terms, relevant inclusions, etc.)

Measurement

- Descriptive statistics: Objectives 1 and 3
 - Number of antibiotics prescribed for upper respiratory tract infections, specifically azithromycin
 - Number of patients prescribed a second round of antibiotic after initial azithromycin therapy.
- Inferential Correlation statistics: Objectives 2 and 4
 - Correlation between number of published articles on antibiotic resistance and rate of antibiotic prescribing
 - Correlation between number of published articles on antibiotic resistance and rate of secondary antibiotic prescribing after initial azithromycin therapy.

PROJECT TIMELINE

Data Collection (Spring 2015-Fall 2016)

Presentatio (Fall 2017)

LIMITATIONS

- Statistical power of the project will be determined by the number of second antibiotic prescriptions found in data collection. The generalizability is limited due to the location of the data collection.
- The charts will not indicate whether the prescriptions were filled.
- Patient population will not be randomized

FUTURE DIRECTIONS

The goal of this study is to define the effectiveness of current education campaigns and to effectively inform the public on antibiotic resistance prevention.

ACKNOWLEDGEMENTS

We would like to thank Dr. Chen for her guidance and encouragement for this project.

REFERENCES

Mossad SB. Upper respiratory tract infections. Cleveland Clinic: Centers for Continuing Education Web site. http://www.clevelandclinicmeded.com/medicalpubs/diseasemanagement/infectious-disease/upper-respiratory-tract-infection/. Published August 2013. Updated 2013. A: 2014.

2014. 2. Clark TV, Medina M, Batham S, Curran MD, Parmar S, Nichobon KG. Adults hospitalised with acute respiratory illness rarely have detectable bacteria in the absence of COPD or pneumonia, viral infection predominates in a large prospective UK sample. J Infect. 2014;96(5):507-515. Accessed: October 19, 2014. Accessed: October 19, 20