

#### **Cedarville University** DigitalCommons@Cedarville

The Research and Scholarship Symposium

The 2014 Symposium

Apr 16th, 11:00 AM - 2:00 PM

#### Evaluating the Bioavailability of Carbamazepine Using a Novel SNEDDS Formulation

Gina M. Mattes Cedarville University, GMattes@cedarville.edu

Zachary A. Wallace Cedarville University, zwallace@cedarville.edu

Derrick L. Chapman Cedarville University, dlchapman@cedarville.edu

Jinwon Byun Cedarville University, jbyun@cedarville.edu

Rebecca A. Kyper Cedarville University, rkyper@cedarville.edu

See next page for additional authors

Follow this and additional works at: http://digitalcommons.cedarville.edu/ research\_scholarship\_symposium



Part of the Pharmacy and Pharmaceutical Sciences Commons

Mattes, Gina M.; Wallace, Zachary A.; Chapman, Derrick L.; Byun, Jinwon; Kyper, Rebecca A.; and Injeti, Elisha R., "Evaluating the Bioavailability of Carbamazepine Using a Novel SNEDDS Formulation" (2014). The Research and Scholarship Symposium. 20.  $http://digital commons.cedarville.edu/research\_scholarship\_symposium/2014/poster\_presentations$ 

This Poster is brought to you for free and open access by DigitalCommons@Cedarville, a service of the Centennial Library. It has been accepted for inclusion in The Research and Scholarship Symposium by an authorized administrator of DigitalCommons@Cedarville. For more information, please contact digitalcommons@cedarville.edu.



Presenters Gina M. Mattes, Zachary A. Wallace, Derrick L. Chapman, Jinwon Byun, Rebecca A. Kyper, and Elisha R. Injeti	

# Effect of Particle Size on Bioavailability of Carbamazepine Administered as a Nanoemulsion

Jinwon Byun, Derrick Chapman, Rebecca Kyper, Gina Mattes, and Zachary Wallace | Dr. Elisha Injeti | Cedarville University

#### Introduction

- Carbamazepine (CBZ) is an anticonvulsant drug primarily used to treat epilepsy, bipolar disorder, trigeminal and glossopharyngeal neuralgia.
- CBZ is a lipophilic and poorly soluble drug with limited ability to diffuse across biological membranes.
- To increase its bioavailability, CBZ is formulated into specialized nanoemulsions called Self Nano-Emulsifying Drug Delivery Systems (SNEDDS).
- SNEDDS are isotropic mixtures of oil, surfactants and cosurfactants that form fine oil-in-water nanoemulsions upon mild agitation, followed by injection into aqueous media of gastrointestinal tract.
- SNEDDS have shown to reduce the particle size of CBZ molecules and improve its solubility and ability to cross the biological membranes<sup>1</sup>.
- However, the *in-vivo* bioavailability of CBZ in SNEDDS hasn't been investigated.

# Objective


• The objective of this study is to study the effect of particle size on *in-vivo* bioavailability of CBZ administered as SNEDDS compared to a standard formulation

# Experimental Methods

- The study will be conducted as a randomized controlled crossover experiment utilizing a rat model.
- Sprague-Dawley rats (n=12) will be divided randomly into two groups of same size.
- One group will be administered CBZ in standard formulation and SNEDDS for other group.
- After administration, blood samples will be collected at 5, 10, 15, 20, 30, 45, 60, 90, and 120 minutes<sup>2</sup>.
- Tail vein incision method will be used to collect blood samples.
- Samples will be stored at -20°C until ready for analysis<sup>2</sup>.
- CBZ in blood samples will be analyzed by reversed phase High Performance Liquid Chromatography (HPLC).



# Data Analysis



• An unpaired t-test will be used to compare the significance between the two sets of data.

### Results and Conclusions

- Bioavailability is expected to increase due to the deceased particle size of SNEDDS formulation.
- Limitations include a small sample size which will lower statistical power of the results.

## References

- 1. Kumar GP, Rambhau D., Aapte S.S. Oral bioavailability enhancement of carbamazepine in healthy human volunteers. *J Pharm Research*. 2011;4(2):361-365.
- 2. Fluttert M, Dalm S, Oitzl MS. A refined method for sequential blood sampling by tail incision in rats. *Laboratory Animals*. 2000;34(4):372-378.