



Cedarville University  
**DigitalCommons@Cedarville**

The Research and Scholarship Symposium

The 2014 Symposium

Apr 16th, 11:00 AM - 2:00 PM

# Baylisascaris procyonis Impacts Raccoon (Procyon lotor) Diets

Matthew E. Ingle

*Cedarville University*, [mingle@cedarville.edu](mailto:mingle@cedarville.edu)

Stephen G. Dunbar

*Loma Linda University*

Jayne L. Bartsch

*Cedarville University*, [jbartsch@cedarville.edu](mailto:jbartsch@cedarville.edu)

Kyle J. Culbertson

*Cedarville University*, [kylejculbertson@cedarville.edu](mailto:kylejculbertson@cedarville.edu)

Taylor A. Fulton

*Cedarville University*, [tfulton@cedarville.edu](mailto:tfulton@cedarville.edu)

*See next page for additional authors*

Follow this and additional works at: [http://digitalcommons.cedarville.edu/research\\_scholarship\\_symposium](http://digitalcommons.cedarville.edu/research_scholarship_symposium)

 Part of the [Animal Sciences Commons](#)

Ingle, Matthew E.; Dunbar, Stephen G.; Bartsch, Jaynee L.; Culbertson, Kyle J.; Fulton, Taylor A.; Guffey, Katherine R.; Juris, Aubrey J.; Nolan, Ashlie N.; Nordquist, Dan P.; Rowlands, Carrie E.; and Sitler, Joshua A., "Baylisascaris procyonis Impacts Raccoon (Procyon lotor) Diets" (2014). *The Research and Scholarship Symposium*. 14.

[http://digitalcommons.cedarville.edu/research\\_scholarship\\_symposium/2014/poster\\_presentations/14](http://digitalcommons.cedarville.edu/research_scholarship_symposium/2014/poster_presentations/14)

This Poster is brought to you for free and open access by DigitalCommons@Cedarville, a service of the Centennial Library. It has been accepted for inclusion in The Research and Scholarship Symposium by an authorized administrator of DigitalCommons@Cedarville. For more information, please contact [digitalcommons@cedarville.edu](mailto:digitalcommons@cedarville.edu).



---

**Presenters**

Matthew E. Ingle, Stephen G. Dunbar, Jaynee L. Bartsch, Kyle J. Culbertson, Taylor A. Fulton, Katherine R. Guffey, Aubrey J. Juris, Ashlie N. Nolan, Dan P. Nordquist, Carrie E. Rowlands, and Joshua A. Sitler



# Baylisascaris procyonis impacts raccoon (Procyon lotor) diets

Ingle, M.E.<sup>1,2</sup>, Dunbar, S.G.<sup>2</sup>, Bartsch, J.L.<sup>1</sup>, Culbertson, K. J.<sup>1</sup>, Fulton, T. A.<sup>1</sup> Guffey, K.R.<sup>1</sup>, Juris, A. J.<sup>1</sup>, Nolan, A.N.<sup>1</sup>, Nordquist, D. P.<sup>1</sup> Rowlands, C.E.<sup>1</sup>, and Sitler, J. A.<sup>1</sup>  
<sup>1</sup> Department of Science and Mathematics. College of Arts and Sciences. Cedarville University.  
<sup>2</sup> Department of Earth and Biological Sciences. School of Medicine, Loma Linda University.

## Introduction


Raccoons (*Procyon lotor*) are the definitive host for raccoon roundworms (*Baylisascaris procyonis*). Raccoon roundworm is responsible for a dangerous neurological disease known as larva migrans encephalopathy. Raccoons are omnivorous animals and rely on various food items. Dietary analyses help determine how a raccoon changes its diet in response to environmental features. Raccoons eat whatever food resource is most convenient and abundant. Parasite infections can potentially affect host eating habits in order to keep the host alive and active longer. In this study, we analyzed the diets of necropsied raccoons from ten townships of Clark and Greene Counties by examining their stomach contents.

## Hypothesis

Raccoons from townships with high prevalence (>60%) have the same vertebrate tissue and plant tissue prevalence as raccoons from townships with low roundworm prevalence (<60%) at necropsy.

## Methods

We categorized stomach contents by separating out plant material, vertebrate tissue, and invertebrate tissue storing each sample separately in an ethanol solution. Before processing, we set the samples out in petri dishes to dry. Our research team used an electronic balance to mass the total stomach contents. We also massed all of the plant content separately to compare and obtain a percentage of plant material in the raccoons' diet. We conducted two  $\chi^2$  tests for equality of distributions.



## Results

County	Township	Number of Raccoon stomachs	Vertebrate	Vertebrate	Plant	Plant tissue
			tissue*	tissue prevalence	Tissue**	prevalence
Greene		119	110	0.924	87	0.731
	Beavercreek	42	41	0.976	24	0.571
	Xenia	27	25	0.926	19	0.704
	Miami	50	44	0.880	44	0.880
	Clark	72	66	0.917	63	0.875
Clark	German	9	8	0.889	8	0.889
	Green	22	21	0.955	18	0.818
	Harmony	16	15	0.938	16	1.000
	Mad River	8	8	1.000	8	1.000
	Moorefield	13	11	0.846	10	0.769
	Springfield	4	3	0.750	3	0.75
	> 0.60 prevalence	101	92	0.911	87	0.861
< 0.60 prevalence	90	84	0.933	63	0.700	

Table 1.1-Prevalence of vertebrate and plant tissues found in stomach contents for the nine townships from Clark and Greene Counties, OH.

	Vertebrate Stomach Contents			Plant Stomach Contents			
	Observed	yes	no	Total	Observed	yes	no
> 60	92	9	101	> 60	87	14	101
< 60	84	6	90	< 60	63	27	90
Total	176	15	191	Total	150	41	191
		Expected		Expected			
> 60	93.06806	7.931937	> 60	79.31937	21.68063		
< 60	82.93194	7.068063	< 60	70.68063	19.31937		

Table 1.2- Observed vs. expected stomach contents in high vs. low prevalence townships.

## Conclusions

- There is no significant difference between raccoons from townships with high prevalence (>60) and those with low prevalence (<60) with regards to proportion of raccoons with vertebrate tissue in the stomach at necropsy.
- Plant tissue prevalence did differ significantly between the two groups ( $\chi^2 = 7.353$ ,  $df = 1$ ,  $p = 0.007$ ), with raccoons from high *B. procyonis* prevalence having significantly higher plant tissue prevalence (0.861 vs 0.700).

## Acknowledgements

We would like to thank Dave Pitstick, Dave Linkhart and Steve Hiller for providing the raccoons for us. We would also like to thank Mindy Vasser, Kara Yutzky and Kelsey Cowen for their work in the field. Finally, we would like to thank Dr. Mark Gathany and Dr. Kristen Page for guidance and encouragement.

## References

- Beasley, J. C., Devault, T. L., & Rhodes Jr., O. E. (2007). Home-range attributes of raccoons in a fragmented agricultural region of northern Indiana. *Journal of Wildlife Management*, 71(3), 844-850.
- Blizzard, E. L., Yabsley, M. J., Beck, M. F., & Harsch, S. (2010). Geographic Expansion of *Baylisascaris procyonis* Roundworms, Florida, USA. *Emerging Infectious Diseases*, 16(11), 1803-1804.
- Page, K. L., Beasley, J. C., Olson, Z. H., Smyser, T. J., Downey, M., Kellner, K. F., . . . Rhodes Jr., O. E. (2011). Reducing *Baylisascaris procyonis* Roundworm Larvae in Raccoon Latrines. *Emerging Infectious Diseases*, 17(1), 90-93.
- Page, K. L., Gehrt, S. D., Titcombe, K. K., & Robinson, N. P. (2005). Measuring prevalence of raccoon roundworm (*Baylisascaris procyonis*): a comparison of common techniques. *Wildlife Society Bulletin*, 33(4), 1406-1412.
- Page, K. L., Swihart, R. K., & Kazacos, K. R. (2001a). Changes in transmission of *Baylisascaris procyonis* to intermediate hosts as a function of spatial scale. *Oikos*, 93, 213-220.
- Page, K. L., Swihart, R. K., & Kazacos, K. R. (2001b). Seed preferences and foraging by granivores at raccoon latrines in the transmission dynamics of the raccoon roundworm (*Baylisascaris procyonis*). *Canadian Journal of Zoology*, 79, 616-622.