



Cedarville University
DigitalCommons@Cedarville

The Research and Scholarship Symposium

The 2016 Symposium

Apr 20th, 11:00 AM - 2:00 PM

Genetic Diversity in Concentration of a Protein Subcomponent in Selected Wheat Lines

Andrew B. Berman

Cedarville University, aberman@cedarville.edu

Kelsey Gentry

Cedarville University, kelseygentry@cedarville.edu

Alexander K. Lee

Cedarville University, aklee@cedarville.edu

Molly Yandrofski

Cedarville University, myandrofski@cedarville.edu

Malorie Young

Cedarville University, mryoung@cedarville.edu

See next page for additional authors

Follow this and additional works at: http://digitalcommons.cedarville.edu/research_scholarship_symposium

 Part of the [Food Chemistry Commons](#), [Genetics Commons](#), and the [Molecular Biology Commons](#)

Berman, Andrew B.; Gentry, Kelsey; Lee, Alexander K.; Yandrofski, Molly; Young, Malorie; and Paris, Robert L., "Genetic Diversity in Concentration of a Protein Subcomponent in Selected Wheat Lines" (2016). *The Research and Scholarship Symposium*. 48.
http://digitalcommons.cedarville.edu/research_scholarship_symposium/2016/poster_presentations/48

This Poster is brought to you for free and open access by DigitalCommons@Cedarville, a service of the Centennial Library. It has been accepted for inclusion in The Research and Scholarship Symposium by an authorized administrator of DigitalCommons@Cedarville. For more information, please contact digitalcommons@cedarville.edu.



Presenters

Andrew B. Berman, Kelsey Gentry, Alexander K. Lee, Molly Yandrofski, Malorie Young, and Robert L. Paris

Research + Scholarship SYMPOSIUM



Determining Concentration of Alpha Gliadin Subcomponent in Wheat

Celiac Disease is a hypersensitive response to gluten caused by HLA-DQ2 or HLA-DQ8 T-cell presentation, initiating destruction of intestinal epithelial cells. Studies indicate that an indigestible fragment of the gluten molecule, alpha-gliadin subcomponent 33-mer, rich in proline and glutamine, is responsible for the hypersensitivity response. Determination of 33 mer concentration in wheat lines would be beneficial to future development of wheat lines with reduced 33 mer concentration. Protein from wheat flour was extracted and subjected to western blot in order to quantify the concentration of 33-mer. This will be a valuable tool for future research efforts focused on identification and development of wheat lines with reduced concentrations of 33-mer. Wheat with reduced 33-mer may be suitable for consumption by individuals with celiac disease.