



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di  
Firenze

## LANGERHANS CELLS AND EPIDERMAL MICROENVIRONMENT

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

LANGERHANS CELLS AND EPIDERMAL MICROENVIRONMENT / N. PIMPINELLI; D. BANI. - In: AMERICAN JOURNAL OF DERMATOPATHOLOGY. - ISSN 0193-1091. - STAMPA. - 11(1989), pp. 188-189.

*Availability:*

This version is available at: 2158/25464 since:

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)


## Letters to the Editor

### Langerhans Cells and Epidermal Microenvironment

#### To the Editor:

We read with great interest the article by Dr. Azizi and co-workers (1) on Langerhans cells (LCs) in basal cell carcinomas (BCCs). In particular, we appreciated the careful usage of fine quantitative and reproducible methods in evaluating number, distribution, and morphology of LCs in BCCs.

Concerning the conclusions, however, some critical points require further discussion. In fact, in the three cases in which both BCC and perilesional sun-exposed skin were examined, no difference was found in the number of LCs/mm<sup>2</sup>, while a highly significant difference was found in the dendricity index. Based upon these findings, the authors conclude that the morphologic and—probably—functional perturbation of LCs observed within BCCs may contribute to the growth of this epithelial tumor; alternatively, they suggest the possibility that the growth of tumor cells produces local perturbation in the morphology of the LCs. We have now to consider the following points: (a) A normally differentiated epidermis (2,3) as well as an epidermoid metaplastic epithelium (4) provide a suitable microenvironment for the differentiation of LCs. (b) Pathologic alterations of the epidermal differentiation, either inherited (ichthyosis) or subsequent to viral infection (warts), are associated with a perturbation in LC morphology (5-7). (c) The density and morphology of LCs in epithelial skin tumors have been reported to be related to the degree of epithelial differentiation (8,9), but not to the intraepithelial T-cell density and the amount of peritumoral infiltrate (8). In particular, the smallest number and the most perturbed morphology of LCs were found in BCCs; through morphometric methods, a significant inverse correlation was demonstrated between mean nuclear area of the epithelial tissue and LC density (8). (d) Immunohistologic and ultrastructural studies performed in different pathology models revealed that LCs acquire their complete and—probably—functional characteristics only following their permanence in the epidermal microenvironment (10,11).



In conclusion, while a correlation between number and differentiation of LCs and degree of differentiation of the epithelial microenvironment seems at present reasonably acceptable, it is not possible to define any relationship between perturbation of LC morphology and distribution and creation of epithelial skin cancers, even from a merely speculative point of view. Further studies are clearly needed to elucidate this problem.

Nicola Pimpinelli, M.D.  
Dermatology Clinic II  
Via della Pergola 58  
50121 Florence, Italy

Daniele Bani, M.D.  
Department of Human Anatomy  
and Histology  
Section of Histology  
University of Florence  
Florence, Italy

#### REFERENCES

1. Azizi F, Bucana C, Goldberg L, Kripke ML. Perturbation of epidermal Langerhans cells in basal cell carcinomas. *Am J Dermatopathol* 1987;9:465-73.
2. Schweizer J, Marks F. A developmental study of the distribution and frequency of Langerhans cells in relation to formation of patterning in mouse tail epidermis. *J Invest Dermatol* 1977;69:198-204.
3. Faure M, Mauduit G, Schmitt D, Kanitakis J, Demidem A, Thivolet J. Growth and differentiation of human epidermal cultures used as auto- and allografts in humans. *Br J Dermatol* 1987;116:161-70.
4. Wong Y, Buck RC. Langerhans cells in epidermoid metaplasia. *J Invest Dermatol* 1971;56:10-7.
5. Kolde G, Happle R. Langerhans cell degeneration in X-linked dominant ichthyosis: a quantitative and ultrastructural study. *Arch Dermatol Res* 1985;277:245-7.
6. Ford GP, Friedman PS, Ross J. Langerhans cells in autosomal dominant ichthyosis vulgaris. *Br J Dermatol* 1983;109:309-11.
7. Chardonnet Y, Viac J, Thivolet J. Langerhans cells in human warts. *Br J Dermatol* 1986;115:669-75.
8. Smolle J, Soyer HP, Ehall R, Bartenstein S, Kerl H. Langerhans cell density in epithelial skin tumors correlates with epithelial differentiation but not with the peritumoral infiltrate. *J Invest Dermatol* 1986;87:477-9.

9. Sudo S, Suito A, Morohashi M. Immunohistochemical study of Langerhans cells in skin tumors. *Dermatologica* 1987; 174:76-83.
10. Murphy GF, Messadi D, Fonferko E, Hancock WW. Phenotypic transformation of macrophages to Langerhans cells in the skin. *Am J Pathol* 1986;123:401-6.
11. Bani D, Moretti S, Pimpinelli N, Giannotti B. The differentiation of monocytes into Langerhans cells in human epidermis. In: *Proceedings of the second international workshop on Langerhans cells*. London-Paris: INSERM/John Libbey, 1988 (in press).

*To the Editor:*

We appreciate the comments of Drs. Pimpinelli and Bani and agree with them completely.

Esther Azizi, M.D.  
Margaret L. Kripke, Ph.D.  
Leonard Goldberg, M.D.  
Corazon Bucana, Ph.D.  
The University of Texas System  
Cancer Center  
Texas Medical Center  
1515 Holcombe Boulevard  
Houston, TX 77030, U.S.A.