



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

1H, 15N and 13C resonance assignments of the cerato-platanin, a phytotoxic protein from *Ceratocystis fimbriata*”, Journal of

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

1H, 15N and 13C resonance assignments of the cerato-platanin, a phytotoxic protein from *Ceratocystis fimbriata*”, Journal of Biomolecular NMR / A.L. OLIVEIRA ; L. PAZZAGLI ; B. PANTERA ; G. CAPPUGI; C.E. BENEDETTI ; A. SPISNI ; T.A. PERTINHEZ. - In: JOURNAL OF BIOMOLECULAR NMR. - ISSN 0925-2738. - STAMPA. - 1(2006), pp. 50-50.

Availability:

This version is available at: 2158/205327 since:

Publisher:

Kluwer Academic Publishers:Journals Department, PO Box 322, 3300 AH Dordrecht Netherlands:011 31 78

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Letter to the Editor

¹H, ¹⁵N and ¹³C resonance assignments of cerato-platanin, a phytotoxic protein from *Ceratocystis fimbriata*
DOI 10.1007/s10858-006-9023-z

Cerato-platanin (CP) is a recently described phytotoxic protein released by ascomycete fungus *C. fimbriata* which is the causative agent of canker stain in a number of plants (Pazzagli et al., 1999). CP elicits defence-related responses, suggesting its involvement in the recognition process and the elicitation of plant defence mechanisms. Since CP functions remain speculative and it shows no structural homology with other proteins, the determination of its 3D structure is of obvious interest. In fact, we expect it will provide new clues on its biological functions and may also disclose some new fold. Based on these considerations we undertook the determination of the 3D solution structure of the recombinant CP ¹⁵N and ¹³C double-labelled. 100% of the backbone atoms and, for the side chains, 92% of ¹³C and 95% of ¹H chemical shifts have been assigned. CSI analysis indicate the presence of seven β -strands and two α -helices. Resonance assignments have been deposited in the BioMagResBank under the accession number 6990.

Reference: Pazzagli et al. (1999) *J. Biol. Chem.*, **274**, 24959–24964.

A.L. Oliveira^{a,b}, L. Pazzagli^c, B. Pantera^c, G. Cappugi^c, C.E. Benedetti^b, A. Spisni^a & T.A. Pertinhez^{a,*}
^a*Dept. Experimental Medicine, University of Parma, Parma, 43100, Italy;* ^b*Center for Molecular Structural Biology, LNL, Campinas, Brazil;* ^c*Dept. Biochemical Sciences, University of Florence, Florence, Italy*

*To whom correspondence should be addressed. E-mail: thelma@unipr.it

Supplementary material is available at <http://dx.doi.org/10.1007/s10858-006-9023-z>