

Parallel Synthesis of an Amide Library Based on the 6,8-Dioxa-3-azabicyclo[3.2.1]octane Scaffold by Direct Aminolysis of Methyl Esters

Fabrizio Machetti,^{*,†} Ilaria Bucelli,[‡] Giovanni Indiani,[‡] C. Oliver Kappe,[§] and Antonio Guarna^{‡,||}

Istituto di chimica dei composti organometallici del CNR c/o Dipartimento di chimica organica "Ugo Schiff", Dipartimento di chimica organica "Ugo Schiff" and Laboratorio di progettazione, sintesi e studio di eterocicli biologicamente attivi (HeteroBioLab), Università degli Studi di Firenze, Via della Lastruccia 13, 50019 Sesto Fiorentino, Firenze, Italy, and Christian Doppler Laboratory for Microwave Chemistry and Institute of Chemistry, Karl-Franzens-University Graz, Heinrichstrasse 28, A-8010 Graz, Austria

Received August 30, 2006

An efficient synthesis of unsubstituted and substituted amides based on the 6,8-dioxa-3-azabicyclo[3.2.1]octane scaffold is described. The reaction, carried out at 60 °C in the absence of solvent, is characterized by its mildness and ease of workup. A library of amides, was synthesized by combination of methyl esters **1–6** with various amines. In addition, the microwave-assisted automated synthesis of the library was compared with the above conventional parallel synthesis. Microwave synthesis significantly decreased the reaction time from hours to minutes.

Introduction

The development of novel scaffolds is one of key tools for the generation of molecular diversity in combinatorial chemistry.^{1,2} Recent communications from our group have described the synthesis of new 6,8-dioxa-3-azabicyclo[3.2.1]octane-based scaffolds, named BTAA (bicycles from tartaric acid and amino acid),³ from readily accessible tartaric acid derivatives and amino aldehydes or amino ketones.^{4,5,6,7} These scaffolds are easy to prepare: they possess modulable stereochemistry and low molecular weight, and they present an unprecedented and fairly rigid core. Most important is their versatility for the generation of chemical diversity from up to six sites (Figure 1).⁵

BTAA's have been used for multiple applications⁸ such as ordered homoligomers⁹ and dipeptide isomers.⁴ To enlarge their potential applications, we have decided to generate molecular diversity by choosing scaffolds substituted in positions C-2, N-3, C-4, and C-5 and to use the carboxylic function on C-7 for further elongation or functionalization. In this context, we planned to diversify the scaffolds through a new amide bond.

Because of the peculiar characteristics,^{10,11,12} ubiquitous nature, biological, pharmacological,¹³ and crop protection




Figure 1.

properties¹⁴ of molecules containing amide bonds, the amide bond¹⁵ has stimulated a great deal of attention.^{16,17,18} In addition, secondary amides are precursors of tertiary amines, which are an extremely important class of compounds from the drug discovery perspective.¹⁹

In this paper, we wish to report the results on the one-step parallel preparation of a series of more than one hundred BTAA amides, by direct aminolysis of the corresponding methyl esters.

Results and Discussion

Preliminary Assays. The direct transformation of esters to amides is an important synthetic tool. However, the application of this transformation to the parallel synthesis of amides is limited for a number of reasons. The aminolysis of esters, unless the esters have good leaving groups,²⁰ in general occurs under harsh conditions requiring high temperature and extended reaction times,^{21,22} the use of high pressure,^{23,24} or the use of strong alkali metal catalysts. These conditions^{25–35} are not suitable for the synthesis of chemical libraries which require high-performing processes and final products in high purity.

Recently, our studies on BTAA's have shown the high reactivity of the methyl ester group of these scaffolds.³⁶ Thus, a methyl amide could be obtained easily by dropwise addition

* To whom correspondence should be addressed. E-mail: fabrizio.machetti@unifi.it

† Istituto di chimica dei composti organometallici del CNR.

‡ Dipartimento di chimica organica "Ugo Schiff".

§ Christian Doppler Laboratory for Microwave Chemistry and Institute of Chemistry.

|| Laboratorio di progettazione sintesi e studio di eterocicli biologicamente attivi.




Figure 2. Time course of the ester aminolysis of BnBTG(O)Me **1** (filled square) and BTGOMe **2** (open square) compared with methyl pentanoate **8** (open circle), methyl *p*-nitrobenzoate **7** (filled triangle), and 2-methoxyacetic acid methyl ester **10** (open triangle) by piperidine, performed solvent free at 40 °C. The conversion of the reaction was monitored by ¹H NMR.

of an ethanol solution of methylamine to methyl ester **1**.³⁷ This result opened the possibility to devise a new methodology for the direct conversion of BTAA methyl ester into a variety of amide derivatives.

Initially, we assessed the reactivity of BTAA methyl esters by comparison with a series of methyl ester species (Figure 2). Direct solvent-free aminolysis experiments were set by heating of the methyl esters in the presence of excess of piperidine at 40 °C. The aminolysis of methyl pentanoate **8** and methyl cyclohexanone-2-carboxylate **9** in those conditions did not lead to any formation of the corresponding piperidine amide, even in case of prolonged reaction time (24 h). In case of methyl *p*-nitrobenzoate **7**,³⁸ the process was extremely slow. On the contrary, BTAA scaffolds **1** and **2** and their structurally similar α -methoxy methyl ester **10** showed high or complete conversion to the corresponding amide after few hours. These results showed the importance of the structure of the scaffold in the enhanced reactivity of the ester.

Because of this encouraging finding, scaffolds **1** and **2** were reacted against a series of amines with different nucleophilicities to understand the utility and limitation of the method.

Synthesis of the Library. The synthesis of the BTAA scaffold-based amide library was achieved by the reaction of a variety of substituted amines (Figure 3) with BTAA methyl esters **1–6** at 60 °C over 16 h, in absence of solvent (Scheme 1). The conditions were set to pursue reaction completion for each amine-scaffold combination. The choice for the amine unit was based on its commercial availability, structural diversity, and melting point, which should be lower than the reaction temperature to give a homogeneous reaction mixture. A simple workup procedure was tailored on the basis of the nature of the amine. When volatile amines (boiling point less than 100 °C) were used, the crude reaction mixtures were concentrated under a gentle stream of nitrogen. For less volatile amines, when the scaffold was a lactam ($R^2 = O$), the reaction mixtures were diluted with MeOH and rapidly filtered through a short column of Amberlyst 15 H⁺ ion-exchange resin to remove the excess amine. Otherwise, when the scaffold was an amine ($R^2 = H_2$), the reaction mixtures were filtered through a short pad of silica gel, eluted





Figure 3. Set of amines {1–31} for the library.

with ethyl acetate; in this way, the amine reagents were rapidly removed from the final amide products. All the amides were obtained with excellent purities (based on HPLC or NMR analysis).

Secondary amides were synthesized according to Scheme 1 and analyzed by HPLC or NMR and MS. (Table 1). The reactions were carried out in parallel with different primary amines either linear or α -branched (cyclopropylamine, cyclohexylamine, and methylbenzylamine) or α -unsaturated (propargylamine). The purities of the resulting amides ranged from 67 to 99%, and the conversion was quantitative except in the case of trifluorethylamine **1–6**{3}, which showed low conversion and consequently low purity of the final products. The reasons were attributed to the low boiling

Scheme 1

**Table 1.** Structure and Analytical Data for Secondary Amides Defined in the Library (Scheme 1)^{a,d}

	R ²	R ³	R ⁴	R ⁵	R ⁷	R _t (min)	purity ^b (conv) ^c		R ²	R ³	R ⁴	R ⁵	R ⁷	R _t (min)	purity ^b (conv) ^c
1{1}	O	Bn	H	H	H	5.5	99	3{12}	O	Bn	Ph	H	MBn	15.5	70
1{2}	O	Bn	H	H	Me	6.5	97	4{3}	O	Bzh	H	Ph	TfE	16.0	69 (67)
1{3}	O	Bn	H	H	TfE	8.4	42 (36)	4{4}	O	Bzh	H	Ph	<i>n</i> Pr	16.2	91
1{4}	O	Bn	H	H	<i>n</i> Pr	9.3	99	4{5}	O	Bzh	H	Ph	Cyp	15.8	99
1{5}	O	Bn	H	H	Cyp	8.1	96	4{6}	O	Bzh	H	Ph	Prg	15.3	67
1{6}	O	Bn	H	H	Prg	5.8	99	4{7}	O	Bzh	H	Ph	<i>n</i> Bu	16.8	96
1{7}	O	Bn	H	H	<i>n</i> Bu	10.8	99	4{8}	O	Bzh	H	Ph	<i>i</i> Am	17.6	99
1{8}	O	Bn	H	H	<i>i</i> Am	11.9	92	4{9}	O	Bzh	H	Ph	Hex	18.2	99
1{9}	O	Bn	H	H	Hex	13.2	93	4{10}	O	Bzh	H	Ph	Cyh	17.9	82
1{10}	O	Bn	H	H	Cyh	12.1	95	4{11}	O	Bzh	H	Ph	Bn	6.5	49 (80)
1{11}	O	Bn	H	H	Bn	11.4	99	4{12}	O	Bzh	H	Ph	MBn	11.2	99
1{12}	O	Bn	H	H	MBn	12.0	99	5{3}	H	Bzh	H	Ph	TfE	—	— (40)
2{3}	H	Bn	H	H	TfE	5.7	23 (21)	5{4}	H	Bzh	H	Ph	<i>n</i> Pr	—	— (99)
2{4}	H	Bn	H	H	<i>n</i> Pr	5.6	76	5{5}	H	Bzh	H	Ph	Cyp	—	— (99)
2{5}	H	Bn	H	H	Cyp	5.6	88	5{6}	H	Bzh	H	Ph	Prg	—	— (50)
2{6}	H	Bn	H	H	Prg	5.4	99	5{7}	H	Bzh	H	Ph	<i>n</i> Bu	—	— (99)
2{7}	H	Bn	H	H	<i>n</i> Bu	7.6	89	5{8}	H	Bzh	H	Ph	<i>i</i> Am	—	— (99)
2{8}	H	Bn	H	H	<i>i</i> Am	8.2	99	5{9}	H	Bzh	H	Ph	Hex	—	— (99)
2{9}	H	Bn	H	H	Hex	5.7	73	5{10}	H	Bzh	H	Ph	Cyh	—	— (99)
2{10}	H	Bn	H	H	Cyh	5.6	82	5{11}	H	Bzh	H	Ph	Bn	—	— (80)
2{11}	H	Bn	H	H	Bn	6.4	90	5{12}	H	Bzh	H	Ph	MBn	—	— (99)
2{12}	H	Bn	H	H	MBn	7.4	68 (95)	6{3}	H	Bn	H	Ph	TfE	—	— (46)
3{3}	O	Bn	Ph	H	TfE	13.9	65 (52)	6{4}	H	Bn	H	Ph	<i>n</i> Pr	—	— (99)
3{4}	O	Bn	Ph	H	<i>n</i> Pr	14.4	86	6{5}	H	Bn	H	Ph	Cyp	—	— (54)
3{5}	O	Bn	Ph	H	Cyp	13.7	90	6{6}	H	Bn	H	Ph	Prg	—	— (99)
3{6}	O	Bn	Ph	H	Prg	13.4	83	6{7}	H	Bn	H	Ph	<i>n</i> Bu	—	— (99)
3{7}	O	Bn	Ph	H	<i>n</i> Bu	14.8	86	6{8}	H	Bn	H	Ph	<i>i</i> Am	—	— (99)
3{8}	O	Bn	Ph	H	<i>i</i> Am	11.2	90	6{9}	H	Bn	H	Ph	Hex	—	— (99)
3{9}	O	Bn	Ph	H	Hex	16.5	84	6{10}	H	Bn	H	Ph	Cyh	—	— (99)
3{10}	O	Bn	Ph	H	Cyh	15.9	67	6{11}	H	Bn	H	Ph	Bn	—	— (80)
3{11}	O	Bn	Ph	H	Bn	15.0	83	6{12}	H	Bn	H	Ph	MBn	—	— (99)

^a Abbreviations: *i*Am, isoamyl; Bn, benzyl; *n*Bu, butyl; Bzh, benzhydryl; Cyp, cyclopentyl; Cyh, cyclohexyl; Hex, hexyl; MBn, (*S*)- α -methylbenzyl; *n*Pr, propyl; Prg, propargyl; TFE, trifluoroethyl. For structural representations of all products, see the Supporting Information.


























^b HPLC purities are given as area percent (UV) at 220 nm. For some C-5-substituted amides, HPLC data are not reliable and are not reported. See ref 6 for additional details. ^c Determined by ¹H NMR. Quantitative if not indicated. ^d Obtained by thermal heating.

point (37–38 °C) of {3}, likely to evaporate from reaction mixture under the conditions set for the aminolysis, and to the strong electron-withdrawing group effect of the trifluoroethyl group, which decreased the nucleophilicity³⁹ of {3}.

Nevertheless, performance of the experiment at lower temperature resulted in no conversion.

Similar results were obtained with either acyclic or alicyclic secondary amines (Table 2), except when hindered

Table 2. Structure and Analytical Data for Tertiary Amides and Amides Derived from Multifunctionalised Amines Defined in the Library (Scheme 1)^{a,d}

Compd	R ²	R ³	R ⁴	R ⁵	R ^{7'}	R ^{7''}	R _t (min)	Purity ^b (Conv) ^c	Compd	R ²	R ³	R ⁴	R ⁵	R ^{7'}	R ^{7''}	R _t (min)	Purity ^b (Conv) ^c	
1{13}	O	Bn	H	H	Me	Me	7.3	94	1{22}	O	Bn	H	H		Me	6.6	78	
1{14}	O	Bn	H	H	Et	Et	5.7	34(25)	2{22}	H	Bn	H	H		Me	6.8	77	
1{15}	O	Bn	H	H	<i>i</i> Pr	<i>i</i> Pr	0	—(0)	3{22}	O	Bn	Bn	H		Me	12.0	56	
1{16}	O	Bn	H	H	<i>t</i> Bu	<i>t</i> Bu	0	—(0)	4{22}	O	Bzh	H	Ph		Me	14.7	47	
1{17}	O	Bn	H	H			9.0	99	5{22}	H	Bzh	H	Ph		Me	—	—(80)	
2{17}	H	Bn	H	H			6.8	99	6{22}	H	Bn	H	Ph		Me	—	—(99)	
3{17}	O	Bn	Bn	H			9.3	99	1{23}	O	Bn	H	H				73	
4{17}	O	Bzh	H	Ph			16.2	91	2{23}	H	Bn	H	H				99	
5{17}	H	Bzh	H	Ph			—	—(99)	3{23}	O	Bn	Bn	H			11.9	82	
6{17}	H	Bn	H	Ph			—	—(99)	4{23}	O	Bzh	H	Ph			9.6	81	
1{18}	O	Bn	H	H			10.8	96	5{23}	H	Bzh	H	Ph			—	99	
2{18}	H	Bn	H	H			8.0	99	6{23}	H	Bn	H	Ph			—	99	
3{18}	O	Bn	Bn	H			13.2	86	1{24}	O	Bn	H	H		H	16.5	51	
4{18}	O	Bzh	H	Ph			17.0	85	2{24}	H	Bn	H	H		H	6.1	99	
5{18}	H	Bzh	H	Ph			—	—(78)	3{24}	O	Bn	Bn	H			H	7.0	67
6{18}	H	Bn	H	Ph			—	—(99)	4{24}	O	Bzh	H	Ph			H	8.7	99
1{19}	O	Bn	H	H			7.4	99	5{24}	H	Bzh	H	Ph		H	—	—(99)	
2{19}	H	Bn	H	H			3.9	83	6{24}	H	Bn	H	Ph		H	—	—(99)	
3{19}	O	Bn	Bn	H			13.1	70	1{25}	O	Bn	H	H			6.02	99	
4{19}	O	Bzh	H	Ph			15.3	99	2{25}	H	Bn	H	H			4.8	99	
5{19}	H	Bzh	H	Ph			—	—(65)	3{25}	O	Bn	Bn	H			11.5	93	
6{19}	H	Bn	H	Ph			—	—(99)	4{25}	O	Bzh	H	Ph			13.9	99(50)	
1{20}	O	Bn	H	H			10.0	90	5{25}	H	Bzh	H	Ph			—	99(70)	
2{20}	H	Bn	H	H			7.3	73	6{25}	H	Bn	H	Ph			—	99(65)	
3{20}	O	Bn	Bn	H			16.4	72	1{26}	O	Bn	H	H			16.2	63	
4{20}	O	Bzh	H	Ph			16.3	99	2{26}	H	Bn	H	H			8.5	99	
5{20}	H	Bzh	H	Ph			—	—(82)	3{26}	O	Bn	Bn	H			9.5	99	
6{20}	H	Bn	H	Ph			—	—(80)	4{26}	O	Bzh	H	Ph			14.4	60	
1{21}	O	Bn	H	H		H	5.4	99	5{26}	H	Bzh	H	Ph			—	—(99)	
2{21}	H	Bn	H	H		H	3.8	63	6{26}	H	Bn	H	Ph			—	—(99)	
3{21}	O	Bn	Bn	H			11.3	99	1{27}	O	Bn	H	H			4.7	30	
4{21}	O	Bzh	H	Ph			13.4	99	2{27}	H	Bn	H	H			7.9	99	
5{21}	H	Bzh	H	Ph		H	—	—(99)	3{27}	O	Bn	Bn	H			9.7	99	
6{21}	H	Bn	H	Ph		H	—	—(99)	4{27}	O	Bzh	H	Ph			10.9	75	
1{30}	O	Bn	H	H	Leucinol	H	10.0	79	5{27}	H	Bzh	H	Ph			—	—(83)	
									6{27}	H	Bn	H	Ph			—	—(74)	

^a For structural representations of all products, see the Supporting Information. ^b HPLC purities are given as area percent (UV) at 220 nm. For some C-5-substituted amides, HPLC data are not reliable and are not reported. See ref 6 for additional details. ^c Determined by ¹H NMR. Quantitative if not indicated. ^d Obtained by thermal heating.

secondary amines {15, 16} and aromatic amines {28, 29} were used. For aromatic amines, no reaction was achieved when the reaction was conducted at higher temperature (100–150 °C). The aminolysis with diamine {21–23} gave only the corresponding monoacetylated amide.

The complete analysis of amides contained in Tables 1 and 2 shows that 82% of the compounds had purities higher than 80% and the average purity was 87%. Analysis of the crude products by NMR indicated that the only contamination of the amide material was from the presence of the unreacted methyl ester.

As an alternative to this method, based on classical synthetic chemistry, microwave technology was sought as an important way to improve the direct aminolysis of BTAA methyl ester scaffolds. Over the past two decades, microwave-assisted organic synthesis (MAOS) has had a significant impact on synthetic chemistry⁴⁰ proving to be beneficial in a vast series of reactions including the aminolysis of ester.^{41,42}

For comparison purposes and with the aim of simplification of the procedures, not only to reduce the reaction times but also to increase the conversion in the case of unreactive amines, we have therefore reinvestigated on a selected number of amines and scaffolds the runs under microwave irradiation using sealed-vessel technology.

The conditions we applied were different than the one used for conventional heating:⁴³ a solvent was added (methanol) and the temperature was higher (150–165 °C). Among different solvents, methanol was chosen because of its polarity (suitable for microwave use) and the absence of a potential transesterification reaction that may occur under these reaction conditions when other alcohols are used. In addition, methanol is easily removed from the reaction mixture.

Microwave (MW) heating using superheated methanol as a solvent led to the complete conversion of methyl esters to the corresponding amides, even in case of incomplete reaction using thermal heating (Table 3, entries 5, 9, 11, 19, and 23). The sterically encumbered diisopropyl {15} and

Table 3. Comparison of Thermal (Neat, 60 °C) versus Microwave Heating (Methanol, 160 °C) for the Aminolysis of **1–6**

entry	scaffold	amine	conversion (%) ^a	
			Δ	MW
1	1	4	100	100
2	4	4	100	100
3	4	6	100	100
4	2	8	100	100
5	5	6	50	100
6	4	8	100	100
7	5	8	100	100
8	2	11	100	100
9	5	11	80	100
10	1	12	100	100
11	2	12	95	100
12	1	15	0	0
13	1	16	0	0
14	4	17	100	100
15	6	17	100	100
16	3	17	100	100
17	2	18	100	100
18	4	18	100	100
19	5	18	78	100
20	6	18	100	100
21	2	19	100	100
22	4	19	100	100
23	5	19	65	100
24	6	19	100	100

^a Determined by ¹H NMR and based on methyl ester.

diterbutyl {16} amines did not react at all (Table 3, entries 12 and 13), as well as the poor nucleophilic aniline{28}. With MW heating, the reaction time was dramatically reduced and the aminolysis of methyl ester **1–6** was completed in minutes (15–20 min).

Conclusions

In summary, we have developed a solvent-free (neat) synthetic method for the parallel synthesis of a set of secondary and tertiary amides based on the 6,8-dioxo-3-azabicyclo[3.2.1]octane scaffolds. The method is mild and compatible with sensitive functional groups on either the methyl ester scaffold or the amine unit. The library resulted in low levels of impurities using simple work up and purification procedures.

The amides obtained have low molecular weight (lower than 500) and variable structure. The availability of a wide variety of primary and secondary amines, diamines, and functionalized amines allows the synthesis of a large and highly diverse small molecule library based on the BTAA scaffolds. With the application of MW heating, the amides were obtained in a shorter time compared to that of conventional heating (15–20 min vs a typical 16 h) and with improved process efficiency.

Experimental Section

General Remarks. All amines were purchased from commercial sources and were used without further purification. Ammonia and methylamine were used as solutions in water (33 and 40%, respectively). Scaffolds **1–6** were prepared following our reported procedure and scaled up when necessary. NMR spectra (CDCl₃ solution, unless

otherwise stated) were recorded on Varian Gemini 200 and Varian Mercury 400 (¹H, 200, 400 MHz) spectrometers. Chemical shifts were determined relative to the residual solvent peak (CHCl₃, 7.24 ppm for ¹H NMR and 77.0 ppm for ¹³C NMR; DMSO-*d*₆, 40.45 ppm for ¹³C). Coupling constants, *J* (in Hz), refer to ³J_{HH}. EI mass spectra were carried out at 30 eV ionizing voltage by direct introduction on a QMD 1000 Carlo Erba instrument. Analytical reversed HPLC experiments were carried out on a Beckman System Gold instrument equipped, using a 5 μm (250 × 4.6 mm) C18 Vydac column monitored at 220 nm. The flow rates were 1 mL min⁻¹. The elution buffer included 0.1% TFA in H₂O (eluant A) and 0.1% TFA in CH₃CN (eluant B), and a linear gradient (30% of B in A to 90% of B in 15 min) was used.

Thermal Experiments. Thermal parallel solution-phase reactions were performed using an homemade laboratory synthesizer containing eight microreaction units supported with temperature control, magnetic stirring, and nitrogen streaming.

Microwave Irradiation Experiments. Microwave reactions were conducted using the Emrys Synthetiser (Biotage AB, Uppsala, Sweden). The microwave instrument comprises a monomode microwave cavity that operates at a frequency of 2.45 GHz with continuous microwave irradiation power from 0 to 300 W. The reaction vials are glass-based, conical, 5 mL closed tubes, sealed with Teflon septa and aluminum crimp tops. The reaction temperatures were measured with a built-in IR sensor and refer to total irradiation times.

Evaluation of Methyl Ester Reactivity. The time courses of the ester aminolysis of BnBTG(O)OMe (**1**) and BnBTGOMe (**2**), compared with methyl ester **7–10** by piperidine, were solvent free (1:10 molar ratio). Methyl ester and amine were mixed in a sealed microreaction vessel and immersed in a temperature-controlled bath at 40 °C for a definite time. Duplicate runs were performed if necessary. The conversion of the reaction was monitored by 400 MHz ¹H NMR integrating the appearance of the 7-H proton of amide scaffolds **1**{18} and **2**{18} (5.12 and 5.0 ppm respectively), while simultaneously integrating the disappearance of 1-H protons (4.74 and 4.82 ppm respectively) of ester scaffolds **1** and **2**, respectively. For methyl ester **10**, the disappearance of its CH₂CO protons (3.96 ppm) was integrated, while the appearance of the CH₂CO protons (4.05 ppm) of the corresponding 1-(methoxyacetyl)piperidine **11** was simultaneously integrated. Spectroscopic data for **11**. ¹H NMR (400 MHz): δ 4.05 (s, 2 H, CH₂CO), 3.51 (t, *J* = 5.6 Hz, 2 H, CH₂N), 3.38 (s, 3 H, CH₃), 3.35 (t, *J* = 5.6 Hz, 2 H, CH₂N), 1.64–1.56 (m, 2 H, Pip-H), 1.56–1.48 (m, 4 H, Pip-H). ¹³C NMR: δ 167.0 (s, C=O), 71.7 (t, CH₂CO), 58.9 (q, CH₃), 45.9 (t, CH₂N), 42.8 (t, CH₂N), 26.4 (t, Pip-C), 25.5 (t, Pip-C), 24.4 (t, Pip-C). GC MS: *m/z* (%) 157 (M⁺, 4), 127 (30), 112 (44), 69 (100). In case of benzoate methyl ester **7**, the aromatic protons and the disappearance of methyl ester protons were monitored.

General Procedure for Acylation of Volatile Amines. GP1. Methyl ester (0.014 mmol) and amine (0.14 mmol) were heated in a sealed microreaction vessel at 60 °C for 16

h. For the workup, the reaction mixture was treated under a stream of nitrogen to afford the final product mainly as a yellow oil.

Representative Analytical Data for Compounds Obtained with Procedure GP1 BnBTG(O)NHMe (1{2}). Yield: 3.9 mg, 100%. $^1\text{H NMR}$: δ 7.39–7.15 (m, 5 H, Ph-H), 6.47 (bs, 1 H, NH), 5.80 (d, $J = 2.2$ Hz, 1 H, 5-H), 5.03 (s, 1 H, 7-H), 4.65 (s, 1 H, 1-H), 4.55 (s, 2 H, PhCH₂), 3.36 (dd, $J = 2.2$ and 12.1 Hz, 1 H, 4-H), 3.10 (d, $J = 12.1$ Hz, 1 H, 4-H), 2.85 (d, $J = 4.8$ Hz, 3 H, CH₃N). $^{13}\text{C NMR}$: δ 168.9 (s, C=O), 165.8 (s, C=O), 135.4 (s, Ph-C), 128.9 (d, 2 C, Ph-C), 128.1 (d, 2 C, Ph-C), 127.9 (d, Ph-C), 99.9 (d, C-5), 79.4 (d, C-1), 77.5 (d, C-7), 50.9 (t, C-4), 48.3 (t, PhCH₂), 26.0 (q, NCH₃). MS: m/z (%) 276 (M⁺, 59), 218 ((M - CONHMe)⁺, 14), 120 (79), 112 (61), 91 ((PhCH₂)⁺, 100), 71 (90). Anal. Calcd for C₁₄H₁₆N₂O₄ (276.29): C, 60.86; H, 5.84; N, 10.14. Found: C, 60.66; H, 5.58; N, 9.96.

BnBTG(O)Pip (1{18}). Yield: 4.7 mg, 100%. $^1\text{H NMR}$: δ 7.17–7.37 (m, 5 H, Ph-H), 5.79 (d, $J = 2.2$ Hz, 1 H, 5-H), 5.12 (s, 1 H, 7-H), 4.89 (s, 1 H, 1-H), 4.68 (part A of AB system $J = 13.9$ Hz, 1 H, PhCH₂N), 4.44 (part B of AB system $J = 13.9$ Hz, 1 H, PhCH₂N), 3.66–3.28 (m, 5 H, 4-H, Pip-H), 3.08 (part B of AB system $J = 12.1$ Hz, 1 H, 4-H), 1.80–1.45 (m, 6 H, Pip-H). $^{13}\text{C NMR}$: δ 166.6 (s, C=O), 165.0 (s, C=O), 135.4 (s, Ph-C), 128.8 (d, 2 C, Ph-C), 128.1 (d, 2 C, Ph-C), 127.8 (d, Ph-C), 99.6 (d, C-5), 77.9 (d, C-1), 76.6 (d, C-7), 51.2 (t, C-4), 48.4 (t, PhCH₂), 46.4 (t, Pip-C), 43.3 (t, Pip-C), 26.3 (t, Pip-C), 25.4 (t, Pip-C), 24.5 (t, Pip-C). IR (CHCl₃): ν 1669 cm⁻¹. MS m/z (%): 330 (M⁺, 30), 218 (5), 166 (25), 112 (100), 91 ((PhCH₂)⁺, 95), 69 (55). Anal. Calcd for C₁₈H₂₂N₂O₄ (330.38): C, 65.44; H, 6.71; N, 8.48. Found: C, 65.63; H, 6.38; N, 8.72.

BnBTGPip (2{18}). Yield: 4.5 mg, 100%. $^1\text{H NMR}$: δ 7.30–7.26 (m, 5 H, Ph-H), 5.51 (s, 1 H, 5-H), 5.00 (s, 1 H, 7-H), 4.97 (s, 1 H, 1-H), 3.80–3.38 (m, 6 H, PhCH₂N, Pip-H), 2.82 (part A of AB system $J = 12.6$ Hz, 1 H, 2-H), 2.72 (part A of AB system $J = 10.8$ Hz, 1 H, 4-H), 2.55 (part B of AB system $J = 10.8$ Hz, 1 H, 4-H), 2.35 (part B of AB system $J = 12.6$ Hz, 1 H, 2-H), 1.75–1.51 (m, 6 H, Pip-H). $^{13}\text{C NMR}$: δ 166.8 (s, C=O), 137.1 (s, Ph-C), 128.6 (d, 2 C, Ph-C), 128.3 (d, 2 C, Ph-C), 127.2 (d, Ph-C), 100.6 (d, C-5), 76.3 (d, C-1), 74.7 (d, C-7), 61.6 (t, PhCH₂), 56.5 (t, C-4), 54.5 (t, C-2), 46.5 (t, Pip-C), 43.3 (t, Pip-C), 26.4 (t, Pip-C), 25.4 (t, Pip-C), 24.6 (t, Pip-C). IR (CDCl₃): ν 1631 cm⁻¹. MS m/z (%): 316 (M⁺, 14), 204 (25), 158 (90), 152 (63), 91 (100). Anal. Calcd for C₁₈H₂₄N₂O₃ (316.40): C, 68.33; H, 7.65; N, 8.85. Found: C, 68.01; H, 7.38; N, 8.59.

BnBTF(O)NHnPr (3{4}). Yield: 5.5 mg, 100%. An analytical sample was obtained after chromatography on silica gel (EtOAc/hexane 2:3). $^1\text{H NMR}$: δ 7.44–7.31 (m, 8 H, Ph-H), 7.09–7.05 (m, 2 H, Ph-H), 6.35 (bs, 1 H, NH), 5.42 (s, 5-H), 5.41 (part A of AB system $J = 15.4$ Hz, 1 H, PhCH₂N), 5.04 (s, 1 H, 7-H), 4.61 (s, 1 H, 1-H), 4.00 (part B of AB system $J = 15.4$ Hz, 1 H, PhCH₂N), 3.34–3.07 (m, 4 H, PhCH₂CH, NCH₂CH₂ and 4-H), 2.82–2.70 (m, 1 H, PhCH₂CH), 1.50–1.42 (m, 2 H, CH₂CH₂CH₃), 0.88–

0.84 (m, 3 H, CH₂CH₃). $^{13}\text{C NMR}$: δ 168.1 (s, C=O), 166.0 (s, C=O), 136.0 (s, 2 C, Ph-C), 129.2 (d, 2 C, Ph-C), 129.0 (d, 2 C, Ph-C), 128.9 (d, 2 C, Ph-C), 128.0 (d, Ph-C), 127.9 (d, 2C, Ph-C), 127.2 (d, Ph-C), 101.0 (d, C-5), 79.7 (d, C-1), 77.6 (d, C-7), 60.5 (d, C-4), 45.7 (t, PhCH₂N), 41.0 (t, CH₂NCO), 36.2 (t, PhCH₂CH), 22.7 (t, CH₂CH₃), 11.2 (q, CH₃). MS m/z (%): 394 (M⁺, 6), 303 (91), 160 (93), 91 (100). Anal. Calcd for C₂₃H₂₆N₂O₄ (394.46): C, 70.03; H, 6.64; N, 7.10. Found: C, 70.42; H, 6.88; N, 6.87.

General Procedure for Acylation of Nonvolatile Diamines and Monoamines with Aminoester Scaffolds. (GP2). Methyl ester (0.014 mmol) and amine (0.14 mmol) were heated in a sealed microreaction vessel at 60 °C for 16 h. For the workup, the reaction mixture was filtered through a pad of silica gel (Pasteur pipet filled with 5 cm of silica gel) and washed with 20 mL of EtOAc (for monoamines) or 20 mL of CH₂Cl₂/MeOH 10:1 (for diamines). Concentration of the solution under a stream of nitrogen afforded the final product mainly as yellow oil.

Representative Analytical Data for Compounds Obtained with Procedure GP2 BnBTKNHCH₂CH₂NH₂ (6{21}). Yield: 3.6 mg, 70%. $^1\text{H NMR}$: δ 7.59–7.50 (m, 2 H, Ph-H), 7.42–7.36 (m, 3 H, Ph-H), 7.34–7.25 (m, 5 H, Ph-H), 6.74 (bs, 1 H, NH), 4.83 (s, 1 H, 7-H), 4.79 (s, 1 H, 1-H), 3.68 (part A of AB system $J = 12.9$ Hz, 1 H, PhCH₂N), 3.55 (part B of AB system $J = 12.9$ Hz, 1 H, PhCH₂N), 3.23 (part A of AB system $J = 5.86$ Hz, 1 H, CONCH₂), 3.18 (part B of AB system $J = 5.86$ Hz, 1 H, CONCH₂), 3.07 (part A of AB system $J = 11.5$ Hz, 1 H, 4-H), 2.93 (part A of AB system $J = 11.9$ Hz, 1 H, 2-H), 2.67 (m, 2 H, CH₂NH₂), 2.57 (part B of AB system $J = 11.9$ Hz, 1 H, 2-H), 2.41 (part B of AB system $J = 11.5$ Hz, 1 H, 4-H), 1.34 (bs, 2 H, NH₂). $^{13}\text{C NMR}$: δ 171.8 (s, C=O), 137.8 (s, Ph-C), 137.2 (s, Ph-C), 129.0 (d, Ph-C), 128.8 (d, 2 C, Ph-C), 128.4 (d, 2 C, Ph-C), 128.3 (d, 2 C, Ph-C), 127.3 (d, Ph-C), 125.1 (d, 2 C, Ph-C), 107.9 (s, C-5), 78.6 (d, C-1), 78.5 (d, C-7), 61.3 (t, CH₂Ph), 60.8 (t, C-4), 54.0 (t, C-2), 41.9 (t, CONHCH₂), 41.3 (t, CH₂NH₂). IR (CHCl₃): ν 3414 (NH), 1667 cm⁻¹. MS m/z (%): 367 (M⁺, 55), 338 (32), 247 (100). Anal. Calcd for C₂₁H₂₅N₃O₃ (367.44): C, 68.64; H, 6.86; N, 11.14. Found: C, 68.42; H, 6.97; N, 11.49.

BnBTGNHCH₂CH₂OH (2{24}). Yield: 3.8 mg, 92%. $^1\text{H NMR}$: δ 7.29–7.26 (m, 5 H, Ph-H), 7.00 (s, 1 H, NH), 5.59 (s, 1 H, 5-H), 4.72 (s, 1 H, 7-H), 4.61 (s, 1 H, 1-H), 3.76–3.71 (m, 2 H, HOCH₂), 3.60–3.44 (m, 4 H, PhCH₂N, HNCH₂), 2.84 (d, $J = 11.7$ Hz, 1 H, 4-H), 2.85 (d, $J = 11.0$ Hz, 2 H, 2-H), 2.52 (part A of AB system $J = 11.0$ Hz, 1 H, 2-H), 2.32 (part B of AB system $J = 11.7$ Hz, 1 H, 4-H). $^{13}\text{C NMR}$: δ 172.0 (C=O), 137.0 (s, Ph-C), 128.6 (d, 2 C, Ph-C), 128.2 (d, 2 C, Ph-C), 127.2 (d, Ph-C), 101.0 (d, C-5), 77.2 (d, C-7), 76.6 (d, C-1), 61.2 (t, CH₂Ph), 60.9 (t, CH₂OH), 55.9 (t, C-4), 54.7 (t, C-2), 41.7 (t, CH₂NH₂). MS m/z (%): 292 (M⁺, 5), 263 (32), 247 (34), 204 (37), 158 (87), 128 (70), 120 (88), 91 (100). Anal. Calcd for C₁₅H₂₀N₂O₄ (292.33): C, 61.63; H, 6.90; N, 9.58. Found: C, 61.79; H, 6.66; N, 9.29.

General Procedure for Acylation of Nonvolatile Monoamines with Amidoester Scaffolds. (GP3). Methyl ester

(0.014 mmol) and amine (0.14 mmol) were heated in a sealed microreaction vessel at 60 °C for 16 h. For the workup, the reaction mixture was then diluted with MeOH and rapidly filtered through a short column of Amberlyst 15 H⁺ ion-exchange resin. Concentration of the solution under a stream of nitrogen afforded the final product mainly as a yellow oil.

Representative Analytical Data for Compounds Obtained with Procedure GP3 BnBTG(O)NHCH₂CH₂OH (1{24}). Yield: 3.8 mg, 88%. An analytical sample was obtained by crystallization from CHCl₃. White solid. ¹H NMR: δ 7.35–7.18 (m, 5 H, Ph-H), 6.85 (s, 1 H, NH), 5.83 (d, *J* = 2.6 Hz, 1 H, 5-H), 5.04 (s, 1 H, 7-H), 4.68 (s, 1 H, 1-H), 4.56 (s, 2 H, PhCH₂N), 3.74 (t, *J* = 5.2 Hz, 2 H, HNCH₂), 3.52–3.39 (m, 3 H, 4-H, HOCH₂), 3.12 (part B of AB system *J* = 12.2 Hz, 1 H, 4-H). ¹³C NMR (DMSO-*d*₆): δ 172.6 (s, C=O), 167.6 (s, C=O), 137.0 (s, Ph-C), 129.1 (d, 2 C, Ph-C), 128.1 (d, 2 C, Ph-C), 127.9 (d, Ph-C), 110.3 (d, C-5), 80.3 (d, C-1), 78.6 (d, C-7), 58.5, (t, CH₂OH), 52.1 (t, C-4), 47.9 (t, PhCH₂), 42.2 (t, CH₂NH₂). MS *m/z* (%): 306 (M⁺, 8), 218 (6), 120 (20), 91 (100). Anal. Calcd for C₁₅H₁₈N₂O₅ (306.31): C, 58.82; H, 5.92; N, 9.15. Found: C, 59.04; H, 5.62; N, 8.88.

BnBTF(O)Eip (3{27}). Yield: 5.1 mg, 74%. ¹H NMR: δ 7.38–7.25 (m, 8 H, Ph-H), 7.07–7.03 (m, 2 H, Ph-H), 5.47 (s, 1 H, 5-H), 5.28 (A part of AB system, *J* = 16.1 Hz, 1 H, PhCH₂N), 5.17 (s, 1 H, 7-H), 4.86 (s, 1 H, 1-H), 4.35–4.05 (m, 3 H, CH₂CH₃, PhCH₂N), 3.39–3.42 (m, 8 H, CH₂CHCO, PhCH₂CH, 4-H, N(CH₂)₂), 2.00–1.58 (m, 4 H, C(CH₂)₂), 1.36–1.18 (m, 3 H, CH₃). ¹³C NMR: 174.0 (s, C=O), 166.8 (s, C=O), 165.5 (s, C=O), 136.1 (s, Ph-C), 135.9 (s, Ph-C), 129.2 (d, 2 C, Ph-C), 129.1 (d, 2 C, Ph-C), 129.0 (d, 2 C, Ph-C), 128.9 (d, Ph-C), 127.8 (d, 2 C, Ph-C), 127.1 (d, Ph-C), 100.9 (d, C-5), 78.4 (d, C-1), 77.6 (d, C-7), 61.1 (t, OCH₂CH₃), 60.7 (d, C-4), 46.2 (t), 44.7 (t), 41.4 (t), 36.5 (t), 29.7 (t, Eip-C), 14.1 (q, OCH₂CH₃). IR (CHCl₃): ν 1665 cm⁻¹. MS *m/z* (%): 492 (M⁺, 5), 401 (95), 262 (39), 161 (48), 91 (100). Anal. Calcd for C₂₈H₃₂N₂O₆ (492.56): δ C, 68.28; H, 6.55; N, 5.69. Found: C, 67.89; H, 6.72; N, 5.38.

General Procedures for Microwave-Assisted Generation of Amide Library. Methyl ester (0.02 mmol) and amine (0.2 mmol) were dissolved in MeOH (600 μL) in a 5 mL conical microwave process vial. The vial was sealed with a Teflon septum and placed into the microwave cavity. After irradiation at 160 °C for 16 min and subsequent gas jet cooling (down to 40 °C), the reaction mixture was concentrated and treated, depending on the case, for the workup described above for the thermal procedures.

Acknowledgment. We thank Università di Firenze and MIUR (COFIN 2004–2006) for financial support. I.B. thanks Università di Firenze for doctoral fellowship. Mrs Brunella Innocenti is acknowledged for their technical support. Financial support by the Ente Cassa di Risparmio di Firenze for the purchase of the 400 NMR instrument is gratefully acknowledged. F.M. deeply thanks Dr Antonio Guidi for helpful discussions. C.O.K. wishes to thank Biotage AB for the Emrys Synthesizer microwave reactor.

Supporting Information Available. Structures and ¹H NMR and MS data of all amides and ¹H NMR and MS spectra for amides 1{2, 24}, 3{4, 27}, 6{21}, and 2{24}. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- (1) For recent selected examples of libraries built around a scaffold, see: (a) Nieto, M. J.; Philip, A. E.; Poupaert, J. H.; McCurdy, C. R. *J. Comb. Chem.* **2005**, *7*, 258–263. (b) Rosania, G. R.; Lee, W. J.; Ding, L.; Yoon, H.-S.; Chang, Y.-T. *J. Am. Chem. Soc.* **2003**, *125*, 1130–1131. (c) Holland, R. J.; Hardcastle, I. R.; Dick, A. G.; Nutley, B. P.; Hayes, A.; Jarman, M. *Tetrahedron Lett.* **2002**, *43*, 719–721. (d) Ding, S.; Gray, N. S.; Wu, X.; Ding, Q.; Schultz, P. G. *J. Am. Chem. Soc.* **2002**, *124*, 1594–1596. (e) Guan, Y.; Green, M. A.; Bergstrom, D. E. *J. Comb. Chem.* **2000**, *2*, 297–300. (f) Organ, M. G.; Mayhew, D.; Cooper, J. T.; Dixon, C. E.; Lavorato, D. J.; Kaldor, S. W.; Siegel, M. G. *J. Comb. Chem.* **2001**, *3*, 64–67. (g) Nilsson, J. W.; Thorstenson, F.; Kvarnstrom, I.; Oprea, T.; Samuelsson, B.; Nilsson, I. *J. Comb. Chem.* **2001**, *3*, 546–553. (h) Hirschmann, R.; Ducry, L.; Smith, A. B., III. *J. Org. Chem.* **2000**, *65*, 8307–8316. (i) Falorni, M.; Giacomelli, G.; Porcheddu, A.; Taddei, M. *Eur. J. Org. Chem.* **2000**, 1669–1675. (j) Couladouros, E. A.; Strongilos, A. T. *Angew. Chem., Int. Ed.* **2002**, *41*, 3677–3680. (k) Gan, Z.; Reddy, P. T.; Quevillon, S.; Couve-Bonnaire, S.; Arya, *Angew. Chem., Int. Ed.* **2005**, *44*, 1366–1368. (l) Farooq Biabani, M. A.; Grover, R. K.; Singh, S. K.; Kumar, S.; Raj, K.; Roy, R.; Kundu, B. *Tetrahedron Lett.* **2001**, *42*, 7119–7121. (m) Rosenbohm, C.; Vanden Berghe, D.; Vlietinck, A.; Wengel, J. *Tetrahedron* **2001**, *57*, 6277–6287. (n) Jain, R.; Kamau, M.; Wang, C.; Ippolito, R.; Wang, H.; Dulina, R.; Anderson, J.; Gange, D.; Sofia, M. J. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2185–2189.
- (2) (a) For natural-product-like scaffolds, see: Boldi, A. M. *Curr. Opin. Chem. Biol.* **2004**, *8*, 281–286. (b) For dihydropyrimidine scaffolds, see: Dallinger, D.; Kappe, C. O. *Pure Appl. Chem.* **2005**, *77*, 155–161. (c) For carbohydrate scaffolds, see: Ohnsmann, J.; Madalinski, M.; Kunz, H. *Chim. Oggi* **2005**, *23*, 20–23; Le, G. T.; Abbenante, G.; Becker, B.; Grathwohl, M.; Halliday, J.; Tometzki, G.; Zuegg, J.; Meutermans, W. *Drug Discovery Today* **2003**, *8*, 701–709. (d) For 1,3,5-triazine scaffolds, see: Giacomelli, G.; Porcheddu, A.; De Luca, L. *Curr. Org. Chem.* **2004**, *8*, 1497–1519. (e) For diketopiperazine scaffolds, see: Fischer, Peter M. J. *Pept. Sci.* **2003**, *9*, 9–35.
- (3) For a detailed explanation of BTAA nomenclature, see ref 4.
- (4) Guarna, A.; Guidi, A.; Machetti, F.; Menchi, G.; Occhiato, E. G.; Scarpi, D.; Sisi, S.; Trabocchi, A. *J. Org. Chem.* **1999**, *64*, 7347–7364.
- (5) Guarna, A.; Occhiato, E. G.; Menchi, G.; Machetti, F. PCT Int. Appl. WO 01 64686, September 7, 2001.
- (6) Guarna, A.; Bucelli, I.; Machetti, F.; Menchi, G.; Occhiato, E. G.; Scarpi, D.; Trabocchi, A. *Tetrahedron* **2002**, *58*, 9865–9870.
- (7) Trabocchi, A.; Menchi, G.; Rolla, M.; Machetti, F.; Bucelli, I.; Guarna, A. *Tetrahedron* **2003**, *59*, 5251–5258.
- (8) Trabocchi, A.; Menchi, G.; Guarna, F.; Machetti, F.; Scarpi, D.; Guarna, A. *Synlett* **2006**, 331–353.
- (9) Machetti, F.; Ferrali, A.; Menchi, G.; Guarna, A. *Org. Lett.* **2000**, *2*, 3987–3990.
- (10) North, M. *Contemp. Org. Synth.* **1994**, *1*, 475.
- (11) Beckwith, A. L. J. In *The Chemistry of Amides: Synthesis of Amides*; Zabicky, J., Ed.; Interscience: New York, 1970; p 96.
- (12) Hudson, D. *J. Org. Chem.* **1988**, *53*, 617–624.

- (13) Negwer, M.; Scharnow, H.-G. In *Organic Chemical Drugs and their Synonyms*, 8th ed; Wiley VCH: Weinheim, Germany, 2001; p 4254.
- (14) *CRC Handbook of Pesticides*; Milne, G. W. A., Ed.; CRC Press: Boca Raton, FL, 1994.
- (15) For a statistical investigation into the structural complementarity of natural products and synthetic compounds, see: Henkel, T.; Brunne, R. M.; Müller, H.; Reichel, F. *Angew. Chem., Int. Ed.* **1999**, *38*, 643–647.
- (16) Bohacek, R. S.; Shakespeare, W. C. In *The Amide Linkage: Structural Significance in Chemistry, Biochemistry, and Materials Science*; Greenberg, A., Breneman, C. M., Liebman, J. F., Eds.; Wiley: New York, 2000; Chapter 12.
- (17) For recent report on amide activity, see: (a) Antitumor. Rauko, P.; Novotny, L.; Doinova, I.; Hunakova, L.; Szekeres, T.; Jayaram, H. N. *Eur. J. Pharm. Sci.* **2001**, *12*, 387–394. (b) Anticonvulsant. Sobol, E.; Bialer, M.; Yagen, B. *J. Med. Chem.* **2004**, *47*, 4316–4326. (c) Antimicrobial. Sener, E. A.; Bingöl, K. K.; Ören, İ.; Arpacı, Ö. T.; Yalçın, İ.; Altanlar, N. *Il Farmaco* **2000**, *55*, 469–476. (d) Appendino, G.; Minassi, A.; Berton, L.; Moriello, A. S.; Cascio, M. G.; De Petrocellis, L.; Di Marzo, V. *J. Med. Chem.* **2006**, *49*, 2333–2338. (e) Antiestrogens. Han, Y.; Giroux, A.; Colucci, J.; Bayly, C. I.; McKay, D. J.; Roy, S.; Xanthoudakis, S.; Vaillancourt, J.; Rasper, D. M.; Tam J.; Nicholson, D. W.; Zamboni, R. J. *Biorg. Med. Chem. Lett.* **2005**, *15*, 1173–1180. (f) Janjic, J. M.; Mu, Y.; Kendall, C.; Stephenson, C. R. J.; Balachandran, R.; Raccor, B. S.; Lu, Y.; Zhu, G.; Xie, W.; Wipf, P.; Day, B. W. *Bioorg. Med. Chem.* **2005**, *13*, 157–164. (f) HIV reverse transcriptase inhibitors. Behforouz, M.; Cai, W.; Stocksdale, M. G.; Lucas, J. S.; Jung, J. Y.; Briere, D.; Wang, A.; Katen, K. S.; Behforouz, N. C. *J. Med. Chem.* **2003**, *46*, 5773–5780. (g) Cannabinoid analogues. Francisco, M. E. Y.; Seltzman, H. H.; Gilliam, A. F.; Mitchell, R. A.; Rider, S. L.; Pertwee, R. G.; Stevenson, L. A.; Thomas, B. F. *J. Med. Chem.* **2002**, *45*, 2708–2719. (h) Neuraminidase inhibitors. Chand, P.; Sudhakar, B. Y.; Bantia, S.; Rowland, S.; Dehghani, A.; Kotian, P. L.; Hutchison, T. L.; Ali, S.; Brouillette, W.; Elkattan, Y.; Lin, T.-H. *J. Med. Chem.* **2004**, *47*, 1919–1929. (i) CCR5 receptor antagonists. Palani, A.; Shapiro, S.; Josien, H.; Bara, T.; Clader, J. W.; Greenlee, W. J.; Cox, K.; Strizzi, J. M.; Baroudy B. M. *J. Med. Chem.* **2002**, *45*, 3143–3160. (j) 5-HT_{1F} receptor agonists. Filla, S. A.; Mathes, B. M.; Johnson, K. W.; Phebus, L. A.; Cohen, M. L.; Nelson, D. L.; Zgombick, J. M.; Erickson, J. A.; Schenck, K. W.; Wainscott, D. B.; Brancheck, T. A.; Schaus, J. M. *J. Med. Chem.* **2003**, *46*, 3060–3071. (k) Hypolipidemic activity. Hernández, D.; Bernal, P.; Cruz, A.; Garciafigueroa, Y.; Garduño, L.; Salazar, M.; Díaz, F.; Chamorro, G.; Tamariz, J. *Drug Dev. Res.* **2004**, *61*, 19–36. (l) Antiproliferative activity against prostate cancer. Gududuru, V.; Hurh, E.; Dalton, J. T.; Miller, D. D. *Biorg. Med. Chem. Lett.* **2004**, *14*, 5289–5293.
- (18) (a) For insecticidal properties, see: Siddiqui, B. S.; Gulzae, T.; Begun, S.; Rasheed, M.; Sattar, F. A. *Helv. Chim. Acta* **2003**, *86*, 2760–2767. (b) Siddiqui, B. S.; Gulzae, T.; Begun, S.; Afshan, F.; Sattar, F. A. *Helv. Chim. Acta* **2004**, *87*, 660–666.
- (19) See ref 10 in Brown, A. R.; Rees, D. C.; Rankovic, Z.; Morphy, J. R. *J. Am. Chem. Soc.* **1997**, *119*, 3288–3295.
- (20) Beckwith, A. L. J. In *The Chemistry of Amides*; Zabicky, J., Ed.; Wiley-Interscience: New York, 1970; pp 96–109. For the application of specific active esters such as 4-nitrophenyl or pentafluorophenyl, respectively, see: a) Bodanszky, M. *Nature*, **1955**, *175*, 685. (b) Kisfaludy, L.; Schön, I. *Synthesis* **1983**, 325–326.
- (21) Roe, E. T.; Scanlan, J. T.; Swern, D. *J. Am. Chem. Soc.* **1949**, *71*, 2215–2218.
- (22) Crossley, R.; Curran, A. C. W.; Hill, D. G. *J. Chem. Soc., Perkin Trans I*, **1976**, 977–982.
- (23) Matsumoto, K.; Hashimoto, S.; Otani, S. *Angew. Chem., Int. Ed.* **1986**, *25*, 565–566.
- (24) Matsumoto, K.; Hashimoto, S.; Uchida, T.; Okamoto, T.; Otani, S. *Chem. Ber.* **1989**, *122*, 1357–1363.
- (25) For trimethylaluminum, see: Dolle, R. E.; Nicolaou, K. C. *J. Am. Chem. Soc.* **1985**, *107*, 1695–1698.
- (26) For bis[bis(trimethylsilyl)amido] tin(II), see: Wang, W.-B.; Roskamp, E. J. *J. Org. Chem.* **1992**, *57*, 6101–6103.
- (27) For lithium aluminum hydride, see: Evans, D. A. *Tetrahedron Lett.* **1969**, *10*, 1573–1576.
- (28) For sodium methoxide, see: (a) Achmatowicz, M.; Jurczak, J. *Tetrahedron: Asymmetry* **2001**, *12*, 111–119. (b) De Feo, R. J.; Strickler, P. D. *J. Org. Chem.* **1963**, *28*, 2915–2917.
- (29) For 2-hydroxypyridine, see: Openshaw, H. I.; Wittaker, N. *J. Chem. Soc. C* **1969**, 89–91.
- (30) For sodium diethylaluminumhydride, see: Sim, T. B.; Yoon, N. M. *Synlett* **1994**, *10*, 827–828.
- (31) For butyllithium, see: Yang, K.-W.; Cannon, J. G.; Rose, J. G. *Tetrahedron Lett.* **1970**, *11*, 1791–1794.
- (32) For methyl magnesium iodide, see: Houghton, R. P.; Williams, C. S. *Tetrahedron Lett.* **1967**, *8*, 3929–3931.
- (33) For sodium hydride, see: Singh, B. *Tetrahedron Lett.* **1971**, *12*, 321–322.
- (34) For potassium amide, see: Levi, E. M.; Mao, C.-L.; Hauser, C. R. *Can. J. Chem.* **1969**, *47*, 3761–3766.
- (35) For other miscellaneous methods, see: (a) Silicon tetrachloride. Chan, H.; Wong, L. T. L. *J. Org. Chem.* **1969**, *34*, 2766–2767. (b) Sodium cyanide. Hogberg, T.; Strom, P.; Ebner, M.; Ramsby, S. *J. Org. Chem.* **1987**, *52*, 2033–2036. (c) Enzymes. West, J. B.; Wong, C.-H. *Tetrahedron Lett.* **1987**, *28*, 1629–1632. (d) Electrochemical method. Arai, K.; Shaw, C. H.; Nozawa, K.; Kawai, K.; Nakajima, S. *Tetrahedron Lett.* **1987**, *28*, 441–442.
- (36) Machetti, F.; Indiani, G.; Bucelli, I.; Guarna, A. C. R. *Chim.* **2003**, 631–633.
- (37) For examples of aminolysis of methyl esters, see: (a) Labelle, M.; Gravel, D. *Chem. Commun.* **1985**, 105–106. (b) Rymbrandt, R. H.; Schmidt, F. L. *J. Med. Chem.* **1971**, *14*, 54–56. (c) Chou, W.-C.; Tan, C.-W.; Chen, S.-F.; Ku, H. *J. Org. Chem.* **1998**, *63*, 10015–10017.
- (38) We choose *p*-nitrobenzo ester rather than other substituted benzoates because the rate of aminolysis of the substituted benzoates increases as the substituents change from a strong electron-donating group to a strong electron-withdrawing group, see: Um, I.-H.; Jeon, S.-E.; Seok, J.-A. *Chem.-Eur. J.* **2006**, *12*, 1237–1243.
- (39) Richard, J. P.; Toteva, M. M.; Crugeiras, J. *J. Am. Chem. Soc.* **2000**, *122*, 1664–1674.
- (40) For recent reviews, see: (a) De La Hoz, A.; Diaz-Ortiz, A.; Moreno, A. *Chem. Soc. Rev.* **2005**, *34*, 164–178. (b) Kappe, C. O. *Angew. Chem., Int. Ed.* **2004**, *43*, 6250–6284. (c) Hayes, B. L. *Aldrichim. Acta* **2004**, *37*, 66–77. (d) Kappe, C. O.; Dallinger, D. *Nat. Rev. Drug Discovery* **2006**, *5*, 51–63.
- (41) For *tert*-butoxide-mediated aminolysis, see: (a) Varma, R. S.; Naicker, K. P. S. *Tetrahedron Lett.* **1999**, *40*, 6177–6180. (b) Zradni, F.-Z.; Hamelin, J.; Derdour, A. *Synth. Commun.* **2002**, *32*, 3525–3531.
- (42) For aminolysis in the absence of base or induced by *tert*-butoxide, see: Perreux, L.; Loupy, A.; Delmotte, M. *Tetrahedron* **2003**, *59*, 2185–2189.
- (43) For technical reasons (minimum filling volume of the reaction vessel = 0.5 mL), it was not feasible to run those reactions under solvent-free conditions.