


Huddersfield]

On: 14 January 2015, At: 00:49

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Thiosilanes Based Delivery of Sulfur Functionalities in Organic Synthesis

Alessandro Degl'innocenti ^a, Antonella Capperucci ^b, Paola Ulivi ^b, Alessandro Mordini ^a, Gianna Reginato ^a & Alfredo Ricci ^{a c}

^a Centro C.N.R. Composti Eterociclici c/o

^b Dipartimento di Chimica Organica, via G. Capponi, 9, 50121, Firenze, Italy

^c Dipartimento di Chimica Organica, v.le Risorgimento 4, 40100, Bologna, Italy

Published online: 23 Sep 2006.

To cite this article: Alessandro Degl'innocenti, Antonella Capperucci, Paola Ulivi, Alessandro Mordini, Gianna Reginato & Alfredo Ricci (1993) Thiosilanes Based Delivery of Sulfur Functionalities in Organic Synthesis, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 74:1-4, 385-386, DOI: [10.1080/10426509308038127](https://doi.org/10.1080/10426509308038127)

To link to this article: <http://dx.doi.org/10.1080/10426509308038127>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing,

systematic supply, or distribution in any form to anyone is expressly forbidden.
Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

THIOSILANES BASED DELIVERY OF SULFUR FUNCTIONALITIES IN ORGANIC SYNTHESIS.


ALESSANDRO DEGL'INNOCENTI^{a*}, ANTONELLA CAPPERUCCI^b, PAOLA ULIVI^b,
ALESSANDRO MORDINI^a, GIANNA REGINATO^a and ALFREDO RICCI^{a,c}

^aCentro C.N.R. Composti Eterociclici c/o ^bDipartimento di Chimica Organica, via G. Capponi, 9, 50121, Firenze, Italy. ^cDipartimento di Chimica Organica, v.le Risorgimento 4, 40100 Bologna, Italy.

Abstract: Thiosilanes have been proved efficient reagents in the transfer of sulfur functionalities onto organic substrates.

Vinyl sulfides have recently emerged as valuable synthetic intermediates¹, taking part in a number of interesting chemical transformations. Nonetheless, their synthesis is not straightforward, although a number of methodologies have been proposed in the literature². Moreover, 1,3-bisphenylthiopropanes have been shown extremely versatile intermediates, being used by Corey³ and Cohen⁴ for some key steps in the total synthesis of some prostaglandin and prostacyclin systems.

We report in this note on a novel synthetic approach to these classes of compounds⁵. We have found that upon treatment of a variety of silyl enol ethers in the presence of equimolar amounts of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ with different silyl sulfides, a clean reaction occurs, leading to the corresponding substituted vinyl derivatives as shown in Scheme 1.




SCHEME 1

This reaction possesses a broad generality, occurring smoothly with different substrates, ranging from linear to cyclic enol ethers, leading to a wide variety of vinyl sulfides, a major advantage being the availability of silyl enol ethers from formally every carbonyl compound possessing an α hydrogen. Moreover, the reaction is regiospecific, as shown in the reaction of

6-methyl-1-cyclohexenyloxytrimethylsilane that affords 6-methyl-1-phenylthio-1-cyclohexene exclusively.

A further valuable extension of this reactivity may be achieved when reacting α -enones with 2 equivalents of PhSSiMe_3 . In this case, a direct and convenient entry to 1,3-bis-phenylthiopropanes can be achieved⁶.

When using propenoylsilane as the reacting enone, the corresponding silyl derivative may be obtained in a stereospecific fashion, and the C-Si bond further functionalized to more valuable compounds or simply desilylated with retention of configuration to afford a stereospecific access to 1,3-bisphenylthiopropane formally deriving from acrolein. Alternatively treatment with *t*-BuLi followed by quench with



SCHEME 2

different electrophiles yields functionalized silyl-bis-phenylthiopropanes which may be conveniently reacted with HgCl_2 to afford a general entry to variously functionalized propenoylsilanes (Scheme 2).

REFERENCES

1. Trost, B.M.; Lavoie, A.C. *J. Am. Chem. Soc.* **1983**, *105*, 5075, and references cited therein.
2. Labiad, B.; Villemin, D. *Synthesis*, **1989**, 143, and references cited therein.
3. Corey, E.J.; Erickson, B.W.; Noyori, R. *J. Am. Chem. Soc.*, **1971**, *93*, 1724.
4. Cohen, T.; Bennet, D.A.; Mura, A.J.Jr. *J. Org. Chem.*, **1976**, *41*, 2506.
5. Degl'Innocenti, A.; Ulivi, P.; Capperucci, A.; Mordini, A.; Reginato, G.; Ricci, A. *Synlett*, **1992**, 499.
6. Sato, T.; Otera, J.; Nozaki, H. *Synlett*, **1991**, 903.