

# JCEM




THE JOURNAL  
OF CLINICAL  
ENDOCRINOLOGY  
& METABOLISM

## TSPY1 Copy Number Variation Influences Spermatogenesis and Shows Differences among Y Lineages

Claudia Giachini, Francesca Nuti, Daniel J. Turner, Ilaria Laface, Yali Xue, Fabrice Daguin, Gianni Forti, Chris Tyler-Smith and Csilla Krausz

J. Clin. Endocrinol. Metab. 2009 94:4016-4022 originally published online Sep 22, 2009; , doi: 10.1210/jc.2009-1029

To subscribe to *Journal of Clinical Endocrinology & Metabolism* or any of the other journals published by The Endocrine Society please go to: <http://jcem.endojournals.org/subscriptions/>



## ***TSPY1* Copy Number Variation Influences Spermatogenesis and Shows Differences among Y Lineages**

Claudia Giachini, Francesca Nuti, Daniel J. Turner, Ilaria Laface, Yali Xue, Fabrice Daguin, Gianni Forti, Chris Tyler-Smith, and Csilla Krausz

Andrology Unit (C.G., F.N., I.L., F.D., G.F., C.K.), Department of Clinical Physiopathology, University of Florence, Florence 50139, Italy; and The Wellcome Trust Sanger Institute (D.J.T., Y.X., C.T.-S.), Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SA, United Kingdom

**Context:** *TSPY1* is a tandemly-repeated gene on the human Y chromosome forming an array of approximately 21–35 copies. The testicular expression pattern and the inferred function of the *TSPY1* protein suggest possible involvement in spermatogenesis. However, data are scarce on *TSPY1* copy number variation in different Y lineages and its role in spermatogenesis.

**Objectives:** We sought to define: 1) the extent of *TSPY1* copy number variation within and among Y chromosome haplogroups; and 2) the role of *TSPY1* dosage in spermatogenic efficiency.

**Materials and Methods:** A total of 154 idiopathic infertile men and 130 normozoospermic controls from Central Italy were analyzed. We used a quantitative PCR assay to measure *TSPY1* copy number and also defined Y haplogroups in all subjects.

**Results:** We provide evidence that *TSPY1* copy number shows substantial variation among Y haplogroups and thus that population stratification does represent a potential bias in case-control association studies. We also found: 1) a significant positive correlation between *TSPY1* copy number and sperm count ( $P < 0.001$ ); 2) a significant difference in mean *TSPY1* copy number between patients and controls ( $28.4 \pm 8.3$  vs.  $33.9 \pm 10.7$ ;  $P < 0.001$ ); and 3) a 1.5-fold increased risk of abnormal sperm parameters in men with less than 33 copies ( $P < 0.001$ ).

**Conclusions:** *TSPY* copy number variation significantly influences spermatogenic efficiency. Low *TSPY1* copy number is a new risk factor for male infertility with potential clinical consequences. (*J Clin Endocrinol Metab* 94: 4016–4022, 2009)

The human testis-specific protein Y-encoded 1 (*TSPY1*) gene was originally identified as a gene on the short arm of the human Y chromosome with a testis-specific expression pattern (1, 2). The majority of *TSPY1* copies are arranged in 20.4 kb tandemly repeated units, each containing one copy of *TSPY1* and one of the *CYorf16* pseudogene transcription unit, forming an approximately 400- to 800-kb cluster (*DYZ5*) corresponding to approximately 20–40 gene copies on proximal Yp (3, 4). Such a structure is highly unusual in the human genome: only 12 protein-coding genes are present

in tandem clusters with more than three copies, and none of the others shows more than 16 copies in the current assembly (5). The *TSPY1* cluster lies within the critical region for the gonadoblastoma locus on the Y chromosome (*GBY*) that predisposes dysgenetic gonads of intersex patients to gonadoblastoma development (6, 7), but the biological role of *TSPY1* is otherwise poorly understood.

*TSPY1* is a member of a protein superfamily that includes SET and NAP, which are activating factors of the replication process, as binding-partners of cyclin B (8, 9). Various

TSPY1 isoforms are generated by alternative splicing events of transcripts originating from the same transcriptional units (10, 11). Full-length TSPY1 is expressed in the normal testis predominantly in spermatogonia and at a low level in primary spermatocytes, indicating a major role in the mitotic division of early germ cells (9). The role of the minor TSPY1 transcripts, which seem to be preferentially expressed in round spermatids, remains to be elucidated (12). Thus, the expression pattern of full-length TSPY1 suggests a potential role in male germ cell development as a proliferation factor during spermatogenesis (2, 9).

The number of TSPY1 gene copies observed in the two main studies ranged from 18 to 47 and from 23 to 64 copies, respectively (13, 14); this variation is likely to be generated by unequal sister chromatid exchange (8, 15). Maintenance of a minimum copy number through selection is suggested by the evolutionary conservation of multiple copies of the gene on the Y chromosomes of other mammals (16, 17) and by limited variation in copy number in humans, with pronounced modes and few excursions to low or high numbers of copies (18). The nature of the putative selective force remains entirely unknown but seems most likely to be related to spermatogenesis.

Thus far, only one study has addressed the question of whether or not TSPY1 copy number variation may influence spermatogenesis and reported an association between increased TSPY1 copy number and infertility (19). The interpretation of these results, however, is difficult for two reasons: 1) the quantification method employed was unable to provide an exact TSPY1 copy number because it was not validated against a gold standard method (*e.g.* pulsed-field gel electrophoresis); and 2) the size of the study population was small, and Y haplogroup (hg) matching was performed only in a minority of cases, so differences might reflect hg stratification.

We have now reinvestigated the role of TSPY1 dosage in spermatogenic efficiency by setting up a validated method able to provide an absolute number of TSPY1 copies and applying it to a large study population. For this purpose we compared TSPY1 copy number between men with normozoospermia (controls) and a heterogeneous group of idiopathic infertile men (cases) with different grades of spermatogenic failure. We controlled for Y lineage effects by determining the hg of participants and thus also provide the largest analysis to date of TSPY1 copy number variation in individuals belonging to six different Y lineages.

## Subjects and Methods

### Subjects

The study population consisted of a total of 284 subjects: 154 idiopathic infertile patients, and 130 normozoospermic men. The

study was approved by the local ethics committee. Infertile patients included in the study were seeking complete andrological diagnostic work-up for couple infertility at the Andrology Unit and the Unit of Physiopathology of Reproduction of the University Hospital Careggi (Florence, Italy). Infertile patients were selected on the basis of a comprehensive andrological examination including medical history, semen analysis, scrotal ultrasound, hormone analysis, karyotype and Y chromosome microdeletion screening. Patients with mono- or bilateral cryptorchidism, varicocele grades 2 and 3, obstructive azoospermia, recurrent infections, iatrogenic infertility, hypogonadotrophic hypogonadism, karyotype anomalies, Y chromosome microdeletions including partial deletions of the AZFc region, and partial AZFc duplications and patients with non-Italian origin were excluded.

According to the three major sperm parameters, the infertile group could be divided as follows: azoospermia in 36 patients; cryptozoospermia (<1 million spermatozoa/ml) in 13 patients; severe oligozoospermia (1–5 million spermatozoa/ml) in 55 patients; moderate oligozoospermia (5–20 million spermatozoa/ml) in 36 patients; astheno and/or teratozoospermia in 14 patients. The mean sperm concentrations in patients and controls were  $4.0 \pm 4.7 \times 10^6$  and  $83.9 \pm 53.6 \times 10^6$  sperm/ml, respectively. The mean total sperm numbers in patients and controls were  $14.5 \pm 19.8 \times 10^6$  and  $271.5 \pm 163.0 \times 10^6$  spermatozoa, respectively. Controls were selected on the basis of normal sperm parameters (sperm count, motility, and morphology) defined according to the World Health Organization criteria (20). Samples were collected using approved protocols, and the informed consent of all individuals was obtained.

### Estimation of TSPY1 copy number

Relative TSPY1 copy number was determined by quantitative PCR, using a region of the single copy PMP22 gene as a control locus as described (21). Absolute numbers of TSPY1 genes were estimated by reference to DNA samples in which copy number was known from size measurement of the hybridizing XbaI fragment in pulsed-field gel analysis (13, 22).

### Detection of the AMELY gene deletion

All patients and controls were screened for AMELY gene deletion by PCR amplification of specific Sequence Tagged Sites: sY70 (GenBank accession no. G66517) and sY276 (GenBank accession no. G38362). The amplified products were run on an agarose gel, and all showed the presence of both fragments, indicating that AMELY was present.

### Y hg definition

To exclude recruitment bias, care was taken in the ethnic and geographic matching of the patients and controls. All patients and controls were asked for their paternal and maternal origin and were included only if they had Central Italian ancestry on both sides. The subjects were genotyped for six binary markers defining eight hgs (including paragroups): hgs A, DE, J, K\* (xN, P), N, P\* (xR1a), R1a, and the remaining Y\* (xA, D, E, J, K) chromosomes. Y chromosome haplotyping was performed as previously described for the YAP, M9, SRY-1532, 92R7, LLY22 g, and 12f2 polymorphisms (23). Polymorphisms were visualized by size or presence/absence of fragments (YAP, 12f2) or restriction enzyme digestion pattern for: M9 (*HinfI*), SRY-1532 (*DraIII*), 92R7 and LLY22 g (*HindIII*). Further analysis showed that most P\* (xR1a) chromosomes fell into R1b1b (M269-de-

**TABLE 1.** Sperm characteristics (expressed as mean  $\pm$  SD of sperm concentration and total sperm number) of patients and controls stratified by Y-chromosome hg

Y hg	Patients (n = 154)			Controls (n = 130)		
	n (%)	Sperm concentration (n $\times$ 10 <sup>6</sup> /ml)	Total sperm no. (n $\times$ 10 <sup>6</sup> )	n (%)	Sperm concentration (n $\times$ 10 <sup>6</sup> /ml)	Total sperm no. (n $\times$ 10 <sup>6</sup> )
Y*(xA, D, E, J, K)	37 (24)	3.2 $\pm$ 3.9	11.0 $\pm$ 15.0	31 (23.9)	87.1 $\pm$ 64.8	257.7 $\pm$ 182.5
P*(xR1a)	76 (49.4)	4.3 $\pm$ 5.3	16.1 $\pm$ 22.7	64 (49.2)	83.5 $\pm$ 53.7	270.0 $\pm$ 160.9
J	20 (13)	3.7 $\pm$ 4.7	13.7 $\pm$ 15.8	12 (9.2)	97.8 $\pm$ 56.3	312.9 $\pm$ 196.5
DE	19 (12.3)	4.3 $\pm$ 3.7	16.3 $\pm$ 20.7	16 (12.3)	84.7 $\pm$ 30.1	298.6 $\pm$ 135.9
K*(xN,P)	1 (0.65)	2.9	17.7	7 (5.4)	47.0 $\pm$ 16.6	213.9 $\pm$ 72.5
R1a	1 (0.65)	0.5	1.7	0 (0)		

rived), but the P\*(xR1a) classification is used here because M269 data are not available for all samples.

### Statistical analysis

Statistical analysis was performed using the statistical package SPSS for Windows (version 15.0.1; SPSS Inc., Chicago, IL). Median and mean values between groups were compared using a nonparametric Mann-Whitney *U* test and Student's *t* test (in case of normal distributions), respectively. Correlation between *TSPY1* copy number and sperm count was ascertained by Spearman's correlation test. The cutoff value of *TSPY1* copy number for the comparison of different subgroups was defined on the basis of receiver operating characteristic curve analysis. Differences in the incidence of subjects with *TSPY1* copy number above and below the selected cutoff between patients and controls were evaluated using the Fisher's exact test. A *P* value of 0.05 was considered statistically significant for each test.

## Results

### Y hg distribution in the study population and sperm count in men belonging to distinct hgs

We performed Y hg analysis in the entire study population. Patients and controls showed a similar Y hg distribution with the exception of the low-frequency hg K\*(xN, P) (Table 1). Hgs A and N were not found in our

study population at all, whereas hg R1a was found only in one subject, who was excluded from subsequent analyses.

The mean sperm concentration of normozoospermic men of hg K\*(xN, P) was significantly lower than of those belonging to hgs Y\* and DE (*P* = 0.04 and *P* = 0.001, respectively). However, when total sperm counts were compared, no significant differences were found between subjects belonging to these Y hgs.

### *TSPY1* copy number variation and Y hgs

The number of *TSPY1* copies varied substantially within each hg (Table 2). Although the variance within different hgs was similar, the mean *TSPY1* copy number differed significantly between the following Y lineages: 1) hg P\*(xR1a) vs. hg J (29.7  $\pm$  8.5 vs. 33.7  $\pm$  11.4 copies; *P* = 0.025); and 2) hg P\*(xR1a) vs. hg DE (29.7  $\pm$  8.5 vs. 34.6  $\pm$  11.1 copies; *P* = 0.005).

### *TSPY1* copy number in patients vs. controls

In our study population, the number of *TSPY1* gene copies ranged from 11 to 72 (from 11 to 54 in patients and from 14 to 72 in controls), but the majority of subjects had a *TSPY1* copy number within the range 21–35 (67% of patients and 62% of controls). We observed a shift toward

**TABLE 2.** *TSPY1* copy number distribution and comparison of mean  $\pm$  SD *TSPY1* copy number in different Y hgs with *P* values referred to hg P\*(xR1a)

Y hg	Study population (n = 284)				Patients (n = 154)		Controls (n = 130)		
	n	<i>TSPY1</i> copy no.	Range	<i>P</i> <sup>a</sup>	n	<i>TSPY1</i> copy no.	n	<i>TSPY1</i> copy no.	<i>P</i> <sup>b</sup>
Y*(xA, D, E, J, K)	68	30.7 $\pm$ 10.3	11–62	ns	37	27.9 $\pm$ 8.8	31	34.1 $\pm$ 11.1	<0.05
P*(xR1a)	140	29.7 $\pm$ 8.5	13–58		76	27.2 $\pm$ 7.3	64	32.6 $\pm$ 9.0	<0.001
J	32	33.7 $\pm$ 11.4	18–72	<0.05	20	30.7 $\pm$ 8.5	12	38.7 $\pm$ 14.1	ns
DE	35	34.6 $\pm$ 11.1	12–66	<0.01	19	32.5 $\pm$ 10.0	16	37.1 $\pm$ 12.2	ns
K*(xN,P)	8	28.4 $\pm$ 12.1	19–56	ns	1	22	7	29.3 $\pm$ 12.8	ns

Comparison of mean *TSPY1* copy numbers between patients and controls is shown separately for each Y hg with the respective *P* values. ns, Not significant.

<sup>a</sup> *P* value relative to mean *TSPY1* copy number in the different Y-hg vs. hg P\*(xR1a).

<sup>b</sup> *P* value relative to mean *TSPY1* copy number in patients vs. controls, within each Y-hg.

higher copy numbers in controls and toward lower copy numbers in patients (Fig. 1A). The mean *TSPY1* copy number was significantly lower in the patient group than in the controls ( $28.4 \pm 8.3$  vs.  $33.9 \pm 10.7$ ;  $P < 0.001$ ). The risk of an individual with less than 33 copies being a patient with impaired sperm parameters rather than a control was significantly increased [odds ratio (OR) = 1.5; 95% confidence interval (CI), 1.2–1.8;  $P < 0.001$ ]. We also calculated the risk for being a man with impaired sperm production after removing the 14 pure astheno/teratozoospermic men (*i.e.* those men who had a normal total sperm count but abnormal motility and/or morphology): OR = 1.5; 95% CI, 1.3–1.8.

To investigate whether or not the significant difference observed in the whole study population was limited to a specific hg, we performed—within each hg—a comparison of the mean *TSPY1* copy number in patients vs. controls (Table 2). The mean *TSPY1* copy number in patients was lower in each hg compared with controls, although a significant difference was observed only in hg P\*(xR1a) ( $27.2 \pm 7.3$  vs.  $32.6 \pm 9.0$  copies;  $P < 0.001$ ) and in hg Y\*(xA, D, E, J, K) ( $27.9 \pm 8.8$  vs.  $34.1 \pm 11.1$  copies;  $P = 0.012$ ). The lack of significance in the remaining Y hgs may be due to the low number of subjects.

Because hg P\*(xR1a) showed the most significant difference between controls and patients we calculated the

OR for this specific subgroup with the same threshold value of *TSPY1* = 33 copies (OR = 1.5; 95% CI, 1.2–2.0;  $P = 0.001$ ). The distribution of *TSPY1* copy number in hg P\*(xR1a) is reported in Fig. 1B.


**Effect of *TSPY1* gene copy number variation on spermatogenesis**

**Correlation between the *TSPY1* copy number and sperm count**


We found a positive correlation between the *TSPY1* copy number and sperm count, both in the whole study population (Rho coefficient = 0.331,  $P < 0.001$ ; and Rho coefficient = 0.315,  $P < 0.001$  for sperm concentration and total sperm count, respectively) and separately in controls (Rho coefficient = 0.282,  $P = 0.001$ ; and Rho coefficient = 0.178,  $P = 0.043$  for sperm concentration and total sperm count, respectively) and in patients (Rho coefficient = 0.190,  $P = 0.018$ ; and Rho coefficient = 0.214,  $P = 0.008$  for sperm concentration and total sperm count, respectively) (Fig. 2).

**Comparison of sperm concentration and total sperm count within subgroups defined on the basis of *TSPY1* copy number**

To investigate further the effect of *TSPY1* copy number variation on spermatogenesis, patients and controls were



**FIG. 1.** *TSPY1* copy number ranges in patients and controls in the whole study population (A) and subjects belonging to hg P\*(xR1a) (B). The majority of subjects have a *TSPY1* copy number within the range 21–35.



**FIG. 2.** Scatter plots between *TSPY1* copy number and total sperm count in the whole study population (A), patients (B), and controls (C). Spearman’s correlation coefficients are reported in the text.

**TABLE 3.** Comparison of the median values (with the respective ranges) of sperm concentration and total sperm count among quartiles I and IV in the two study populations (patients and controls)

	I Quartile			IV Quartile			<i>P</i> <sup>a</sup>	<i>P</i> <sup>b</sup>
	<i>TSPY1</i> range (n)	Sperm concentration (n × 10 <sup>6</sup> /ml)	Total sperm no. (n × 10 <sup>6</sup> )	<i>TSPY1</i> range (n)	Sperm concentration (n × 10 <sup>6</sup> /ml)	Total sperm no. (n × 10 <sup>6</sup> )		
Patients	11–22 (n = 37)	0.9 (0–14.8)	1.7 (0–74)	33–54 (n = 38)	4.0 (0–27)	14.3 (0–135)	0.012	0.007
Controls	14–25 (n = 29)	53 (28–188)	224.0 (37.8–570)	40–72 (n = 36)	80.0 (22–329)	285.7 (51–730.8)	0.001	0.041

<sup>a</sup> *P* value obtained comparing the median sperm concentration, within patients and controls, between quartiles I and IV.


<sup>b</sup> *P* value obtained comparing the median total sperm count, within patients and controls, between quartiles I and IV.

grouped into quartiles on the basis of their *TSPY1* copy number. The comparison of the median sperm concentration and total sperm count within quartiles is reported in Table 3. Sperm concentration and total sperm count were higher in the fourth quartile (with the highest *TSPY1* copy numbers) than in the first quartile, and reached statistical significance for sperm concentration in both groups.

We compared sperm concentration and total sperm count within patients and controls, above and below the threshold value used for the calculation of OR, *i.e.* *TSPY1* = 33. The analysis further confirmed that subjects with less than 33 *TSPY1* copy number have a significantly lower sperm production (Fig. 3).

## Discussion

The Y chromosome is enriched for genes involved in spermatogenesis, and its structure predisposes to it deletions



**FIG. 3.** Comparison of the median values of sperm concentration and total sperm number above and below the selected threshold value (33 *TSPY1* copies) used for the calculation of OR, within patients (A) and controls (B). \*, *P* < 0.05; \*\*, *P* = 0.001.

and duplications, and thus to variation in the copy number of these genes. During the past 15 yr, much emphasis has been given to the *AZF* (azoospermia factor) regions on the long arm of the Y chromosome. The complete loss of *AZF* region(s) by deletion causes impairment of spermatogenesis and represents the most frequent known molecular genetic cause of male infertility (3, 24). In addition, partial loss of the *AZFc* region (gr/gr deletion) is a significant genetic risk factor for oligozoospermia (25, 26). We now provide evidence for a significant effect of the copy number of *TSPY1*, a multicopy gene situated on Yp, on spermatogenesis.

Because the Y chromosome shows marked population stratification and data in the literature concerning *TSPY1* copy number variation in different Y lineages are scarce, we first investigated whether the different Y hgs that are common in the Italian population showed different *TSPY1* copy numbers. Based on the analysis of 47 Y hgs, Repping *et al.* (14) had previously observed little variation in *TSPY1* copy number within and between Y hgs, whereas an earlier analysis limited to the few hgs that could be identified in 1994 observed more variation within hgs (13). We now report substantial variation in the number of *TSPY1* copies within the six common Y hgs and paragroups present in our study population. The variation was especially high in hg J (from 18 to 72 copies), although this was not significantly greater than in the other Y lineages. In all hgs we observed a peak of frequencies for *TSPY1* copy number in the range 21–35, which may indicate an optimal range of copy number. Interestingly, hg P (the most frequent hg in Central Italy) contained significantly lower *TSPY1* copy numbers than hgs J and DE; a *P vs. DE* difference is in line with the observations of Mathias *et al.* (13) on equivalent hgs then named 1 and 4, which were made using a different method (pulsed-field gel electrophoresis) on a smaller study population (20 subjects belonging to hg DE *vs.* 31 hg P).

Our finding indicates that the analysis of *TSPY1* copy number variation in the context of case-control association studies is susceptible to stratification bias. This observation implies that data from unmatched study pop-

ulations may not provide reliable information. The distribution of Y hgs in controls and patients in our study was similar, leading us to two major conclusions: 1) no Y hg predisposed to impaired spermatogenesis to a detectable extent; and 2) *TSPY1* copy number differences between controls and cases were independent of the Y background. Concerning the former, the relationship between Y background and reproductive fitness has been controversial, which may partly be related to study design biases (small sample size, inappropriate inclusion criteria for patients and controls) (27). This is the first study of a large European sample in which Y hg distribution was compared between normozoospermic controls and idiopathic infertile men. The lack of a Y hg predisposing to impaired sperm production in our study population, together with the lack of significant differences in total sperm counts of men belonging to different Y hgs, provides strong evidence on the lack of a significant Y hg effect in the Italian population.

Concerning our second conclusion, we found a significantly lower number of *TSPY1* copies in infertile men with abnormal sperm parameters compared with men with normozoospermia. As emphasized above, this difference was not related to a difference in the prevalence of hgs DE or J—the hgs that have a higher mean *TSPY1* copy number than P—in the control group. We observed a trend toward lower *TSPY1* copy number in the infertile group within each hg, suggesting that the contraction or expansion responsible for the observed differences between cases and controls occurs in all Y lineages. Accordingly to our analysis, the risk of having abnormal sperm parameters was increased 1.5-fold for a man with fewer than 33 *TSPY1* copies. Other evidence for a significant contribution of *TSPY1* dosage to spermatogenesis included: 1) positive correlation between sperm count and *TSPY1* copy number in the whole study population and separately in controls and cases; and 2) significant differences in sperm production in men with more and less than 33 copies. Given the link between *TSPY1* copy number and sperm count on the one hand and the difference in mean *TSPY1* copy number between hgs on the other, we would expect to see a *TSPY1*-dependent relationship between hgs and sperm count. However, such an effect is small and is obscured by the large variation in *TSPY1* within each hg, and would only be detected in substantially larger samples.

The maintenance of the evolutionarily unstable *TSPY1* array across mammalian species suggested that low copy number must be disadvantageous, and an effect on spermatogenesis was likely on the basis of the observed expression pattern and inferred function of the *TSPY1* protein. Spermatogenesis requires the concerted action of thousands of genes, all contributing to the efficiency of

spermatogenesis to a different extent. Mutations in genes essential for spermatogenesis inevitably lead to impaired spermatogenesis, whereas mutations in genes acting as tuners/modulators of the efficiency of spermatogenesis may not necessarily lead to clinically overt conditions (28, 29). Infertility is a complex disorder. Hence, instead of highly penetrant mutations in single genes (until now very few causative mutations have been identified), the interaction of multiple factors, both genetic and environmental, with individually small effects is expected to play a major etiopathogenic role. We have now provided evidence for a link between low *TSPY1* copy number and impaired spermatogenesis. Low *TSPY1* copy number therefore represents a new genetic risk factor for male infertility with potential clinical consequences and should be taken into consideration in the context of a multigenic approach to idiopathic infertility.

## Acknowledgments

We thank Grazia Matera, Francesca Gensini, and Valeri Tchepelev for technical assistance, and Vieri Boldi for expert help in statistical analysis. We also thank the Italian Ministry of Education and Research and Telethon-Italy for financial support.

Address all correspondence and requests for reprints to: Csilla Krausz, M.D., Ph.D., Andrology Unit, Department of Clinical Physiopathology, Viale Pieraccini 6, 50139 Firenze, Italy. E-mail: c.krausz@dfc.unifi.it.

This work was supported by the Italian Ministry of Education and Research (Grant PRIN 2008-2010, to C.K.) and by Telethon-Italy (Grant GGP08204, to C.K.). D.J.T., Y.X., and C.T.-S. were supported by The Wellcome Trust.

Disclosure Summary: The authors have nothing to disclose.

## References

1. Arnemann J, Epplen JT, Cooke HJ, Sauermann U, Engel W, Schmidtke J 1987 A human Y-chromosomal DNA sequence expressed in testicular tissue. *Nucleic Acids Res* 15:8713–8724
2. Zhang JS, Yang-Feng TL, Muller U, Mohandas TK, de Jong PJ, Lau YF 1992 Molecular isolation and characterization of an expressed gene from the human Y chromosome. *Hum Mol Genet* 1:717–726
3. Skaletsky H, Kuroda-Kawaguchi T, Minx PJ, Cordum HS, Hillier L, Brown LG, Repping S, Pyntikova T, Ali J, Bieri T, Chinwalla A, Delhaunty A, Delhaunty K, Du H, Fewell G, Fulton L, Fulton R, Graves T, Hou SF, Latrielle P, Leonard S, Mardis E, Maupin R, McPherson J, Miner T, Nash W, Nguyen C, Ozersky P, Pepin K, Rock S, Rohlffing T, Scott K, Schultz B, Strong C, Tin-Wollam A, Yang SP, Waterston RH, Wilson RK, Rozen S, Page DC 2003 The male-specific region of the human Y chromosome is a mosaic of discrete sequence classes. *Nature* 423:825–837
4. Tyler-Smith C, Taylor L, Müller U 1988 Structure of a hypervariable tandemly repeated DNA sequence on the short arm of the human Y chromosome. *J Mol Biol* 203:837–848
5. Warburton PE, Hasson D, Guillem F, Lescale C, Jin X, Abrusan G 2008 Analysis of the largest tandemly repeated DNA families in the human genome. *BMC Genomics* 9:533

6. Lau YF 1999 Gonadoblastoma, testicular and prostate cancers, and the TSPY gene. *Am J Hum Genet* 64:921–927
7. Tsuchiya K, Reijo R, Page DC, Distechi CM 1995 Gonadoblastoma: molecular definition of the susceptibility region on the Y chromosome. *Am J Hum Genet* 57:1400–1407
8. Li Y, Lau YF 2008 TSPY and its X-encoded homologue interact with cyclin B but exert contrasting functions on cyclin-dependent kinase 1 activities. *Oncogene* 27:6141–6150
9. Schnieders F, Dörk T, Arnemann J, Vogel T, Werner M, Schmidtke J 1996 Testis-specific protein, Y-encoded (TSPY) expression in testicular tissues. *Hum Mol Genet* 5:1801–1807
10. Honecker F, Stoop H, de Krijger RR, Chris Lau YF, Bokemeyer C, Looijenga LH 2004 Pathobiological implications of the expression of markers of testicular carcinoma in situ by fetal germ cells. *J Pathol* 203:849–857
11. Lau YF, Lau HW, Kömüves LG 2003 Expression pattern of a gonadoblastoma candidate gene suggests a role of the Y chromosome in prostate cancer. *Cytogenet Genome Res* 101:250–260
12. Kido T, Lau YF 2005 A Cre gene directed by a human TSPY promoter is specific for germ cells and neurons. *Genesis* 42:263–275
13. Mathias N, Bayès M, Tyler-Smith C 1994 Highly informative compound haplotypes for the human Y chromosome. *Hum Mol Genet* 3:115–123
14. Repping S, van Daalen SK, Brown LG, Korver CM, Lange J, Marszalek JD, Pyntikova T, van der Veen F, Skaletsky H, Page DC, Rozen S 2006 High mutation rates have driven extensive structural polymorphism among human Y chromosomes. *Nat Genet* 38:463–467
15. Smith GP 1976 Evolution of repeated DNA sequences by unequal crossover. *Science* 191:528–535
16. Guttenbach M, Müller U, Schmid M 1992 A human moderately repeated Y-specific DNA sequence is evolutionarily conserved in the Y chromosome of the great apes. *Genomics* 13:363–367
17. Murphy WJ, Pearks Wilkerson AJ, Raudsepp T, Agarwala R, Schäffer AA, Stanyon R, Chowdhary BP 2006 Novel gene acquisition on carnivore Y chromosomes. *PLoS Genet* 2:e43
18. Tyler-Smith C 2008 An evolutionary perspective on Y-chromosomal variation and male infertility. *Int J Androl* 31:376–382
19. Vodicka R, Vrtel R, Dusek L, Singh AR, Krizova K, Svacinova V, Horinova V, Dostal J, Oborna I, Brezinova J, Sobek A, Santavy J 2007 TSPY gene copy number as a potential new risk factor for male infertility. *Reprod Biomed Online* 14:579–587
20. World Health Organization 1999 WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction. 3rd ed. Cambridge, UK: Cambridge University Press; 4–33
21. Jobling MA, Lo IC, Turner DJ, Bowden GR, Lee AC, Xue Y, Carvalho-Silva D, Hurles ME, Adams SM, Chang YM, Kraaijenbrink T, Henke J, Guanti G, McKeown B, van Oorschot RA, Mitchell RJ, de Knijff P, Tyler-Smith C, Parkin EJ 2007 Structural variation on the short arm of the human Y chromosome: recurrent multigene deletions encompassing amelogenin Y. *Hum Mol Genet* 16:307–316
22. Oakey R, Tyler-Smith C 1990 Y chromosome DNA haplotyping suggests that most European and Asian men are descended from one of two males. *Genomics* 7:325–330
23. Rosser ZH, Zerjal T, Hurles ME, Adojaan M, Alavantic D, Amorim A, Amos W, Armenteros M, Arroyo E, Barbujani G, Beckman G, Beckman L, Bertranpetit J, Bosch E, Bradley DG, Brede G, Cooper G, Còrte-Real HB, de Knijff P, Decorte R, Dubrova YE, Evgrafov O, Gilissen A, Glisic S, Gölge M, Hill EW, Jeziorowska A, Kalaydjieva L, Kayser M, Kivisild T, Kravchenko SA, Krumina A, Kucinskis V, Lavinha J, Livshits LA, Malaspina P, Maria S, McElreavey K, et al. 2000 Y-chromosomal diversity in Europe is clinal and influenced primarily by geography, rather than by language. *Am J Hum Genet* 67:1526–1543
24. Krausz C, Degl'Innocenti S 2006 Y chromosome and male infertility: update, 2006. *Front Biosci* 11:3049–3061
25. Giachini C, Laface I, Guarducci E, Balercia G, Forti G, Krausz C 2008 Partial AZFc deletions and duplications: clinical correlates in the Italian population. *Hum Genet* 124:399–410
26. Repping S, Skaletsky H, Brown L, van Daalen SK, Korver CM, Pyntikova T, Kuroda-Kawaguchi T, de Vries JW, Oates RD, Silber S, van der Veen F, Page DC, Rozen S 2003 Polymorphism for a 1.6-Mb deletion of the human Y chromosome persists through balance between recurrent mutation and haploid selection. *Nat Genet* 35:247–251
27. Krausz C, Quintana-Murci L, Forti G 2004 Y chromosome polymorphisms in medicine. *Ann Med* 36:573–583
28. Nuti F, Krausz C 2008 Gene polymorphisms/mutations relevant to abnormal spermatogenesis. *Reprod Biomed Online* 16:504–513
29. Tyler-Smith C, Krausz C 2009 The will-o'-the-wisp of genetics—hunting for the azoospermia factor gene. *N Engl J Med* 360:925–927