



Withholding Irrigation During the Establishment Phase Affected Growth and Physiology of Norway Maple (*Acer platanoides*) and Linden (*Tilia* spp.)

Alessio Fini, Francesco Ferrini, Piero Frangi, Gabriele Amoroso, and Riccardo Piatti

Abstract. The aim of this work was to investigate the drought tolerance of different *Tilia* species and of different cultivars of *Acer platanoides* grown during the establishment phase, and to evaluate irrigation effect on their growth and physiology. In winter 2004–2005, 168 trees [8–10 cm (3–4 in) circumference] of *Tilia platyphyllos*, *T. cordata*, *T. × europaea*, *T. tomentosa*, *Acer platanoides* ‘Summershade’, *A. platanoides* ‘Deborah’, and *A. platanoides* ‘Emerald Queen’ were planted in the field. Eighty-four plants were irrigated with a drip irrigation system (4 l/h) and eighty-four were not. Height, trunk diameter, and shoot elongation were measured at the end of the growing season in 2005, 2006, and 2007. Leaf gas exchange and chlorophyll fluorescence were measured monthly during the 2006 and 2007 growing seasons. Leaf greenness index content was measured in 2006 and 2007. Results indicate that *T. tomentosa* and *T. cordata* are more drought tolerant during establishment than *T. platyphyllos*, while *Acer platanoides* ‘Summershade’ is less drought tolerant during establishment than the cultivars ‘Emerald Queen’ and ‘Deborah’.

Key Words. *Acer platanoides*; Chlorophyll Fluorescence; Drought Avoidance; Leaf Gas Exchange; *Tilia* spp.; Water Stress.

Global environmental conditions have changed during the last century and based on current trends, temperature will rise by about 1°–3.5°C (1.8°–6.3°F) over the next 70 years, and rainfall will also be affected by a decrease in the frequency and an increase in intensity of rainy events (UNEP/IUC 1999). Water limitation may prove to be a critical constraint to primary productivity of plants under future scenarios of more arid climates due to climate change (Fisher et al. 2001). When referred to shade trees planted in the urban environment and peri-urban forests and recreational areas, plant productivity may correspond with the ability of single plants or plant communities to provide benefits to the inhabitants. Healthy, long-lived trees provide environmental, ecological, economic, social, cultural, and aesthetic benefits to the community (Akbari 2002; Brack 2002; Fini and Ferrini 2007; Nowak et al. 2007; Elmendorf 2008; Escobedo et al. 2008).

Mortality rate in the urban environment is usually very high and ranges from 10% to 50% with water stress playing a major role, especially where pavements, soil compaction, and small planting pits prevent infiltration into the root zone (Kaushal and Aussenac 1989; Miller and Miller 1991; Whitlow et al. 1992; Pauleit et al. 2002). This threat is very dangerous in the first years after planting, when mortality can be 50% in the first year, and up to 34% in the second (Gilbertson and Bradshaw 1985; Nowak et al. 1990). Irrigation is an important factor to increase plant survival and quality during the establishment phase, but landscape water consumption is highly visible and provides a prime target for water restrictions and subsequent regulation (Scheiber et al. 2007). Despite the need of saving water, water restrictions during landscape establishment can be detrimental to plants which have not had enough time to develop a sufficient root system to compensate for evapotranspirational losses (Montague et al. 2000). One way to increase water efficiency is to irrigate trees until they are fully established

and then terminate irrigation unless there are periods of extreme drought. Establishment time can be estimated by comparing leaf gas exchange and growth rates of newly planted stressed (nonirrigated) and unstressed (irrigated) trees (Scheiber et al. 2007).

Another strategy to reduce drought-related transplant losses is to plant species/cultivars that show a certain degree of drought tolerance during the establishment phase. Even within a genus or a species, great differences in water needs for establishment can be found among species/cultivars. A previous work ranked drought tolerance of *Fraxinus* genotypes on the basis of drought-induced changes in chlorophyll fluorescence, chlorophyll content, and carbon assimilation (Percival et al. 2006).

Few studies have assessed drought tolerance among maples and lindens: most were performed on container-grown plants (Abrams and Kubiske 1990; Zwack et al. 1998; Fini et al. 2008) and none of them addressed drought tolerance during the establishment phase. This is very surprising because maples and lindens are widely-used species for the urban environment in Italy and Europe. Four linden species are commonly used for urban forestry: 1) *T. cordata* Mill., a native European species widespread from Spain to Northern Greece to Southern Finland which forms climax forest in plain and mountain areas up to 1500 m (0.93 mi) above sea level; 2) *T. platyphyllos* Scop., a native European species which grows widely from Northern Spain to Caucasus and to Southern Sweden and forms climax forest preferentially in mountain areas, up to 1600 m (1 mile) above sea level; 3) *T. × europaea* (syn. *T. × vulgaris*) L. is the natural hybrid between *T. cordata* and *T. platyphyllos*; 4) *T. tomentosa* Moench, a native of Southern Europe and Asia. Norway maple (*Acer platanoides* L.), and Sycamore maple (*Acer pseudoplatanus* L.) are the most commonly used maple species in European and Italian towns. A previous study found that *A. platanoides* is more

advisable for urban plantings than *A. pseudoplatanus* (Fini et al. 2008). Several Norway maple cultivars are available for sale, but some such as ‘Deborah’, ‘Emerald Queen’, and ‘Summershade’ are particularly appreciated due to reddish colour of new sprouts, intense green colour of foliage, and heat tolerance, respectively.

The aim of this work was to: 1) investigate the effects of irrigation on growth and physiology of *Tilia* and *Acer platanoides* during the establishment phase; 2) evaluate whether differences in drought tolerance during the establishment phase existed within *Tilia* genus; and 3) evaluate whether differences in drought tolerance during the establishment phase existed among three *Acer platanoides* cultivars. This information can be useful to urban foresters and arborists in selecting the right species and cultivars within a desired genus.

MATERIALS AND METHODS

Plant Material and Growing Conditions

In winter 2004–2005, 168 uniform, 2.5 to 3 m (8.3 to 9.9 ft) tall, 3 to 4 cm (1.2 to 1.6 in) diameter [measured at 1.3 m (4.3 ft) height], balled and burlapped [size of the root ball was approximately 30 cm (12 in) in diameter] *Tilia platyphyllos*, *T. cordata*, *T. × europaea*, *T. tomentosa*, *Acer platanoides* ‘Summershade’, *A. platanoides* ‘Deborah’, and *A. platanoides* ‘Emerald Queen’ were planted in an experimental plot located at the Fondazione Minoprio (Como, Italy, 45°44’N, 9°04’E). Mean annual temperature and rainfall of the experimental site, calculated over the last 20 years, are 12.99°C (55.38°F) and 1086 mm (47.76 in) respectively. Despite of relatively high rainfall, some drought spells are likely to occur in the study site in summer and especially in July, which is the driest and the warmest month.

Trees were planted in a loam soil (51% sand, 40% silt, 9% clay). Planting holes were two times the width of the root-ball and deep enough to position the root flare exactly at the soil level.

Trees of the different species or cultivars were planted in a randomized complete block with three replicates. Each block was made up of four plants per species/cv. and irrigation treatment. All plants were irrigated in 2005 to avoid transplant shock. Starting in 2006, 12 plants per species/cultivar (irrigated; 84 plants for the whole experiment) were irrigated with a drip irrigation system (4 l/h) in order to maintain constant soil moisture and for 12 plants per species/cv. (nonirrigated) irrigation was withheld throughout the experiment. Irrigation was provided to irrigated plants when evapotranspiration exceeded precipitation, which occurred from May to the end of August 2006, and from June 20 to the end of September 2007 (Figure 1). Effective evapotranspiration (ETE) can be calculated as the product of reference evapotranspiration (ET_0) and a crop coefficient (Kc). ET_0 was measured with a Class A pan evaporimeter. Kc was estimated to be 0.5 to 0.8 for many landscape trees and shrubs including maple and linden (Daugherty 2002; Pardossi et al. 2004). Since we only mattered that irrigated plants experienced no water stress, we overestimated Kc and we considered a Kc = 1 for effective evapotranspiration determination as proposed by Pardossi et al. (2004). The difference between rainfall and ETE gave the water balance of the trees. Two irrigation events per week were performed on irrigated plants. Irrigation time was the time needed to recover to 0 (when negative) the difference between rainfall and ETE. Time needed to achieve this goal varied, generally, from 3 (May) to 15 (July) hours per week.




Figure 1. Net balance between rainfall and ETo measured weekly from the first of June to the 20th of September 2007.

Data Collection

Shoot elongation was measured on 20 randomly chosen shoots on first order lateral branches per plant on two plants per species/cultivar per irrigation treatment per replicate at the end of the growing season in 2006 and 2007. Trunk diameter was measured on all plants at 1.3 m (4.3 ft) above the root flare.

Leaf gas exchange (net photosynthesis, A, $\mu\text{mol m}^{-2} \text{s}^{-1}$; transpiration, E, $\text{mmol m}^{-2} \text{s}^{-1}$; stomatal conductance, g_s , $\text{mmol m}^{-2} \text{s}^{-1}$) was measured in 2006 and 2007 from June to September using the CIRAS-2 portable infrared gas analyzer (PP Systems, Hertfordshire, UK). The readings were taken between 8.00 h and 11.00 h (maple), and from 14.00 h to 17.00 h (linden) on sunny days under fixed CO_2 concentration (360 ppm) and saturating irradiance ($1300 \mu\text{mol m}^{-2} \text{s}^{-1}$, provided by a built-in red LED radiation source) on two plants per species/cultivar per irrigation treatment and replicate. Five fully expanded leaves per plant were selected from the outer portion of the crown and at different heights and checked for gas exchange. On July 20 and 27, 2007, leaf gas exchange was measured continuously for the entire day, to assess the daily trend of gas exchange. Data are presented in Tables 5 and 6 as the average between the two sampling dates. Water Use Efficiency (WUE) was calculated as the ratio between A and E as reported in a previous work (Ferrini et al. 2008).

Leaf greenness index was measured with a chlorophyll meter (SPAD-502, Minolta, Sakai, Osaka, Japan) on six leaves per species per treatment and replicate at the end of July in 2006 and at the beginning of September in 2007. A recent work on three tree species shows that SPAD values of leaf chlorophyll content can also be a good indicator of leaf N content and leaf carotenoids (Percival 2008).

Chlorophyll fluorescence was measured on six leaves per species per treatment and replicate with a Plant Efficiency Analyser Chlorophyll Fluorimeter (Handy Pea, Hansatech Ins., King's Lynn, Norfolk, U.K.) in July 2006 and in June, July, and August 2007. Fluorescence values were obtained by placing leaves in darkness for 30 minutes by attaching light-exclusion clips to the leaf surface of whole trees. F_0 (minimal fluorescence); F_v/F_m and F_v/F_0 were measured after exposing the leaf to a saturating light ($3000 \mu\text{mol m}^{-2} \text{s}^{-1}$). These parameters are reliable

indicators of the occurrence of environmental stresses, including water stress, on PSII of several woody and herbaceous species (Angelopoulos et al. 1996; Maxwell and Johnson 2000; Percival and Fraser 2001; Percival et al. 2003, Percival 2005; Lazár 2006; Li et al. 2006; Percival et al. 2006; Yamada et al. 2006). F_0 , the minimal fluorescence, is a measure of the stability of the light harvesting complex. F_v/F_m is the maximum quantum yield of the PSII. F_v/F_0 is an estimate of the maximum primary yield of photochemistry of PSII. In this study, chlorophyll fluorescence was used to evaluate the occurrence of water stress within the genus *Tilia* and within the *Acer platanoides* species. Presence of stress was identified by changes in F_0 or decrease in F_v/F_m and F_v/F_0 in nonirrigated plants if compared to irrigated plants of the same species/cultivar. Comparison between mean annual leaf gas exchange of irrigated and nonirrigated plants of the same species were measured to identify the drought tolerance strategy adopted by the different species/cultivars.

Statistical Analysis

All data were subjected to one- or two-way analysis of variance (ANOVA) using SPSS statistical package for Windows (SPSS Inc., Chicago, IL). The two genera were always analyzed independently. Effects of irrigation and species/cultivar were analyzed with a mixed model two-way ANOVA, where irrigation was the fixed effect and species/cultivar the random effect. When no significant interaction between factors was found, differences among species/cultivars within a genus/species were tested with Duncan's multiple range test ($P \leq 0.05$ and $P \leq 0.01$). Parameters which showed significant interaction between factors were plotted separately in order to compare each level of factor A (species/cultivar) for each level of factor B (irrigation) (Chew 1976). Data on leaf gas exchange were analyzed per single sampling date, merged together, and processed again to obtain an average value on annual basis. Data of daily trend of A and E were analyzed with repeated measures General Linear Model.

RESULTS AND DISCUSSION

Over the three years of the experiment, no plant died for water stress. Some signs of wilting were found in July 2007 (the driest month in the two years) in nonirrigated *T. platyphyllos*. By the end of August 2006, nonirrigated *Acer platanoides* 'Deborah' shed over half of the leaves, while no leaf shedding was observed in irrigated *Acer platanoides* 'Deborah' and in the other cultivars of Norway maple and linden species.

Effects of Species/Cultivar and Irrigation on Growth, Gas Exchange and Chlorophyll Content

Among the linden species, *T. cordata* and *T. x europaea* had, respectively, the highest shoot growth in 2006 and 2007 (Table 1; Figure 2). In 2006, *T. platyphyllos* and *T. tomentosa* were slower growing than both the former species and in 2007 were slower growing than *T. x europaea*. Trunk diameter growth had a trend similar to shoot growth in 2006, and no difference among the species appeared in 2007 (data not shown). In 2006, *T. cordata* had the highest mean annual A and WUE (Table 1). High gas exchange was found also in *T. tomentosa*, while *T. platyphyllos* had the lowest gas exchange in both years. In 2006 and 2007, *T. cordata* and *T. tomentosa* had the highest chlorophyll con-

tent respectively, while *T. platyphyllos* had the lowest. No difference among the species appeared for WUE in 2007. An increase in shoot growth and leaf gas exchange between 2006 and 2007 was observed for *T. x europaea*, which in 2006 had lower A, WUE, and chlorophyll content especially when compared to *T. cordata* and *T. tomentosa*, while it ranked very high in 2007.

Irrigation increased shoot growth and transpiration in both years (Table 1). Net photosynthesis was increased by irrigation only in 2006. WUE and chlorophyll content were unaffected by irrigation. Significant interaction between factors was found for shoot extension in 2006, and for E in 2006 and 2007 (Figure 2). In 2006, shoot growth was increased by irrigation in all species but *T. x europaea*. In 2006, *T. cordata* had the highest E both in non irrigated and irrigated environments.




Figure 2. Means separation of those parameters [shoot growth in 2006, transpiration (E, $\text{mmol s}^{-1} \text{m}^{-2}$) in 2006 and 2007], which were significantly affected by species \times irrigation interaction in *Tilia*. In this case, Duncan's Multiple Range Test was used to compare the species for each level of irrigation separately.

Table 1. Shoot extension (cm), Net Photosynthesis (A, $\mu\text{mol m}^{-2} \text{s}^{-1}$), Transpiration (E, $\text{mmol m}^{-2} \text{s}^{-1}$), Water Use Efficiency (WUE) and leaf greenness index (Chl, Unit SPAD) in *Tilia* spp grown under different irrigation regimes. Data are reported as annual means (five sampling dates in 2006, five in 2007), and subjected to two-way ANOVA.

	Shoot extension (cm)		A ($\mu\text{mol m}^{-2} \text{s}^{-1}$)		E ($\text{mmol m}^{-2} \text{s}^{-1}$)		WUE ($\mu\text{mol CO}_2 / \text{mmol H}_2\text{O}$)		Chl (Unit SPAD)	
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007
	Species									
<i>T. platyphyllos</i>	30.2	45.4b	4.6d	8.1b	1.9	2.8	2.4c	2.9	38.6d	43.5c
<i>T. cordata</i>	66.4	50.2b	9.8a	10.3a	3.2	3.4	3.1a	3.0	49.1a	50.8b
<i>T. × europaea</i>	35.9	58.6a	5.9c	10.5a	2.4	3.4	2.5c	3.1	43.4c	51.9b
<i>T. tomentosa</i>	29.9	42.9b	8.2b	10.3a	3.0	3.4	2.8b	3.0	46.9b	55a
P	**	**	**	**	**	**	**	N.S.	**	**
	Irrigation									
Irrigated	43.1	53.4	7.3	10.1	2.7	3.4	2.7	3.0	44.0	50.3
Nonirrigated	38.2	43.1	6.6	9.5	2.4	3.1	2.8	3.1	45.0	50.3
P	*	**	*	N.S.	**	**	N.S.	N.S.	N.S.	N.S.
	Species × Irrigation									
P	**	N.S.	N.S.	N.S.	**	*	N.S.	N.S.	N.S.	N.S.

Different letters within the same column indicate statistically differences among species at $P < 0.05$ (*) or $P < 0.01$ (**) using Duncan's MRT.

T. tomentosa had lower E than *T. cordata* when irrigation was withheld, but E did not differ between the two species in well-watered conditions. *T. platyphyllos* and *T. × europaea* had lower E than the former two cultivars when irrigation was withheld. Irrigation increased E in *T. × europaea* but not in *T. platyphyllos*. In 2007, *T. cordata* had higher E than *T. platyphyllos* in non irrigated conditions. In the presence of irrigation, *T. tomentosa*

showed higher E than *T. × europaea*. Among the species studied, the lowest E was measured in *T. platyphyllos* (Figure 2).

In 2006, *Acer platanoides* 'Deborah' had higher shoot extension, mean annual A, and leaf chlorophyll content than the other two cultivars (Table 2). No differences were found for WUE and stem diameter (data not shown). In 2007, cultivar 'Deborah' demonstrated faster growth than the other cultivars. There were no differences in A and chlorophyll content among the cultivars in 2007. Irrigation had no effect on shoot extension, trunk diameter growth (data not shown) and chlorophyll content in 2006 (Table 2), but increased mean annual A and E and decreased WUE. In 2007, irrigation reduced shoot growth. In 2007, mean annual A was unaffected by irrigation and mean annual E was increased. In spite of this, no difference in WUE was found between irrigated and nonirrigated plants. Significant interaction between factors was found for E in 2006 and 2007. In 2006, nonirrigated 'Deborah' maple had higher E than nonirrigated 'Emerald Queen' which, in turn, had higher E than nonirrigated 'Summershade'. Where irrigation was applied, 'Deborah' had higher E than the other two cultivars (Figure 3). In 2007, nonirrigated 'Deborah' had higher E than the other two cultivars. In the presence of irrigation, differences among cultivars disappeared in 2007.




Figure 3. Means separation of those parameters [transpiration (E, $\text{mmol s}^{-1} \text{m}^{-2}$) in 2006 and 2007], which were significantly affected by cultivar × irrigation interaction in *Acer platanoides*. In this case, Duncan's Multiple Range Test was used to compare the cultivars for each level of irrigation separately.

Species/Cultivar Strategies to Cope with Water Stress

Drought tolerance of the different species was evaluated comparing chlorophyll fluorescence of different individuals of the same species/cultivar grown in well-watered or water-shortage conditions (Angelopoulos et al. 1996; Li et al. 2006). Chlorophyll content was not affected by drought treatment in linden species in 2006 and 2007 (Table 3).

F_0 is the minimal fluorescence of dark-adapted leaves and this value was used to quantify the detrimental effects of drought on leaf tissues. In *T. cordata*, *T. × europaea* and *T. tomentosa*, F_0 was unaffected by water shortage (Table 3). Contrary to the other species, nonirrigated *T. platyphyllos* had higher F_0 on July 12, 2007 if compared to irrigated plants. Increases in F_0 have been reported in *Olea*, *Quercus*, and some *Acer* and *Fraxinus* genotypes in response to water and other environmental and chemical stresses (Angelopoulos et al. 1996; Percival et al. 2003; Percival 2005; Percival et al.

Table 2. Shoot extension (cm), Net Photosynthesis (A. $\mu\text{mol m}^{-2} \text{s}^{-1}$), Transpiration (E. $\text{mmol m}^{-2} \text{s}^{-1}$), Water Use Efficiency (WUE) and leaf chlorophyll content (Chl, Unit SPAD) in different cultivars of *Acer platanoides* grown under different irrigation regimes. Data are reported as annual means (five sampling dates in 2006, five in 2007), and subjected to two-way ANOVA.

	Shoot extension (cm)		A ($\mu\text{mol m}^{-2} \text{s}^{-1}$)		E ($\text{mmol m}^{-2} \text{s}^{-1}$)		WUE ($\mu\text{mol CO}_2 / \text{mmol H}_2\text{O}$)		Chl (Unit SPAD)		
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007	
	Cultivar										
'Summershade'	30.1b	48.4c	6.3c	8.4	2.2	2.8	2.9	3.0	35.3c	42.8	
'Deborah'	45.7a	130.9a	8.9a	8.5	3.0	3.1	3.0	2.7	42.8a	44.5	
'Emerald Queen'	23.7c	95.9b	7.6b	8.8	2.5	2.8	3.0	3.1	39.3b	43	
P	**	**	**	N.S.	**	*	N.S.	N.S.	**	N.S.	
	Irrigation										
Irrigated	32.4	83.0	7.9	8.9	2.8	3.1	2.8	2.9	39.8	43.9	
Nonirrigated	34.4	100.3	7.3	8.2	2.3	2.8	3.2	2.9	38.6	42.9	
P	N.S.	**	**	N.S.	**	**	*	N.S.	N.S.	N.S.	
	Species \times Irrigation										
P	N.S.	N.S.	N.S.	N.S.	*	*	N.S.	N.S.	N.S.	N.S.	

Different letters within the same column indicate statistically differences among cultivars at $P < 0.05$ (*) or $P < 0.01$ (**), using Duncan's MRT.

2006). From a theoretical point of view, an increase in F_0 can be interpreted as a stress-induced reduction of the rate constant of energy trapping by PSII centers, which result from a physical dissociation of light harvesting complexes from PSII reaction center (Armond et al. 1980; Havaux 1993). Despite the possibility of being influenced by the difference in optical properties between control and stressed leaves (Angelopoulos et al. 1996), drought induced variations of F_0 were the best predictors ($r = 0.88$) of grain yield of barley plants exposed to water-stressing conditions (Li et al. 2006). Moreover, Percival and Sheriffs (2002), in a study on drought tolerance, found that F_0 was an earlier responding parameter than F_v/F_m and F_v/F_0 to water stress in some woody perennials.

F_v/F_m is a quantitative measure of the maximal photochemical efficiency of PSII (Willits and Peet 2001). F_v/F_m was lower in July and August 2007 in nonirrigated *T. platyphyllos* if compared to irrigated ones. F_v/F_m was unaffected by drought treatment in *T. cordata* and *T. tomentosa* (Table 3). In August 2007, F_v/F_m was higher in nonirrigated *T. tomentosa* if compared to irrigated plants. Reductions of F_v/F_m have been observed in drought-sensitive barley and *Fraxinus* genotypes in response to water stress (Li et al. 2006; Percival et al. 2006).

F_v/F_0 is the maximum primary yield of photochemistry of PSII. F_v/F_0 was not affected by withholding irrigation in *Tilia cordata*, *T. \times europaea*; and *T. tomentosa*. *T. platyphyllos* showed a significant decline of F_v/F_0 in nonirrigated conditions (Table 3). Similar results were obtained in water-stressed woody perennials and in drought-sensitive barley genotypes (Percival and Sheriffs 2002; Li et al. 2006).

With regard to leaf gas exchange, the different *Tilia* species responded to water shortage in different ways (Table 4). In *T. platyphyllos*, which had the lowest photosynthetic potential and WUE, no difference appeared between irrigated and nonirrigated plants for mean annual A, E, and g_s . In *T. cordata*, water shortage affected only mean annual E in 2006. In *T. \times europaea*, water shortage reduced mean annual E, g_s and A in 2006, but differences were not found in 2007. In particular for this species, a 14% decrease of g_s caused a 16% and 17% reduction of E and A, respectively. The greater decline in A than g_s observed in 2006 for this species suggests that non-stomatal factors limited photosynthesis of nonirrigated plants, in agreement with that reported by previous research (Davies and Kozlowski 1977; Gazal and Kubiske 2004). In *T. tomentosa*, water shortage affected leaf gas exchange in both years. Respectively

Table 3. Chlorophyll content and chlorophyll fluorescence (F_0 , F_v/F_m and F_v/F_0) in *Tilia* spp. grown with or without irrigation.

Irrigation	Chl content		F_0 Jul/06	Jun/07	Jul/07	Aug/07	F_v/F_m				F_v/F_0			
	2006	2007					Jul/06	Jun/07	Jul/07	Aug/07	Jul/06	Jun/07	Jul/07	Aug/07
	<i>T. platyphyllos</i>													
Irrigated	38.7	44.4	247.5	206.9	220.0	209.3	0.763	0.774	0.750	0.760	3.23	3.45	2.99	3.15
Nonirrigated	38.6	42.6	257.0	205.4	239.0	215.4	0.748	0.774	0.720	0.740	3.15	3.45	2.69	2.88
P	N.S.	N.S.	N.S.	N.S.	**	N.S.	N.S.	N.S.	*	*	N.S.	N.S.	*	*
	<i>T. cordata</i>													
Irrigated	42.5	51.5	300.0	224.3	241.8	224.5	0.796	0.778	0.758	0.765	3.95	3.58	3.18	3.29
Nonirrigated	44.3	52.3	275.5	217.4	246.4	222.7	0.793	0.767	0.744	0.759	3.84	3.31	2.93	3.16
P	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.
	<i>T. \times europaea</i>													
Irrigated	42.5	51.5	285.8	211.5	239.1	219.8	0.777	0.776	0.749	0.755	3.59	3.47	3.05	3.11
Nonirrigated	44.3	52.3	293.0	219.8	239.5	216.4	0.785	0.778	0.717	0.744	3.67	3.51	2.62	2.95
P	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.
	<i>T. tomentosa</i>													
Irrigated	45.5	56.3	322.0	238.0	271.4	253.7	0.797	0.749	0.718	0.733	4.00	3.10	2.42	2.80
Nonirrigated	48.3	53.7	330.8	237.0	270.3	248.7	0.798	0.751	0.736	0.758	4.00	3.13	2.84	3.14
P	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	*	N.S.	N.S.	N.S.	N.S.

* and ** indicate statistical differences at $P < 0.05$ (*) or $P < 0.01$ (**), between irrigated and nonirrigated plants of the same species.

Table 4. Net Photosynthesis (A, $\mu\text{mol m}^{-2} \text{s}^{-1}$), Transpiration (E, $\text{mmol m}^{-2} \text{s}^{-1}$) and Stomatal conductance (gs, $\text{mmol m}^{-2} \text{s}^{-1}$) in *Tilia* spp., grown with or without irrigation. Data are reported as means of the values obtained from June to September (five sampling dates in 2006, five in 2007), and subjected to analysis of variance.

Treatment	A ($\mu\text{mol m}^{-2} \text{s}^{-1}$)		E ($\text{mmol m}^{-2} \text{s}^{-1}$)		Gs ($\text{mmol m}^{-2} \text{s}^{-1}$)	
	2006	2007	2006	2007	2006	2007
	<i>T. platyphyllos</i>					
Irrigated	4.4	8.1	1.8	2.8	59.4	140.6
Nonirrigated	4.5	8.1	1.9	2.8	62.9	139.0
P	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.
	<i>T. cordata</i>					
Irrigated	10.1	10.6	3.4	3.5	124.0	187.6
Nonirrigated	9.2	10.1	3.0	3.2	114.9	171.9
P	N.S.	N.S.	*	N.S.	N.S.	N.S.
	<i>T. × europaea</i>					
Irrigated	6.2	10.5	2.5	3.4	85.0	187.1
Nonirrigated	5.1	10.6	2.1	3.4	73.0	192.8
P	**	N.S.	**	N.S.	**	N.S.
	<i>T. tomentosa</i>					
Irrigated	8.5	11.2	3.3	3.9	124.0	223.1
Nonirrigated	7.4	9.4	2.7	3.0	102.0	172.8
P	*	**	*	**	**	**

* and ** indicate statistical differences at $P < 0.05$ (*) or $P < 0.01$ (**) between irrigated and nonirrigated plants of the same species.

Table 5. Chlorophyll content and chlorophyll fluorescence (F_0 , F_v/F_m and F_v/F_0) in different cultivars of *Acer platanoides* grown with or without irrigation.

Irrigation	Chl content		F_0				F_v/F_m				F_v/F_0			
	2006	2007	Jul/06	Jun/07	Jul/07	Aug/07	Jul/06	Jun/07	Jul/07	Aug/07	Jul/06	Jun/07	Jul/07	Aug/07
	‘Summershade’													
Irrigated	35.3	43.0	238.0	243.7	257.0	225.5	0.775	0.775	0.738	0.736	3.50	3.47	2.83	2.83
Nonirrigated	35.3	42.6	270.0	240.7	288.0	247.3	0.778	0.768	0.701	0.719	3.52	3.33	2.47	2.63
P	N.S.	N.S.	*	N.S.	*	*	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	*	*
	‘Deborah’													
Irrigated	44.6	45.9	237.0	244.6	266.7	235.7	0.782	0.764	0.733	0.732	3.70	3.24	2.80	2.75
Nonirrigated	41.1	43.6	235.4	239.8	266.6	236.4	0.784	0.765	0.71	0.713	3.60	3.30	2.62	2.68
P	*	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.
	‘Emerald Queen’													
Irrigated	39.3	43.4	231.8	222.3	269.0	228.9	0.779	0.774	0.722	0.731	3.60	3.46	2.52	2.73
Nonirrigated	39.4	42.6	236.2	232.4	261.2	241.9	0.778	0.768	0.708	0.713	3.60	3.33	2.62	2.64
P	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.

* and ** indicate statistical differences at $P < 0.05$ (*) or $P < 0.01$ (**) between irrigated and nonirrigated plants of the same cultivar.

Table 6. Net Photosynthesis (A, $\mu\text{mol m}^{-2} \text{s}^{-1}$), Transpiration (E, $\text{mmol m}^{-2} \text{s}^{-1}$) and Stomatal Conductance (gs, $\text{mmol m}^{-2} \text{s}^{-1}$) in different cultivars of *Acer platanoides*, grown with or without irrigation. Data are reported as means of the values obtained from June to September (five sampling dates in 2006, five in 2007), and subjected to analysis of variance.

Treatment	A ($\mu\text{mol m}^{-2} \text{s}^{-1}$)		E ($\text{mmol m}^{-2} \text{s}^{-1}$)		Gs ($\text{mmol m}^{-2} \text{s}^{-1}$)	
	2006	2007	2006	2007	2006	2007
	‘Summershade’					
Irrigated	6.6	8.9	2.6	3.1	118.2	175.2
Nonirrigated	6.1	7.8	1.9	2.5	93.0	148.9
P	N.S.	N.S.	**	**	**	N.S.
	‘Deborah’					
Irrigated	9.5	9.0	3.2	3.1	163.4	179.0
Nonirrigated	8.4	8.6	2.7	3.2	147.0	194.3
P	**	N.S.	**	N.S.	**	N.S.
	‘Emerald Queen’					
Irrigated	7.8	8.9	2.6	3.0	124.7	174.5
Nonirrigated	7.5	8.1	2.4	2.7	117.5	124.7
P	N.S.	N.S.	N.S.	*	N.S.	N.S.

* and ** indicate statistical differences at $P < 0.05$ (*) or $P < 0.01$ (**) between irrigated and nonirrigated plants of the same cultivar.

for 2006 and 2007, a reduction in g_s of 17% and 22% decreased E by 19% and 22%, and reduced A by 12% and 16%. WUE was never affected by water regime in *Tilia* species (data not shown).

Chlorophyll content of maple cultivars was unaffected by irrigation except for 'Deborah' in 2006, when irrigated plants had higher leaf chlorophyll content than nonirrigated plants (Table 5).

F_0 , Fv/Fm, and Fv/F₀ were unaffected by water regime in cultivars 'Deborah' and 'Emerald Queen'. In July and August 2007,

nonirrigated 'Summershade' had higher F_0 and lower Fv/F₀ than in the irrigated treatment, although no difference was found for Fv/Fm. Considering the different maple cultivars, water shortage never affected mean annual A, except for 'Deborah' in 2006, when irrigated plants fixed 11% more atmospheric carbon than nonirrigated ones (Table 6). No difference was found in 2007.

If compared to irrigated plants, nonirrigated maples had lower mean annual E in both years (cv. 'Summershade'); only in 2006




Figure 4. Daily trend of net assimilation (A, $\mu\text{mol m}^{-2} \text{s}^{-1}$) and transpiration (E, $\text{mmol m}^{-2} \text{s}^{-1}$) in the four linden species grown in well watered and water shortage conditions. Data are the average between the two sampling dates (July 20 and 27, 2007). Single asterisk (*) and double-asterisks (**) in the right-top corner of each graph indicate statistically differences at $P < 0.05$ (*) or $P < 0.01$ (**), using ANOVA test for the factors I (irrigation) and T (time).




Figure 5. Daily trend of net assimilation (A, $\mu\text{mol m}^{-2} \text{s}^{-1}$) and transpiration (E, $\text{mmol m}^{-2} \text{s}^{-1}$) in the three Norway maple cultivars grown in well watered and water shortage conditions. Data are the average between the two sampling dates (July 20 and 27, 2007). Single asterisk (*) and double-asterisks (**) in the right-top corner of each graph indicate statistically differences at $P < 0.05$ (*) or $P < 0.01$ (**), using ANOVA test for the factors I (irrigation) and T (time).

(cv. 'Deborah') or only in 2007 (cv. 'Emerald Queen'). g_s was affected by water regime only in 2006 in the cultivars 'Summershade' and 'Deborah'.

Daily gas exchange trend, measured on July 20 and 27, 2007, differed among species/cultivar (Figure 4; Figure 5). A was not affected by water regime except for *T. tomentosum*, which showed higher carbon assimilation under irrigated conditions (Figure 4). A was affected by sampling time in all linden species, generally reaching the minimum between 12.00 and 14.00 h. Some recovery was observed at 18.00 h only in irrigated *T. platyphyllos* and nonirrigated *T. cordata* and *T. tomentosum*. E was not influenced by water conditions in *T. platyphyllos* and *T. × europaea*. Irrigated *T. tomentosum* and *T. cordata* transpired more water than nonirrigated plants of the same species, but in littleleaf linden significant differences between irrigation treatments appeared only at 9.00 h. In irrigated *T. tomentosum*, E remained fairly constant during the day, while in nonirrigated silver linden a significant drop in E occurred at 12.00 h and 14.00 h. In this species, E recovered completely at 18.00 h.

In *Acer platanoides* 'Summershade' and 'Emerald Queen', irrigation affected gas exchange, with irrigated plants having higher A and E than nonirrigated plants for the entire day (Figure 3). On the contrary, in *Acer platanoides* 'Deborah' irrigation treatment had no influence on A and E throughout the day. Sampling time influenced A and E in all cultivars (Figure 5). A reached its maximum at 9.00 h in all cultivars. In *Acer platanoides*, 'Summershade' and 'Deborah', A kept declining during the day, reaching its minimum at 17.00 h. In *Acer platanoides*, 'Emerald Queen' a drop in A was observed at 11.00 h, then there was a subsequent recovery until 15.00 h and a final drop at 17.00 h. E was also influenced by sampling time in all the cultivars. In *Acer platanoides*, 'Summershade' the increase in E observed from 9.00 h to 13.00 h was followed by a drop in the afternoon. This drop was particularly evident in nonirrigated *Acer platanoides* 'Summershade'. No clear trend was observed in the 'Emerald Queen' cultivar.

CONCLUSION

Norway maple establishment was promoted by irrigation for two years after transplant. After this period, irrigation is probably not needed anymore. Linden establishment was promoted by irrigation also in the third growing season after transplant, while more research is needed to determine for how long *Tilia* should be irrigated after planting in the landscape. Only in *T. platyphyllos* and *Acer* ‘Summershade’ did water shortage cause a significant increase in F_0 and decline in the primary yield of PSII. In *T. platyphyllos*, F_v/F_m was significantly decreased by drought. These effects on chlorophyll fluorescence parameters have been related to the occurrence of water or other environmental and chemical stresses. On this basis, these two species were considered the least tolerant to water stress during the establishment phase. Different responses to water shortage were found in the different species and cultivars. *T. platyphyllos* had lower gas exchange than the other species, even in well-watered conditions and stomatal conductance, carbon assimilation and transpiration were unaffected by water shortage. This strategy was also found in beech seedlings from dry (rainfall 580–680 mm/year) provenances (Pueke et al. 2002), but in the case of linden, it failed to prevent disruption of PSII under nonirrigated conditions. *T. cordata*, *T. × europaea*, *Acer* ‘Deborah’ and ‘Emerald Queen’ showed, in 2007, higher gas exchange which was little affected by water shortage. This “water spending” strategy may depend on a better access to soil water thanks to a more developed root system (Levitt 1972; Mediavilla and Escudero 2004). This is also confirmed by the fact that before full establishment (2006, two years from transplant) effects of water shortage on leaf gas exchange were observed—especially in *T. × europaea* and *Acer platanoides* ‘Deborah’. *T. tomentosa*—showed high gas exchange, but, in response to water shortage, stomatal conductance and transpiration were reduced. A similar strategy was found on *Quercus coccinea* and classified as “drought avoidance based on water saving” by Sakcali and Ozturk (2004).

The results obtained in this experiment may be a useful tool for species/cultivar selection for urban forestry. Among lindens, *T. tomentosa* seems particularly adapted to those situations where root development and root access to water is limited by small planting pits (e.g., street trees). *T. cordata*, which showed the highest gas exchange and water use efficiency in the year following transplant, seems well suited to perform well where access to underground water is not restricted. Differences among Norway maple cultivars seem smaller than those observed among lindens (no significant differences in F_v/F_m). Regardless, in the establishment phase, *A. platanoides* ‘Deborah’ and ‘Emerald Queen’ seem better performers than *Acer* ‘Summershade’.

Acknowledgments. This work has been undertaken as a research project with the title “Tecniche eco-compatibili di gestione del vivaismo e del verde ornamentale” financed by Regione Lombardia - Agricultural Department, according to the Plan of Research and Development 2004.

LITERATURE CITED

- Abrams, M.D., and M.E. Kubiske. 1990. Photosynthesis and water relations during drought in *Acer rubrum* L. genotypes from contrasting sites in central Pennsylvania. *Functional Ecology* 4:727–733.
- Akbari, H. 2002. Shade trees reduce building energy use and CO₂ emissions from power plants. *Environmental Pollution* 116:S119–S126.
- Angelopoulos, K., B. Dichio, and C. Xiloyannis. 1996. Inhibition of photosynthesis in olive trees (*Olea europaea* L.) during water stress and rewatering. *Journal of Experimental Botany* 47(301):1093–1100.
- Armond, P.A., O. Björkman, and L.A. Staehelin. 1980. Dissociation of supramolecular complexes in chloroplast membranes: a manifestation of heat damage to the photosynthetic apparatus. *Biochimica et Biophysica Acta* 601:433–442.
- Brack, C.L. 2002. Pollution mitigation and carbon sequestration by an urban forest. *Environmental Pollution* 116:S195–S200.
- Chew, V. 1976. Uses and abuses of Duncan’s Multiple Range Test. *Proc. Fla. State Hort. Soc.* 89:251–253.
- Daugherty, R. 2002. Untitled. Front Range Community College, Westminster, CO, www.timberlinelandscaping.com/Rev'dPlants-FRCC-TL/TreesPDFs.
- Davies, W.J., and T.T. Kozlowski. 1977. Variations among woody plants in stomatal conductance and photosynthesis during and after growth. *Plant and Soil* 46:435–444.
- Elmendorf, W. 2008. The importance of trees and nature in community: a review of the relative literature. *Arboriculture & Urban Forestry* 34(3):152–156.
- Escobedo, F.J., J.E. Wagner, D.J. Nowak, C.L. De la Maza, M. Rodriguez and D.E. Crane. 2008. Analyzing the cost-effectiveness of Santiago, Chile’s policy of using urban forest to improve air quality. *Journal of Environmental Management* 86:148–157.
- Ferrini, F., A. Fini, P. Frangi, and G. Amoroso. 2008. Mulching of ornamental trees: effects on growth and physiology. *Arboriculture & Urban Forestry* 34(3):157–162.
- Fini, A., and F. Ferrini. 2007. Influenza dell’ambiente urbano sulla fisiologia e la crescita degli alberi. *Italus Hortus* 14(1):9–24.
- Fini, A., G.B. Mattii, and F. Ferrini. 2008. Physiological responses to different irrigation regimes for shade trees grown in container. *Advances in Horticultural Sciences* 22(1):13–20.
- Fisher, G., M. Shah, H. van Veuiluzen, and F.O. Nachtergaele. 2001. Global agro-ecological assessment for agriculture in the 21st Century. Laxenburg, Austria: IIASA and FAO.
- Gazal R.M., and M.E. Kubiske. 2004. Influence of initial root length on physiological responses of cherrybark oak and Schumard oak seedlings to field drought conditions. *Forest Ecology and Management* 189:295–305.
- Gilbertson, P., and A.D. Bradshaw. 1985. Tree survival in the cities: the extent and nature of the problem. *Arboricultural Journal* 9:131–142.
- Havaux, M. 1993. Characterization of thermal damage to photosynthetic electron transport system in potato leaves. *Plant Science* 94:19–33.
- Kaushal, P., and G. Aussenac. 1989. Transplant shock in corsican pine and cedar of atlas seedlings: internal water deficits, growth and root regeneration. *Forest Ecology and Management* 27:29–40.
- Lazár, D. 2006. The polyphasic chlorophyll a fluorescence rise measured under high intensity of exciting light. *Functional Plant Biology* 33:9–30.
- Levitt, J. 1972. Responses of plants to environmental stresses. Academic Press, New York, 697 pp.
- Li, R., P. Guo, M. Baum, S. Grando, and S. Ceccarelli. 2006. Evaluation of chlorophyll content and fluorescence parameters as indicators of drought tolerance in barley. *Agricultural Sciences in China* 5(10):751–757.
- Maxwell, K., and G.N. Johnson. 2000. Chlorophyll fluorescence - a practical guide. *Journal of Experimental Botany* 51:659–668.
- McPherson, E.G. 2003. A benefit-cost analysis of ten street tree species in Modesto, California, U.S. *Journal of Arboriculture* 29(1):1–8.

- Mediavilla, S., and A. Escudero. 2004. Stomatal responses to drought of mature trees and seedlings of two co-occurring Mediterranean oaks. *Forest Ecology and Management* 187:281–294.
- Miller, R.H., and R.W. Miller. 1991. Planting survival of selected street tree taxa. *Journal of Arboriculture* 17(7):185–191.
- Montague, T., R. Kjelgren, and L. Rupp. 2000. Gas exchange and growth of two transplanted, field grown tree species in an arid climate. *HortScience* 35:763–768.
- Nowak, D.J., J.R. McBride, and R.A. Beatty. 1990. Newly planted street tree growth and mortality. *Journal of Arboriculture* 16(5):124–129.
- Nowak, D.J., R. Hoehn, and D.E. Crane. 2007. Oxygen production by urban trees in the United States. *Arboriculture and Urban Forestry* 33(3):220–226.
- Pardossi, A., L. Incrocci, and P. Marzialetti. 2004. La razionalizzazione dell'irrigazione nel florovivaismo: una sintesi. In ARSIA (Eds.), *Uso razionale delle risorse nel florovivaismo: l'acqua*, Florence, Italy, 285 pp.
- Pauleit, S., N. Jones, G. Garcia-Martin, J.L. Garcia-Valdecantos, L.M. Rivière, L. Vidal-Beaudet, M. Bodson, and T.B. Raundrup. 2002. Tree establishment practice in towns and cities – Results from a European survey. *Urban Forestry & Urban Greening* 1:83–96.
- Percival, G.C., and G.A. Fraser. 2001. Measurement of the salinity and freezing tolerance of *Crataegus* genotypes using chlorophyll fluorescence. *Journal of Arboriculture* 27(5):233–245.
- Percival, G.C., and C.N. Sheriffs. 2002. Identification of drought-tolerance woody perennials using chlorophyll fluorescence. *Journal of Arboriculture* 28(5):215–223.
- Percival, G.C., G.A. Fraser, and G. Oxenham. 2003. Foliar salt tolerance of *Acer* genotypes using chlorophyll fluorescence. *Journal of Arboriculture* 29(2):61–65.
- Percival, G.C. 2005. The use of chlorophyll fluorescence to identify chemical and environmental stress in leaf tissue of three oak (*Quercus*) species. *Journal of Arboriculture* 31(5):215–227.
- Percival, G.C., I.P. Keary, and S. AL-Habsi. 2006. An assessment of the drought tolerance of *Fraxinus* genotype for urban landscape planting. *Urban Forestry & Urban Greening* 5(1):17–27.
- Percival, G.C., I.P. Keary, and K. Noviss. 2008. The potential of a chlorophyll content SPAD meter to quantify nutrient stress in foliar tissue of sycamore (*Acer pseudoplatanus*), English oak (*Quercus robur*), and european beech (*Fagus sylvatica*). *Arboriculture & Urban Forestry* 34(2):89–100.
- Pueke A.D., C. Schraml, W. Hartung, and H. Rennenberg. 2002. Identification of drought-sensitive beech ecotypes by physiological parameters. *New Phytologist* 154:373–387.
- Sakcali M.S., and M. Ozturk. 2004. Eco-physiological behaviour of some Mediterranean plants as suitable candidates for reclamation of degraded areas. *Journal of Arid Environments* 57:1–13.
- Scheiber, S.M., E.F. Gilman, M. Paz, and K.A. Moore. 2007. Irrigation affects landscape establishment of Burford holly, pittosporum and sweet viburnum. *HortScience* 42(2):344–348.
- UNEP/IUC. 1999. United Nations Framework Convention on Climate Change Information kit. Climate Change information sheet. Williams M. (Eds.), Geneva, Chatelaine, Switzerland, 40 pp.
- Willits D.H., and M.M. Peet. 2001. Using chlorophyll fluorescence to model leaf photosynthesis in greenhouse pepper and tomato. *Acta Horticulturae* 507:311–315.
- Whitlow, T.H., N.L. Bassuk, and D.L. Reichert. 1992. A 3-year study of water relations of urban street trees. *Journal of Applied Ecology* 29(2):436–450.
- Yamada, M., D. Hidaka, and H. Fukamachi. 1996. Heat tolerance in leaves of tropical fruit crops as measured by chlorophyll fluorescence. *Scientia Horticulturae* 67:39–48.
- Zwack, J.A., W.R. Graves, and A.M. Townsend. 1998. Leaf water relations and plant development of three Freeman maple cultivars subjected to drought. *Journal of American Society of Horticulture Science* 123(3):371–375.

Alessio Fini

*Dipartimento di Ortoflorofrutticoltura
Viale delle Idee, 30
Sesto Fiorentino (FI), 50019 Italy
alessio.fini@unifi.it*

*Francesco Ferrini (corresponding author)
Dipartimento di Ortoflorofrutticoltura
Viale delle Idee, 30
Sesto Fiorentino (FI), 50019 Italy
francesco.ferrini@unifi.it*

*Piero Frangi
Fondazione Minoprio
Centro Mirt
Viale Raimondi, 54
Vertemate con Minoprio (CO), 22070 Italy
frangi@fondazioneminoprio.it*

*Gabriele Amoroso
Fondazione Minoprio
Centro Mirt
Viale Raimondi, 54
Vertemate con Minoprio (CO), 22070 Italy
g.amoroso@fondazioneminoprio.it*

*Riccardo Piatti
Fondazione Minoprio
Centro Mirt
Viale Raimondi, 54
Vertemate con Minoprio (CO), 22070 Italy
r.piatti@fondazioneminoprio.it*

Zusammenfassung. Das Ziel dieser Arbeit war, die Trockenheitsresistenz von verschiedenen Lindenarten und verschiedenen Ahorn-Kultivaren während der Anpflanzphase zu untersuchen und den Bewässerungseffekt auf ihr Wachstum und Physiologie zu bewerten. Im Winter 2004/05 wurden 168 Bäume mit Stammumfang von 8-10 cm (*Tilia platyphyllos*, *T. cordata*, *T. × europaea*, *T. tomentosa*, *Acer platanoides* 'Summershade', *A. platanoides* 'Deborah', und *A. platanoides* 'Emerald Queen' im Feld ausgepflanzt. 84 Pflanzen wurden mit einer Tröpfchenbewässerung (4l/h) bewässert und 84 erhielten kein Wasser. Höhe, Stammdurchmesser und Triebblängen wurden am Ende der Wachstumsperiode 2005, 2006 und 2007 gemessen. Blattgasaustausch und Chlorophyllfluoreszenz wurden monatlich während der Wachstumsperiode in 2006 und 2007 gemessen. Der Blattgrün-Index-Gehalt wurde in 2006 und 2007 gemessen. Die Resultate zeigen, dass *T. tomentosa* and *T. cordata* mehr stressresistent sind während der Anwachsphase als *T. platyphyllos*, während *Acer platanoides* 'Summershade' weniger stressresistent während der Anwachsphase ist als die Kultivare 'Emerald Queen' and 'Deborah'.

Resumen. La intención de este trabajo fue investigar la tolerancia a la sequía de especies de Tilia y de diferentes cultivares de *Acer platanoides* crecidos durante la fase de establecimiento, y evaluar el efecto del riego en su fisiología y crecimiento. En el invierno de 2004-2005, 168 árboles [8-10 cm (3-4 pulg) circunferencia] de *Tilia platyphyllos*, *T. cordata*, *T. × europaea*, *T. tomentosa*, *Acer platanoides* 'Summershade', *A. platanoides* 'Deborah', y *A. platanoides* 'Emerald Queen' fueron plantados en el campo. Ochenta y cuatro plantas fueron regadas con sistema por goteo (4 l/h) y otros 84 no lo fueron. Se midió altura, diámetro del tronco y elongación de los brotes al final de la estación de crecimiento en 2005, 2006 y 2007. El intercambio de gases en las hojas y la fluorescencia de clorofila fueron medidas mensualmente durante las estaciones de crecimiento de 2006 y 2007. El índice de vigor contenido en las hojas fue medido en 2006 y 2007. Los resultados indican que *T. tomentosa* y *T. cordata* son más tolerantes a la sequía durante el establecimiento que *T. platyphyllos*, mientras que *Acer platanoides* 'Summershade' es menos tolerante a la sequía durante el establecimiento que los cultivares 'Emerald Queen' y 'Deborah'.