

SNYDER GEOMETRY AND QUANTUM FIELD THEORY

P. VALTANCOLI

*Dipartimento di Fisica, Polo Scientifico Università di Firenze and INFN,
Sezione di Firenze, Via G. Sansone 1, 50019 Sesto Fiorentino, Italy
valtancoli@fi.infn.it*

Received 2 July 2012

Revised 6 November 2012

Accepted 16 November 2012

Published 11 December 2012

We find that, in presence of the Snyder geometry, the notion of translational invariance needs to be modified, allowing a momentum dependence of this symmetry. This step is necessary to build the maximally localized states and the Feynman rules of the corresponding quantum field theory.

Keywords: Noncommutative geometry; minimal position uncertainty.

PACS number: 11.10.Nx

1. Introduction

At the Planck scale we expect that the gravitational effects of the test particle's energy modify the structure of space-time, inducing a finite limit Δx_0 on the possible resolution of distances. String theory, the leader candidate to describe the quantum gravitational phenomena, suggests also a certain type of correction to the uncertainty relation¹⁻⁵

$$\Delta x \Delta p \geq \frac{\hbar}{2} (1 + \beta (\Delta p)^2), \quad \beta > 0, \quad (1.1)$$

implying a nonzero minimal uncertainty Δx_0 . This uncertainty relation can be modeled by introducing a correction term to the commutation relations:

$$[\hat{x}, \hat{p}] = i\hbar(1 + \beta \hat{p}^2). \quad (1.2)$$

In the case of n -dimensions, the simplest generalization of this algebra gives rise to the Snyder geometry.⁶ Snyder introduced it as a UV regularization of quantum field theory but he did not relate it to the notion of minimal position uncertainty.

In this paper, we study in detail the implications that this algebra induces on quantum field theory. First, we need to modify the translational invariance, which is broken by this algebra, in such a way as to make covariant with it.

This concept allows us to build the maximally localized states which carry the maximal quantum information compatible with this algebra. These states are then necessary to compute the basic Feynman rules, as the vertex and the propagator. In particular the propagator $G(\xi - \eta)$ is no more singular in the limit $\xi \rightarrow \eta$, avoiding the UV divergencies, typical of quantum field theory.

2. The Algebra

Snyder algebra is simply obtained modifying the representation of the position operator in the following way (for simplicity we work with an Euclidean metric)

$$\hat{x}^\mu = i\hbar[\delta^{\mu\nu} + \beta p^\mu p^\nu] \frac{\partial}{\partial p^\nu}, \quad (2.1)$$

from which we obtain the following commutation relations:

$$[\hat{x}^\mu, \hat{p}^\nu] = i\hbar[\delta^{\mu\nu} + \beta \hat{p}^\mu \hat{p}^\nu], \quad [\hat{x}^\mu, \hat{x}^\nu] = i\hbar\beta[\hat{p}^\nu \hat{x}^\mu - \hat{p}^\mu \hat{x}^\nu]. \quad (2.2)$$

These define a possible n -dimensional generalization of the unique modification of one-dimensional quantum mechanics,

$$[\hat{x}, \hat{p}] = i\hbar[1 + \beta \hat{p}^2], \quad (2.3)$$

inducing a nonzero minimal uncertainty on the position measure

$$(\Delta x)^2 \geq \beta \hbar^2. \quad (2.4)$$

In the Euclidean n -dimensional case, each spatial dimension contributes with an analogous uncertainty giving rise to the following formula:

$$(\Delta x)^2 \geq n\beta \hbar^2. \quad (2.5)$$

Therefore, the physical meaning of the Snyder geometry is to introduce a spatial hypercube of side-length $\sqrt{\beta}\hbar$, which is not accessible to the position measurements. It results in a very efficient regularization of the UV divergencies of quantum field theory.

3. The ρ Variables

It is convenient to introduce a new representation of the Snyder algebra, given in terms of a new variable ρ^μ living on a compact hypersphere of radius $\frac{1}{\sqrt{\beta}}$:

$$\hat{x}^\mu = i\hbar\sqrt{1 - \beta\rho^2} \frac{\partial}{\partial \rho^\mu}, \quad 0 < \rho^2 < \frac{1}{\beta}, \quad \hat{p}^\mu = \frac{\rho^\mu}{\sqrt{1 - \beta\rho^2}}. \quad (3.1)$$

This representation will help us, in the following, to simplify all the calculations. It works also in the one-dimensional case:

$$\hat{x} = i\hbar\sqrt{1 - \beta\rho^2} \frac{\partial}{\partial \rho}, \quad 0 < \rho < \frac{1}{\sqrt{\beta}}, \quad \hat{p} = \frac{\rho}{\sqrt{1 - \beta\rho^2}}, \quad (3.2)$$

where it faithfully reproduces all the results reported in Ref. 7.

The operators \hat{x}^μ and \hat{p}^μ are symmetric with respect to the following scalar product:

$$\langle \psi(\rho) | \phi(\rho) \rangle = \int d^n \rho \frac{\psi^*(\rho) \phi(\rho)}{\sqrt{1 - \beta \rho^2}}. \quad (3.3)$$

In the momentum representation, the operator \hat{x}^μ is again symmetric with the following choice:

$$\hat{x}^\mu = i\hbar \left[\frac{\partial}{\partial p^\mu} + \beta p^\mu p^\nu \frac{\partial}{\partial p^\nu} + \beta \left(\frac{n+1}{2} \right) p^\mu \right]. \quad (3.4)$$

The corresponding scalar product is now

$$\langle \psi(p) | \phi(p) \rangle = \int d^n p \psi^*(p) \phi(p). \quad (3.5)$$

The operator \hat{x}^μ does not admit a unique self-adjoint extension, which is of course a consequence of the minimal uncertainty of the position measurement.

4. Modifying the Translation Invariance

Snyder geometry is compatible with the rotation group but not with the translation one. The fact that the coordinates \hat{x}^μ do not commute is an obstacle to the study of maximally localized states compatible with it. This is simply because the eigenvalue equation

$$\hat{x}^\mu \psi_\xi(\rho) = \xi^\mu \psi_\xi(\rho) \quad (4.1)$$

does not make sense, since the coordinates \hat{x}^μ cannot be simultaneously diagonalized, and the combination $(\hat{x}^\mu - \xi^\mu)$ is not covariant with respect to the commutation rules.

Our contribution is modifying the concept of translation in order to make it covariant with respect to the Snyder algebra, which is considered also in Ref. 8 (with a different result), and to make sense of the eigenvalue equation for \hat{x}^μ .

Our strategy is to define a momentum dependent operator analogous to the translation

$$\hat{x}_\xi^\mu = \xi^\mu f(\rho^2) + \rho^\mu (\rho \cdot \xi) g(\rho^2), \quad (4.2)$$

imposing that the combination

$$\hat{\hat{x}}^\mu = \hat{x}^\mu - \hat{x}_\xi^\mu \quad (4.3)$$

satisfies to the same commutation rules of the Snyder algebra. Moreover, we require that for $n = 1$ this operator reduces to the simple translation

$$\begin{aligned} 2f'(\rho^2) - g(\rho^2) + \frac{\beta}{1 - \beta \rho^2} f(\rho^2) &= 0, \\ f(\rho^2) + \rho^2 g(\rho^2) &= 1 \end{aligned} \quad (4.4)$$

This system can be resolved by introducing the series

$$f(\rho^2) = 1 + \sum_{n=1}^{\infty} \beta^n f_n(\rho^2)^n, \quad (4.5)$$

$$\rho^2 g(\rho^2) = \sum_{n=1}^{\infty} \beta^n g_n(\rho^2)^n,$$

from which

$$f_n = -g_n = -\frac{(2n-2)!!}{(2n+1)!!}. \quad (4.6)$$

These series can be resummed to give

$$f(\rho^2) = 1 - \rho^2 g(\rho^2) = 1 + \frac{1}{2} \int_0^1 dx \ln[1 - 4x(1-x)\beta\rho^2]. \quad (4.7)$$

An alternative representation for $f(\rho^2)$ is

$$f(\rho^2) = \sqrt{\frac{1-\beta\rho^2}{\beta\rho^2}} \arcsin \sqrt{\beta\rho^2}. \quad (4.8)$$

Coming back to the momentum variables we obtain

$$f(p^2) = \frac{1}{\sqrt{\beta p^2}} \arctan \sqrt{\beta p^2} = \int_0^1 dx \frac{1}{1 + \beta p^2 x^2}, \quad (4.9)$$

from which we can give the final representation for the translation operator

$$\hat{x}_\xi^\mu = \int_0^1 dx \left(\frac{\xi^\mu + \beta p^\mu (p \cdot \xi) x^2}{1 + \beta p^2 x^2} \right). \quad (4.10)$$

For $n = 1$ it is obvious that $\hat{x}_\xi^\mu \rightarrow \xi$.

By construction, the combination $\hat{x}^\mu - \hat{x}_\xi^\mu$ is covariant with respect to the commutation rules. Now we want to show that the following analogue of the eigenvalue equation,

$$\hat{x}^\mu \psi_\xi = \hat{x}_\xi^\mu \psi_\xi, \quad \langle \psi_\xi | \psi_\xi \rangle = 1, \quad (4.11)$$

can be solved. In fact the solution in the momentum coordinates is simply

$$\psi_\xi(p) = \frac{c}{(1 + \beta p^2)^{\frac{n+1}{4}}} e^{\frac{\xi \cdot p}{i\hbar\sqrt{\beta p^2}} \arctan \sqrt{\beta p^2}} \quad (4.12)$$

that, translated in the ρ variables, results in

$$\psi_\xi(\rho) = \left[\left(\frac{\beta}{\pi} \right)^{\frac{n}{2}} \frac{\Gamma(\frac{n+1}{2})}{\sqrt{\pi}} \right]^{\frac{1}{2}} e^{\frac{\xi \cdot \rho}{i\hbar\sqrt{\beta\rho^2}} \arcsin \sqrt{\beta\rho^2}}. \quad (4.13)$$

Let us note that the solution (4.12) is the obvious generalization for $n \neq 1$ of the state studied in Ref. 7. We will need this wave function to define the maximally localized states and the analogue of the Fourier transform.

Obviously, the scalar product of two wave functions with two independent translations is different from zero as in the case $n = 1$, due to the fuzziness of space–time:

$$\langle \psi_\xi | \psi_\eta \rangle = c^2 \int \frac{d^n \rho}{\sqrt{1 - \beta \rho^2}} e^{i \frac{(\xi - \eta) \cdot \rho}{\hbar \sqrt{\beta \rho^2}} \arcsin \sqrt{\beta \rho^2}}. \quad (4.14)$$

The angular part can be computed as

$$\int d\Omega e^{-i \frac{\alpha \cdot \rho}{\sqrt{\rho^2}} \arcsin \sqrt{\beta \rho^2}} = (2\pi)^{\frac{n}{2}} \frac{J_{\frac{n}{2}-1}(\alpha \arcsin \sqrt{\beta \rho^2})}{(\alpha \arcsin \sqrt{\beta \rho^2})^{\frac{n}{2}-1}}, \quad (4.15)$$

where $\alpha = \frac{|\xi - \eta|}{\hbar \sqrt{\beta}}$ and J_ν is the Bessel function of the first kind.

Introducing the following change of variables $\rho = \frac{1}{\sqrt{\beta}} \sin x$ the radial part of the integral (4.14) results in

$$\langle \psi_\xi | \psi_\eta \rangle = \left(\frac{2}{\alpha} \right)^{\frac{n}{2}} \frac{\alpha}{\sqrt{\pi}} \Gamma\left(\frac{n+1}{2} \right) \int_0^{\frac{\pi}{2}} dx \frac{(\sin x)^{n-1}}{x^{\frac{n}{2}-1}} J_{\frac{n}{2}-1}(\alpha x). \quad (4.16)$$

Unfortunately this integral cannot be computed for generic n . For $n = 1$ we reobtain

$$\langle \psi_\xi | \psi_\eta \rangle = \frac{\sin\left[\frac{\pi}{2}\alpha\right]}{\frac{\pi}{2}\alpha} \quad (4.17)$$

and for $n = 3$

$$\langle \psi_\xi | \psi_\eta \rangle = \frac{1}{\pi\alpha} \left[-\text{Si} \left[\frac{\pi}{2}(\alpha - 2) \right] + 2 \text{Si} \left[\frac{\pi}{2}\alpha \right] - \text{Si} \left[\frac{\pi}{2}(\alpha + 2) \right] \right]. \quad (4.18)$$

5. Maximally Localized States

After modifying the concept of translation, we are ready to build the maximally local states, i.e. the physical states with which we will build the Feynman rules of the theory. To do this, it is necessary to study the following equation:

$$(\hat{x}^\mu - \hat{x}_\xi^\mu + ik\hat{p}^\mu)\psi_k(p^2) = 0, \quad (5.1)$$

choosing the parameter k in order to minimize the uncertainty associated to $\psi_k(p^2)$. For simplicity we resolve this problem in the origin and then we apply the translation wave function discussed in Sec. 4 to define the generic case.

In the origin, Eq. (5.1) written in the variables ρ implies

$$\frac{\partial}{\partial \rho^\mu} \psi_k(\rho^2) = -\frac{k}{\hbar} \frac{\rho^\mu}{(1 - \beta \rho^2)} \psi_k(\rho^2), \quad \langle \psi_k | \psi_k \rangle = 1 \quad (5.2)$$

that is resolved by

$$\psi_k(\rho^2) = \left[\left(\frac{\beta}{\pi} \right)^{\frac{n}{2}} \frac{\Gamma\left(\frac{k}{\beta\hbar} + \frac{n+1}{2} \right)}{\Gamma\left(\frac{k}{\beta\hbar} + \frac{1}{2} \right)} \right]^{\frac{1}{2}} (1 - \beta \rho^2)^{\frac{k}{2\beta\hbar}}. \quad (5.3)$$

The indetermination on this state results in

$$(\Delta x)^2|_k = \langle \psi_k | \hat{x}^2 | \psi_k \rangle = \frac{n}{2\beta} \frac{k^2}{\left(\frac{k}{\beta\hbar} - \frac{1}{2}\right)} \rightarrow k = \beta\hbar. \quad (5.4)$$

The minimum is obtained for $k = \beta\hbar$ and the value coincides with what we discussed at the beginning of the paper:

$$(\Delta x)^2|_{\beta\hbar} = n\beta\hbar^2. \quad (5.5)$$

The translated wave function can be obtained simply by multiplying it with the solution to the modified eigenvalue problem (4.11):

$$\psi^{ml}(\rho) = \left[\left(\frac{\beta}{\pi}\right)^{\frac{n}{2}} \frac{\Gamma(\frac{n+3}{2})}{\Gamma(\frac{3}{2})} \right]^{\frac{1}{2}} (1 - \beta\rho^2)^{\frac{1}{2}} e^{\frac{\xi \cdot \rho}{i\hbar\sqrt{\beta\rho^2}} \arcsin \sqrt{\beta\rho^2}}. \quad (5.6)$$

This is our final wave function describing the maximal localization in the neighborhood of the point ξ^μ .

6. Feynman Rules

The maximally localized states are the building blocks to define the Feynman rules of the quantum field theory based on the Snyder algebra. In particular, we need to compute the vertex and the propagator (see also Ref. 9)

$$\begin{aligned} \tilde{\delta}(\xi^{ml}, \eta^{ml}) &= \langle \xi^{ml} | \eta^{ml} \rangle, \\ G(\xi^{ml}, \eta^{ml}) &= \frac{\hbar^2}{(\Delta x_0)^2} \langle \xi^{ml} | \frac{1}{p^2 + m^2} | \eta^{ml} \rangle, \quad (\Delta x_0)^2 = n\beta\hbar^2. \end{aligned} \quad (6.1)$$

The vertex is just the scalar product of two maximally localized states:

$$\tilde{\delta}(\xi^{ml}, \eta^{ml}) = c^2 \int_0^{\frac{1}{\sqrt{\beta}}} d\rho \rho^{n-1} \sqrt{1 - \beta\rho^2} \int d\Omega e^{-i\frac{\alpha \cdot \rho}{\sqrt{\beta\rho^2}} \arcsin \sqrt{\beta\rho^2}}, \quad (6.2)$$

where

$$c^2 = \left(\frac{\beta}{\pi}\right)^{\frac{n}{2}} \frac{\Gamma(\frac{n+3}{2})}{\Gamma(\frac{3}{2})}, \quad |\alpha| = \frac{|\xi - \eta|}{\hbar\sqrt{\beta}}. \quad (6.3)$$

The final result is the integral

$$\begin{aligned} \tilde{\delta}(\xi^{ml}, \eta^{ml}) &= \left(\frac{2}{\alpha}\right)^{\frac{n}{2}} \frac{\Gamma(\frac{n+3}{2})}{\Gamma(\frac{3}{2})} \alpha \int_0^{\frac{\pi}{2}} dx \\ &\times \left[\frac{(\sin x)^{n-1} - (\sin x)^{n+1}}{x^{\frac{n}{2}-1}} \right] J_{\frac{n}{2}-1}(\alpha x). \end{aligned} \quad (6.4)$$

For $n = 1$, it is exactly solvable in terms of elementary functions

$$\tilde{\delta}(\xi^{ml}, \eta^{ml})|_{n=1} = \frac{\sin \frac{\pi}{2} \alpha}{\pi} \left[\frac{1}{\left(\frac{\alpha}{2}\right) - \left(\frac{\alpha}{2}\right)^3} \right], \quad (6.5)$$

confirming what is computed in Ref. 7. The propagator is instead

$$G(\xi^{ml}, \eta^{ml}) = \frac{1}{n\beta} \langle \xi^{ml} | \frac{1}{p^2 + m^2} | \eta^{ml} \rangle. \quad (6.6)$$

Due to the identity

$$\frac{1}{p^2 + m^2} = \frac{1}{(1 - \beta m^2)^2} \frac{1}{\rho^2 + \frac{m^2}{1 - \beta m^2}} - \frac{\beta}{1 - \beta m^2}, \quad (6.7)$$

the propagator can be divided in two contributions:

$$G(\xi^{ml}, \eta^{ml}) = G_0(\xi^{ml}, \eta^{ml}) - \frac{1}{n(1 - \beta m^2)} \tilde{\delta}(\xi^{ml}, \eta^{ml}). \quad (6.8)$$

The core term is obviously

$$G_0(\xi^{ml}, \eta^{ml}) = \left(\frac{2}{\alpha}\right)^{\frac{n}{2}} \frac{\Gamma\left(\frac{n+3}{2}\right)}{\Gamma\left(\frac{3}{2}\right)} \frac{\alpha}{n(1 - \beta m^2)^2} \times \int_0^{\frac{\pi}{2}} dx \left[\frac{(\sin x)^{n-1} - (\sin x)^{n+1}}{x^{\frac{n}{2}-1} \left(\sin^2 x + \frac{\beta m^2}{1 - \beta m^2}\right)} \right] J_{\frac{n}{2}-1}(\alpha x). \quad (6.9)$$

This integral cannot be computed for $m^2 \neq 0$. In the massless case we obtain instead

$$G_0(\xi^{ml}, \eta^{ml})|_{m^2=0} = \left(\frac{2}{\alpha}\right)^{\frac{n}{2}} \frac{\Gamma\left(\frac{n+3}{2}\right)}{\Gamma\left(\frac{3}{2}\right)} \frac{\alpha}{n} \times \int_0^{\frac{\pi}{2}} dx \left[\frac{(\sin x)^{n-3} - (\sin x)^{n-1}}{x^{\frac{n}{2}-1}} \right] J_{\frac{n}{2}-1}(\alpha x). \quad (6.10)$$

In the limit $\alpha \rightarrow 0$ (i.e. $\xi \rightarrow \eta$ and β fixed) we obtain the interesting result

$$G_0(\xi^{ml}, \eta^{ml})|_{m^2=0}(\alpha \rightarrow 0) = \frac{n+1}{n(n-2)} \quad (6.11)$$

and for the complete propagator

$$G(\xi^{ml}, \eta^{ml})|_{m^2=0}(\alpha \rightarrow 0) = \frac{3}{n(n-2)}. \quad (6.12)$$

From this formula we note that the propagator, as defined in (6.1), makes sense only for $n \geq 3$, while for $n = 2$ the integral diverges logarithmically. In the limiting case $n = 3$ for $\alpha \rightarrow 0$ we obtain simply the identity, while for $\alpha \rightarrow \infty$ ($\beta \rightarrow 0$, $|\xi - \eta| = \text{fixed}$) the propagator assumes the usual form

$$G(\xi^{ml}, \eta^{ml})|_{m^2=0}(\alpha \rightarrow \infty) \propto \frac{1}{|\xi - \eta|}. \quad (6.13)$$

The physical meaning of this result is interesting. The UV divergencies of quantum field theory arise because the propagator is a singular function in the limit $\xi \rightarrow \eta$, i.e. it is a distribution due to the locality property.

Usually in the perturbative expansion we find products of propagators and the products of distributions are generally ill defined, as in the case

$$[G(\xi - \eta)]^2 = \int \frac{d^4p}{(2\pi)^4} e^{ip \cdot (\xi - \eta)} \int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 - m^2)((p - k)^2 - m^2)}, \quad (6.14)$$

and we need to introduce subtractions to make sense of the perturbative result. Instead in the case of the Snyder geometry the propagator is finite in the limit $\xi \rightarrow \eta$ and the product of propagators is a well-defined function. The theory is regularized in its roots (modifying the quantization rules) without the need of introducing extra subtractions.

7. Conclusions

In summary, the Snyder algebra represents the simplest generalization of one-dimensional quantum mechanics with a nonzero minimal position uncertainty. It introduces a hypercube of side-length $\sqrt{\beta}\hbar$, not accessible to the measurements.

We have succeeded to modify the translational invariance in order to make it covariant with the noncommutative algebra. This step is fundamental to define the maximally localized states that realize the highest physical information accessible.

Thanks to these states we can build the Feynman rules like the vertex and the propagator. Unlike ordinary quantum field theory, the propagator is no more a distribution, i.e. singular in the limit $\xi \rightarrow \eta$, but finite. This property assures that the UV divergencies of quantum field theory are absent and there is no need of ad hoc subtractions.

Although the UV divergencies are absent, this does not exclude the possibility of other type of divergencies, due to the UV-IR mixing (as discussed in Ref. 10), which is a typical property of noncommutative field theories.

Then we discuss the limits of our technique. The first limit we see is that our scheme is strongly dependent on our Euclidean choice of the space-time metric. Frankly we do not know if our formulas can be analytically continued to the Minkowskian case.¹¹ The second limit is that the introduction of a nonlocality of space-time is in conflict with the principle of unitarity (which is strictly related to the locality principle). Since with the fuzziness of space-time we have no control of the Planck scale at the measurement level, we expect that the requirement of unitarity must be limited to the laboratory energy scale. Work is in progress in this direction.

References

1. P. K. Townsend, *Phys. Rev. D* **15**, 2795 (1976).
2. D. J. Gross and P. F. Mende, *Nucl. Phys. B* **303**, 407 (1988).

3. D. Amati, M. Ciafaloni and G. Veneziano, *Phys. Lett. B* **216**, 41 (1989).
4. K. Konishi, G. Paffuti and P. Provero, *Phys. Lett. B* **234**, 276 (1990).
5. M. Maggiore, *Phys. Lett. B* **319**, 83 (1993).
6. H. S. Snyder, *Phys. Rev.* **71**, 38 (1947).
7. A. Kempf, G. Mangano and R. B. Mann, *Phys. Rev. D* **52**, 1108 (1995), arXiv:hep-th/9412167.
8. R. Banerjee, S. Kulkarni and S. Samanta, *J. High Energy Phys.* **0605**, 077 (2006), arXiv:hep-th/0602151.
9. A. Kempf and G. Mangano, *Phys. Rev. D* **55**, 7909 (1997), arXiv:hep-th/9612084.
10. C. N. Yang, *Phys. Rev.* **72**, 874 (1947).
11. A. Bose, arXiv:1011.5234v1 [physics.gen-ph].