

# *Hippotherium malpassii* (Equidae, Mammalia) from the latest Miocene (late Messinian; MN13) of Monticino gypsum quarry (Brisighella, Emilia-Romagna, Italy)

Lorenzo ROOK & Raymond L. BERNOR

L. Rook, Dipartimento di Scienze della Terra, Università di Firenze, Via G. La Pira 4, I-50121 Firenze, Italy; lorenzo.rook@unifi.it  
R.L. Bernor, College of Medicine, Department of Anatomy, Laboratory of Evolutionary Biology, Howard University, Washington, DC 20059, USA; rbernor@howard.edu

**KEY WORDS** - *Hippotherium*, dentition, postcranial skeleton, Messinian, Brisighella, Italy.

**ABSTRACT** - We describe the *Hippotherium* record from the latest Miocene (MN13) vertebrate faunal assemblage of the Monticino gypsum quarry (also known as Brisighella). This small sample would appear to be attributable to a single species of hipparionine horse. The referral to the recently described species *Hippotherium malpassii* Bernor, Kaiser, Nelson & Rook, 2011 would appear to be the best one possible. The Monticino gypsum quarry specimens correspond to the size of *Hippotherium malpassii* and show a suite of morphological features that allow their specific attribution. Formal description of the Brisighella hipparion specimens augments the previous knowledge of the Monticino gypsum quarry vertebrate fauna, one of the best-known latest Messinian fossil assemblages of continental Europe.

**RIASSUNTO** - [*Hippotherium malpassii* (Equidae, Mammalia) dal Miocene terminale (Messiniano superiore; MN13) della Cava di gessi del Monticino (Brisighella, Emilia-Romagna, Italia)] - Vengono descritti i resti fossili di *Hippotherium* della associazione faunistica del Miocene terminale (MN13 nella biocronologia continentale a mammiferi dell'Europa) proveniente della Cava di gessi del Monticino (nota anche come fauna di Brisighella). Il campione è rappresentato da un numero limitato di resti, tutti riferibili ad una singola specie di ippoterio. Il campione è attribuito alla specie *Hippotherium malpassii* Bernor, Kaiser, Nelson & Rook, 2011, recentemente descritta sulla base dell'abbondante campione fossile proveniente dal bacino di Baccinello-Cinigiano, in provincia di Grosseto. In base alle dimensioni, i resti della Cava di gessi del Monticino corrispondono alla taglia della specie *Hippotherium malpassii*. Questa attribuzione è inoltre supportata dalla presenza di una serie di tratti morfologici caratteristici. La descrizione formale dell'ippoterio di Brisighella espande la nostra conoscenza sulla fauna della Cava di gessi del Monticino, una delle associazioni faunistiche a mammiferi del Messiniano terminale meglio conosciute in Europa. A conclusione della descrizione morfologica del piccolo campione di *Hippotherium* di Brisighella viene infine discussa la peculiare distribuzione del record fossile dei perissodattili nel quadro della località a mammiferi del Miocene Superiore della Penisola Italiana.

## INTRODUCTION

The occurrence of a rich late Messinian fossil vertebrate assemblage in karst sediments associated with the Messinian evaporites in the Monticino gypsum quarry near Brisighella (Emilia-Romagna, Central Italy; Fig. 1) was first reported by Costa et al. (1986). According to local and regional geological constraints (Marabini & Vai, 1989; Vai, 1989), the vertebrate assemblage is correlated with the late Messinian Stage and European Land Mammal Age (ELMA) MN13. The vertebrate fauna has been the subject of a number of descriptive papers (De Giuli et al., 1988; De Giuli, 1989; Kotsakis, 1989; Kotsakis & Masini, 1989; Masini, 1989; Masini & Thomas, 1989; Torre, 1989; Rook et al., 1991; Rook, 1992; Masini & Rook, 1993; Rook & Masini, 1994; Delfino, 2002; Rook, 2009; Gallai & Rook, 2011).

Soon after the 1985 discovery of the paleontological site within the gypsum quarry, a proposal was advanced to create an open-air geo-paleontological museum at Brisighella (Vai, 2007). Between 2001-2004, the first preservation activities at the site were undertaken thanks to an agreement between the Regional Administration (Regione Emilia-Romagna) and the Town Administration (Comune di Brisighella), and an open-air museum of the quarry area was realized in 2005-2006. A geo-park with a 14.5 hectare environmental refuge was soon created for

a variety of didactic activities, and opened to the general public in 2007 (Sami, 2007a, b).

In this paper we report on the unpublished fossil remains of a tridactyl horse (*Hippotherium*) from the Monticino gypsum quarry. This taxon, represented by limited fossil material, has not been yet formally described, although its recovery dates back to the time of the Brisighella fauna's discovery (De Giuli et al., 1988). The identification of this sample has important biochronologic and biogeographic implications. We also provide a discussion of the significance of the hipparion record in the Messinian of Italy.

## GEOLOGICAL SETTING

The exposed succession within the Monticino gypsum quarry (Fig. 1) is composed, at the base, of a series of tilted gypsum layers with interbedded clays of the early Messinian Gessoso-Solfifera Formation. The gypsum was slightly eroded and subjected to a karst dissolution cycle that took place following the tectonic tilting of the strata. The deposits that infill the net of fissures intersecting the gypsum and the depressions on the upper surface of the gypsum sequence consist of marly-clays, fine grained conglomerates and sand lenses derived from the Colombacci Formation (grey pelites and variegated




Fig. 1 - Location map (left) and outline of the geological section outcropping in the Monticino gypsum quarry (center), with formal geological units (Formations) and chronologic setting of the exposed sediments (right). The fossils vertebrates of the Monticino gypsum quarry faunal assemblages are found in the Colombacci Fmt., the deposition of which is tightly placed in time by the constraints of the geological setting. The Colombacci Fmt. is deposited on the karstic surface of the Messinian gypsum (thereafter the infra-Messinian tectonic phase that affected the Gessoso-solfifera Fmt.), and is capped as an angular conformity by the earliest Pliocene marine clays of the Argille Azzurre Fmt. (Zanclean). Modified after Marabini & Vai (1989).

marly clays with whitish to yellow thin evaporitic marly limestone; latest Messinian in age). They contain partially articulated and fragmented skeletons of large mammals, and very abundant microvertebrate remains. Several fossiliferous collecting points have been identified along the Monticino gypsum quarry outcrop (Fig. 2), some of which represent successive sampling of the same fissure that were brought to light by the progress of the quarrying activity. The Gessoso-Solfifera Formation is unconformably overlain by a thin discontinuous layer of greenish to blackish clay (0 to 1 m thick) that is considered equivalent to the later depositional phases of the Colombacci Formation and is correlated with the late Messinian. The succession is capped by up to 20 m of Early Pliocene marine gray marly clays (Argille Azzurre Formation) whose base falls in the *Sphaeroidinellopsis* Zone, and which overlie, with apparent conformity, the Colombacci Formation. We refer the reader to Vai (1989) for a more complete discussion of the geological setting of the Romagna Apennine, and to Marabini & Vai (1989) for further details on the geology of the Monticino gypsum quarry.




Fig. 2 - Scheme of the fossil collecting sites within the Monticino gypsum quarry outcrop. Sites that yielded *Hippotherium* are highlighted by circled numbers on a coloured background (after a sketch by M. Sami, modified from Rook & Delfino, 2007).

## MATERIALS AND METHODS

### Studied Material

The described specimens from Monticino gypsum quarry are housed in the MSF, whereas the comparative sample includes material from Baccinello V3, housed at the IGF and NHMB (see acronyms below).

### Institutional Acronyms

“MSF” indicates the Museo Civico di Scienze Naturali “Malmerendi”, Faenza, “IGF” denotes the Museo di Storia Naturale (Sezione di Geologia e Paleontologia) of the Università di Firenze, and “NHMB” refers to the Basel Naturhistorisches Museum (Switzerland).

### Fossil Site Acronyms

“BRS” indicates the fossiliferous locality (=Brisighella); the number following the acronym (e.g., BRS 25) indicates the single karst fissure (Fig. 2), whereas when a second number follows (e.g., BRS 25/1) it indicates the specimen field catalogue number. “BCB V3” refers to Baccinello V3 (latest Messinian, Tuscany; Benvenuti et al., 2001; Rook et al., 2011).

### Measurements

We have measured the upper and lower dentitions using standard equid measurements published by Eisenmann et al. (1988) and Bernor et al. (1997). The third metacarpal (MCIII) is too fragmentary to measure, but its general size is comparable to *Hippotherium malpassii*. Measurements of the Brisighella hipparion specimens are provided in Appendix, while teeth metrical comparison and species descriptive statistics are offered in Tab. 1. Measurement numbers (M1, M2, M3, etc.) refer to those published by Eisenmann et al. (1988; and rounded to 0.1 mm) for postcranium (also in Bernor et al., 1997), whereas tooth measurement numbers refer to those published by Bernor et al. (1997), Bernor & Franzen (1997) and Bernor & Harris (2003). The osteological nomenclature has been adapted from Nickel et al. (1986). Getty (1982) was also consulted for morphological identification and comparison.

BRS			<i>H. malpassii</i> from Baccinello V3					
P <sup>4</sup>	MSF119	MSF120	n	Mean	Min	Max	STD	95%CI
M1	23.0	23.6	8	25.4	21.7	27.8	2.16	±1.49
M2	19.8	19.7	7	21.1	19.4	23.6	1.78	±1.32
M3	22.7	21.5	8	22.6	20.9	25.1	1.19	±0.83
M4	20.8	20.3	6	24.3	22.7	26.2	2.10	±1.68
M5	49.1	(47.0)	7	35.3	23.3	48.1	10.10	±7.48
M10	6.4	6.0	8	5.7	4.0	7.2	1.22	±0.85
M11	3.6	4.1	8	4.4	3.4	5.6	0.74	±0.51
M <sub>2</sub>	MSF122		n	Mean	Min	Max	STD	95%CI
M1	20.8		9	22.5	20.7	25.1	1.70	±1.11
M2	19.5		6	21.2	19.7	22.4	1.06	±0.85
M3	13.8		10	12.2	11.3	13.4	0.67	±0.42
M4	6.7		10	6.8	4.7	8.4	1.16	±0.72
M5	7.7		10	8.4	6.4	10.6	1.30	±0.81
M6	13.6		8	11.4	8.2	14.9	2.24	±1.55
M7	11.9		7	12.4	9.3	13.3	1.41	±1.05
M8	11.4		10	10.7	9.5	12.1	0.95	±0.59
M9	10.3		8	10.0	8.8	11.7	0.95	±0.66
M10	32.3		7	35.8	25.6	45.7	7.88	±5.84

Tab. 1 - *Hippotherium* P<sup>4</sup> and M<sub>2</sub> (see Appendix for metrical variables explanation) from Monticino gypsum quarry compared to descriptive statistics of *Hippotherium malpassii* from type locality (Baccinello V3).

#### Taxonomic remarks

The genus nomen *Hipparion* has been used in a variety of ways by different authors. Here we follow the systematics recently provided by Bernor et al. (1996, 1997, 2011) for hipparionine horse superspecific taxa.

### SYSTEMATIC PALEONTOLOGY

Suborder HIPPOMORPHA Wood, 1937

Superfamily EQUOIDEA Hay, 1902

Family EQUIDAE Gray, 1821

Subfamily EQUINAE Steinmann & Doderlein, 1890

Tribe HIPPARIONINI Quinn, 1955

Genus *Hippotherium* Kaup, 1832

*Hippotherium malpassii* Bernor, Kaiser, Nelson & Rook, 2011  
(Figs 3-5)

1988 *Hipparion* sp. DE GIULI ET AL., p. 68.

1993 *Hipparion* sp. MASINI & ROOK, p. 80.

2004 *Hipparion* sp. ROOK & DELFINO, p. 194.

2007 *Hipparion* sp. ROOK & DELFINO, p. 118.

**Material examined** - The Monticino gypsum quarry *Hippotherium* sample comprises five specimens: MFS119, left P<sup>4</sup> (Fig. 3a); MFS120, left P<sup>4</sup> (Fig. 3b); BRS 15/1, upper molar; MSF122, left M<sub>2</sub> (Fig. 4); MSF121, fragmentary left Metacarpal III (Fig. 5).

**Site** - All the described material has been collected from sites within the Monticino gypsum quarry, near Brisighella (RA), specifically from three (possibly five) single fissures: BRS 1 (MSF119), BRS 15 (BRS15/1), and BRS 25 (MSF122), while two erratic specimens (MSF120 and MSF121) possibly come from fissures BRS 3 and BRS 5 (Fig. 2).

**Description** - The Monticino gypsum quarry hipparion sample includes two very similar left P<sup>4</sup>s, MSF119 (crown height = 49.1 mm) and MSF122 (crown height = 47.0 mm), that are in early middle wear (Fig. 3). Both specimens exhibit the following salient features: pre- and postfossettes have complex mesial and distal borders, but in particular the distal border of the prefossette; they have a very large, single pli; they have lenticular shaped protocones; the protocone is placed well lingual of the hypocone; the hypoglyph is deeply incised; the parastyle and mesostyle are narrow loops. The plications of the pre- and postfossettes are not as complex as some of the Baccinello sample, but this may be due to their relatively early stage of wear. Both of these teeth have paracone labial enamel borders that are high and round and equal to score #2 of Muhlbachler et al. (2011) and Wolf et al. (2012), which has the effect of producing a deep-wide groove across the mesostyle-protocone portion of the crown. As such, it may be inferred that this horse had a low abrasive diet (= large browse component) and was likely a mixed-feeder at the browsing end of the browser-grazer spectrum.

The lower molar, MSF122 (Fig. 4), is a left M<sub>2</sub> with a crown height of 32.3 mm. This specimen is in middle adult wear and has the following salient characters: the metaconid and metastylid are round; pre- and postflexids are labiolingually compressed and have simple margins; linguaflexid has a deep U-shaped morphology; ectoflexid is deep, fully penetrating the metaconid-metastylid junction; protoconid and metaconid have rounded margins; there is no evidence of an ectostylid or a pli caballinid; protoconid raises only about 23 mm from the base of the crown and is pointed at its tip.

There is a fragmentary metacarpal III (Fig. 5) that includes a portion of the proximal end and the fragmentary anterior two-thirds of the shaft. The specimen is too fragmentary to measure but, in general, would appear to have dimensions and proportions similar to the Baccinello sample of *Hippotherium malpassii* (see Appendix).

As an assemblage, this would appear to be material of a single species of hipparionine horse. The referral to *Hippotherium malpassii* would appear to be the best one possible. It is the size of *Hippotherium malpassii* type material (Tab. 1; Fig. 6) and has a constellation of morphological features that best fit this taxon: there is no ectostylid or pli caballinid typical of Late Miocene - early Pliocene *Eurygnathohippus*; neither the metaconid nor metastylid is pointed as is often the case in *Eurygnathohippus*; it is neither very large as in *Sivalhippus* nor very small as in *Cremohipparion*; the MCIII does not have either massive proportions as is often seen in *Sivalhippus*, nor extremely gracile proportions as seen in *Hipparion* or *Cremohipparion*. In sum, the Brisighella hipparion is best referred to *Hippotherium malpassii* recently described from Baccinello (Bernor et al., 2011).





Fig. 3 - Left P<sup>4</sup> of *Hippotherium malpassii* from Monticino gypsum quarry (Brisighella). a) MSF119 in distal (a1), mesial (a2), occlusal (a3), labial (a4), and lingual (a5) views. b) MSF120 in distal (b1), mesial (b2), occlusal (b3), labial (b4), and lingual (b5) views. Figures a5 and b5 illustrate the broad labio-lingual groove across the mesostyle-protocone portion of the tooth. Scale bar = 3 cm.

CONCLUSIONS

The Monticino gypsum quarry (Brisighella) karst fissure fillings complex yielded very few horse remains documented from five collecting sites (BRS 1, BRS 3,

BRS 5, BRS 15, BRS 25), described herein for the first time. On the basis of morphological and morphometric comparisons we attribute the Brisighella sample to *Hippotherium malpassii*, a species recently described from the Late Miocene (Messinian) of Baccinello V3 in southern Tuscany (Bernor et al., 2011).

The Italian Late Miocene *Hippotherium* record, although relatively poor, contributes to our knowledge of the peculiar paleobiogeographic evolution of the Italian Peninsula. During Late Turolian (Late Miocene, MN12 time equivalent), no equid (nor any perissodactyls) are known in Italy either from the so-called V2 faunal assemblages of the Tusco-Sardinian bioprovince (Monte Bamboli in Tuscany and Fiume Santo in Sardinia; Rook et al., 1999, 2006; Bernor et al., 2001; Chesi et al., 2009) or from the faunas of the Abruzzi Apulian bioprovince (Abbazzi et al., 1996; Rook et al., 2006).

The faunal turnover that characterized the continental peninsular Italy during the latest Miocene led to the extinction of all taxa typical of the Tusco-Sardinian area and was characterized by the dispersal of a faunas composed by continental European taxa, characterizing the latest Miocene Italian mammal assemblages, along the northern Apennines, from Piedmont (Ciabot Cagna,



Fig. 4 - Left M<sub>2</sub> of *Hippotherium malpassii* from Monticino gypsum quarry (Brisighella). MSF122 in occlusal (a), labial (b), lingual (c), mesial (d), and distal (e) views. Scale bar = 3 cm.



Fig. 5 - Left metacarpal III of *Hippotherium malpassii* from Monticino gypsum quarry (Brisighella). MSF121 in cranial (a), caudal (b), proximal (c), lateral (d), and medial (e) views. Scale bar = 5 cm.

Verduno and Moncucco; Cavallo et al., 1993; Angelone et al., 2011; Colombero & Pavia, 2013) to Romagna (Brisighella, this paper) and southern Tuscany (Borro Strolla, Baccinello V3, etc.; Rook et al., 2006; Abbazzi et

al., 2008). Perissodactyla were part of this renewed fauna, including tapirs (Rook & Rustioni, 1991), rhinos (Hürzeler & Engesser, 1976; De Giuli et al., 1998) and hipparions (Bernor et al., 2011; this paper). The MN13 record of the *Hippotherium* record in Italy is the youngest known for this lineage in Eurasia.

In the light of the renewed interest in the Italian *Hippotherium* record (Bernor et al., 2011; this paper), a revision of the hipparion sample from Casino - only in part figured by Forsyth Major (1875, 1877) and by Pirlot (1956), and generally referred to as part of the “*Hipparion gracile* group” (among others Alberdi, 1986; Forsten, 2002) - would be most welcome, given its possible identification as representing a form close to the species *Hippotherium malpassii*.




Fig. 6 - Bivariate plot of P<sup>4</sup> measurements. Protocone width (M11) is plotted versus protocone length (M10). BRS specimens (asterisks) are compared with a sample of Late Miocene Old World hipparions: Baccinello V3, Italy (open triangle); Gaiselberg, Austria (solid star); Höwenegg, Germany (cross circled); Sinap, Turkey (cross boxed); Baltavar, Hungary (diamond). Comparative data derived from Bernor et al. (1997, 2011).



## ACKNOWLEDGMENTS

We wish to thank Tonino Benericetti from Zattaglia for the patient and careful activity on the Monticino fossiliferous locality, as well as Gian Paolo Costa, and Marco Sami (Faenza) for constructive discussions. Finally we would like to acknowledge the Cultural Heritage service of the Town Administration for facilitating access to fossil material in the collections of the Faenza Museum. This contribution is part of a broader research program on Late Neogene vertebrate evolution developed at the University of Florence (coordinator L.R.). We would like to acknowledge financial support for his research from the National Science Foundation: EAR0125009, EAR113175 (to RLB) and BCS-0321893 (grant made to F.C. Howell and T.D. White; RHOI, University of California at Berkeley). Thanks are extended to M. Delfino, D.M. Alba and an anonymous reviewer for their constructive comments, which greatly ameliorated this short paper.

## REFERENCES

- Abbazzi L., Benvenuti M., Boschian G., Dominici S., Masini F., Mezzabotta C., Rook L., Valleri G. & Torre D. (1996). The Neogene and Pleistocene succession, and the mammal faunal assemblages of an area between Apricena and Poggio Imperiale (Foggia, Italy). *Memorie della Società Geologica Italiana*, 51: 383-402.
- Abbazzi L., Benvenuti M., Ceci M.E., Esu D., Faranda C., Rook L. & Tangocci F. (2008). The end of the Lago-Mare time in the SE Valdelsa Basin (Central Italy): interference between local tectonism and regional sea-level rise. *Geodiversitas*, 30: 611-639.
- Alberdi M.T. (1986). The Pliocene species of *Hipparion* and their biostratigraphical meanings. *Géobios*, 19: 517-522.
- Angelone C., Colombero S., Esu D., Giuntelli P., Marcolini F., Pavia M., Trenkwalder S., Van Den Hoek Ostende L.W., Zunino M. & Pavia G. (2011). Moncucco Torinese, a new postevaporitic Messinian fossiliferous site from Piedmont (NW Italy). *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 259: 89-104.
- Benvenuti M., Papini M. & Rook L. (2001). Mammal biochronology, UBSU and paleoenvironment evolution in a post-collisional basin: evidence from the Late Miocene Baccinello-Cinigiano basin in southern Tuscany. Italy. *Bollettino della Società Geologica Italiana*, 120 (1995): 97-118.
- Bernor R.L., Fahlbusch V., Andrews P., De Bruijn H., Fortelius M., Rögl F., Steininger F.F. & Werdelin L. (1996). The evolution of Western Eurasian Neogene mammal faunas: a chronologic, systematic, biogeographic, and paleoenvironmental synthesis. In Bernor R.L., Fahlbusch V. & Mittman H.W. (eds), *The Evolution of Western Eurasian Neogene Mammal Faunas*, Columbia University Press, New York: 449-469.
- Bernor R.L., Fortelius M. & Rook L. (2001). Evolutionary biogeography and paleoecology of the “*Oreopithecus bambolii* Faunal Zone” (Late Miocene, Tusco-Sardinian Province). *Bollettino della Società Paleontologica Italiana*, 40: 139-148.
- Bernor R.L. & Franzen J. (1997). The hipparionine horses from the Turolian Age (Late Miocene) locality of Dorn Dürkheim, Germany. *Courier Forschungsinstitut Senckenberg*, 197: 117-185.
- Bernor R.L. & Harris J. (2003). Systematics and evolutionary biology of the Late Miocene and Early Pliocene hipparionine horses from Lothagam, Kenya. In Leakey M. & Harris J. (eds), *Lothagam: The Dawn of Humanity in Eastern Africa*. Columbia University Press, New York: 387-438.
- Bernor R.L., Kaiser T.M., Nelson S.V. & Rook L. (2011). Systematics and paleobiology of *Hippotherium malpassii* n. sp. (Equidae, Mammalia) from the latest Miocene of Baccinello V3 (Tuscany, Italy). *Bollettino della Società Paleontologica Italiana*, 50: 175-208.
- Bernor R.L., Tobien H., Hayek L.-A. & Mittmann H.-W. (1997). The Höwenegg hipparionine horses: systematics, stratigraphy, taphonomy and paleoenvironmental context. *Andrias*, 10: 1-230.
- Cavallo O., Sen S., Rage J.-C. & Gaudant J. (1993). Vertébrés messiniens du Faciès à congéries de Ciabòt Cagna, Corneliano d'Alba (Piémont, Italie). *Rivista Piemontese di Storia Naturale*, 14: 3-22.
- Chesi F., Delfino M. & Rook L. (2009). Late Miocene *Mauremys* (Testudines, Geoemydidae) from Tuscany (Italy): evidence of terrapin persistence after a mammal turnover. *Journal of Paleontology*, 83: 379-388.
- Colombero S. and Pavia G. (2013). *Centralomys benericettii* (De Giuli, 1989) (Mammalia, Rodentia): a latest Messinian murid from Northern and Central Italy. New data from Piedmont. *Bollettino della Società Paleontologica Italiana*, 52: 27-34.
- Costa G.P., Colalongo M.L., De Giuli C., Marabini S., Masini F., Torre D. & Vai G.B. (1986). Latest Messinian vertebrate fauna preserved in a paleokarst-neptunian dyke setting. *Le Grotte d'Italia*, 12: 221-235.
- De Giuli C. (1989). The rodents from the Brisighella latest Miocene fauna. *Bollettino della Società Paleontologica Italiana*, 28: 197-212.
- De Giuli C., Masini F., Torre D., Benericetti A., Costa G.P., Fosella M. & Sami M. (1988). The mammal fauna of Monticino Quarry. In De Giuli C. & Vai G.B. (eds), *Fossil Vertebrates in the Lamone Valley Romagna Appennines*. Field Trip Guidebook. Faenza: 65-69.
- Delfino M. (2002). Erpetofaune italiane del Neogene e del Quaternario. 382 pp. PhD Thesis, Università degli Studi di Modena e Reggio Emilia.
- Eisenmann V., Alberdi M.T., De Giuli C. & Staesche U. (1988). Volume I: Methodology. In Woodburne M. & Sondaar P. (eds), *Studying Fossil Horses*. Collected papers after the “New York International Hipparion Conference, 1981”. Brill, Leiden: 1-71.
- Forsten A. (2002). Latest *Hipparion* Christol, 1832 in Europe. A review of the Pliocene *Hipparion crassum* Gervais group and other finds (Mammalia, Equidae). *Geodiversitas*, 24: 465-486.
- Forsyth Major C.I. (1875). Considerazioni sulla fauna dei Mammiferi pliocenici e postpliocenici della Toscana. II. La Fauna dei Mammiferi del Pliocene inferiore (Orizzonte di Casino). *Memorie della Società Toscana di Scienze Naturali*, 1: 223-245.
- Forsyth Major C.I. (1877). Considerazioni sulla fauna dei Mammiferi pliocenici e postpliocenici della Toscana. III. Cani fossili del Val d'Arno superiore e della Valle d'Era. *Memorie della Società Toscana di Scienze Naturali*, 3: 207-227.
- Gallai G. & Rook L. (2011). *Propotamochoerus provincialis* (Gervais, 1859) (Suidae, Mammalia) from the latest Miocene (late Messinian; MN13) of Monticino quarry (Brisighella, Emilia-Romagna, Italy). *Bollettino della Società Paleontologica Italiana*, 50: 29-34.
- Getty R. (1982). *The Anatomy of Domestic Animals*. Saunders Company, Philadelphia: 1-1211.
- Gray J.E. (1821). On the natural arrangement of vertebrate animals. *London Medical Repository*, 15: 296-310.
- Hay O.P. (1902). Bibliography and catalogue of the fossil Vertebrata of North America. *U.S. Geological Survey Bulletin*, 179: 1-868.
- Hürzeler J. & Engesser B. (1976). Les faunes des mammifères néogènes du Bassin de Baccinello (Grosseto, Italie). *Comptes Rendus de l'Académie de Sciences de Paris, sér. 2D*, 283: 333-336.
- Kaup J.J. (1832). Description d'ossements fossiles de mammifères inconnus jusqu'à présent qui se trouvent au Muséum grand ducal de Darmstadt. 119 pp. Meyer, Darmstadt.
- Kotsakis T. (1989). Late Turolian amphibians and reptiles from Brisighella (Northern Italy). *Bollettino della Società Paleontologica Italiana*, 28: 271-276.
- Kotsakis T. & Masini F. (1989). Late Turolian bats from Brisighella (Northern Italy). *Bollettino della Società Paleontologica Italiana*, 28: 281-285.

- Marabini S. & Vai G.B. (1989). Geology of the Monticino quarry, Brisighella, Italy. Stratigraphic implications of its late Messinian mammal fauna. *Bollettino della Società Paleontologica Italiana*, 28: 369-382.
- Masini F. (1989). *Prolagus sorbinii* nov. sp. a new Ochotonid (Mammalia, Lagomorpha) from the Messinian of Italy. *Bollettino della Società Paleontologica Italiana*, 28: 295-306.
- Masini F. & Rook L. (1993). *Hystrix primigenia* (Mammalia, Rodentia) from the Late Messinian of the Monticino gypsum quarry (Faenza, Italy). *Bollettino della Società Paleontologica Italiana*, 32: 79-87.
- Masini F. & Thomas H. (1989). *Samotragus occidentalis* nov. sp. a new bovid from the late Messinian of Italy. *Bollettino della Società Paleontologica Italiana*, 28: 307-316.
- Mihlbachler M.C., Rivals F., Solounias N. & Semprebon G.M. (2011). Dietary change and evolution of horses in North America. *Science*, 331: 1178-1181.
- Nickel R., Schummer A., Seiferle E., Wilkens H., Wille K.-H. & Frewein J. (1986). The Anatomy of the Domestic Animals, Volume I: The Locomotor System. 499 pp. Verlag Paul Parey-Springer Verlag, Berlin.
- Pirlot P.L. (1956). Les formes européennes du genre *Hipparion*. *Memorias y Comunicaciones del Instituto Geológico, Diputación Provincial de Barcelona*, 14: 1-121.
- Quinn J.H. (1955). Miocene Equidae of the Texas Gulf Coastal Plain. *Bureau of Economic Geology, University of Texas Publications*, 5516: 1-102.
- Rook L. (1992). "*Canis*" *monticinensis* sp. nov. a new Canidae (Carnivora, Mammalia) from the late Messinian of Italy. *Bollettino della Società Paleontologica Italiana*, 30: 151-156.
- Rook L. (2009). The wide ranging genus *Eucyon* Tedford & Qiu, 1996 (Mammalia, Carnivora, Canidae) in Mio-Pliocene of the Old World. *Geodiversitas*, 31: 723-743.
- Rook L., Abbazzi L. & Engesser B. (1999). An overview on the Italian Miocene land mammal faunas. In Agustí J., Rook L. & Andrews P. (eds), *The Evolution of Neogene Terrestrial Ecosystems*. Cambridge University Press, Cambridge: 191-205.
- Rook L. & Delfino M. (2004). I vertebrati fossili di Brisighella nel quadro dei popolamenti continentali del Mediterraneo durante il Neogene. *Ravenna Studi e Ricerche*, 11 (2003): 179-207.
- Rook L. & Delfino M. (2007). La fauna preistorica di Brisighella e i popolamenti continentali del Mediterraneo durante il Miocene Superiore. In Sami M. (ed.), *Il Parco geologico Cava Monticino, Brisighella. Una guida, una storia*. Regione Emilia Romagna, Tipografia Carta Bianca, Faenza: 97-124.
- Rook L., Ficarelli G. & Torre D. (1991). Messinian carnivores from Italy. *Bollettino della Società Paleontologica Italiana*, 30: 7-22.
- Rook L., Gallai G. & Torre D. (2006). Lands and endemic mammals in the Late Miocene of Italy: Constraints for paleogeographic outlines of Tyrrhenian area. *Palaeogeography Palaeoclimatology Palaeoecology*, 238: 263-269.
- Rook L. & Masini F. (1994). *Orycteropus* cf. *gaudryi* (Mammalia, Tubulidentata) from Late Miocene of the Monticino Quarry (Faenza, Italia). *Bollettino della Società Paleontologica Italiana*, 28: 369-374.
- Rook L., Oms O., Benvenuti M. & Papini M. (2011). Magnetostratigraphy of the Late Miocene Baccinello-Cinigiano basin (Tuscany, Italy) and the age of *Oreopithecus bambolii* faunal assemblages. *Palaeogeography, Palaeoclimatology Palaeoecology*, 305: 286-294.
- Rook L. & Rustioni M. (1991). *Tapirus* cf. *arvernensis* from the Late Turolian Baccinello V3 Faunal Assemblage (Grosseto, Italy). *Bollettino della Società Paleontologica Italiana*, 30: 325-327.
- Sami M. (2007a). Cava Monticino: i fossili ce ne raccontano il passato. In Sami M. (ed.), *Il Parco geologico Cava Monticino, Brisighella. Una guida, una storia*. Regione Emilia Romagna, Tipografia Carta Bianca, Faenza: 71-94.
- Sami M. (2007b). Il Parco geologico Cava Monticino, Brisighella. Una guida, una storia. Regione Emilia Romagna, Tipografia Carta Bianca, Faenza: 1-224.
- Steinmann G. & Döderlein L. (1890). *Elemente der Paläontologie*. Wilhelm Engelmann, Leipzig: 1-848.
- Torre D. (1989). *Plioiviverrops faventinus* nov. sp. a new carnivore of late Messinian age. *Bollettino della Società Paleontologica Italiana*, 28: 323-327.
- Vai G.B. (1989). A field trip guide to the Romagna Appennine Geology. The Lamone Valley. *Bollettino della Società Paleontologica Italiana*, 28: 343-367.
- Vai G.B. (2007). Storia del Geoparco di Brisighella. In Sami M. (ed.), *Il Parco geologico Cava Monticino, Brisighella. Una guida, una storia*. Regione Emilia Romagna, Tipografia Carta Bianca, Faenza: 13-26.
- Wolf D., Semprebon G.M. & Bernor R.L. (2012). New observations on the paleodiet of the Late Miocene Höwenegg (Hegau, Germany) *Hippotherium primigenium*. *Bollettino della Società Paleontologica Italiana*, 51: 185-191.
- Wood H.E. (1937). Perissodactyl suborders. *Journal of Mammalogy*, 18: 1-106.

Manuscript received 5 July 2012

Revised manuscript accepted 2 August 2013

Published online 30 October 2013

Editor Massimo Delfino

