



Memory retrieval of inhibitory avoidance requires histamine H₁ receptor activation in the hippocampus

Roberta Fabbri^{a,b}, Cristiane Regina Guerino Furini^a, Maria Beatrice Passani^c, Gustavo Provensi^b, Elisabetta Baldi^d, Corrado Bucherelli^d, Ivan Izquierdo^{a,1}, Jociane de Carvalho Myskiw^{a,1,2}, and Patrizio Blandina^{b,1,2}

^aMemory Center, Brain Institute of Rio Grande do Sul, Pontifical Catholic University of Rio Grande do Sul, 90610-000 Porto Alegre, RS, Brazil; ^bDipartimento di Neuroscienze, Psicologia, Area del Farmaco e Salute del Bambino, Sezione di Farmacologia e Tossicologia, Università di Firenze, 50139 Florence, Italy; ^cDipartimento di Scienze della Salute, Sezione di Farmacologia e Chemioterapia, Università di Firenze, 50139 Florence, Italy; and ^dDipartimento di Medicina Sperimentale e Clinica, Università di Firenze, 50134 Florence, Italy

Contributed by Ivan Izquierdo, March 30, 2016 (sent for review March 2, 2016; reviewed by Michel Baudry and Benno Roozendaal)

Retrieval represents a dynamic process that may require neuro-modulatory signaling. Here, we report that the integrity of the brain histaminergic system is necessary for retrieval of inhibitory avoidance (IA) memory, because rats depleted of histamine through lateral ventricle injections of α -fluoromethylhistidine (a-FMHis), a suicide inhibitor of histidine decarboxylase, displayed impaired IA memory when tested 2 d after training. a-FMHis was administered 24 h after training, when IA memory trace was already formed. Infusion of histamine in hippocampal CA1 of brain histamine-depleted rats (hence, amnesic) 10 min before the retention test restored IA memory but was ineffective when given in the basolateral amygdala (BLA) or the ventral medial prefrontal cortex (vmPFC). Intra-CA1 injections of selective H₁ and H₂ receptor agonists showed that histamine exerted its effect by activating the H₁ receptor. Noteworthy, the H₁ receptor antagonist pyrilamine disrupted IA memory retrieval in rats, thus strongly supporting an active involvement of endogenous histamine; 90 min after the retention test, c-Fos-positive neurons were significantly fewer in the CA1s of a-FMHis-treated rats that displayed amnesia compared with in the control group. We also found reduced levels of phosphorylated cAMP-responsive element binding protein (pCREB) in the CA1s of a-FMHis-treated animals compared with in controls. Increases in pCREB levels are associated with retrieval of associated memories. Targeting the histaminergic system may modify the retrieval of emotional memory; hence, histaminergic ligands might reduce dysfunctional aversive memories and improve the efficacy of exposure psychotherapies.

memory | retrieval | inhibitory avoidance | histamine | H1 receptor

Memory determines the uniqueness of our personal history and is decisive for each individual to survive and prosper. It is a multistate process that includes acquisition, consolidation, and retrieval (1). Whereas considerable advancement has been made toward understanding the specific brain structures (e.g., amygdala, prefrontal cortex, and hippocampus) and molecular mechanisms (receptors and signaling pathways) that underlie acquisition and consolidation (2, 3), the understanding of retrieval has lagged behind, although it is ultimately the only possible measure of memory (4–6). Indeed, retrieval is not simply a static readout of stored information; rather, it represents a dynamic process that can be studied separately from either acquisition or consolidation, with which it shares similar mechanisms (1, 7, 8). Furthermore, retrieval can elicit specific processes that modify the recalled memory (9). In this regard, protein synthesis, a necessary step in the transfer of a labile short-term memory into a stable long-term memory (LTM) (2), is required to enable retrieval, because infusion of protein synthesis inhibitors in the amygdala 10 min before retrieval impaired fear memory expression (10). An interesting question is whether neuromodulatory signaling is required for not only memory acquisition and consolidation (11) but also, retrieval. There is evidence that adrenergic signaling is required for the

retrieval of various types of hippocampus-dependent memory in mice (12) as well as in humans (13). Intrahippocampal blockade of metabotropic glutamate or alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors immediately before retrieval impaired LTM expression of an inhibitory avoidance (IA) response (7), thus suggesting that glutamatergic signaling in hippocampus is necessary for retrieval of an aversive memory. In contrast, cholinergic and dopaminergic signaling contributes to acquisition and/or consolidation but is generally not required for retrieval (6). The histaminergic system modulates memory consolidation in many cognitive tasks from IA to fear conditioning and object recognition (14–16). We recently showed that rats were able to consolidate an IA memory only when the brain histaminergic system was intact (17). Histamine within the brain is synthesized from histidine by histidine-decarboxylase solely in hypothalamic tuberomammillary nucleus neurons (18), which are organized into functionally distinct circuits (19), display selective control mechanisms (19, 20), and impinge on different brain regions, including amygdala, prefrontal cortex, and hippocampus, that are responsible for many forms of learning (21).

Based on the above findings, it is conceivable that histaminergic signaling is required for memory retrieval as well. Consequently, this study was specifically designed to address this question using an IA task, an associative learning paradigm that has largely contributed

Significance

Several neurotransmitters contribute to memory formation by modulating selectively acquisition, consolidation, and/or retrieval. Integrity of the brain histamine system is necessary for the consolidation of inhibitory avoidance (IA) memory. Here, we report that cerebral histamine depletion also impairs retrieval of IA in rats and blunts retrieval-induced c-Fos activation and cAMP-responsive element binding protein phosphorylation in the CA1 region of the hippocampus. Histamine infusion into the CA1 restores IA retrieval in histamine-depleted rats by targeting brain histamine H₁ receptors. Our study uncovers previously unidentified mechanisms involved in memory retrieval and may offer possible targets for eventual pharmacotherapies to treat dysfunctional aversive memories, including phobias, panic attacks, and posttraumatic stress disorders, as well as improve the efficacy of exposure psychotherapies.

Author contributions: R.F., C.R.G.F., M.B.P., I.I., J.d.C.M., and P.B. designed research; R.F., C.R.G.F., G.P., E.B., C.B., and J.d.C.M. performed research; R.F., C.R.G.F., M.B.P., I.I., J.d.C.M., and P.B. analyzed data; and M.B.P., I.I., J.d.C.M., and P.B. wrote the paper.

Reviewers: M.B., Western University of Health Sciences; and B.R., Donders Institute, Radboud University.

The authors declare no conflict of interest.

¹To whom correspondence may be addressed. Email: izquier@terra.com.br, jociane_carvalho@hotmail.com, or patrizio.blandina@unifi.it.

²J.d.C.M. and P.B. contributed equally to this work.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1604841113/-DCSupplemental.

to the understanding of memory processes (22). We first examined whether brain histamine depletion through administration of the histidine-decarboxylase suicide inhibitor α -fluoromethylhistidine (a-FMHs) (23) in the lateral ventricle (LV) 24 h after training affected IA retrieval. Then, we tested whether the local infusion of histamine in the basolateral amygdala (BLA), the ventral medial prefrontal cortex (vmPFC), or the CA1 region of the dorsal hippocampus overcame a-FMHs-elicited impairment of retrieval, and we characterized the type of histamine receptor involved. We also investigated changes in neuronal activity by assessing after retrieval the pattern of c-Fos expression as well as cyclic adenosine monophosphate (cAMP)-responsive element binding protein (CREB) phosphorylation in BLA, vmPFC, and CA1 region of histamine-depleted rats. Finally, we investigated the ability of histamine to restore retrieval after administration of protein synthesis inhibition in the CA1 region.

Results

Depletion of Histamine Impairs Retrieval Independently of Consolidation.

We recently showed that intra-LV infusion of a-FMHs (5 μ g/ μ L) quickly and fully suppressed the release of brain histamine measured by microdialysis, which was restored to control levels after about 48 h (17). Thus, to investigate the role of endogenous histamine in retrieval, we examined the performance of rats infused into the LV with a-FMHs 24 h after the IA training session. The retention test was carried out 48 h after training. Latencies were compared with those of rats treated with a-FMHs 24 h before training and animals given equivalent infusions of saline (controls). Latencies of all groups during IA training did not differ. Fig. 1 shows the stepdown latency during testing of rats treated with a-FMHs and control. One-way ANOVA performed at the retention test revealed a significant difference across groups ($F_{2,41} = 62.89$; $P < 0.0001$). Additional analysis with Bonferroni's multiple




Fig. 1. Effect of histamine acute depletion through a-FMHs administration on IA task. The schematic drawing shows the sequence of behavior procedures and treatments. Rats were implanted with an infusion cannula in the LV and distributed to three groups: one group received saline (SAL) 24 h before and after IA training (control), a second group received a-FMHs 24 h before IA training and SAL 24 h after training, and the third group received SAL 24 h before IA training and a-FMHs 24 h after training. Data are expressed as means \pm SEMs of 10–12 animals for each group. **** $P < 0.0001$ vs. control (one-way ANOVA and Bonferroni's MCT).

comparisons tests (MCT) showed that latencies of all rats treated with a-FMHs either 24 h before or 24 h after IA training were significantly shorter than those of controls. Further analysis with Bonferroni's MCT showed that latencies of all rats treated with a-FMHs, either 24 h before or after IA training, were significantly shorter than those of controls. As a-FMHs effect on histamine synthesis persists for about 48 h (22), it is conceivable that rats receiving a-FMHs prior to IA training lacked integrity of the histaminergic system during the consolidation process, whereas those treated 24 h after training, during the retrieval. These findings confirm that histamine depletion impairs the consolidation of IA memory (17) and suggest that it also worsens IA-LTM expression by influences on memory retrieval mechanisms. To further test these hypotheses, we investigated whether intracerebral administration of histamine reversed the amnesic effect of histamine depletion caused by a-FMHs.

Effects of Histamine Infusion into the BLA, vmPFC, or CA1 on a-FMHs-Induced Amnesia.

Histamine (1 μ g/ μ L) was infused bilaterally into the BLA (Fig. 2A), the vmPFC (Fig. 2B), or the CA1 (Fig. 2C) 10 min before the retention test of rats given a-FMHs 24 h before or after IA training. The retention test was carried out 48 h after IA training. Controls received equivalent infusions of saline. Latencies of all groups during IA training did not differ. One-way ANOVA performed at the retention test revealed a significant difference across groups (BLA: $F_{2,24} = 58.76$; $P < 0.0001$; vmPFC: $F_{2,40} = 41.05$; $P < 0.0001$; CA1: $F_{2,27} = 34.11$; $P < 0.0001$). Additional analysis with Bonferroni's MCT showed that rats treated with a-FMHs before training and histamine in the BLA (Fig. 2A), the vmPFC (Fig. 2B), or the CA1 (Fig. 2C) displayed latencies significantly shorter than respective controls. Also, latencies of rats infused with a-FMHs 24 h after training and histamine into the BLA (Fig. 2A) or the vmPFC (Fig. 2B) were significantly shorter than those of corresponding controls. Conversely, latencies of rats treated with a-FMHs intra-LV 24 h after training and intra-CA1 histamine did not differ from their controls (Fig. 2C). Taken together, these results indicate that histamine never rescued IA-LTM of rats given a-FMHs before training. When infused into the BLA or the vmPFC of animals treated with a-FMHs after training, histamine did not restore IA-LTM. Conversely, it antagonized amnesia when given into the CA1 region. These findings suggest that animals treated with a-FMHs after IA training formed a memory of this experience and support a crucial role for histamine neurotransmission in the CA1 during retrieval.

Intra-CA1 Infusion of Histamine Did Not Reverse Amnesia of IA-LTM Elicited by Intra-CA1 Administration of Anisomycin.

To verify whether the amnesia induced by a-FMHs given 24 h before IA training shared features with the memory impairment of rats unable to consolidate memory, we investigated IA-LTM in animals that received bilateral infusion of anisomycin (80 μ g/ μ L), a protein synthesis inhibitor, in the CA1 immediately after training, saline or a-FMHs intra-LV 24 h after training, and saline or histamine (1 μ g/ μ L) into the CA1 10 min before the retention test (Fig. 2D). Controls received comparable infusions of saline. Latencies of all groups during the IA training did not differ. One-way ANOVA performed on the retention test revealed a significant group effect ($F_{3,55} = 98.10$; $P < 0.0001$). As shown in Fig. 2D, Bonferroni's MCT analysis showed no difference among the three groups of rats given anisomycin independently of the subsequent treatments, but all displayed latencies significantly shorter than controls. Thus, anisomycin-treated rats showed a substantial memory impairment that was not reversed by histamine administration in the CA1.

Histamine H_1 Receptor Activation Is Required for IA-LTM Retrieval.

Histaminergic fibers innervate the hippocampus through the fornix and a caudal route (24), and histamine modulates hippocampal functions, including memory processes, through interactions




Fig. 2. Effect of histamine infusion into the BLA, vmPFC, or CA1 on a-FMHis- or anisomycin-induced amnesia. The schematic drawings show the sequence of behavior procedures and treatments. Rats were implanted with infusion cannula in the LV to administer a-FMHis or saline (SAL) and a second cannula bilaterally in the (A) BLA, (B) vmPFC, or (C and D) CA1. Data are expressed as means \pm SEMs of 10–12 animals for each group. **** P < 0.0001 vs. respective controls (one-way ANOVA and Bonferroni's MCT). ANI, anisomycin; HA, histamine.

with H_1 and H_2 receptors (25–27). To characterize the histamine receptor type involved in the reversal of a-FMHis-induced amnesia (Fig. 2C), rats were infused with a-FMHis into the LV 24 h after IA training and 2-(2-pyridyl)ethylamine (PEA; 1.2 $\mu\text{g}/\mu\text{L}$) or dimaprit (DIM; 2.3 $\mu\text{g}/\mu\text{L}$) into the CA1 10 min before the retention test (Fig. 3A). PEA selectively activates H_1 receptors, whereas DIM is an H_2 agonist with no H_1 activity (28). Controls received equivalent infusions of saline. One-way ANOVA performed at the retention test displayed a significant difference in latency across groups ($F_{2,42} = 118.1$; $P < 0.0001$), and Bonferroni's MCT analysis indicated that rats infused with DIM had significantly shorter latencies than those infused with saline or PEA. Hence, DIM did not reverse a-FMHis-induced amnesia. Controls and PEA-treated rats displayed similar levels of latency during the retention test, thus suggesting that the H_1 receptor contributed to the full reinstatement of IA-LTM expression in animals rendered amnesic by a-FMHis. To further investigate the role of H_1 receptors, we examined the effects of pyrilamine (20 $\mu\text{g}/\mu\text{L}$), an H_1 antagonist (29) infused in the CA1 10 min before the retention test. During IA training, there were no significant differences among the groups in their latencies (Fig. 3B). However, on the retention test, rats infused with pyrilamine displayed shorter latencies

than those infused with saline (unpaired t test; $P < 0.05$) (Fig. 3B), thus indicating that blockade of H_1 receptors in the CA1 impaired IA-LTM expression. This finding extends the observations with PEA, strongly suggesting an active involvement of endogenous histamine in IA-LTM retrieval.

Effect of Histamine Depletion on c-Fos Expression After IA-LTM Retrieval in Rat BLA, vmPFC, and CA1. In an attempt to clarify how brain histamine deficit may influence IA-LTM retrieval, we measured c-Fos protein expression in three brain regions of rats given saline or a-FMHis into the LV 24 h after IA training and subjected to a retention test 48 h after training. Rats were euthanized 90 min after the retention test. No differences in c-Fos expression were found in the BLA or the vmPFC of rats treated with either saline or a-FMHis (Fig. 4A and B). Conversely, neurons immunopositive for c-Fos were significantly fewer in the CA1 of a-FMHis-treated rats compared with in the control group (unpaired t test; $P < 0.0001$) (Fig. 4C).

Effect of Histamine Depletion on Levels of pCREB After IA-LTM Retrieval in Rat BLA, vmPFC, and CA1. CREB is a crucial mediator in the formation of IA-LTM (17, 30), and an increase in




Fig. 4. c-Fos expression in the CA1 of rats trained and tested for IA memory is blunted in a-FMHis-treated rats. The schematic drawing shows the sequence of procedures and treatment. Brain coronal sections show the effect of exposure to retention test on c-Fos protein expression in the (A) BLA, (B) vmPFC, and (C) CA1 of rats given intra-LV infusions of saline (SAL) or a-FMHis 24 h after IA training. Data are expressed as means \pm SEM of three to four rats for each group. (Scale bars: A, 100 μ m; B, 500 μ m; C, Upper, 100 μ m; C, Lower, 500 μ m.) **** $P < 0.0001$ (unpaired t test).

investigated in this study. There is much evidence that activation of H_2 receptors potentiates consolidation in aversive tasks (14, 27, 44), including IA (26, 45). The hippocampus expresses H_1 and H_2 receptors (46, 47), and both facilitate cAMP accumulation (48). Hippocampal infusion with the H_2 receptor agonist DIM 10 min before the retention test failed to restore a-FMHis-induced amnesia, whereas rats injected with the H_1 receptor agonist PEA showed latencies similar to those of saline-injected controls. Furthermore, administering the H_1 antagonist pyrilamine 10 min before retention test in rats with normal levels of histamine completely disrupted IA retention. This result strongly supports the notion that histamine signaling through H_1 receptors is essential for IA-LTM retrieval. Our results are in agreement with previous reports showing that the H_1 receptor is implicated in memory processes, because H_1 receptor KO mice showed impaired memory performance in the Barnes maze (49) and the radial maze task (50). Accordingly, long-term potentiation in the CA1 was significantly reduced in H_1 -KO compared with WT mice (49). Studies using c-fos-tTA reporter mice have shown that the same neurons in the amygdala, hippocampus, and cortex that are active during contextual fear learning are reactivated when memory is retrieved (51, 52). Recently, it has been reported that CA1 neurons that were engaged during contextual fear

learning reduced levels of pCREB and were responsible for re-instating memory representations that occurred during learning in the cortex at the time of retrieval (53). When those specific CA1 neurons were silenced, specific cortical reactivation was reduced, and mice became unable to retrieve a previously formed contextual fear memory (53). We may speculate that IA-LTM shares these features with contextual fear memory and that H_1 receptor activation is necessary for reactivating the CA1 neurons engaged during learning, thus being responsible for reinstatement of cortical-specific activity necessary for retrieval. Consistent with this idea, given that gene expression in c-fos-tTA mice largely recapitulated endogenous c-Fos expression (52), we found that, 90 min after exposure to the retention test, c-Fos-positive neurons were significantly fewer in the CA1 of a-FMHis-treated rats that also displayed amnesia compared with those in the same region of animals given saline and displaying intact IA-LTM retention. Previous studies supported this contention (53, 54). Immediately after the retention test, we found reduced levels of pCREB in the CA1 of a-FMHis-treated animals compared with in controls. CREB phosphorylation represents a crucial step for the consolidation of LTM (30), and increases in pCREB levels associated with retrieval of fear conditioning as well as spatial memory in different brain regions,




Fig. 5. Effect of histamine depletion on levels of CREB phosphorylation in the BLA, vmPFC, and CA1 of rats trained and tested for IA memory. The schematic drawing displays the sequence of procedures and treatments. Rats were implanted with an infusion cannula in the LV. Representative immunoblots and densitometric quantification are shown. Data are expressed as means \pm SEMs of four rats. ** $P < 0.01$ vs. respective saline (SAL; one-way ANOVA and Bonferroni's MCT).

including the hippocampus, have been reported (55, 56). We suggest that increased levels of pCREB are required for the retrieval of IA-LTM through activation of H_1 receptors. Consistently, earlier reports indicate that histamine signaling is necessary to trigger CREB phosphorylation in the hippocampus (17, 57). The hippocampus is involved in both the aversive component of the IA and its contextual aspect as reviewed in the work by Izquierdo et al. (58).

Taken together, this study suggests (i) that the integrity of the histaminergic system in the CA1 region of the hippocampus is crucial for IA-LTM retrieval, (ii) that histamine-depleted rats do not retrieve IA-LTM and display a reduced number of c-Fos-positive cells as well as a lower level of pCREB in CA1 after retrieval compared with controls, and (iii) that blockade of H_1 receptors in the CA1 of normal rats impairs the retrieval of IA-LTM 2 d after training. Advances in the understanding of mechanisms underlying IA memory may help in the search for treatments of psychiatric diseases, such as posttraumatic stress disorder or obsessive-compulsive disorder. Here, we provide evidence that targeting the histaminergic system may modify the encoding, consolidation, and retrieval of emotional memory.

Materials and Methods

Animals. Male Wistar rats (3 mo old; 300–330 g) purchased from Centro de Modelos Biologicos Experimentais of the Pontifical Catholic University of Rio Grande do Sul (our regular provider) were used. Animals were housed four to a cage with water and food ad libitum under a 12-h light/dark cycle (lights on at 7:00 AM). The temperature of the animals' room was maintained at $23 \text{ }^\circ\text{C} \pm 1 \text{ }^\circ\text{C}$. All procedures were in accordance with the NIH's *Guide for the Care and Use of Laboratory Animals* (59) and approved by Animal Committee on Ethics in the Care and Use of Laboratory Animals of the Pontifical Catholic University of Rio Grande do Sul.

Surgery. At least 1 wk after their arrival, animals were anesthetized (75 mg/kg ketamine plus 10 mg/kg xylazine i.p.) and placed on a stereotaxic frame

(Kopf). A stainless steel cannula (22 gauge) was implanted in the LV (anterior, -0.9 mm ; lateral, -1.5 mm ; ventral, -2.6 mm from Bregma) (60) and fixed to the skull by using dental cement. Animals were also implanted bilaterally with 22-gauge guide cannulas 1 mm above the CA1 region of the dorsal hippocampus (anterior, -4.2 mm ; lateral, $\pm 3.0 \text{ mm}$; ventral, -1.8 mm from Bregma), the BLA (anterior, -2.4 mm ; lateral, $\pm 5.1 \text{ mm}$; ventral, -7.5 mm from Bregma), or the vmPFC (anterior, $+3.2 \text{ mm}$; lateral, $\pm 0.8 \text{ mm}$; ventral, -2.0 mm from Bregma) (60). Cannula placements were verified postmortem as described in detail in *SI Materials and Methods*. Animals were allowed 7 d to recover from surgery before behavioral procedures. All rats were handled once daily for 3 consecutive d, and all behavioral procedures was conducted between 8:00 and 11:00 AM.

IA Task. The apparatus consisted of a Plexiglas box ($50 \times 25 \times 25 \text{ cm}$) with a floor made of parallel 1-mm-caliber bronze bars spaced 0.8 cm apart and a wood platform (5-cm high, 8-cm wide, and 25-cm long) on the left extreme of the box. For the IA training session, animals were gently placed on the platform facing the left rear corner. When stepping down and placing their four paws on the grid, animals received a 2-s 0.5-mA scrambled foot shock and then, were immediately withdrawn from the training box. After 48 h, animals were placed again on the platform as described for a retention test without the foot shock. In the retention test, the stepdown latency was 300 s. Latency to stepdown was measured with an automated stopwatch.

Drugs and Infusion Procedures. At the time of drug microinfusions, animals were gently restrained by hand, and an injection needle (30 gauge) was fitted tightly into the guide, extending 1 mm from the tip of the guide cannulas. The injection needle was connected to a 10- μL Hamilton microsyringe, and the infusions were performed at a rate of 0.5 $\mu\text{L}/30 \text{ s}$. The injection needle was left in place for an additional 60 s to minimize backflow. It was then carefully withdrawn and placed on the other side, where the procedure was repeated. Infusions into the BLA, vmPFC, or CA1 were performed 10 min before the retention test. Infusions into the LV were performed 24 h either before or after IA training. The drugs used were histamine (1 $\mu\text{g}/\mu\text{L}$), DIM (2.3 $\mu\text{g}/\mu\text{L}$), PEA (1.2 $\mu\text{g}/\mu\text{L}$), pyrilamine (20 $\mu\text{g}/\mu\text{L}$), and anisomycin (80 $\mu\text{g}/\mu\text{L}$) purchased from Sigma-Aldrich. The a-FMHs (5 $\mu\text{g}/\mu\text{L}$) was synthesized at Abbott Laboratories. The doses were chosen among those found to be effective in previous papers and had no effects on locomotion or exploration activity (26, 61). The volume of the drugs infused was 0.5 μL per side in the BLA and 1 μL per side into the CA1, vmPFC, and LV. Control groups received equal volumes of sterile saline (0.9%).

Immunohistochemistry. Ninety minutes after the retention test, animals were deeply anesthetized with chloral hydrate (400 mg/kg i.p.) and perfused transcardially with saline followed by 4% (vol/vol) paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). Brains were postfixed in the same solution overnight (4 $^\circ\text{C}$) and cryoprotected in 30% (wt/vol) sucrose in phosphate buffer. Forty-micrometer-thick sections were then processed for standard immunostaining (details in *SI Materials and Methods*).

Western Blotting Analysis. Animals were killed immediately after the IA retention test, the brain was dissected out on ice, and the amygdala, the vmPFC, and the CA1 were immediately isolated. The pooled structures (left and right) were individually homogenized in 200 μL ice-cold lysis buffer containing protease and phosphatase inhibitors [50 mM Tris-HCl (pH 7.5), 50 mM NaCl, 10 mM EGTA, 5 mM EDTA, 2 mM sodium pyrophosphate, 4 mM *p*-nitrophenyl phosphate, 1 mM Na_3VO_4 , 1.1 mM PMSF, 20 $\mu\text{g}/\mu\text{L}$ leupeptin, 50 $\mu\text{g}/\mu\text{L}$ aprotinin, 0.1% SDS] and centrifuged at $13.8 \times g$ at 4 $^\circ\text{C}$ for 15 min. The following procedure is described in detail in *SI Materials and Methods*. For each sample, a ratio of p^{Ser133} -CREB/CREB densities was calculated, and then, all of the individual rates were expressed as a percentage of the average of ratios obtained from the control group.

Data and Statistical Analysis. Statistical analysis was performed by using Prism Software (GraphPad). Data are expressed as means \pm SEMs. IA latencies and number of c-Fos-positive nuclei as well as the pCREB/CREB ratio were analyzed with unpaired *t* test or one-way ANOVA. The source of the detected significances was determined by Bonferroni's multiple comparison posthoc test. *P* values less than 0.05 were considered statistically significant. The number of rats per group is indicated in the figures.

ACKNOWLEDGMENTS. We thank Dr. Alessia Costa for technical support and Dr. M. Cowart for the synthesis of a-FMHs. Work was supported by research grants from the National Council of Research of Brazil, the CNPq, the Compagnia di San Paolo (Italy), and the Ente Cassa di Risparmio di Firenze (Italy).

1. Abel T, Lattal KM (2001) Molecular mechanisms of memory acquisition, consolidation and retrieval. *Curr Opin Neurobiol* 11(2):180–187.
2. McGaugh JL (2000) Memory—a century of consolidation. *Science* 287(5451):248–251.
3. Johansen JP, Cain CK, Ostroff LE, LeDoux JE (2011) Molecular mechanisms of fear learning and memory. *Cell* 147(3):509–524.
4. Szapiro G, et al. (2002) Molecular mechanisms of memory retrieval. *Neurochem Res* 27(11):1491–1498.
5. Brown R, Silva AJ (2004) Molecular and cellular cognition; the unraveling of memory retrieval. *Cell* 117(1):3–4.
6. Thomas SA (2015) Neuromodulatory signaling in hippocampus-dependent memory retrieval. *Hippocampus* 25(4):415–431.
7. Szapiro G, et al. (2000) Participation of hippocampal metabotropic glutamate receptors, protein kinase A and mitogen-activated protein kinases in memory retrieval. *Neuroscience* 99(1):1–5.
8. Vianna MRM, Szapiro G, McGaugh JL, Medina JH, Izquierdo I (2001) Retrieval of memory for fear-motivated training initiates extinction requiring protein synthesis in the rat hippocampus. *Proc Natl Acad Sci USA* 98(21):12251–12254.
9. Kim J, Kwon J-T, Kim HS, Josselyn SA, Han JH (2014) Memory recall and modifications by activating neurons with elevated CREB. *Nat Neurosci* 17(1):65–72.
10. Lopez J, Gamache K, Schneider R, Nader K (2015) Memory retrieval requires ongoing protein synthesis and NMDA receptor activity-mediated AMPA receptor trafficking. *J Neurosci* 35(6):2465–2475.
11. Atherton LA, Dupret D, Mellor JR (2015) Memory trace replay: The shaping of memory consolidation by neuromodulation. *Trends Neurosci* 38(9):560–570.
12. Murchison CF, et al. (2004) A distinct role for norepinephrine in memory retrieval. *Cell* 117(1):131–143.
13. Sterpenich V, et al. (2006) The locus coeruleus is involved in the successful retrieval of emotional memories in humans. *J Neurosci* 26(28):7416–7423.
14. Benetti F, Baldi E, Bucherelli C, Blandina P, Passani MB (2013) Histaminergic ligands injected into the nucleus basalis magnocellularis differentially affect fear conditioning consolidation. *Int J Neuropsychopharmacol* 16(3):575–582.
15. da Silveira CK, Furini CR, Benetti F, Monteiro SdaC, Izquierdo I (2013) The role of histamine receptors in the consolidation of object recognition memory. *Neurobiol Learn Mem* 103:64–71.
16. Baldi E, et al. (2005) The H3 receptor protean agonist proxyfan enhances the expression of fear memory in the rat. *Neuropharmacology* 48(2):246–251.
17. Benetti F, et al. (2015) Histamine in the basolateral amygdala promotes inhibitory avoidance learning independently of hippocampus. *Proc Natl Acad Sci USA* 112(19):E2536–E2542.
18. Green JP, Prell GD, Khandelwal JK, Blandina P (1987) Aspects of histamine metabolism. *Agents Actions* 22(1–2):1–15.
19. Blandina P, Munari L, Provensi G, Passani MB (2012) Histamine neurons in the tuberomammillary nucleus: A whole center or distinct subpopulations? *Front Syst Neurosci* 6(2012):33.
20. Munari L, Provensi G, Passani MB, Blandina P (2013) Selective brain region activation by histamine H₃ receptor antagonist/inverse agonist ABT-239 enhances acetylcholine and histamine release and increases c-Fos expression. *Neuropharmacology* 70: 131–140.
21. Panula P, Nuutinen S (2013) The histaminergic network in the brain: Basic organization and role in disease. *Nat Rev Neurosci* 14(7):472–487.
22. Izquierdo I, Medina JH (1997) Memory formation: The sequence of biochemical events in the hippocampus and its connection to activity in other brain structures. *Neurobiol Learn Mem* 68(3):285–316.
23. Garbarg M, Barbin G, Rodergas E, Schwartz JC (1980) Inhibition of histamine synthesis in brain by alpha-fluoromethylhistidine, a new irreversible inhibitor: In vitro and in vivo studies. *J Neurochem* 35(5):1045–1052.
24. Panula P, Pirvola U, Auvinen S, Airaksinen MS (1989) Histamine-immunoreactive nerve fibers in the rat brain. *Neuroscience* 28(3):585–610.
25. Haas HL, Sergeeva OA, Selbach O (2008) Histamine in the nervous system. *Physiol Rev* 88(3):1183–1241.
26. da Silva WC, Bonini JS, Bevilacqua LRM, Izquierdo I, Cammarota M (2006) Histamine enhances inhibitory avoidance memory consolidation through a H2 receptor-dependent mechanism. *Neurobiol Learn Mem* 86(1):100–106.
27. Giovannini MG, et al. (2003) Improvement in fear memory by histamine-elicited ERK2 activation in hippocampal CA3 cells. *J Neurosci* 23(27):9016–9023.
28. Panula P, et al. (2015) International Union of Basic and Clinical Pharmacology. XCVIII. Histamine Receptors. *Pharmacol Rev* 67(3):601–655.
29. Ganellin CR (1982) *Pharmacology of Histamine Receptors*, eds Ganellin C, Parson ME (Wright, Bristol, England), pp 10–102.
30. Carlezon WA, Jr, Duman RS, Nestler EJ (2005) The many faces of CREB. *Trends Neurosci* 28(8):436–445.
31. Bernabeu R, et al. (1997) Involvement of hippocampal cAMP/cAMP-dependent protein kinase signaling pathways in a late memory consolidation phase of aversively motivated learning in rats. *Proc Natl Acad Sci USA* 94(13):7041–7046.
32. Viola H, et al. (2000) Phosphorylated cAMP response element-binding protein as a molecular marker of memory processing in rat hippocampus: Effect of novelty. *J Neurosci* 20(23):RC112.
33. Izquierdo I, et al. (1997) Sequential role of hippocampus and amygdala, entorhinal cortex and parietal cortex in formation and retrieval of memory for inhibitory avoidance in rats. *Eur J Neurosci* 9(4):786–793.
34. Hotte M, et al. (2006) D₁ receptor modulation of memory retrieval performance is associated with changes in pCREB and pDARPP-32 in rat prefrontal cortex. *Behav Brain Res* 171(1):127–133.
35. Bambah-Mukku D, Travaglia A, Chen DY, Pollonini G, Alberini CM (2014) A positive autoregulatory BDNF feedback loop via C/EBPβ mediates hippocampal memory consolidation. *J Neurosci* 34(37):12547–12559.
36. Taubenfeld SM, Milekic MH, Monti B, Alberini CM (2001) The consolidation of new but not reactivated memory requires hippocampal C/EBPβ. *Nat Neurosci* 4(8): 813–818.
37. Bekinschtein P, et al. (2007) Persistence of long-term memory storage requires a late protein synthesis- and BDNF- dependent phase in the hippocampus. *Neuron* 53(2): 261–277.
38. Bekinschtein P, Weisstaub N (2014) Role of PFC during retrieval of recognition memory in rodents. *J Physiol Paris* 108(4–6):252–255.
39. Milekic MH, Brown SD, Castellini C, Alberini CM (2006) Persistent disruption of an established morphine conditioned place preference. *J Neurosci* 26(11):3010–3020.
40. Quevedo J, et al. (1999) Two time windows of anisomycin-induced amnesia for inhibitory avoidance training in rats: Protection from amnesia by pretraining but not pre-exposure to the task apparatus. *Learn Mem* 6(6):600–607.
41. Inda MC, Muravieva EV, Alberini CM (2011) Memory retrieval and the passage of time: From reconsolidation and strengthening to extinction. *J Neurosci* 31(5): 1635–1643.
42. Black AH, Nadel L, O’Keefe J (1977) Hippocampal function in avoidance learning and punishment. *Psychol Bull* 84(6):1107–1129.
43. Garín-Aguilar ME, Medina AC, Quirarte GL, McGaugh JL, Prado-Alcalá RA (2014) Intense aversive training protects memory from the amnesic effects of hippocampal inactivation. *Hippocampus* 24(1):102–112.
44. Bonini JS, et al. (2011) Histamine facilitates consolidation of fear extinction. *Int J Neuropsychopharmacol* 14(9):1209–1217.
45. Gianlorenço AC, Canto-de-Souza A, Mattioli R (2013) Intra-cerebellar microinjection of histamine enhances memory consolidation of inhibitory avoidance learning in mice via H2 receptors. *Neurosci Lett* 557(Pt B):159–164.
46. Vizuete ML, et al. (1997) Detailed mapping of the histamine H₂ receptor and its gene transcripts in guinea-pig brain. *Neuroscience* 80(2):321–343.
47. Palacios JM, Wamsley JK, Kuhar MJ (1981) The distribution of histamine H₁-receptors in the rat brain: An autoradiographic study. *Neuroscience* 6(1):15–37.
48. Baudry M, Martres MP, Schwartz J-C (1975) H₁ and H₂ receptors in the histamine-induced accumulation of cyclic AMP in guinea pig brain slices. *Nature* 253(5490): 362–364.
49. Dai H, et al. (2007) Selective cognitive dysfunction in mice lacking histamine H1 and H2 receptors. *Neurosci Res* 57(2):306–313.
50. Zlomuzica A, Ruocco LA, Sadile AG, Huston JP, Dere E (2009) Histamine H1 receptor knockout mice exhibit impaired spatial memory in the eight-arm radial maze. *Br J Pharmacol* 157(1):86–91.
51. Reijmers LG, Perkins BL, Matsuo N, Mayford M (2007) Localization of a stable neural correlate of associative memory. *Science* 317(5842):1230–1233.
52. Liu X, et al. (2012) Optogenetic stimulation of a hippocampal engram activates fear memory recall. *Nature* 484(7394):381–385.
53. Tanaka KZ, et al. (2014) Cortical representations are reinstated by the hippocampus during memory retrieval. *Neuron* 84(2):347–354.
54. Tayler KK, Tanaka KZ, Reijmers LG, Wiltgen BJ (2013) Reactivation of neural ensembles during the retrieval of recent and remote memory. *Curr Biol* 23(2):99–106.
55. Izumi T, et al. (2011) Retrieval of conditioned fear activates the basolateral and intercalated nucleus of amygdala. *J Neurosci Res* 89(5):773–790.
56. Zhou G, Xiong W, Zhang X, Ge S (2013) Retrieval of consolidated spatial memory in the water maze is correlated with expression of pCREB and Egr1 in the hippocampus of aged mice. *Dement Geriatr Cogn Dis Extra* 3(1):39–47.
57. Munari L, et al. (2015) Brain histamine is crucial for selective serotonin reuptake inhibitors’ behavioral and neurochemical effects. *Int J Neuropsychopharmacol* 18(10): pyv045.
58. Izquierdo I, Furini CRG, Myskiw JC (2016) Fear memory. *Physiol Rev* 96(2):695–750.
59. National Institutes of Health (1985) *Guide for the Care and Use of Laboratory Animals* (National Institutes of Health, Bethesda, MD).
60. Paxinos G, Watson C (1998) *The Rat Brain in Stereotaxic Coordinates* (Academic, New York).
61. Passani MB, et al. (2001) Histamine H3 receptor-mediated impairment of contextual fear conditioning and in-vivo inhibition of cholinergic transmission in the rat basolateral amygdala. *Eur J Neurosci* 14(9):1522–1532.
62. de Carvalho Myskiw J, Benetti F, Izquierdo I (2013) Behavioral tagging of extinction learning. *Proc Natl Acad Sci USA* 110(3):1071–1076.
63. Fiorenza NG, Rosa J, Izquierdo I, Myskiw JC (2012) Modulation of the extinction of two different fear-motivated tasks in three distinct brain areas. *Behav Brain Res* 232(1):210–216.
64. Myskiw JC, et al. (2008) On the participation of mTOR in recognition memory. *Neurobiol Learn Mem* 89(3):338–351.
65. Provensi G, et al. (2014) Satiety factor oleylethanolamide recruits the brain histaminergic system to inhibit food intake. *Proc Natl Acad Sci USA* 111(31):11527–11532.

Supporting Information

Fabbri et al. 10.1073/pnas.1604841113

SI Materials and Methods

Correct Cannula Placements. Correct cannula placement was verified by infusing a 4% (wt/vol) methylene blue solution over 30 s into the CA1 region of the dorsal hippocampus, the vmPFC (both 1 μ L per side), or the BLA (0.5 μ L per side) 2 d after the behavioral procedures. Rats were euthanized 30 min later with an overdose of anesthetic. Brains were removed and stored in formalin. The spread of the dye was taken as an estimate of drug infusions in the same animal. Cannula placements were considered correct when the spread was 1 mm³ (62–64) or less from the intended infusion sites, which occurred in 98% of the rats. Only behavioral data from animals with cannulas placed correctly were analyzed.

Immunohistochemistry Procedure. Ninety minutes after IA retention test, to fully assess *c-Fos* expression, rats were deeply anesthetized with chloral hydrate (400 mg/kg i.p.) and perfused transcardially with cold physiological saline followed by 4% (vol/vol) paraformaldehyde in 0.1 M phosphate buffer (PB; pH 7.4). Brains were postfixed in the same solution overnight (4 °C) and cryoprotected in 30% (wt/vol) sucrose in PB; 40- μ m-thick sections were cut on a cryostat and collected in PB. Sections were preincubated in 0.75% H₂O₂ in PB for 30 min and 0.2% BSA for 30 min and then incubated overnight in rabbit *c-Fos* primary antibody (1:5,000; Sigma-Aldrich) at 4 °C. The immunoreactive product was detected with the avidin-biotin peroxidase system (Vectastain Kit; Vector Laboratories). After washing, sections were mounted on gelatin-coated slides, dehydrated, coverslipped, and observed using an Olympus BX40 Microscope equipped with a Nikon DS-F1 Camera. *c-Fos*-immunopositive nuclei were counted bilaterally using the ImageJ software (NIH) on four to five sections per region per rat and normalized to a 1-mm² area according to the work by Provensi et al. (65). Atlas coordinates relative to Bregma (60) for the sections analyzed were from –3.8 to –4.4 mm for hippocampal CA1, from –2.12 to –2.75 mm for the BLA, and from 2.7 to 1.95 mm for the vmPFC. All regions analyzed receive histaminergic fibers. Statistics were calculated on the average values from four to five sections of individual regions for each animal.

pCREB Experiments and Western Blotting Analysis. For the experiments aimed to evaluate pCREB levels, male Wistar rats were infused with saline or a-FMHs through a cannula into the LV 24 h after the IA training session. Immediately after the IA retention test, animals were killed, rat brains were dissected out on ice, and the BLA, the CA1 region of the hippocampus, and the vmPFC were isolated immediately. Pools of structures (left and right) were individually homogenized in 200 μ L ice-cold lysis buffer containing protease and phosphatase inhibitors [50 mM Tris-HCl (pH 7.5), 50 mM NaCl, 10 mM EGTA, 5 mM EDTA 2 mM sodium pyrophosphate, 4 mM *p*-nitrophenyl phosphate, 1 mM Na₃VO₄, 1.1 mM PMSF, 20 μ g/ μ L leupeptin, 50 μ g/ μ L aprotinin, 0.1% SDS] and centrifuged at 13.8 \times *g* at 4 °C for 15 min. Supernatants were collected, and levels of total protein were quantified using Pierce BCA Protein Assay (Thermo Scientific). Aliquots of protein homogenates were diluted with a mix of lysis buffer and loading buffer two times [50 mM Tris (pH 6.8), 100 mM DTT, 10% (vol/vol) glycerol, 1% bromophenol blue, 2% (vol/vol) SDS] and boiled for 10 min. Aliquots containing 50 μ g total proteins were resolved by electrophoresis on a 10% SDS/PAGE and transferred on PVDF membranes (Immobilon Transfer Membranes; Millipore). Membranes were then blocked in Tris-buffered saline (pH 7.6) containing 0.1% Tween 20 (TBS-T) and 5% (wt/vol) nonfat dry milk (Bio-Rad Laboratories) for 2 h at room temperature and incubated overnight at 4 °C with mAbs against pCREB (Ser133; Cell Signaling Technology) or CREB (Cell Signaling Technology) diluted 1:1,000 in TBS-T containing 5% (wt/vol) BSA or 5% (wt/vol) nonfat dry milk, respectively. Immunodetection was performed with secondary antibodies (anti-rabbit IgG conjugated to HRP; Cell Signaling Technology) diluted 1:5,000 in TBS-T containing 1% nonfat dry milk. Blots were washed in TBS-T, and then, reactive bands were detected by using ECL (Luminata Crescendo; Millipore). Quantitative densitometry was assessed using QuantityOne software (Bio-Rad Laboratories). A ratio of pSer133-CREB/CREB densities was calculated of all samples, and all of the individual rates were expressed as percentages of the average of ratios obtained from the control group.