

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2018 September 21; 24(35): 3965-4092



### EDITORIAL

- 3965 Role of endoscopic therapy in early esophageal cancer  
*Malik S, Sharma G, Sanaka MR, Thota PN*
- 3974 Biomarkers for hepatocellular carcinoma: What's new on the horizon?  
*Ocker M*

### REVIEW

- 3980 Pediatric hepatocellular carcinoma  
*Khanna R, Verma SK*
- 4000 Changing role of histopathology in the diagnosis and management of hepatocellular carcinoma  
*Rastogi A*

### MINIREVIEWS

- 4014 Endoscopy in inflammatory bowel disease: Role in diagnosis, management, and treatment  
*Spiceland CM, Lodhia N*
- 4021 Biosimilars in paediatric inflammatory bowel disease  
*Sieczkowska-Golub J, Jarzebicka D, Oracz G, Kierkus J*

### ORIGINAL ARTICLE

#### Basic Study

- 4028 Adiponectin affects the mechanical responses in strips from the mouse gastric fundus  
*Idrizaj E, Garella R, Castellini G, Mohr H, Pellegata NS, Francini F, Ricca V, Squecco R, Baccari MC*
- 4036 Daikenchuto (Da-Jian-Zhong-Tang) ameliorates intestinal fibrosis by activating myofibroblast transient receptor potential ankyrin 1 channel  
*Hiraishi K, Kurahara LH, Sumiyoshi M, Hu YP, Koga K, Onitsuka M, Kojima D, Yue L, Takedatsu H, Jian YW, Inoue R*

#### Retrospective Cohort Study

- 4054 Portosplenomesenteric vein thrombosis in patients with early-stage severe acute pancreatitis  
*Ding L, Deng F, Yu C, He WH, Xia L, Zhou M, Huang X, Lei YP, Zhou XJ, Zhu Y, Lu NH*

#### Retrospective Study

- 4061 Serum anti-*Helicobacter pylori* antibody titer and its association with gastric nodularity, atrophy, and age: A cross-sectional study  
*Toyoshima O, Nishizawa T, Sakitani K, Yamakawa T, Takahashi Y, Yamamichi N, Hata K, Seto Y, Koike K, Watanabe H, Suzuki H*



**Observational Study**

- 4069 Real-life chromoendoscopy for dysplasia surveillance in ulcerative colitis

*Klepp P, Tollisen A, Røseth A, Cvancarova Småstuen M, Andersen SN, Vatn M, Moum BA, Brackmann S*

**Randomized Clinical Trials**

- 4077 Usefulness of the clip-flap method of endoscopic submucosal dissection: A randomized controlled trial

*Ban H, Sugimoto M, Otsuka T, Murata M, Nakata T, Hasegawa H, Inatomi O, Bamba S, Andoh A*

**CASE REPORT**

- 4086 Infant cholestasis patient with a novel missense mutation in the *AKR1D1* gene successfully treated by early adequate supplementation with chenodeoxycholic acid: A case report and review of the literature

*Wang HH, Wen FQ, Dai DL, Wang JS, Zhao J, Setchell KDR, Shi LN, Zhou SM, Liu SX, Yang QH*

**ABOUT COVER**

Editorial board member of *World Journal of Gastroenterology*, Amedeo Lonardo, MD, Doctor, Senior Lecturer, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena 41126, Italy

**AIMS AND SCOPE**

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

**INDEXING/ABSTRACTING**

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports® cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35<sup>th</sup> among 80 journals in gastroenterology and hepatology (quartile in category Q2).

**EDITORS FOR THIS ISSUE**

**Responsible Assistant Editor:** *Xiang Li*  
**Responsible Electronic Editor:** *Shu-Yu Yin*  
**Proofing Editor-in-Chief:** *Lian-Sheng Ma*

**Responsible Science Editor:** *Xue-Jiao Wang*  
**Proofing Editorial Office Director:** *Ze-Mao Gong*

**NAME OF JOURNAL**

*World Journal of Gastroenterology*

**ISSN**

ISSN 1007-9327 (print)  
 ISSN 2219-2840 (online)

**LAUNCH DATE**

October 1, 1995

**FREQUENCY**

Weekly

**EDITORS-IN-CHIEF**

**Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States**

**EDITORIAL BOARD MEMBERS**

All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

**EDITORIAL OFFICE**

Ze-Mao Gong, Director  
*World Journal of Gastroenterology*  
 Baishideng Publishing Group Inc  
 7901 Stoneridge Drive, Suite 501,  
 Pleasanton, CA 94588, USA  
 Telephone: +1-925-2238242  
 Fax: +1-925-2238243  
 E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
 Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

**PUBLISHER**

Baishideng Publishing Group Inc  
 7901 Stoneridge Drive, Suite 501,  
 Pleasanton, CA 94588, USA  
 Telephone: +1-925-2238242  
 Fax: +1-925-2238243  
 E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
 Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

**PUBLICATION DATE**

August 28, 2018

**COPYRIGHT**

© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

**SPECIAL STATEMENT**

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

**INSTRUCTIONS TO AUTHORS**

Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

**ONLINE SUBMISSION**

<http://www.f6publishing.com>

## Basic Study

**Adiponectin affects the mechanical responses in strips from the mouse gastric fundus**

Eglantina Idrizaj, Rachele Garella, Giovanni Castellini, Hermine Mohr, Natalia S Pellegata, Fabio Francini, Valdo Ricca, Roberta Squecco, Maria Caterina Baccari

Eglantina Idrizaj, Rachele Garella, Fabio Francini, Roberta Squecco, Maria Caterina Baccari, Department of Experimental and Clinical Medicine, Section of Physiological Sciences, University of Florence, Florence 50134, Italy

Giovanni Castellini, Valdo Ricca, Psychiatry Unit, Department of Health Sciences, University of Florence, Florence 50134, Italy

Hermine Mohr, Natalia S Pellegata, Institute for Diabetes and Cancer, Helmholtz Zentrum München, Neuherberg 85764, Germany

ORCID number: Eglantina Idrizaj (0000-0002-2756-6552); Rachele Garella (0000-0003-3194-7603); Giovanni Castellini (0000-0003-1265-491X); Hermine Mohr (0000-0002-3061-1166); Natalia S Pellegata (0000-0002-8000-7784); Fabio Francini (0000-0002-9255-1824); Valdo Ricca (0000-0002-9291-2124); Roberta Squecco (0000-0002-6534-3675); Maria Caterina Baccari (0000-0003-4665-1426).

**Author contributions:** Idrizaj E and Garella R contributed equally to this work; Idrizaj E and Garella R performed the functional experiments; Idrizaj E and Mohr H performed the PCR analysis; Castellini G, Ricca V, Squecco R and Baccari MC designed the research study; Pellegata NS and Francini F contributed to design the research study; Idrizaj E, Garella R and Baccari MC analyzed the data; Baccari MC wrote the paper; Idrizaj E, Garella R, Castellini G, Mohr H, Pellegata NS, Francini F, Ricca V, Squecco R and Baccari MC critically revised the manuscript.

**Supported by** the Florence University (No. RTD CO 090101010107; RICATEN14) and Fondazione CRF (No. 2017.0777).

**Institutional animal care and use committee statement:** The experimental protocol was designed in compliance with the guidelines of the European Communities Council Directive 2010/63/UE and the recommendations for the care and use of laboratory animals approved by the Animal Care Committee of the University of Florence, Italy, with authorization from the Italian Ministry of Health nr. 787/2016-PR.

**Conflict-of-interest statement:** No conflicts of interest, financial

or otherwise, are declared by the authors.

**Data sharing statement:** No additional data are available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Correspondence to:** Maria Caterina Baccari, PhD, Professor, Department of Experimental and Clinical Medicine, Section of Physiological Sciences, University of Florence, Viale G.B. Morgagni 63, Florence 50134, Italy. [mcaterina.baccari@unifi.it](mailto:mcaterina.baccari@unifi.it)  
Telephone: +39-55-2751600  
Fax: +39-55-4379506

**Received:** June 8, 2018

**Peer-review started:** June 8, 2018

**First decision:** July 3, 2018

**Revised:** July 12, 2018

**Accepted:** July 22, 2018

**Article in press:** July 22, 2018

**Published online:** September 21, 2018

**Abstract****AIM**

To investigate whether the adipocytes derived hormone adiponectin (ADPN) affects the mechanical responses in strips from the mouse gastric fundus.

**METHODS**

For functional experiments, gastric strips from the fundal

region were cut in the direction of the longitudinal muscle layer and placed in organ baths containing Krebs-Henseleit solution. Mechanical responses were recorded *via* force-displacement transducers, which were coupled to a polygraph for continuous recording of isometric tension. Electrical field stimulation (EFS) was applied *via* two platinum wire rings through which the preparation was threaded. The effects of ADPN were investigated on the neurally-induced contractile and relaxant responses elicited by EFS. The expression of ADPN receptors, Adipo-R1 and Adipo-R2, was also evaluated by touchdown-PCR analysis.

### RESULTS

In the functional experiments, EFS (4-16 Hz) elicited tetrodotoxin (TTX)-sensitive contractile responses. Addition of ADPN to the bath medium caused a reduction in amplitude of the neurally-induced contractile responses ( $P < 0.05$ ). The effects of ADPN were no longer observed in the presence of the nitric oxide (NO) synthesis inhibitor L-N<sup>G</sup>-nitro arginine (L-NNA) ( $P > 0.05$ ). The direct smooth muscle response to methacholine was not influenced by ADPN ( $P > 0.05$ ). In carbachol precontracted strips and in the presence of guanethidine, EFS induced relaxant responses. Addition of ADPN to the bath medium, other than causing a slight and progressive decay of the basal tension, increased the amplitude of the neurally-induced relaxant responses ( $P < 0.05$ ). Touchdown-PCR analysis revealed the expression of both Adipo-R1 and Adipo-R2 in the gastric fundus.

### CONCLUSION

The results indicate for the first time that ADPN is able to influence the mechanical responses in strips from the mouse gastric fundus.

**Key words:** Adiponectin; Adiponectin receptors; Gastric motility; Nitric oxide; Non-adrenergic, non-cholinergic neurotransmission

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Evidence exists that some white adipose-tissue derived hormones that are involved in the regulation of food intake also influence the motor responses of the gastrointestinal tract. In this view, adiponectin (ADPN) too has been reported to influence food intake but no data concerning its effects on the gastric mechanical activity are available. The present results indicate for the first time that ADPN is able to influence the mechanical responses in strips from the mouse gastric fundus. It could be speculated that these peripheral effects on motor phenomena might represent an additional mechanism engaged by the hormone to control food intake.

Idrizaj E, Garella R, Castellini G, Mohr H, Pellegata NS, Francini F, Ricca V, Squecco R, Baccari MC. Adiponectin affects the mechanical responses in strips from the mouse gastric fundus. *World J Gastroenterol* 2018; 24(35): 4028-4035 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i35/4028.htm>

## INTRODUCTION

White adipose tissue is recognized as an endocrine organ<sup>[1]</sup>, being able to secrete many substances, known as adipokines, that are involved in a wide range of functions<sup>[1,2]</sup>. Among them, accumulating evidence suggests that adipose-tissue derived hormones, such as leptin and adiponectin (ADPN), play an important role in the systemic metabolic homeostasis, also acting as key neuromodulators of food intake<sup>[1-6]</sup>. ADPN is a peptide hormone mainly produced by white adipose tissue although its expression has been reported also in other organs of different animal species<sup>[7]</sup>, including the mouse stomach<sup>[8]</sup>. ADPN exerts its effects mainly through the activation of two seven transmembrane domain receptors, Adipo-R1 and Adipo-R2<sup>[9]</sup>, whose expression has been revealed, other than in the brain, in a variety of mammalian peripheral tissues<sup>[7]</sup>. Particularly, Adipo-R2 mRNA expression has been reported in the gastrointestinal tract of rats<sup>[10]</sup>.

Interestingly, many substances that act centrally to modulate food intake also influence the activity of gastrointestinal smooth muscle<sup>[11-14]</sup>, whose motor phenomena represent a source of peripheral signals involved in the control of feeding behavior through the gut-brain axis<sup>[15]</sup>.

All the above knowledge are consistent with the possibility that ADPN too could affect gastrointestinal motor responses, although no data are present in the literature concerning these effects.

On this ground, in the present study we investigated whether ADPN influences the mechanical responses in strips from the mouse gastric fundus. The expression of both the ADPN receptors was also evaluated by touchdown-PCR analysis.

## MATERIALS AND METHODS

### Animals

Experiments were carried out on 8-12 wk old C57BL/6 female mice (Charles River, Lecco, Italy). The animals were fed standard laboratory chow and water, and were housed under a 12-h light/12-h dark (12 L : 12 D) photoperiod and controlled temperature (21 °C ± 1 °C). The experimental protocol was designed in compliance with the guidelines of the European Communities Council Directive 2010/63/UE and the recommendations for the care and use of laboratory animals approved by the Animal Care Committee of the University of Florence, Italy, with authorization from the Italian Ministry of Health nr. 787/2016-PR. The mice were killed by prompt cervical dislocation to minimize animal suffering.

### Mechanical experiments

As previously published by our research group<sup>[16,17]</sup>, the

**Table 1 PCR components**

Reagents	Volume
Forward primer (0.01 pmol/L)	1 µL
Reverse primer (0.01 pmol/L)	1 µL
GOTaq colorless Master Mix sample (Promega <sup>®</sup> )	12.5 µL
H <sub>2</sub> O	9.5 µL
cDNA template	1 µL

**Table 2 Touchdown-PCR setup**

Step	Temperature (°C)	Time	Note
1	94	3 min	-
2	94	20 s	-
3	64	30 s	-0.5 °C per cycle
4	72	35 s	Repeat steps 2-4 for 12 cycles
5	94	20 s	-
6	58	30 s	-
7	72	35 s	Repeat steps 5-7 for 25 cycles
8	72	2 min	-
9	4		Hold

stomach was rapidly dissected from the abdomen and two full-thickness strips (2 mm × 10 mm) were cut in the direction of the longitudinal muscle layer from each fundal region. One end of each strip was tied to a platinum rod while the other was connected to a force displacement transducer (Grass model FT03) by a silk thread for continuous recording of isometric tension. The transducer was coupled to a polygraph (7K; Grass Instrument). Muscle strips were mounted in 5 mL double-jacketed organ baths containing Krebs-Henseleit solution, which consisted of 118 mmol/L NaCl, 4.7 mmol/L KCl, 1.2 mmol/L MgSO<sub>4</sub>, 1.2 mmol/L KH<sub>2</sub>PO<sub>4</sub>, 25 mmol/L NaHCO<sub>3</sub>, 2.5 mmol/L CaCl<sub>2</sub>, and 10 mmol/L glucose pH 7.4, and bubbled with 95% O<sub>2</sub>/5% CO<sub>2</sub>. Prewarmed water (37 °C) was circulated through the outer jacket of the tissue bath *via* a constant-temperature circulator pump. The temperature of the Krebs-Henseleit solution in the organ bath was maintained within ± 0.5 °C.

Electrical field stimulation (EFS) was applied *via* two platinum wire rings (2 mm diameter, 5 mm apart) through which the preparation was threaded. Electrical impulses (rectangular waves, 80 V, 4-16 Hz, 0.5 ms, for 15 s) were provided by a Grass model S8 stimulator. Strips were allowed to equilibrate for 1 h under an initial load of 0.8 g. During this period, repeated and prolonged washes of the preparations with Krebs-Henseleit solution were done to avoid accumulation of metabolites in the organ baths.

**Drugs**

The following drugs were used: acetyl-β-methylcholine chloride (methacholine), recombinant full-length mouse adiponectin (ADPN), atropine sulphate, carbachol (CCh), guanethidine sulphate, L-N<sup>G</sup>-nitro arginine (L-NNA), tetrodotoxin (TTX). All drugs were obtained from Sigma Chemical (St. Louis, MO, United States). Solutions were prepared on the day of the experiment, except for TTX

and ADPN, for which stock solutions were kept stored at -20 °C.

Drug concentrations are referred as final bath concentrations. For ADPN we started from 20 pmol/L, *i.e.*, the dose previously reported to be effective in *in vitro* gastric preparations<sup>[8]</sup>. Direct smooth muscle contractions were obtained by addition of methacholine to the bath medium. The interval between two subsequent applications of methacholine was no less than 30 min, during which repeated and prolonged washes of the preparations with Krebs-Henseleit solution were performed. In order to investigate the effects of ADPN in non-adrenergic, non-cholinergic (NANC) condition, guanethidine and CCh were added to the bath medium, to rule out the adrenergic and the cholinergic influences, respectively. When contraction elicited by CCh reached a stable plateau phase, EFS or drugs were applied.

**Polymerase chain reaction for ADPN receptors**

Aiming to assess the presence of mRNA for ADPN receptors in gastric fundal tissue, mouse tissues were snap-frozen in liquid nitrogen and stored at -80 °C until use.

Then, from frozen tissue sections, RNA was extracted using RNeasy Mini Kit (Qiagen, Hilden, Germany) following the manufacturer’s instructions. Polymerase chain reaction (PCR) was performed for AdipoR1 and AdipoR2 receptors and the following primers were used:

AdipoR1 forward 5’-CAG AGA AGC TGA CAC AGT GGA G-3’ and reverse 5’-GTC CCT CCC AGA CCT TAT ACA C-3’; AdipoR2 forward 5’-GGA CTC CAG AGC CAG ATA TAC G-3’ and reverse 5’-ACT CTT CCA TTC GTT CCA TAG C-3’.

The mixtures listed in Table 1 were prepared in thin walled 0.2 mL tubes and analysed by Biometra PCR cyclo. PCRs were run with the program reported in Table 2.


Three gastric fundus were evaluated and a negative control without cDNA template as well as a positive control containing cDNA of inguinal fat pad were analyzed in parallel. The positive control was chosen since both Adipo-R1 and Adipo-R2 are well expressed in adipocytes which are the most relevant cell types for these receptors<sup>[18]</sup>.

**Data collection**

Amplitude of contractile responses is expressed as percentage of the muscular contraction evoked by 2 µmol/L methacholine, assumed as 100% or as absolute values (g). Amplitude of contractile responses to methacholine was measured 30 s after a stable plateau phase was reached. Relaxant responses are expressed as percentage decrease relative to the muscular tension induced by 1 µmol/L CCh just before obtaining relaxations. Amplitude values of EFS-induced relaxations refer to the maximal peak obtained during the stimulation period.

**Statistical analysis**

Statistical analysis was performed by means of



**Figure 1** Effects of adiponectin on the neurally-induced contractile responses in strips from the mouse gastric fundus. A: Typical tracing showing the contractile responses to EFS (left hand panel). ADPN (20 nmol/L) decreases the amplitude of the neurally-induced contractile responses (right hand panel); B: Bar charts showing the influence of ADPN (20 nmol/L) on the mean amplitude of the EFS-induced contractile responses in the absence (left hand panel) and in the presence (right hand panel) of L-NNA (200  $\mu$ mol/L). Note that, in the presence of L-NNA, ADPN no longer decreases the amplitude of the neurally-induced excitatory responses in the whole range of stimulation frequency employed. Amplitude of contractile responses is expressed as percentage of the muscular contraction evoked by 2  $\mu$ mol/L methacholine, assumed as 100%. All values are means  $\pm$  SE of 6 strips from 3 mice. <sup>a</sup> $P < 0.05$  vs the control; <sup>b</sup> $P < 0.05$  and  $P > 0.05$  vs the control and vs L-NNA, respectively (Student's *t*-test plus ANOVA and Newman-Keuls post-test). ADPN: Adiponectin; L-NNA: L-N<sup>G</sup>-nitro arginine.

Student's *t*-test to compare two experimental groups or one-way ANOVA followed by Newman-Keuls posttest when more than two groups were compared. Values were considered significantly different with  $P < 0.05$ . Results are given as means  $\pm$  SE. The number of muscle strip preparations is designated by *n* in the results.

## RESULTS

### Mechanical experiments

At basal tension EFS (4-16 Hz) induced ( $n = 22$ ) contractile responses, whose amplitude increased by increasing the stimulation frequency (Figure 1A). As previously observed in the mouse gastric fundus<sup>[16,17]</sup>, the EFS-evoked contractile responses were abolished by TTX (1  $\mu$ mol/L) ( $P < 0.05$ ) or atropine (1  $\mu$ mol/L) ( $P < 0.05$ ), thus indicating that they were neurally-induced and cholinergic in nature.


Addition of ADPN to the bath medium caused, at 20 nmol/L ( $n = 12$ ), a statistically significant reduction ( $P < 0.05$ ) in amplitude of the neurally-induced excitatory

responses in the whole range of stimulation frequency employed (Figure 1A and B). The effects of ADPN were already appreciable 10 min after its addition to the bath medium and persisted up to 60 min (longer time not observed). Addition of the nitric oxide (NO) synthesis inhibitor L-NNA (200  $\mu$ mol/L) to the bath medium ( $n = 6$ ) caused an increase in amplitude of the EFS-induced contractile responses (Figure 1B). In the presence of L-NNA, the depressant effects of ADPN on the neurally-induced contractile responses were no longer observed (Figure 1B).


Addition of the muscarinic receptors agonist methacholine (2  $\mu$ mol/L) to the bath medium ( $n = 4$ ) caused, after 10-15 s of contact time, a sustained contracture which reached a plateau phase (mean amplitude 0.99 g  $\pm$  0.2 g) that persisted until washout. The amplitude of the direct muscular response elicited by methacholine was not influenced by 20 nmol/L ADPN (mean amplitude 0.98  $\pm$  0.3;  $P > 0.05$ ).

The effects of ADPN on the fundal strips were also tested in NANC conditions. As previously observed<sup>[16]</sup>,





**Figure 2** Effects of adiponectin on the neurally-induced relaxant responses in strips from the mouse gastric fundus. A: Typical tracing showing the relaxant responses to EFS (left hand panel). ADPN (20 nmol/L) increases the amplitude of the neurally-induced relaxant responses (right hand panel); B: Bar chart showing the influence of ADPN on the mean amplitude of the EFS-induced relaxant responses. Note that ADPN increases the amplitude of the neurally-induced relaxation in the whole range of stimulation frequency employed. Relaxant responses are expressed as percentage decrease relative to the muscular tension induced by 1  $\mu$ mol/L CCh just before obtaining relaxations. All values are means  $\pm$  SE of 6 strips from 3 mice. <sup>a</sup> $P < 0.05$  vs the control (Student's *t*-test). ADPN: Adiponectin.



**Figure 3** Expression of adiponectin receptors, Adipo-R1 and Adipo-R2, in murine gastric fundus. Typical PCR analysis recorded in the same GF revealed the expression of both AdipoR1 (left hand) and AdipoR2 (right hand) receptors. Positive controls were derived from inguinal fat pad cDNA (Fat) and NC without cDNA template (water). Marker: 100 bp marker (NEB). GF: Gastric fundus; NC: Negative control.

18 strips examined, a slight (mean amplitude 0.15 g  $\pm$  0.02 g) and progressive decay of the basal tension that was not influenced by 1  $\mu$ mol/L TTX ( $P > 0.05$ ). In CCh precontracted strips, EFS (4-16 Hz) elicited ( $n = 16$ ), during the stimulation period, relaxant responses (Figure 2A) whose amplitudes were increased following the addition of ADPN (20 nmol/L) to the bath medium (Figure 2A and B).

### Touchdown-PCR analysis

By touchdown-PCR analysis ADPN receptors expression, analysed using specific primer pairs for Adipo-R1 and Adipo-R2, has been detected in cDNA in three murine gastric fundus. Two bands corresponding to Adipo-R1 and Adipo-R2 were identified in agarose gel electrophoresis. The positive controls were derived from inguinal fat pad cDNA. A typical experiment from a gastric fundus is shown in Figure 3.

CCh (1  $\mu$ mol/L) induced ( $n = 18$ ) a contractile response (mean amplitude 1.1 g  $\pm$  0.3 g) that reached a plateau phase which persisted until washout. Addition of ADPN (20 nmol/L) to the bath medium caused, in 14 out of the

## DISCUSSION

The results of the present study indicate for the first time that ADPN is able to influence the motor responses

in strips from the mouse gastric fundus in which we revealed the expression of both Adipo-R1 and Adipo-R2. Particularly, the ability of ADPN to reduce the amplitude of the neurally-induced contractile responses, without affecting the direct muscular response to methacholine, indicates that the hormone exerts a neuromodulatory action. Furthermore, the lack of effects of ADPN on the contraction to methacholine, other than excluding non-specific effects, demonstrated that muscle responsiveness was not compromised.

It is known that gastric motor responses are the result of a balance between nervous excitatory and inhibitory influences on the smooth muscle and that during EFS both excitatory cholinergic and NANC inhibitory nervous fibres are activated. Thus, the reduction in amplitude of the neurally-induced contractile responses by ADPN, observed in the present experiments, may be ascribable to either a minor activation of the excitatory component or to a major nervous inhibitory influence exerted on the smooth muscle. In this view, NO is considered the main NANC inhibitory neurotransmitter released during EFS to cause gastrointestinal relaxation<sup>[19,20]</sup>. Actually, the increase in amplitude of the EFS-induced contractile responses by the NO synthesis inhibitor L-NNA supports the removal of a nitroergic inhibitory nervous influence. Notably, the observation that ADPN, in the presence of L-NNA, is no longer able to depress the amplitude of the neurally-induced excitatory responses strongly indicates that the nitroergic neurotransmission is involved in the depressant effects of the hormone. This is further confirmed by the ability of ADPN to increase the amplitude of the EFS-induced relaxant responses (elicited by EFS during the stimulation period) whose nitroergic nature has been reported in the mouse gastric fundus<sup>[16,21]</sup>. Actually, NO appears to be a shared target pathway in the hormonal control of gastrointestinal motility<sup>[16,21-24]</sup> and thus ADPN effects too may occur, at least in part, through NO. In this view, the ability of the hormone to increase NO synthase expression has been reported in vascular tissues<sup>[25,26]</sup>.

However, in the present study, the observation that ADPN is also able to cause a TTX-insensitive decay of the basal tension indicates that the hormone, in addition to its neuromodulatory effects, exerts a direct muscular action. Experiments are in progress to better clarify these direct effects of ADPN on gastric smooth muscle.

From a physiological point of view, the ability of ADPN to decrease the amplitude of the neurally-induced contractile responses and to enhance that of the relaxant ones, coupled to a decay of the basal tension, suggests that the hormone effects are directed to facilitate gastric relaxation, so increasing the distension of the gastric wall and in turn the organ capacity. Thus, it could be speculated that the effects of ADPN represent a peripheral satiety signal that might contribute to the anorexigenic effects of the hormone<sup>[4,27]</sup>. In this view, the importance of the gastrointestinal tract in controlling food intake is well recognized<sup>[28]</sup> and the ability of signals generated in this apparatus to influence food intake through the vagal afferent fibers is well established<sup>[29,30]</sup>. In this regard,

the presence of ADPN receptors in both mucosal and muscular vagal afferents in the gastric antrum of mice has been reported<sup>[8]</sup>, suggesting a stomach-vagal-brain pathway for ADPN to modulate satiety signals. Thus, the observed effects of ADPN on gastric motor responses might be regarded as a reinforcing peripheral mechanism engaged by ADPN to regulate food intake.

In conclusion, the present study demonstrates for the first time that ADPN, likely involving the NO pathway, is able to affect the mechanical responses in strips from the mouse gastric fundus. This may furnish the basis to better explore the link between the peripheral and the central effects of ADPN in sending satiety signals and, thus, in the regulation of food intake<sup>[31]</sup>. Interestingly, clinical evidence of an increased NO production in patients affected by eating disorders, such as anorexia nervosa and bulimia nervosa, has been reported<sup>[32]</sup>. Thus, even if caution is needed when transferring data obtained in animal to human, the present results might represent a contribution to consider ADPN as a potential candidate for both novel therapeutic strategies in eating disorders and diagnostic tools.

## ARTICLE HIGHLIGHTS

### Research background

The adipose-tissue derived hormone adiponectin (ADPN) has been reported to have a role as a key neuromodulator of food intake. It is known that gastrointestinal motor phenomena represent a source of peripheral signals involved in the control of feeding behavior. However, no data are available concerning on the effects of ADPN on gastrointestinal motility. This novel information could highlight an additional mechanism engaged by the hormone in the control of food intake.

### Research motivation

The finding that ADPN could influence gastric motility might represent an initial step to regard the peripheral effects of the hormone as possible signals involved in the control of food intake. This aspect, that certainly deserves to be deeper investigated, could lead to consider ADPN as a potential candidate for both novel therapeutic strategies in eating disorders and diagnostic tools.

### Research objectives

The main objective of the present study was to investigate the influence of ADPN on the mechanical responses and the expression of its receptors in the mouse gastric fundus.

### Research methods

For the mechanical experiments, longitudinal muscle strips from the mouse gastric fundus were connected to force displacement transducers for continuous recording of isometric tension. The expression of ADPN receptors in gastric fundal tissue was revealed by touchdown- polymerase chain reaction (PCR) analysis.

### Research results

The present results highlight that, in the mouse gastric strips, ADPN induces inhibitory effects by decreasing the amplitude of the neurally-induced contractile responses, enhancing that of the relaxant ones and causing a decay of the basal tension. Some of these actions appear to be mediated, at least in part, by nitric oxide although further studies are needed to better clarify the mechanism through which the hormone exerts its effects.

### Research conclusions

The results of the present study indicate for the first time that ADPN is able to exert inhibitory effects on the mechanical responses of the mouse gastric

fundus in which we revealed the expression of ADPN receptors. It could be hypothesized that the hormone effects may be directed to facilitate muscle relaxation, so increasing the distension of the gastric wall that represents a peripheral satiety signal. In this view, the results of the present study may furnish the basis to better explore the link between peripheral and central effects of ADPN in the regulation of food intake.

### Research perspectives

The results of the present basic research might furnish a contribution to consider ADPN as a potential candidate for both novel therapeutic strategies in eating disorders and diagnostic tools.

## ACKNOWLEDGMENTS

We thank Mr. Adrio Vannucchi for preparation of the Figures 1 and 2.

## REFERENCES

- 1 **Harwood HJ Jr.** The adipocyte as an endocrine organ in the regulation of metabolic homeostasis. *Neuropharmacology* 2012; **63**: 57-75 [PMID: 22200617 DOI: 10.1016/j.neuropharm.2011.12.010]
- 2 **Fasshauer M, Blüher M.** Adipokines in health and disease. *Trends Pharmacol Sci* 2015; **36**: 461-470 [PMID: 26022934 DOI: 10.1016/j.tips.2015.04.014]
- 3 **Blüher M, Mantzoros CS.** From leptin to other adipokines in health and disease: facts and expectations at the beginning of the 21st century. *Metabolism* 2015; **64**: 131-145 [PMID: 25497344 DOI: 10.1016/j.metabol.2014.10.016]
- 4 **Coope A, Milanski M, Araújo EP, Tambascia M, Saad MJ, Geloneze B, Velloso LA.** AdipoR1 mediates the anorexigenic and insulin/leptin-like actions of adiponectin in the hypothalamus. *FEBS Lett* 2008; **582**: 1471-1476 [PMID: 18394428 DOI: 10.1016/j.febslet.2008.03.037]
- 5 **Stern JH, Rutkowski JM, Scherer PE.** Adiponectin, Leptin, and Fatty Acids in the Maintenance of Metabolic Homeostasis through Adipose Tissue Crosstalk. *Cell Metab* 2016; **23**: 770-784 [PMID: 27166942 DOI: 10.1016/j.cmet.2016.04.011]
- 6 **Wang ZV, Scherer PE.** Adiponectin, the past two decades. *J Mol Cell Biol* 2016; **8**: 93-100 [PMID: 26993047 DOI: 10.1093/jmcb/mjw011]
- 7 **Thundiyil J, Pavlovski D, Sobey CG, Arumugam TV.** Adiponectin receptor signalling in the brain. *Br J Pharmacol* 2012; **165**: 313-327 [PMID: 21718299 DOI: 10.1111/j.1476-5381.2011.01560.x]
- 8 **Kentish SJ, Ratcliff K, Li H, Wittert GA, Page AJ.** High fat diet induced changes in gastric vagal afferent response to adiponectin. *Physiol Behav* 2015; **152**: 354-362 [PMID: 26074203 DOI: 10.1016/j.physbeh.2015.06.016]
- 9 **Yamauchi T, Kamon J, Ito Y, Tsuchida A, Yokomizo T, Kita S, Sugiyama T, Miyagishi M, Hara K, Tsunoda M, Murakami K, Ohteki T, Uchida S, Takekawa S, Waki H, Tsuno NH, Shibata Y, Terauchi Y, Froguel P, Tobe K, Koyasu S, Taira K, Kitamura T, Shimizu T, Nagai R, Kadowaki T.** Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 2003; **423**: 762-769 [PMID: 12802337 DOI: 10.1038/nature01705]
- 10 **González CR, Caminos JE, Gallego R, Tovar S, Vázquez MJ, Garcés MF, Lopez M, García-Caballero T, Tena-Sempere M, Nogueiras R, Diéguez C.** Adiponectin receptor 2 is regulated by nutritional status, leptin and pregnancy in a tissue-specific manner. *Physiol Behav* 2010; **99**: 91-99 [PMID: 19887079 DOI: 10.1016/j.physbeh.2009.10.015]
- 11 **Florian V, Caroline F, Francis C, Camille S, Fabielle A.** Leptin modulates enteric neurotransmission in the rat proximal colon: an in vitro study. *Regul Pept* 2013; **185**: 73-78 [PMID: 23816465 DOI: 10.1016/j.regpep.2013.06.010]
- 12 **Okumura T, Nozu T.** Role of brain orexin in the pathophysiology of functional gastrointestinal disorders. *J Gastroenterol Hepatol* 2011; **26** Suppl 3: 61-66 [PMID: 21443712 DOI: 10.1111/j.1440-1746.2011.06626.x]
- 13 **Squecco R, Garella R, Luciani G, Francini F, Baccari MC.** Muscular effects of orexin A on the mouse duodenum: mechanical and electrophysiological studies. *J Physiol* 2011; **589**: 5231-5246 [PMID: 21911618 DOI: 10.1113/jphysiol.2011.214940]
- 14 **Yarandi SS, Hebbar G, Sauer CG, Cole CR, Ziegler TR.** Diverse roles of leptin in the gastrointestinal tract: modulation of motility, absorption, growth, and inflammation. *Nutrition* 2011; **27**: 269-275 [PMID: 20947298 DOI: 10.1016/j.nut.2010.07.004]
- 15 **Camilleri M.** Peripheral mechanisms in appetite regulation. *Gastroenterology* 2015; **148**: 1219-1233 [PMID: 25241326 DOI: 10.1053/j.gastro.2014.09.016]
- 16 **Garella R, Idrizaj E, Traini C, Squecco R, Vannucchi MG, Baccari MC.** Glucagon-like peptide-2 modulates the nitrergic neurotransmission in strips from the mouse gastric fundus. *World J Gastroenterol* 2017; **23**: 7211-7220 [PMID: 29142468 DOI: 10.3748/wjg.v23.i40.7211]
- 17 **Squecco R, Garella R, Francini F, Baccari MC.** Influence of obestatin on the gastric longitudinal smooth muscle from mice: mechanical and electrophysiological studies. *Am J Physiol Gastrointest Liver Physiol* 2013; **305**: G628-G637 [PMID: 23989009 DOI: 10.1152/ajpgi.00059.2013]
- 18 **Blüher M, Fasshauer M, Kralisch S, Schön MR, Krohn K, Paschke R.** Regulation of adiponectin receptor R1 and R2 gene expression in adipocytes of C57BL/6 mice. *Biochem Biophys Res Commun* 2005; **329**: 1127-1132 [PMID: 15752770 DOI: 10.1016/j.bbrc.2005.02.081]
- 19 **Min YW, Hong YS, Ko EJ, Lee JY, Ahn KD, Bae JM, Rhee PL.** Nitrgenic Pathway Is the Main Contributing Mechanism in the Human Gastric Fundus Relaxation: An In Vitro Study. *PLoS One* 2016; **11**: e0162146 [PMID: 27589594 DOI: 10.1371/journal.pone.0162146]
- 20 **Rand MJ.** Nitrgenic transmission: nitric oxide as a mediator of non-adrenergic, non-cholinergic neuro-effector transmission. *Clin Exp Pharmacol Physiol* 1992; **19**: 147-169 [PMID: 1325878 DOI: 10.1111/j.1440-1681.1992.tb00433.x]
- 21 **Vannucchi MG, Garella R, Cipriani G, Baccari MC.** Relaxin counteracts the altered gastric motility of dystrophic (mdx) mice: functional and immunohistochemical evidence for the involvement of nitric oxide. *Am J Physiol Endocrinol Metab* 2011; **300**: E380-E391 [PMID: 21081707 DOI: 10.1152/ajpendo.00375.2010]
- 22 **Abot A, Lucas A, Bautzova T, Bessac A, Fournel A, Le-Gonidec S, Valet P, Moro C, Cani PD, Knaut C.** Galanin enhances systemic glucose metabolism through enteric Nitric Oxide Synthase-expressed neurons. *Mol Metab* 2018; **10**: 100-108 [PMID: 29428595 DOI: 10.1016/j.molmet.2018.01.020]
- 23 **Idrizaj E, Garella R, Francini F, Squecco R, Baccari MC.** Relaxin influences ileal muscular activity through a dual signaling pathway in mice. *World J Gastroenterol* 2018; **24**: 882-893 [PMID: 29491682 DOI: 10.3748/wjg.v24.i8.882]
- 24 **Squecco R, Garella R, Idrizaj E, Nistri S, Francini F, Baccari MC.** Relaxin Affects Smooth Muscle Biophysical Properties and Mechanical Activity of the Female Mouse Colon. *Endocrinology* 2015; **156**: 4398-4410 [PMID: 26360621 DOI: 10.1210/en.2015-1428]
- 25 **Chen H, Montagnani M, Funahashi T, Shimomura I, Quon MJ.** Adiponectin stimulates production of nitric oxide in vascular endothelial cells. *J Biol Chem* 2003; **278**: 45021-45026 [PMID: 12944390 DOI: 10.1074/jbc.M307878200]
- 26 **Nour-Eldine W, Ghantous CM, Zibara K, Dib L, Issaa H, Itani HA, El-Zein N, Zeidan A.** Adiponectin Attenuates Angiotensin II-Induced Vascular Smooth Muscle Cell Remodeling through Nitric Oxide and the RhoA/ROCK Pathway. *Front Pharmacol* 2016; **7**: 86 [PMID: 27092079 DOI: 10.3389/fphar.2016.00086]
- 27 **Qi Y, Takahashi N, Hileman SM, Patel HR, Berg AH, Pajvani UB, Scherer PE, Ahima RS.** Adiponectin acts in the brain to decrease body weight. *Nat Med* 2004; **10**: 524-529 [PMID: 15077108 DOI: 10.1038/nm1029]
- 28 **Monteiro MP, Batterham RL.** The Importance of the Gastrointestinal Tract in Controlling Food Intake and Regulating

- Energy Balance. *Gastroenterology* 2017; **152**: 1707-1717.e2 [PMID: 28193513 DOI: 10.1053/j.gastro.2017.01.053]
- 29 **Bray GA**. Afferent signals regulating food intake. *Proc Nutr Soc* 2000; **59**: 373-384 [PMID: 10997653 DOI: 10.1017/S0029665100000422]
- 30 **Cummings DE**, Overduin J. Gastrointestinal regulation of food intake. *J Clin Invest* 2007; **117**: 13-23 [PMID: 17200702 DOI: 10.1172/JCI30227]
- 31 **Wilson JL**, Enriori PJ. A talk between fat tissue, gut, pancreas and brain to control body weight. *Mol Cell Endocrinol* 2015; **418** Pt 2: 108-119 [PMID: 26316427 DOI: 10.1016/j.mce.2015.08.022]
- 32 **Vannacci A**, Ravaldi C, Giannini L, Rotella CM, Masini E, Faravelli C, Ricca V. Increased nitric oxide production in eating disorders. *Neurosci Lett* 2006; **399**: 230-233 [PMID: 16495002 DOI: 10.1016/j.neulet.2006.01.060]

**P- Reviewer:** Beales ILP **S- Editor:** Gong ZM **L- Editor:** A  
**E- Editor:** Yin SY





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>



ISSN 1007-9327

