

COMMUNICATION

Direct biocatalysed synthesis of first sulfur-, selenium- and tellurium- containing L-ascorbyl hybrid derivatives with radical trapping and GPx-like properties

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Damiano Tanini,* Beatrice Lupori, Gianni Malevolti, Moira Ambrosi, Pierandrea Lo Nostro, Antonella Capperucci*

6-O-L-ascorbyl selenoesters, thioesters and telluroesters can be efficiently and directly prepared from L-ascorbic acid and suitable functionalised chalcogenoesters through lipase-catalysed transesterification reactions. Novel synthesised L-ascorbyl derivatives exhibited remarkable chain breaking and glutathione peroxidase-like activities.

L-ascorbic acid (vitamin C) is a powerful water soluble antioxidant essential for the correct functioning of the body, being involved in a number of biological processes ranging, amongst others, from the biosynthesis of collagen¹ and catecholamine to the modulation of neurotransmission.²

The enhanced concentration of harmful reactive oxygen species (ROS) has long been related with the onset of several human diseases such as cancer, immune disorders, cystic fibrosis, and neurodegenerative diseases.³ Liposoluble vitamin E and hydrosoluble vitamin C, together with phenolic compounds, carotenoids, and trace elements such as zinc and selenium are the main exogenous defences against oxidative stress. In this context, the design and the development of novel antioxidants have been attracting growing interest over the last decades.⁴ Particularly, owing to their capability to mimic the glutathione peroxidase enzymatic activity, the synthesis of chalcogen-containing antioxidants has recently attracted growing attention. Furthermore, organochalcogenides can possess anticancer, antibacterial, and enzyme inhibitor activities.⁵ The functionalization of bioactive natural products with chalcogens represents an effective strategy to modulate or improve their biological properties. For example, the introduction of chalcogen-containing moieties onto natural compounds such as tocopherols,^{6,7} tocotrienols,⁷ retinol,⁸ hydroxytyrosol,⁹ chrysin,¹⁰ quercetin,^{10a} and resveratrol^{10c} have been extensively studied. However, whilst sulfur-, selenium-, or tellurium-functionalised vitamin E and vitamin A derivatives have been reported, to the best of our knowledge the synthesis of

chalcogen-containing vitamin C derivatives has never been described. The paucity of such results is reasonably due to the instability of the L-ascorbic acid core, which easily undergoes oxidation of the enediol moiety and ring opening of the lactone ring. Furthermore, several alcohol protecting group strategies cannot be applied since the conditions of the endgame protecting group cleavage are often not compatible with chalcogens or with the L-ascorbic acid core. For example, because the presence of chalcogens poisons the Pd-catalyst,¹¹ the most commonly employed benzyl protecting group cannot be cleaved under mild Pd/C catalysed hydrogenation conditions. Additionally, harsh bases- or acid-mediated deprotection procedures cause severe products decomposition.

We sought to approach this problem from a different perspective and evaluated the possibility of applying lipase biocatalysed transesterifications¹² of chalcogen-containing esters and L-ascorbic acid. This approach would ideally allow a straightforward access to chalcogen-containing L-ascorbic acid derivatives, without requiring tedious and detrimental protection/deprotection steps. Lipase B catalysed transesterification is indeed a versatile tool to synthesise L-ascorbyl esters under green and mild conditions.¹³ However, to the best of our knowledge, its application to chalcogen-containing acyl donors has never been described and its feasibility and functional group tolerance were not obvious.

We commenced our studies by establishing the optimal conditions required to promote the biocatalysed transesterification reaction of β -selenoester **2a**, prepared through seleno-Michael addition from benzeneselenol **1a** and methyl acrylate (see ESI), with L-ascorbic acid **3**.

According to a literature survey, the optimal temperature for the enzyme activity was established to be 45°C. Polar solvents, such as acetone and tertiary alcohols are the media of choice for lipase-catalysed L-ascorbic acid esterifications.^{12,13} In our hands, acetone proved to be the most effective, plausibly owing to the high capability to dissolve both **2a** and **3**. Evaluation of different solvents commonly used for Lipase B catalysed reactions, such as *tert*-butanol and 2-methyl-1-butanol gave lower yields. Thus, we investigated the effect of

University of Florence, Department of Chemistry "Ugo Schiff", Via della Lastruccia 13, I-50019 Sesto Fiorentino, Italy

Electronic supplementary information (ESI) available: Full experimental details and products characterisation. See DOI: 10.1039/x0xx00000x

the reaction stoichiometry and the amount of lipase. We found that, whilst poor yields were achieved by using 500 U/mmol of enzyme (Table 1, entries 1-3), a doubling of the enzyme amount brought about a significant yield improvement (Table 1, entry 4).

The optimal reaction time was found to be 48 h, as partial decomposition of the reaction product was observed after 72 h (Table 1, entry 5). On the other hand, shorter times resulted in much lower yields (Table 1, entry 6). Pleasingly, we found that by using an excess of β -selenoester **2a** the desired 6-O-L-ascorbyl ester **4a** was formed in rather good yield for this type of biocatalysed transformations (45%, Table 1, entry 8).

Remarkably, a significant improvement in yield was achieved upon performing the reaction in the presence of 4 A molecular sieves (Table 1, entries 9 and 10). Particularly, by using an excess of **2a** under these conditions, the ascorbyl derivative **4a** was formed in 74% yield, which represents an excellent results for lipase-catalysed transesterification reactions. The striking effect of zeolites can be reasonably ascribed to the trapping of both MeOH and H₂O. Indeed, the removal of methanol produced by the transesterification displaces the equilibrium of the reaction toward the formation of **4a**. Furthermore, by trapping water, molecular sieves hamper the competitive lipase-catalysed hydrolysis of **4a**.


Table 1. Optimization of the biocatalysed synthesis of selenium-containing 6-O-ascorbyl ester **4a**.

Entry	2a (Equiv.)	Lipase B (U/mmol)	Time (h)	Yield (%) ^a	Entry	2a (Equiv.)	Lipase B (U/mmol)	Time (h)	Yield (%) ^a
1	1.0	500	48	12	6	0.3	1000	24	<10
2	0.5	500	48	<10	7	1.0	1000	48	26
3	0.3	500	48	<10	8	3.0	1000	48	45
4	0.3	1000	48	22	9	3.0	1000	48	74 ^c
5	0.3	1000	72	14 ^b	10	0.3	1000	48	32 ^c

^aIsolated yield is reported; ^bPartial decomposition of **4a** was observed; ^c4 A molecular sieves (300 mg) were added. See ESI for details.

Having identified optimal reaction conditions, we then proceeded to investigate the scope of the transformation with respect to different chalcogen-containing esters. Thus, a large variety of differently substituted selenium-, sulfur-, and tellurium-containing esters (Schemes 1 and S1, ESI) was synthesised as reported in the Scheme 1. β -Arylseleno- and β -alkylseleno-esters **2a-g** were smoothly achieved through a novel seleno-Michael addition¹⁴ involving suitable aryl- or alkyl-selenols¹⁵ **1** and methyl acrylate. The reaction occurred under very mild conditions in the presence of Al₂O₃ (Schemes 1 and S1, *via A*). Furthermore, α -selenoesters **2h,i** and γ -selenoester **2j** were easily obtained by exploiting the reactivity of selenols **1** with methyl bromoacetate and ethyl 4-bromobutyrate, respectively (Schemes 1 and S1, *via B*). Similarly, variously substituted and functionalised α -, β -, and γ -thioesters **6a-i** were prepared from the corresponding aryl or alkyl thiols and suitable electrophiles (Scheme S1, *via A* and *via*

B). Disulfide **6j**, bearing two ester functions and four sulfur atoms, was synthesised from 1,9-nonanethiol through a two-step procedure involving a thio-Michael addition and a DCF (dicyanofumarate) mediated oxidation sequence (see ESI for details).¹⁶ Finally, β -aryltelluroesters **8a,b** were prepared from the corresponding ditellurides and methyl 3-bromopropionate (Scheme 1 and S1).




Via B, reagents and conditions. *Conditions i*: R/ArSH **1** or R/ArSeH **5** (1.0 eq.), Cs₂CO₃ (1.0 eq.), TBAI (1.0 eq.), dry DMF, 0°C, bromoester (1.1 eq.), then 0°C to r.t. for 4h; *Conditions ii*: ArTeTeAr **7** (1.0 eq.), NaBH₄ (3.0 eq.), 0°C for 30 min, bromoester (2.2 eq.), then 0°C to r.t. for 4h. See ESI for details

Scheme 1. Synthesis of selenoesters **2**, thioesters **6**, and telluroesters **8**.

Having in hands a wide range of differently substituted and functionalised chalcogen-containing esters, we then explored the scope of the lipase-catalysed transesterification reaction with L-ascorbic acid (Scheme 2). Under the optimised reaction conditions, *o*-, *m*-, and *p*-methoxy phenylselenoesters **2b-d** gave selectively the corresponding 6-O-ascorbyl arylselenoalkanoates **4b-d** in good yield. β -Selenoester **2e**, bearing a *p*-F-C₆H₄ ring, also reacted efficiently with **3** to afford compound **4e**. The biocatalysed transesterification reaction was also successfully applied to the synthesis of L-ascorbyl derivatives **4f,g** bearing different alkylseleno moieties, including the functionalised glycidol derivative (**4g**). Furthermore, methyl and ethyl α - and γ -selenoesters **2h-j** were efficiently transesterified with **3** yielding the corresponding 6-O-L-ascorbyl esters **4h-j** in rather good yields, therefore demonstrating the versatility of the biocatalysed approach towards the synthesis of variously functionalised homologous L-ascorbyl selenoesters.

Next, we turned our attention to evaluating the generality of such reaction with respect to sulfur-containing esters. β -Arylthioesters **6a-d** were smoothly converted into the corresponding sulfurated 6-O-ascorbyl derivatives bearing a phenyl ring (**9a**) or *o*- (**9b**), *m*- (**9c**), and *p*- (**9d**) bromo-substituted benzenes bonded to the S atom. This methodology could also be applied to more interesting highly functionalised alkyl sulfides. The hydroxy-substituted S-alkyl β -thioester **6e** and the enantioenriched epichlorohydrin derivatives (*S*)-**6f** and (*R*)-**6f** were successfully transferred onto the L-ascorbic acid core, affording the corresponding functionalised ester **9e** and the enantioenriched derivatives **9f** and **9g**, containing three controlled stereogenic centers and the further functionalisable chlorinated chain. Additionally, chiral enantioenriched *N*-Tosyl amino-substituted β -thioester **6g**, synthesised from L-valine,^{14,15a} could be efficiently employed to access the enantioenriched S,N-containing 6-O-L-ascorbyl ester **9h**.



Scheme 2. Biocatalysed direct synthesis of selenium-, sulfur-, and tellurium-containing 6-*O*-ascorbyl esters **4**, **9**, **10** from L-ascorbic acid. Isolated yields are reported.

Furthermore, also α - and γ -thioesters **9i** and **9j** were conveniently achieved treating **3** with **6h** and **6j**, respectively, under the above described conditions. Remarkably, the disulfide **6j** was successfully used in a double biocatalysed transesterification to afford the potentially valuable *bola type* bis-ascorbyl ester **9k**, bearing two L-ascorbyl moieties and four sulfur atoms. Indeed, owing to the unique physicochemical properties of *bola type* structures, novel bifunctional chalcogen-containing amphiphilic ascorbic acid derivatives could find wide application both in polymers and material sciences and in medicinal chemistry for the drug delivery of lipophilic molecules.^{12a,17}

Finally, in order to enlarge the scope of this methodology to organotellurium derivatives, we evaluated the reactivity of β -aryltelluroesters **8a,b** with **3**. Under the optimised reaction conditions, 6-*O*-L-ascorbyl aryltelluroalkanoates **10a,b** were directly and selectively formed in good yield, thus highlighting the remarkably broad scope of the procedure and the possibility to use the lipase-catalysed approach with all chalcogens.

Owing to the presence of both the vitamin C free enediol moiety and the selenium or tellurium atom, ascorbyl derivatives **4** and **10** can exhibit both chain breaking and catalytic antioxidant activities, therefore representing excellent antioxidant candidates. Having developed a convenient procedure to access novel chalcogen-containing L-ascorbic acid derivatives, we wished to investigate their antioxidant properties. Pleasingly, according to the DPPH

assay, all the synthesised 6-*O*-ascorbyl esters exhibited remarkable chain breaking activity,¹⁸ leading to a rapid free radical quenching (complete decolouration of ethanolic DPPH solution occurred within 3 seconds) and, thus, showing radical scavenger properties comparable to those of L-ascorbic acid, which represents the most powerful hydrophilic antioxidant. Furthermore, the catalytic thiol peroxidase like activity of selected L-ascorbyl derivatives was also pursued according to literature reported methods using dithiothreitol (DTT)¹⁹ or glutathione (GSH)¹⁹ as substrates (see Scheme S2 and Fig. S2, S3). We were delighted to discover that all tested selenium- and tellurium-containing vitamin C derivatives behave as catalysts in promoting the reduction of hydrogen peroxide in the presence of a thiol cofactor (Table 2). Intriguingly, when the DTT oxidation assay is accomplished using 1.0 eq. of hydrogen peroxide ca. 20% of DTT remained unreacted, thus demonstrating that the reaction of H₂O₂ with the enediol moiety occurs faster with respect to the oxidation of the chalcogen atom, which represents the first step of the GPx-like catalytic cycle. On the other hand, complete DTT oxidation was achieved upon using an excess (1.2 eq.) of H₂O₂.²⁰ As can be noticed (Table 2 and Figure 1), under these conditions all the tested compounds exhibited catalytic thiol-peroxidase like properties, being able to promote the oxidation of both DTT and GSH. Intriguingly, the β -alkylselenoester **4g** showed higher catalytic activity with respect to the arylseleno-substituted analogues **4a-c,e**. Furthermore, according to both assays, L-ascorbyl β -aryltelluroalkanoates **10a,b** behaved as more effective catalysts with respect to the similar selenium-containing derivatives **4a-e**. Furthermore, particularly interesting are the results obtained through the GSH/GR/NADPH coupled test, that better reproduces the cellular environment (Figure 1). Indeed, under these conditions the novel synthesised L-ascorbyl derivatives exhibited GPx-like activity, thus demonstrating the effective enhanced antioxidant properties of these novel amphiphilic systems, thus offering new opportunities for their potential wide application in chemistry, biology, and materials science.

Table 2. Thiol-peroxidase like activity of selenium- and tellurium-containing 6-*O*-L-ascorbyl esters **4** and **10** according to DTT and GSH/GR methods

Entry	Compound	DTT (T ₅₀) ^{a,b}	GSH/GR (T ₅₀) ^{a,c}
1	4a	3405 (\pm 268)	43 (\pm 5)
2	4b	4270 (\pm 325)	48 (\pm 7)
3	4c	4046 (\pm 314)	45 (\pm 5)
4	4e	3862 (\pm 362)	53 (\pm 4)
5	4g	1346 (\pm 151)	37 (\pm 3)
6	4h	2846 (\pm 116)	42 (\pm 6)
7	10a	654 (\pm 104)	18 (\pm 3)
8	10b	386 (\pm 93)	14 (\pm 3)

^aT₅₀ is the time required, in seconds, to halve the initial thiol concentration after the addition of H₂O₂; data in parenthesis are the experimental error. ^bDTT oxidation was monitored by the mean of ¹H NMR spectroscopy; 10 mol% of **4** and 1 mol% of **10** were used. ^cNADPH consumption was monitored by UV spectroscopy (340 nm).




Figure 1. NADPH-coupled GPx assay. Reaction conditions: $[\text{NADPH}]_0 = 0.3 \text{ mM}$, $[\text{GSH}]_0 = 1.0 \text{ mM}$, $[\text{H}_2\text{O}_2]_0 = 2.5 \text{ mM}$, $[\text{GR}] = 4 \text{ units per mL}$, $[\text{catalyst}] = 0.1 \text{ mM}$ in pH 7.4 phosphate buffer at ambient temperature. The mean \pm SD values of three separate experiments are reported.

Preliminary physicochemical characterisation of these novel amphiphilic antioxidants showed that most of them behave as semicrystalline solids. The evolution of the crystal and of the amorphous fraction depends on the thermal history of the sample, a common behaviour of polymers. The DSC curves obtained for **9c** are reported in Fig. S1 (see ESI). The crystalline fraction in **9c** melted at about $75 \text{ }^\circ\text{C}$ producing an endothermic peak with $\Delta H = 21.6 \text{ J/g}$. When the melt was slowly cooled to $-20 \text{ }^\circ\text{C}$ (at $5 \text{ }^\circ\text{C/min}$), crystallisation occurred producing an exothermic peak at about $35.5 \text{ }^\circ\text{C}$ with $\Delta H = 28.2 \text{ J/g}$. The crystalline fraction was found to melt between $40 \text{ }^\circ\text{C}$ and $100 \text{ }^\circ\text{C}$ upon heating. Instead, when the sample was cooled quickly down to $-20 \text{ }^\circ\text{C}$ (at $50 \text{ }^\circ\text{C/min}$), no crystallisation was shown, and a cold crystallisation exothermic peak was detected in the following heating cycle followed by a broad endothermic melting peak. Remarkably, while this behaviour is typical for polymers, it is rather unusual for such small molecules as those synthesized in the present work and may represent an opportunity for their application in material science.

In summary, we have found a convenient, green, mild, and direct route to synthesise novel chalcogen-containing L-ascorbyl derivatives. Such compounds would not have been accessible through classic methodology. Owing to their enhanced chain breaking and catalytic antioxidant properties, these conjugate molecules represent potential valuable systems in biology and in medicinal chemistry.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- R. S. Pinnell, *Yale J Biol Med.*, 1985, **58**, 553.
- R. Figueroa-Méndez and S. Rivas-Arancibia, *Front. Physiol.*, 2015, **6**, 397.
- (a) J. K. Andersen, *Nat. Rev. Neurosci.*, 2004, **5**, S18; (b) R. C. S. Seet, C.-Y. J. Lee, E. C. H. Lima, J. J. H. Tan, A.M. L. Quek, W.-L. Chong, W.-F. Looi, S.-H. Huang, H. Wang, Y.-H. Chand and B. Halliwell, *Free Radical Biol. Med.*, 2010, **48**, 560.
- (a) S. Yoshida, F. Kumakura, I. Komatsu, K. Arai, Y. Onuma, H. Hojo, B. G. Singh, K. I. Priyadarsini and M. Iwaoka, *Angew. Chem., Int. Ed.*, 2011, **50**, 2125; (b) C. W. Nogueira, G. Zeni and J. B. T. Rocha, *Chem. Rev.*, 2004, **104**.

- (a) A. R. Patra, S. S. Roy, A. Basu, A. Bhuniya, A. Bhattacharjee, S. Hajra, U. H. Sk, R. Baral and S. Bhattacharya, *Sci. Rep.*, 2018, **8**, 2194; (b) A. Angeli, D. Tanini, A. Capperucci and C. T. Supuran, *Bioorg. Chem.*, 2018, **76**, 268; (c) A. Angeli, D. Tanini, A. Capperucci, G. Malevolti, F. Turco, M. Ferraroni, C. T. Supuran, *Bioorg. Chem.*, 2018, **81**, 642; (d) A. Angeli, D. Tanini, A. Capperucci and C. T. Supuran, *ACS Med. Chem. Lett.*, 2017, **8**, 1213.
- (a) V. P. Singh, J. F. Poon and L. Engman, *Org. Lett.*, 2013, **15**, 6274.
- P. S. Vraha, C. Drouza, M. P. Rikkou, A. D. Odysseos and A. D. Keramidas, *Bioorg. Med. Chem.*, 2006, **14**, 2684.
- S. C. Welch and M. Gruber, *J. Med. Chem.*, 1979, **22**, 1532.
- G. Rodríguez-Gutiérrez, F. Rubio-Senent, A. Gómez-Carretero, I. Maya, J. Fernández-Bolaños, G. G. Duthie and B. de Roos, *Eur. J. Nutr.*, 2018, doi: 10.1007/s00394-018-1733-y.
- (a) S. F. Fonseca, D. B. Lima, D. Alves, R. G. Jacob, G. Perin, E. J. Lenardão and L. Savegnago, *New J. Chem.*, 2015, **39**, 3043; (b) I. L. Martins, C. Charneira, V. Gandin, J. L. Ferreira da Silva, G. C. Justino, J. P. Telo, A. J. S. C. Vieira, C. Marzano and A. M. M. Antunes, *J. Med. Chem.*, 2015, **58**, 4250; (c) D. Tanini, L. Panzella, R. Amorati, A. Capperucci, E. Pizzo, A. Napolitano, S. Menichetti and M. d'Ischia, *Org. Biomol. Chem.*, 2015, **13**, 5757.
- P. Albers, J. Pietsch and S. F. Parker, *J. Mol. Catal. A: Chem.*, 2001, **173**, 275.
- (a) C. Dolle, P. Magrone, S. Riva, M. Ambrosi, E. Fratini, N. Peruzzi and P. Lo Nostro, *J. Phys. Chem. B*, 2011, **115**, 11638; (b) G. Carrea and S. Riva, *Angew. Chem. Int. Ed.*, 2000, **39**, 222
- (a) K. Nott, A. Brognaux, G. Richard, P. Laurent, A. Favrelle, C. Jérôme, C. Blecker, J. P. Wathélet, M. Paquot and M. Deleu, *Prep. Biochem. Biotechnol.*, 2012, **42**, 348; (b) S. Park, F. Viklund, K. Hulta and R. J. Kazlauskas, *Green Chemistry*, 2003, **5**, 715
- D. Tanini, S. Scarpelli, E. Ermini, A. Capperucci, *Adv. Synth. Catal.*, 2019, DOI 10.1002/adsc.201900168
- (a) D. Tanini, C. Tiberi, C. Gellini, P. R. Salvi, A. Capperucci, *Adv. Synth. Catal.*, 2018, **360**, 3367; (b) A. Angeli, D. Tanini, A. Nocentini, A. Capperucci, M. Ferraroni, P. Gratterer and C. T. Supuran, *Chem. Commun.*, 2019, **55**, 648.
- G. Mlostoń, A. Capperucci, D. Tanini, R. Hamera-Fałdyga and H. Heimgartner, *Eur. J. Org. Chem.*, 2017, 6831
- S. Palma, R.H. Manzo, P. Lo Nostro and D. Allemandi, *Int. J. Pharm.* 2007, **345**, 26.
- (a) E. Tempestini, M. Bucci, V. Mastromartino, M. Gori, D. Tanini, M. Ambrosi, E. Fratini, A. Capperucci and P. Lo Nostro, *ChemPhysChem* 2017, **18**, 1400; (b) P. Lo Nostro, G. Capuzzi, A. Romani and N. Mulinacci, *Langmuir*, 2000, **16**, 1744.
- (a) D. Tanini, A. Grechi, L. Ricci, S. Dei, E. Teodori, A. Capperucci, *New J. Chem.*, 2018, **42**, 6077; (b) F. Kumakura, B. Mishra, K. I. Priyadarsini and M. Iwaoka, *Eur. J. Org. Chem.*, 2010, 440.
- A reasonable catalytic cycle is the following. Ring opening of the oxidised lactone ring cannot be ruled out. See: J. C. Deutsch, *Anal. Biochem.*, 1998, **255**, 1.

