

## The importance of the nursery pathway for the spread of *Phytophthora* species to natural ecosystems in Europe.

Thomas Jung<sup>1,2</sup>, Leszek Orlikowski<sup>3</sup>, Beatrice Henricot<sup>4</sup>, Paloma Abad-Campos<sup>5</sup>, A. G. Aday<sup>6</sup>, Olga Aguín Casal<sup>7</sup>, Jozsef Bakonyi<sup>8</sup>, Santa Olga Cacciola<sup>9</sup>, Thomas Cech<sup>10</sup>, D. Chavarriaga<sup>11</sup>, Tamara Corcobado<sup>12</sup>, Alfredo Cravador<sup>1</sup>, T. Decourcelle<sup>13</sup>, Geoffrey Denton<sup>5</sup>, Stefanos Diamandis<sup>14</sup>, H. T. Doğmuş-Lehtijärvi<sup>7</sup>, A. Franceschini<sup>15</sup>, Beatrice Ginetti<sup>16</sup>, Milka Glavendekić<sup>17</sup>, Jarkko Hantula<sup>18</sup>, Günther Hartmann<sup>19</sup>, Maria Herrero<sup>20</sup>, D. Ivic<sup>21</sup>, Marilia Horta Jung<sup>1</sup>, Arija Lilja<sup>18</sup>, Nenad Keca<sup>17</sup>, Vlodimir Kramarets<sup>22</sup>, A. Lyubenova<sup>23</sup>, Helena Machado<sup>24</sup>, Gaetano Magnano di San Lio<sup>9</sup>, P. J. Mansilla Vázquez<sup>7</sup>, Benoit Marçais<sup>25</sup>, Irina Matsiakh<sup>22</sup>, Ivan Milenkovic<sup>17</sup>, Salvatore Moricca<sup>16</sup>, Jan Nechwatal<sup>26</sup>, Christer Olsson<sup>27</sup>, Tomasz Oszako<sup>28</sup>, Antonella Pane<sup>9</sup>, E. J. Paplomatas<sup>29</sup>, Cristina Pintos Varela<sup>7</sup>, Simone Prospero<sup>30</sup>, C. Rial Martínez<sup>7</sup>, Daniel Rigling<sup>30</sup>, Cecile Robin<sup>13</sup>, A. Rytönen<sup>18</sup>, Maria E. Sánchez<sup>31</sup>, Bruno Scanu<sup>15</sup>, Alexandra Schlenzig<sup>32</sup>, Jörg Schumacher<sup>33</sup>, S. Slavov<sup>23</sup>, Alejandro Solla<sup>12</sup>, Eduardo Sousa<sup>24</sup>, Jan Stenlid<sup>27</sup>, Venche Talgø<sup>20</sup>, Z. Tomic<sup>21</sup>, Panos Tsopelas<sup>34</sup>, Andrea Vannini<sup>35</sup>, Anna M. Vettrai<sup>35</sup>, Marcel Wenneker<sup>36</sup>, Steve Woodward<sup>11</sup>, A.V. Sanz Ros<sup>37</sup>, Ana Pérez-Sierra<sup>38</sup>

<sup>1</sup>Phytophthora Research Centre, Mendel University in Brno, 613 00 Brno, Czech Republic; <sup>2</sup>Phytophthora Research and Consultancy, 83131 Nußdorf, Germany; <sup>3</sup>Institute of Horticulture, Konstytucji 3 Maja 1/3, 96-100 Skierniewice, Poland; <sup>4</sup>Royal Horticultural Society, Wisley, Woking, Surrey GU23 6QB, UK; <sup>5</sup>Grupo de Investigación en Hongos Fitopatógenos, Instituto Agroforestal Mediterráneo, Universidad Politécnica de Valencia, Camino de Vera s/n, 46022 Valencia, Spain; <sup>6</sup>Süleyman Demirel University, Faculty of Forestry, 32260 Isparta, Turkey; <sup>7</sup>EXCMA, Diputación Provincial de Pontevedra, Servicio Agrario, Estación Fitopatología “do Areiro”, Subida a la Robleda S/N, 36153 Pontevedra, Spain; <sup>8</sup>Plant Protection Institute, Hungarian Academy of Sciences, Herman Ottó út 15, 1022 Budapest, Hungary; <sup>9</sup>Department of Agriculture, Food and Environment (Di3A), University of Catania, via S. Sofia 100, 95123 Catania, Italy; <sup>10</sup>Federal Research and Training Centre for Forests, Natural Hazards and Landscape (BFW), Seckendorff-Gudent-Weg 8, A-1131 Vienna, Austria; <sup>11</sup>Institute of Biological and Environmental Sciences, University of Aberdeen, Cruickshank Building, Aberdeen AB24 3UU, UK; <sup>12</sup>Ingeniería Técnica Forestal, Universidad de Extremadura, Avenida Virgen del Puerto 2, 10600 Plasencia, Spain; <sup>13</sup>INRA, UMR BioGeCo, 69 route d'Arcachon, 33612 Cestas Cedex, France; <sup>14</sup>NAGREF-Forest Research Institute, 570 06 Vassilika-Thessaloniki, Greece; <sup>15</sup>Dipartimento di Agraria, Sezione di Patologia vegetale ed Entomologia (SPaVE), Università degli Studi di Sassari, Viale Italia, 39, 07100 – Sassari, Italy; <sup>16</sup>Department of Agricultural Biotechnology, Section of Plant Protection, University of Florence, Piazzale delle Cascine 28, I-50144 Florence, Italy; <sup>17</sup>Faculty of Forestry, University of Belgrade, Kneaza Viseslava 1, 11030 Belgrade, Serbia; <sup>18</sup>Finnish Forest Research Institute, Vantaa Research Centre, P.O. Box 18, 01301 Vantaa, Finland; <sup>19</sup>Lower Saxony Forest Research Station, Grätzelstrasse 2, D-37079 Göttingen, Germany; <sup>20</sup>Norwegian Institute for Agricultural and Environmental Research (Bioforsk), Plant Health and Plant Protection Division, Høgskoleveien 7, 1432 Ås, Norway; <sup>21</sup>Institute for Plant Protection, Croatian Centre for Agriculture, Food and Rural Affairs, Rim 98, 10000 Zagreb, Croatia; <sup>22</sup>National Forestry University of

Ukraine, Forestry Department, General Chuprynki Str. 103, 79057 Lviv, Ukraine; <sup>23</sup>Agrobiointitute, Centre of Excellence in Plant Biotechnology, 8 Dragan Tsankov Blvd., 1164 Sofia, Bulgaria; <sup>24</sup>Instituto Nacional de Recursos Biológicos, Unidade de Silvicultura e Produtos Florestais, Quinta do Marquês, 2780-159 Oeira, Portugal; <sup>25</sup>INRA, Nancy Université, UMR 1136 Interactions Arbres/Microorganismes, IFR 110, F-54280 Champenoux, France; <sup>26</sup>Bavarian State Institute of Agriculture (LfL), D-85354 Freising, Germany; <sup>27</sup>Department of Forest Mycology and Pathology, Swedish University of Agricultural Sciences, PO Box 7026, 750 07, Uppsala, Sweden; <sup>28</sup>Department of Forest Phytopathology, Forest Research Institute, ul. Braci Lesnej 3, 05-090 Raszyn, Poland; <sup>29</sup>Laboratory of Plant Pathology, Agricultural University of Athens, 75 Iera Odos, 11855 Athens, Greece; <sup>30</sup>Swiss Federal Institute for Forest, Snow and Landscape Research WSL, CH-8903 Birmensdorf, Switzerland; <sup>31</sup>Departamento Agronomía, ETSIAM, Universidad de Córdoba. Apdo. 3048, 14080-Córdoba, Spain; <sup>32</sup>Scottish Government-SASA, Plant Health, Roddinglaw Rd, Edinburgh, EH12 9FJ, UK; <sup>33</sup>University of Applied Sciences, Eberswalde, Germany; <sup>34</sup>NAGREF-Institute of Mediterranean Forest Ecosystems, Terma Alkmanos, 11528 Athens, Greece; <sup>35</sup>Department for Innovation in Biological, Agro-Food and Forest Systems, University of Tuscia, Via S. Camillo de Lellis, 01100 Viterbo, Italy; <sup>36</sup>Wageningen UR/Applied Plant Research, Sector Flower bulbs, Tree nursery & Fruit, Lingewal 1, 6668 LA Randwijk, 6670 AE Zetten, the Netherlands; <sup>37</sup>Centro de Sanidad Forestal de Calabazanos, 34190, Palencia, Spain; <sup>38</sup>Forest Research, Alice Holt Lodge, Farnham, Surrey GU10 4LH, UK.

An analysis of incidence of *Phytophthora* spp. in 732 European nurseries producing forest transplants, larger specimen trees, landscape plants and ornamentals, plus 2525 areas in which trees and shrubs were planted, is presented based on work conducted by 38 research groups in 23 European countries between 1972 and 2013. Forty-nine *Phytophthora* taxa were recorded in 670 nurseries (91.5%); within these nurseries, 1614 of 1992 nursery stands (81.0%) were infested, although most affected plants appeared healthy. In forest and landscape plantings, 56 *Phytophthora* taxa were recovered from 1667 of 2525 tested sites (66.0%). Affected plants frequently showed symptoms such as crown thinning, chlorosis and dieback caused by extensive fine root losses and/or collar rot. Many well-known highly damaging host–*Phytophthora* combinations were frequently detected but 297 and 407 new *Phytophthora*–host associations were also observed in nurseries and plantings, respectively. On average, 1.3 *Phytophthora* species/taxa per infested nursery stand and planting site were isolated. At least 47 of the 68 *Phytophthora* species/taxa detected in nurseries and plantings were exotic species several of which are considered well established in both nurseries and plantings in Europe. Seven known *Phytophthora* species/taxa were found for the first time in Europe, while 10 taxa had not been previously recorded from nurseries or plantings; in addition, 5 taxa were first detections on woody plant species. Seven *Phytophthora* taxa were previously unknown to science. The reasons for these failures of plant biosecurity in Europe, implications for forest and semi-natural ecosystems and possible ways to improve biosecurity are discussed.