

PROF. OLIVER PFAAR (Orcid ID : 0000-0003-4374-9639)  
DR. NIKOLAOS G PAPAPOULOS (Orcid ID : 0000-0002-4448-3468)  
PROF. IOANA AGACHE (Orcid ID : 0000-0001-7994-364X)  
PROF. CLAUS BACHERT (Orcid ID : 0000-0003-4742-1665)  
DR. VICTORIA CARDONA (Orcid ID : 0000-0003-2197-9767)  
PROF. THOMAS B CASALE (Orcid ID : 0000-0002-3149-7377)  
PROF. STEPHEN R. DURHAM (Orcid ID : 0000-0001-5264-6207)  
PROF. WYTSKE WJ FOKKENS (Orcid ID : 0000-0003-4852-229X)  
DR. ROY GERTH VAN WIJK (Orcid ID : 0000-0002-9608-8742)  
DR. TARI HAAHTELA (Orcid ID : 0000-0003-4757-2156)  
DR. TOMOHISA IINUMA (Orcid ID : 0000-0002-9940-5520)  
DR. JÖRG KLEINE-TEBBE (Orcid ID : 0000-0002-2862-7353)  
PROF. MAREK LESZEK KOWALSKI (Orcid ID : 0000-0002-8442-2774)  
DR. DÉSIRÉE ERLINDA LARENAS-LINNEMANN (Orcid ID : 0000-0002-5713-5331)  
PROF. KARIN C. LØDRUP CARLSEN (Orcid ID : 0000-0002-9257-1198)  
PROF. OLGA LOURENÇO (Orcid ID : 0000-0002-8401-5976)  
DR. JOAQUIM MULLOL (Orcid ID : 0000-0003-3463-5007)  
PROF. HAE-SIM PARK (Orcid ID : 0000-0003-2614-0303)  
PROF. GIOVANNI PASSALACQUA (Orcid ID : 0000-0002-5139-3604)  
PROF. GRAHAM C ROBERTS (Orcid ID : 0000-0003-2252-1248)  
DR. DERMOT RYAN (Orcid ID : 0000-0002-4115-7376)  
DR. MOHAMED H SHAMJI (Orcid ID : 0000-0003-3425-3463)  
PROF. ULRICH WAHN (Orcid ID : 0000-0002-5723-6132)  
PROF. TORSTEN ZUBERBIER (Orcid ID : 0000-0002-1466-8875)

Article type : EAACI Position Paper

## 2019 ARIA Care pathways for allergen immunotherapy

**Short informative title** Many guidelines or national practice guidelines have been produced but many are complex and none propose care pathways.

**Short title: 2019 ARIA in allergen immunotherapy**

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/all.13805

This article is protected by copyright. All rights reserved.

J Bousquet (1, 2)\*, O Pfaar (3)\*, A Togias (4)\*\*, HJ Schünemann (5), I Ansoategui (6), NG Papadopoulos (7), I Tsiligianni (8), I Agache (9), JM Anto (10-13), C Bachert (14), A Bedbrook (1), KC Bergmann (15), S Bosnic-Anticevich (16), I Bosse (17), J Brozek (5), M Calderon (18), GW Canonica (19), L Caraballo (20), V Cardona (21), T Casale (22), L Cecchi (23), DK Chu (5), E Costa (24), AA Cruz (25), W Czarlewski (27), SR Durham (26), G Du Toit (27), M Dykewicz (28), M Ebisawa (29), JL Fauquert (30), M Fernandez-Rivas (31), WJ Fokkens (32), J Fonseca (33), JF Fontaine (34), R Gerth van Wijk (35), T Haahtela (36), S Halken (37), PW Hellings (38), D Ierodiakonou (8), T Iinuma (39), JC Ivancevich (40), L Jacobsen (41), M Jutel (42), I Kaidashev (43), M Khaitov (44), O Kalayci (45), J Kleine Tebbe (46), L Klimek (46), ML Kowalski (47), P Kuna (48), V Kvedariene (49), S La Grutta (50), D Larenas-Linemann (51), S Lau (52), D Laune (53), L Le (54), K Lodrup Carlsen (55), O Lourenço (56), HJ Malling (57), G Marien (58), E Menditto (59), G Mercier (60), J Mullol (61), A Muraro (62), R O'Hehir (63), Y Okamoto (39), GB Pajno (64), HS Park (65), P Panzner (66), G Passalacqua (67), N Pham-Thi (68), G Roberts (69), C Rolland (70), N Rosario (71), D Ryan (72), B Samolinski (73), M Sanchez-Borges (74), G Scadding (75), MH Shamji (76), A Sheikh (77), GJ Sturm (78), A Todo Bom (79), S Toppila-Salmi (80), M Valentin-Rostan (81), A Valiulis (82), E Valovirta (83), MT Ventura (84), U Wahn (85), S Walker (86), D Wallace (87), S Wasserman (88), A Yorgancioglu (89), T Zuberbier (15) and the ARIA Working Group

\*: These two authors contributed equally to the paper

\*\* : Dr. Togias' co-authorship of this publication does not constitute endorsement by the US National Institute of Allergy and Infectious Diseases or by any other United States government agency

1. MACVIA-France, Fondation partenariale FMC VIA-LR, Montpellier, France.
2. INSERM U 1168, VIMA : Ageing and chronic diseases Epidemiological and public health approaches, Villejuif, Université Versailles St-Quentin-en-Yvelines, UMR-S 1168, Montigny le Bretonneux, France, Euforea, Brussels, Belgium, and Charité, Universitätsmedizin Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Comprehensive Allergy Center, Department of Dermatology and Allergy, Berlin, Germany
3. Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Germany.
4. Division of Allergy, Immunology, and Transplantation (DAIT), National Institute of Allergy and Infectious Diseases, NIH, Bethesda, US.
5. Department of Health Research Methods, Evidence and Impact, Division of Immunology and Allergy, McMaster University, Hamilton, ON, Canada.
6. Hospital Quirónsalud Bizkaia, Bilbao, Spain.
7. Division of Infection, Immunity & Respiratory Medicine, Royal Manchester Children's Hospital, University of Manchester, Manchester, UK, and Allergy Department, 2nd Pediatric Clinic, Athens General Children's Hospital "P&A Kyriakou," University of Athens, Athens, Greece

8. Clinic of Social and Family Medicine, Faculty of Medicine, University of Crete, Crete, Greece.
9. Faculty of Medicine, Transylvania University, Brasov, Romania.
10. ISGloBAl, Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain.
11. IMIM (Hospital del Mar Research Institute), Barcelona, Spain.
12. Universitat Pompeu Fabra (UPF), Barcelona, Spain.
13. CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain.
14. Upper Airways Research Laboratory, ENT Dept, Ghent University Hospital, Ghent, Belgium.
15. Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin and Berlin Institute of Health, Comprehensive Allergy-Centre, Department of Dermatology and Allergy, member of GA<sup>2</sup>LEN, Berlin, Germany.
16. Woolcock Institute of Medical Research, University of Sydney and Woolcock Emphysema Centre and Local Health District, Glebe, NSW, Australia.
17. Allergist, La Rochelle, France.
18. Quebec Heart and Lung Institute, Laval University, Québec City, Quebec, Canada.
19. Calderon. Imperial College London - National Heart and Lung Institute, Royal Brompton Hospital NHS, London, UK.
20. Personalized Medicine Clinic Asthma & Allergy, Humanitas University, Humanitas Research Hospital, Rozzano, Milan, Italy
21. Institute for Immunological Research, University of Cartagena, Campus de Zaragocilla, Colombia, and Foundation for the Development of Medical and Biological Sciences (Fundemeb), Cartagena, Colombia.
22. Allergy Section, Department of Internal Medicine, Hospital Vall d'Hebron & ARADyAL research network, Barcelona, Spain.
23. Division of Allergy/Immunology, University of South Florida, Tampa, USA
24. SOS Allergology and Clinical Immunology, USL Toscana Centro, Prato, Italy.
25. UCIBIO, REQUIMTE, Faculty of Pharmacy, and Competence Center on Active and Healthy Ageing of University of Porto (AgeUPNetWork), University of Porto, Portugal
26. ProAR – Nucleo de Excelencia em Asma, Federal University of Bahia, Brasil and WHO GARD Planning Group, Brazil.
27. Medical Consulting Czarlewski, Levallois, France.
28. Allergy and Clinical Immunology Section, National Heart and Lung Institute, Imperial College London, UK.
29. Guy's and St Thomas' NHS Trust, Kings College London, UK.
30. Section of Allergy and Immunology, Saint Louis University School of Medicine, Saint Louis, Missouri, USA.
31. Clinical Research Center for Allergy and Rheumatology, Sagami-hara National Hospital, Sagami-hara, Japan.
32. Unité de pneumo-allergologie de l'enfant, pôle pédiatrique, CHU de Clermont-Ferrand-Estaing, Clermont-Ferrand, France.
33. Allergy Department, Hospital Clinico San Carlos, IdiSSC, Madrid, Spain.
34. Department of Otorhinolaryngology, Academic Medical Centres, AMC, Amsterdam, the Netherlands.

35. CINTESIS, Center for Research in Health Technology and Information Systems, Faculdade de Medicina da Universidade do Porto; and Medida, Lda Porto, Portugal
36. Allergist, Reims, France.
37. Department of Internal Medicine, section of Allergology, Erasmus MC, Rotterdam, the Netherlands.
38. Skin and Allergy Hospital, Helsinki University Hospital, and University of Helsinki, Helsinki, Finland.
39. Hans Christian Andersen Children's Hospital, Odense University Hospital, Odense, Denmark.
40. Dept of Otorhinolaryngology, Univ Hospitals Leuven, Belgium, and Academic Medical Center, Univ of Amsterdam, The Netherlands and Euforea, Brussels, Belgium.
41. Dept of Otorhinolaryngology, Chiba University Hospital, Chiba, Japan.
42. Servicio de Alergia e Immunologia, Clinica Santa Isabel, Buenos Aires, Argentina.
43. Allergy Learning and Consulting, Copenhagen, Denmark.
44. Department of Clinical Immunology, Wrocław Medical University, Poland.
45. Ukrainian Medical Stomatological Academy, Poltava, Ukraine.
46. National Research Center, Institute of Immunology, Federal Medicobiological Agency, Laboratory of Molecular immunology, Moscow, Russian Federation.
47. Pediatric Allergy and Asthma Unit, Hacettepe University School of Medicine, Ankara, Turkey.
48. Allergy & Asthma Center Westend, Berlin, Germany.
49. Department of Immunology and Allergy, Healthy Ageing Research Center, Medical University of Lodz, Poland and Sach's Childrend and Youth Hospital, Södersjukhuset, Stockholm, Sweden
50. Division of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Poland.
51. Institute of Biomedical Sciences, Department of Pathology, Faculty of Medicine, Vilnius University and Institute of Clinical medicine, Clinic of Chest diseases and Allergology, faculty of Medicine, Vilnius University, Vilnius, Lithuania
52. Institute of Biomedicine and Molecular Immunology (IBIM), National Research Council (CNR), Palermo, Italy.
53. Center of Excellence in Asthma and Allergy, Médica Sur Clinical Foundation and Hospital, México City, Mexico.
54. Department of Pediatric Pneumology and Immunology, Charité Universitätsmedizin, Berlin, Germany.
55. KYomed INNOV, Montpellier, France.
56. University of Medicine and Pharmacy, Hochiminh City, Vietnam.
57. Oslo University Hospital, Department of Paediatrics, Oslo, and University of Oslo, Faculty of Medicine, Institute of Clinical Medicine, Oslo, Norway.
58. Faculty of Health Sciences and CICS – UBI, Health Sciences Research Centre, University of Beira Interior, Covilhã, Portugal.
59. Danish Allergy Centre, University of Copenhagen, Copenhagen, Denmark.
60. EUFOREA, Bussels, Belgium.
61. CIRFF, Center of Pharmacoeconomics, University of Naples Federico II, Naples, Italy.
62. Unité Médico-Economie, Département de l'Information Médicale, University Hospital, Montpellier, France

63. Rhinology Unit & Smell Clinic, ENT Department, Hospital Clínic; Clinical & Experimental Respiratory Immunoallergy, IDIBAPS, CIBERES, University of Barcelona, Spain.
64. Food Allergy Referral Centre Veneto Region, Department of Women and Child Health, Padua General University Hospital, Padua, Italy.
65. Department of Allergy, Immunology and Respiratory Medicine, Alfred Hospital and Central Clinical School, Monash University, Melbourne, Victoria, Australia.
66. Department of Pediatrics, Allergy Unit, University of Messina, Messina, Italy.
67. Department of Allergy and Clinical Immunology, Ajou University School of Medicine, Suwon, South Korea.
68. Department of Immunology and Allergology, Faculty of Medicine in Pilsen, Charles University in Prague, Pilsen, Czech Republic.
69. Allergy and Respiratory Diseases, Ospedale Policlinico San Martino -University of Genoa, Italy
70. Allergy department, Pasteur Institute, Paris, France.
71. David Hide Centre, St Mary's Hospital, Isle of Wight and University of Southampton, Southampton, UK.
72. Association Asthme et Allergie, Paris, France.
73. Hospital de Clinicas, University of Parana, Brazil.
74. Honorary Clinical Research Fellow, Allergy and Respiratory Research Group, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Medical School, Edinburgh, UK
75. Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Poland.
76. Allergy and Clinical Immunology Department, Centro Medico-Docente La Trinidad, Caracas, Venezuela.
77. The Royal National TNE Hospital, University College London, UK.
78. Immunomodulation and Tolerance Group, Imperial College London, and Allergy and Clinical Immunology, Imperial College London, London, UK.
79. The Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK.
80. Department of Dermatology and Venerology, Medical University of Graz, Graz, Austria and Outpatient Allergy Clinic Reumannplatz, Vienna, Austria.
81. Imunoalergologia, Centro Hospitalar Universitário de Coimbra and Faculty of Medicine, University of Coimbra, Portugal.
82. Skin and Allergy Hospital, Helsinki University Hospital and University of Helsinki, Helsinki, Finland.
83. Allergist, Montevideo, Uruguay.
84. Vilnius University Institute of Clinical Medicine, Clinic of Children's Diseases, and Institute of Health Sciences, Department of Public Health, Vilnius, Lithuania; European Academy of Paediatrics (EAP/UEMS-SP), Brussels, Belgium.
85. Department of Lung Diseases and Clinical Immunology, University of Turku and Terveystalo allergy clinic, Turku, Finland.
86. University of Bari Medical School, Unit of Geriatric Immunoallergy, Bari, Italy.
87. Pediatric Department, Charité, Berlin, Germany.

88. Asthma UK, Mansell street, London, UK.

89. Nova Southeastern University, Fort Lauderdale, Florida, USA.

90. Department of Medicine, Clinical Immunology and Allergy, McMaster University, Hamilton, Ontario, Canada.

91. Department of Pulmonary Diseases, Celal Bayar University, Faculty of Medicine, Manisa, Turkey

## Address for correspondence

Professor Jean Bousquet

CHU Montpellier, 371 Avenue du Doyen Gaston Giraud, 34295 Montpellier Cedex 5, France

Tel +33 611 42 88 47 jean.bousquet@orange.fr

## Conflicts of interest:

Dr Claus Bachert reports personal fees from Uriach.

Dr Bosnic-Anticevich reports personal fees from TEVA, Boehringer Ingelheim, Sanofi, AstraZeneca, GSK, grants from TEVA, MEDA, outside the submitted work.

Dr Bousquet reports personal fees and other from Chiesi, Cipla, Hikma, Menarini, Mundipharma, Mylan, Novartis, Sanofi-Aventis, Takeda, Teva, Uriach, other from Kyomed, outside the submitted work.

Dr. Cardona reports personal fees from Allergopharma, personal fees from ALK, personal fees from Diater, personal fees from Leti, personal fees from Thermofisher, outside the submitted work.

Dr. Casale reports grants and personal fees from Stallergenes, outside the submitted work.

Dr. Cecchi reports personal fees from Menarini, Malesci, ALK, outside the submitted work.

Dr. Cruz reports grants and personal fees from GSK, personal fees and non-financial support from Boehringer Ingelheim, Astrazeneca, MEDA, personal fees from Novartis, Boston Scientific, EUROFARMA, non-financial support from CHIESI, outside the submitted work.

Dr. Durham reports personal fees from Adiga, Anergis, ALK, Allergopharma, MedicalUpdate GmBC, UCB, outside the submitted work.

Dr Du Toit reports personal fees from Stallergenes, outside the submitted work. Dr. Ebisawa reports personal fees from DBV Technologies, Mylan EPD, maruho, Shionogi & CO., Ltd., Kyorin Pharmaceutical Co., Ltd., Thermofisher Diagnostics Pfizer, Beyer, Nippon Chemifar, Takeda Pharmaceutical Co., Ltd., MSD, outside the submitted work.

Dr. Gerth van Wijk reports personal fees from ALK Abello, Allergopharma, outside the submitted work.

Dr. Haahtela reports personal fees from Orion Pharma, Mundipharma, during the conduct of the study.

Dr. Hellings reports grants from Mylan, other from Stallergenes, Allergopharma, Sanofi, during the conduct of the study; grants from Mylan.

Dr. Ivancevich reports personal fees from Eurofarma Argentina, Faes Farma, other from Sanofi, Laboratorios Casasco, outside the submitted work.

Dr. Kvedariene reports personal fees from GSK, Berlin Chemie Menarini, non-financial support from StallergenGreer, Mylan, AstraZeneca, outside the submitted work.

Larenas Linnemann reports personal fees from GSK, Astrazeneca, MEDA, Boehringer Ingelheim, Novartis, Grunenthal, UCB, Amstrong, Siegfried, DBV Technologies, MSD, Pfizer. grants from Sanofi, Astrazeneca, Novartis, UCB, GSK, TEVA, Chiesi, Boehringer Ingelheim, outside the submitted work.

Dr Okamoto reports personal fees from Shionogi Co. Ltd., Torii Co. Ltd., GSK, MSD, Kyowa Co. Ltd., from Eizai Co. Ltd., grants and personal fees from Kyorin Co. Ltd., Tiho Co. Ltd., grants from Yakuruto Co. Ltd., Yamada Bee Farm, outside the submitted work.

Dr Papadopoulos reports personal fees from Abbvie Novartis, Faes Farma, BIOMAY, HAL, Nutricia Research, Menarini, Novartis, MEDA, MSD, Omega Pharma, Danone, grants from Menarini, outside the submitted work

Dr. Panzner reports personal fees from ALK, AstraZeneca, Novartis, Stallergenes, outside the submitted work.

Dr. Pfaar reports grants and personal fees from ALK-Abelló, Allergopharma, Stallergenes Greer, HAL Allergy Holding B.V./HAL Allergie GmbH, Bencard Allergie GmbH/Allergy Therapeutics, Lofarma, ASIT Biotech Tools S.A., Laboratorios LETI/LETI Pharma, Anergis S.A., grants from Biomay, Nuvo, Circassia, personal fees from Novartis Pharma, MEDA Pharma, Mobile Chamber Experts (a GA2LEN Partner), Pohl-Boskamp, Indoor Biotechnologies, grants from Glaxo Smith Kline, outside the submitted work.

Dr. Scadding reports personal fees from ALK, Sequirus, outside the submitted work.

Dr. Sheikh reports grants from EAACI, outside the submitted work.

Dr. Todo-Bom reports grants and personal fees from Teva, Mundipharma, personal fees from AstraZeneca, GSK, Novartis, grants from Bial, grants from Leti, outside the submitted work.

Dr. Wallace reports other from ALK, outside the submitted work.

Waserman reports personal fees from Merck, GSK, Novartis, Behring, Shire, Sanofi, Barid Aralez, Mylan Meda, PEDIAPHARM outside the submitted work.

Dr. Zuberbier reports and Organizational affiliations: Committee member: WHO-Initiative "Allergic Rhinitis and Its Impact on Asthma" (ARIA). Member of the Board: German Society for Allergy and Clinical Immunology (DGAKI). Head: European Centre for Allergy Research Foundation (ECARF) ; Secretary General: Global Allergy and Asthma European Network (GA<sup>2</sup>LEN). Member: Committee on Allergy Diagnosis and Molecular Allergology, World Allergy Organization (WAO).

## Abstract

Allergen immunotherapy (AIT) is a proven therapeutic option for the treatment of allergic rhinitis and/or asthma. Many guidelines or national practice guidelines have been produced but the evidence-based method varies, many are complex and none propose care pathways. This paper reviews care pathways for AIT using strict criteria and provides simple recommendations that can be used by all stakeholders including health professionals. The decision to prescribe AIT for the patient should be individualized and based on the relevance of the allergens, the persistence of symptoms despite appropriate medications according to guidelines as well as on the availability of good-quality and efficacious extracts. Allergen extracts cannot be regarded as generics. Immunotherapy is selected by specialists for stratified patients. There are no currently available validated biomarkers that can predict AIT success. In adolescents and adults, AIT should be reserved for patients with moderate/severe rhinitis or for those with moderate asthma who, despite appropriate pharmacotherapy and adherence, continue to exhibit exacerbations that appear to be related to allergen exposure, except in some specific cases. Immunotherapy may be even more advantageous in patients with multimorbidity. In children, AIT may prevent asthma onset in patients with rhinitis. mHealth tools are promising for the stratification and follow up of patients.

## Key words

Allergen immunotherapy, asthma, rhinitis, children, stratification, mHealth

## Abbreviations

AIT: Allergen immunotherapy

AR: Allergic rhinitis

ARIA: Allergic Rhinitis and its Impact on Asthma

CDSS: Clinical decision support system

CRD: Chronic respiratory disease

DB-PC-RCT: Double-blind, placebo-controlled, randomized trial

EIP on AHA: European Innovation Partnership on Active and Healthy Ageing

EIT: European Institute for Innovation and Technology

EU: European Union

GRADE: Grading of Recommendations Assessment, Development and Evaluation

ICER: Incremental cost effectiveness ratio

ICP: Integrated care pathway

JA-CHRODIS: Joint Action on Chronic Diseases and Promoting Healthy Ageing across the Life Cycle

MACVIA: Fighting chronic diseases for active and healthy ageing

MASK-air<sup>®</sup>: (formerly Allergy Diary)

MASK: Mobile Airways Sentinel Network

NICE: National Institute for Health and Clinical Excellence (UK)

PCP: Primary health care professional

QALY: Quality-adjusted life year

QOL: Quality-of-life

RCT: Randomized controlled trials

RWE: Real-world evidence

SmPC: Summary of product characteristics

SCIT: Subcutaneous immunotherapy

SCUAD: Severe Chronic Upper Airway Disease

SLIT: Sublingual immunotherapy

WHO: World Health Organization

## Introduction

In all societies, the burden and cost of allergic diseases are increasing rapidly and “change management” strategies are needed to support the transformation of the health care system for integrated care. As an example for allergic disease care, the newest ARIA (Allergic Rhinitis and its Impact on Asthma) project (ARIA phase 4) (1, 2) and POLLAR (Impact of Air POLLution on Asthma and Rhinitis, EIT Health) (3) are proposing digitally-enabled, integrated, person-centred care for rhinitis and asthma multimorbidity embedding environmental exposure (2, 4).



Integrated Care Pathways (ICPs) are structured multi-disciplinary care plans detailing the key steps of patient care (5). They promote the translation of guideline recommendations into local protocols and their application to clinical practice (6, 7). ICPs should integrate recommendations from clinical practice guidelines, but they usually enhance recommendations by combining interventions iteratively, integrate quality assurance and offer recommendation on the coordination of care.

Allergen immunotherapy (AIT) is a proven therapeutic option for the treatment of allergic rhinitis and/or asthma for many standardized products by sublingual (SLIT) or sub-cutaneous (SCIT) routes (8-14). Studies using prescription databases have recently found that the efficacy demonstrated in double-blind, placebo-controlled, randomized clinical trials (DB-PC-RCT) translates into real life (15). In most countries, AIT is more expensive than other medical treatments for allergic rhinitis (AR) and should therefore be considered in patients within a stratified medicine approach (16). Many international and national guidelines on AIT (8-14, 17) have been produced but the evidence-based method varies, many are complex and none propose ICPs.

The aim of the present publication is to develop the ARIA ICPs for both SCIT and SLIT that were proposed by a EAACI Task Force (18).

## **1. Development of the document**

The original draft of this document was prepared by JB and was circulated to several authors for comments. A questionnaire (Annex 1) was also circulated and the answers were collected.

The document and the questionnaire answers were reviewed during a meeting including ARIA, EIT Health (POLLAR: Impact of Air POLLution on Asthma and Rhinitis) (3), the European Innovation Partnership on Active and Healthy Ageing (19) and the Global Alliance against Chronic Respiratory Diseases (GARD, WHO Alliance) with the participation of major allergy societies and patient's organizations (Paris, December 3, 2018). The meeting was carried out with the support of many organizations (Figure 1).

The final document was approved by the members of the Paris study group and the ARIA working group.

A Pocket Guide has been developed and includes the major recommendations of this document in a simple format. It is to be used digitally and in paper form to guide clinical practice for all stakeholders including patients, all health care providers and policy makers.

## 2. Gaps in AIT knowledge

AIT is an effective treatment but there are many gaps including those identified by AIRWAYS ICPs (19, 20) (Table 1) (21). Some of these gaps are the basis for the development of ARIA ICPs for AIT.

## 3. Allergens to be used

### a. Relevant extract

The decision to prescribe AIT for the patient should be based on allergen relevance and on the persistence of clinical symptoms, despite appropriate medications according to guidelines, as well as on the availability of good-quality extracts.

Adequate quality is essential for any medicinal product to be eligible for marketing (10, 22). Only regulated, standardized allergen extracts that are efficacious and safe should be used for AIT (23, 24).

### b. Extrapolation to untested products

AIT products have to show efficacy and safety in line with regulatory requirements (25). Allergen extracts cannot be regarded as generics. In the EU, each individual product (individual product or mixtures), with the exceptions made by EMA (European Medicines Agency) or PEI (Paul Ehrlich Institute), must prove its efficiency (23).

In some cases, exceptions related to the concept of homologous groups defining allergens with a significant clinical important cross-reactivity can be accepted without specific clinical documentation. These homologous groups include a range of pollen allergen extracts and house dust mites which are defined in the respective EMA guides for allergen products (23).

There is no evidence that mixing different allergens will have the same effect as separately administering individual allergens. Mixing different allergens can result in a dilutional effect - under dosing of the treatment and potential specific allergen degradation - due to enzymatic activity of certain allergens (26). The risk of allergic side effects can increase, especially by the degradation, when a new batch is used (27). Therefore, the EMA has recommended only to use mixed allergen products of allergens represented by the allergen sources from homologous groups (23).

### **c. Named Patient Products**

In many countries, Named Patient Products (NPP) are used by practitioners in an effort to apply precision treatment to patients. However, this practice requires appropriate confirmatory trials and real-world evidence since clinical data with some allergens cannot be directly extrapolated to NPP practice. NPPs are marketed on exception from the European legislation on allergen extracts (14, 28).

### **d. Polysensitized patients**

Allergic diseases are among the most complex and diverse diseases. Patients are often sensitized (IgE) to many allergens (polysensitization), but not all of these sensitizations may be clinically relevant. Therefore, it is important to treat the allergies that give rise to allergic symptoms and not the sensitizations potentially irrelevant for the patient. There is a broad range of evidence for the clinical efficacy of single extracts in polysensitized patients (29-31).

Instead of mixing extracts, the different allergens can be applied separately (12). However, it has been proposed without data that two extracts can be given with a 30-minute interval in two different injection spots. By doing so, each allergen can be monitored separately for the local reaction and potential systemic side effects.

In general, the question regarding the efficacy of poly- compared to oligo- or mono-allergen immunotherapy in polysensitized patients has not been addressed in carefully designed clinical trials. A recent report from an NIH-sponsored international workshop on aeroallergen immunotherapy outlines trial concepts to address this important knowledge gap (32).

## **4. Stratification of allergic patients for AIT**

### **a. Concept of patient stratification**

Precision medicine aims to customize healthcare with medical decisions, practices, and/or products tailored to the individual patient. It also refers to the tailoring of medical treatment to the clinical and social characteristics of each patient and not necessarily to genomics (33). The stratification of patients into subpopulations is the basis of clinical decision-making for increased diagnostic and treatment efficacy (34, 35). Patient stratification also integrates cost trends and social determinant risk models to match the patient to the right care management. This model applies to AIT (36).

In non-life threatening diseases with a very high prevalence, such as allergic diseases, patient stratification is required (i) to identify the best candidates for intervention through complex care management, (ii) to reduce the amount of time and resources needed to match the right patient to a care management programme and (iii) to optimize costs as some therapeutic interventions cannot be administered to all patients. Patient stratification may also help to improve the patient's engagement (37).

Molecular diagnosis, when used with other tools and patients' clinical records, can help clinicians better to select the most appropriate patients and allergens for AIT (38) and, in some cases, predict the risk of adverse reactions (39). The pattern of sensitization to allergens could potentially predict the efficacy of allergen immunotherapy provided that these immunotherapy products contain a sufficient amount of these allergens. Nevertheless, multiplex assay remains a third-level approach, not to be used as screening method in current practice (39).

VAS may be useful also for monitoring AIT effectiveness and medication use as it is easy-to-use and also validated for AR control of severity. It has been used as secondary endpoint in both adult and pediatric trials (40, 41)

The role of precision medicine in selecting an AIT regimen was proposed by an expert meeting (36) (Table 2).

The Flow of Precision Medicine approach in allergic disease has been adapted (Figure 2) from (16) and (36). In some instances, AIT can be offered to patients whose AR is controlled by pharmacotherapy such as those who may develop thunderstorm-induced asthma (42, 43). AIT should also be considered even in moderate AR, particularly (but not necessarily only) in patients who have had asthma exacerbations during the pollen season and who live in geographically at-risk regions.

#### **b. Biomarkers in AIT**

Biomarkers - clinical or laboratory characteristics that reflect biological processes - are essential for monitoring the health of patients. They include clinical signs identified by physical examination, biological assays, mHealth outcomes, genomic indices, and others that can be objectively measured and used as indicators of pathophysiological processes (44). They can be used individually or in combination but they need further studies.

There are currently no validated genetic or blood biomarkers for predicting or monitoring the efficacy of AIT at an individual patient level although several candidates have been investigated (45). Biomarkers associated with mHealth and a clinical decision support system (CDSS) (46) may change the scope of AIT as they will help monitor the patient's disease control (47, 48) for (i) patient

stratification, (ii) clinical trials and real-world evidence, (iii) monitoring efficacy and safety of targeted therapies (a critical process for identifying appropriate reimbursement) and (iv) implementation of stopping rules (Figure 3). Clinical stopping rules should be developed for AIT, similarly to what is currently considered for biologics in severe asthma, as a guidance for continuing or stopping treatment after a short (early stopping rule) or long (late stopping rule) period. As an example, a global treatment evaluation after 16 weeks is used as an early stopping rule for omalizumab treatment (49, 50).

### **c. ARIA**

In ARIA 2008 (16), it was indicated that DB-PC-RCTs have confirmed the efficacy of SCIT and SLIT. However, trial-based clinical efficacy is one of the many factors in a clinician's decision-making process for the use of AIT, especially since AIT RCTs are designed to fulfill regulatory demands for marketing authorization (51). Before starting AIT, it is essential to appreciate the relative value of pharmacotherapy and AIT and the degree of disease control achieved using pharmacotherapy. Furthermore, it is important to consider the rest of the patient's medical history as well as his/her social and geographical environment. The indications for SCIT in ARIA 2008 were similar to those published in 1998 (52) and 2001 (53).

## **5. Economic burden of allergic rhinitis and asthma, and cost-effectiveness of AIT**

Allergic diseases place a huge burden on society in terms of high prevalence, morbidity, direct costs (health service and drugs prescribed) and indirect costs, in particular those related to presenteeism (54). In addition, allergies have an impact on learning and performance at all levels of education (55, 56). Better care for allergies based on guideline-based treatment would allow substantial savings for Europe's economy (55).

Patients with allergic diseases may not understand the benefits of treatment, and adherence to treatment is poor (47). A substantial proportion of AR patients can be managed by appropriate pharmacological treatment (1). However, a subset of patients (10 to 20%) is poorly controlled and is ascribed to SCUAD (severe chronic upper airway disease) (57, 58, 59). Patients with asthma tend to incur higher rhinitis costs.

The cost-effectiveness of AIT should be considered for ICPs. However, it varies widely between countries and, in some countries such as Japan, the costs of AIT and pharmacotherapy are similar, whereas in the EU, acquisition costs of AIT are higher than pharmacotherapy. A health technology

assessment examined the comparative costs of SLIT and SCIT using the UK cost model (60). A benefit from both SCIT and SLIT compared with placebo was consistently demonstrated, but the extent of this effectiveness in terms of clinical benefit was considered unclear. The study concluded that both SCIT and SLIT may be cost-effective from around six years compared with standard treatment using the National Institute for Health and Care Excellence (NICE) cost-effectiveness threshold of £20,000-30,000 per quality-adjusted life years (QALY) (60, 61). A systematic overview showed that the cost-effectiveness of AIT is limited and of low methodological quality, but suggests that AIT may be cost-effective for people with AR with or without asthma (62). This systematic overview suggested that SLIT and SCIT would be considered cost-effective using the NICE cost-effectiveness threshold of £20 000 per QALY (63). Many of these studies were based on assumptions of the preventive effect of AIT using prediction models such as the Markov's model. These costs should be compared to biologics in the treatment of severe asthma. Although many limitations were identified, NICE proposed that Omalizumab (64), Mepolizumab (65) or Reslizumab (66) are likely to be cost-effective in severe asthma at the threshold of £30,000 per QALY.

However, the cost model of NICE may be questioned as it was developed for diseases impairing mobility or for severe diseases and does not take indirect costs (e.g. presenteeism) into account. Furthermore, it neglects the potential savings outside the UK health care system which may not be generalizable.

## **6. Safety**

### **a. Subcutaneous immunotherapy (SCIT)**

A typical reaction (local reaction) is redness and swelling at the injection site immediately or several hours after the injection. Sometimes, sneezing, nasal congestion or hives can occur (systemic reactions) (67). Serious reactions to injections are very rare but require immediate medical attention. Symptoms of an anaphylactic reaction can include swelling in the throat, wheezing or tightness in the chest, nausea and dizziness. Most serious reactions develop within 30 minutes after the injections and it is therefore recommended patients wait in their doctor's office for at least 30 minutes after an injection.

### **b. Sublingual immunotherapy (SLIT)**

Allergen drops or tablets have a more favorable safety profile than injections. SLIT can be administered at home after the first dose is > administered under the supervision of a physician. The large majority of adverse events are local (mouth itching, lip swelling, nausea) and spontaneously subside after the first days of administration. The severity of local side effects is graded according to

Accepted Article

persistence and impact on the quality of life (68). In some countries excluding Europe, SLIT tablets include a warning about possible severe allergic reactions and adrenaline-auto injectors are > routinely recommended. This is not the case in Europe although in the rare event that a general allergic reaction occurs after SLIT then the risk/benefit should be reassessed and a decision made whether to continue SLIT and if appropriate whether a rescue auto-injector should be provided

## 7. Patient's views

The patient's perspective should always be considered to enable a customized approach in shared decision making. There are contrasting real-life studies assessing the level of knowledge, perceptions, expectations and satisfaction about AIT. In two European studies, there was a relatively high degree of patient's perception and satisfaction that corresponded well with the physician's views (69, 70). However, most studies report a lack of information from allergic patients and every effort should be made to improve communication leading to increased patient knowledge and increased patient satisfaction (71, 72). Many AR patients have never heard of AIT (72).

Adherence to allergen immunotherapy (AIT) is crucial for its efficacy. SCIT requires regular (often monthly) visits, while SLIT is performed with a daily intake of allergen tablets or drops at home. Non-adherence to an AIT schedule and premature discontinuation are common problems (73). Results from various studies have concluded in controversial results on the rate of AIT adherence. Evidence-based communication, strategy-patient-centred care, motivational interviewing, and shared-decision making underscore the importance of taking time to establish trust, understand patient concerns and priorities, and involve the patient in decisions regarding AIT (74). A well-organized allergologist's time schedule not only increases safety but also offers the possibility of close follow-up and an increase in patient loyalty (73)

Information from a medical, economical and legal perspective illustrates the importance of the effort for evidence. From the medicolegal standpoint, the application of current medical knowledge, in combination with care for the patient's welfare, should drive daily medical practice. Medical criteria need to be prioritized over economic aspects, as physicians need to choose treatments according to the commonly acknowledged professional standards. Furthermore, the physician has the obligation to inform the patient about treatment options according to professional standards, detailing application ways, benefits and risks of available treatments/drugs, and to involve him/her in the decision (75).

## 8. Pharmacist's views

Self-medication to treat AR symptoms is common, and most patients self-manage their AR with few interactions with their physician (76). Community pharmacists are the most accessible health professionals for the public and AR is one of the most common diseases managed by pharmacists. They play an essential role in the management of pharmacotherapy, counselling, disease prevention and primary care. In particular, with the availability of non-prescribed medications (OTC) in the pharmacy, the community pharmacy is often the first stop for AR management (77, 78).

AR treatment encompasses three different aspects: avoidance of allergen exposure, pharmacotherapy and immunotherapy. The pharmacist intervention can specifically tackle the first two and might be an opportunity for patient education in terms of avoidance of allergen exposure, disease information, and medication use, especially medication administration and adherence. However, products for allergen immunotherapy are available in the pharmacies of many countries and the pharmacist must be well-informed about this treatment. Moreover, the pharmacist might play an important role in educating patients about the commitments involved in immunotherapy and its risks. For example, if patients miss several doses of immunotherapy, they may have to restart it. It is therefore important for patients to know what is expected up front and the pharmacist can play a significant role in providing this information.

## 9. General Practitioner's views

In many countries, the diagnosis and management of allergic disorders take place almost exclusively in primary care that has an essential role in the diagnosis and management of allergic diseases (79, 80). The continuous, easy to access, and holistic role of primary care can support the identification of allergic patients, reassure early diagnosis, and regularly follow up allergic patients for assessment of disease control, treatment adjustments and shared-decision making that is patient-centred. However, few general practitioners (GPs) receive formal undergraduate or postgraduate training in allergy (81, 82). Although considered important (80, 83, 84), there are minimal requirements for training and certification of subspecialists in allergy (85). Therefore, it is important for GPs to have access to training and evidence-based primary care allergy guidelines (86). Although some attempts of ICPs have been made (87), close collaboration with specialists for proper and time-efficient referral of cases will be beneficial for the patient and the health care system. Clear referral criteria and pathway plans should be created, implemented and validated by national circumstances and by cost-efficiency evaluation (88). Furthermore, GPs play a major role in patient education, self-medication and shared decision making (34, 88, 89), borrowing good practices from the management of other chronic diseases. Greater patient adherence to AIT is reported if AIT is provided by a GP rather than a



specialist (90). SCIT could also be done in primary care and although it is associated with some risks, these can be minimized when given by trained GPs that carefully select patients in an appropriate environment with available primary care facilities for treating systemic anaphylactic reactions (91-94).

## **10. Practical approach for patient stratification in AIT**

Shared decision making is required for AIT. Patients should be informed about all possible treatment options, benefits and drawbacks of AIT including its duration. Moreover, patients should know whether AIT is covered by their health system or insurance company and whether it will generate partial out-of-pocket costs or will need to be fully covered out-of-pocket.

Although biologics in severe asthma and AIT in allergic diseases target two different populations, costs per QALY, at least in some European countries, appear to be similar between AIT and biologics. This indicates that AIT should be reserved for stratified rhinitis patients insufficiently responsive to pharmacologic treatment (e.g. SCUAD (57)) who have been evaluated and guided with respect to adherence to pharmacotherapy. For asthma, a similar recommendation applies, but AIT should not be considered for severe asthma patients who are candidates for biologics. This recommendation is in line with the indications for a house dust mite tablet recently approved by the European Medicines Agency (95).

### **a. Rhinitis and rhinoconjunctivitis in adolescents and adults**

The selection of pharmacotherapy for AR patients depends on several factors, including age, predominant symptoms, severity, AR control, patient preferences and cost. Allergen exposure and resulting symptoms vary, e.g., based upon seasonal exposure or change in environment, making it necessary to make adjustments to therapy. CDSS may be beneficial by assessing disease control (96). They should be based on the best evidence algorithms to aid patients and health care professionals to jointly decide on the treatment and its step-up or step-down strategy depending on AR control (shared decision making).

The treatment of AR also requires consideration of (i) the phenotype (rhinitis, conjunctivitis and/or asthma) and severity of symptoms, (ii) the relative efficacy of the treatment, (iii) speed of onset of action of treatment, (iv) current treatment, (v) historic response to treatment, (vi) patient's preference, (vi) interest to self-manage and (viii) resource use. Guidelines and various statements by experts for AR pharmacotherapy usually propose the approach summarized in Table 3 (8, 97, 98).

All recommended medications are considered to be safe at the usual dosage except first generation oral H<sub>1</sub>-antihistamines which should be avoided (99). Notably, despite guidelines, the practice of prescribing both an INCS and an oral H<sub>1</sub>-antihistamine is globally common.

For step-up and step-down management, a simple algorithm was devised by MACVIA, but its applicability varies depending on the availability of medications and resources in different countries (Figure 4) (100).

Algorithms inherently result from combining individual decision nodes that represent separate recommendations. To be fully validated, the algorithm needs to be tested as a complete management plan and compared to alternative plans to explore if the combination of these separate recommendations leads to more benefit than harm when applied in practice. A large scale mobile technology study (47), a speed of onset study (101) and new recommendations supported the algorithm (97, 98).

#### **b. Asthma in adolescents and adults**

An algorithm is not yet available for asthma. Uncontrolled asthma is a contra-indication for AIT (102).

GINA (Global Initiative for Asthma) has endorsed SLIT for house dust mite-asthma (103). From the SmPC for the approved SLIT house dust mite tablet (95): (i) the patient should not have had a severe asthma exacerbation within the last 3 months of AIT initiation; (ii) in patients with asthma and experiencing an acute respiratory tract infection, initiation of treatment should be postponed until the infection has resolved; (iii) AIT is not indicated for the treatment of acute exacerbations and patients must be informed of the need to seek medical attention immediately if their asthma deteriorates suddenly; (iv) mite AIT should initially be used as an add-on therapy to controller treatment and reduction in asthma controllers should be performed gradually under the supervision of a physician according to management guidelines.

No other AIT product has been approved for asthma in the EU.

#### **c. Multimorbidity**

Multimorbidity, the co-existence of more than one allergic disease in the same patient, is very common in allergic diseases and over 85% of patients with asthma also have AR. On the other hand, only 20-30% of patients with AR have asthma. AR multimorbidity increases the severity of asthma (104).

An advantage of AIT is that it can control many aspects of multimorbidity including AR, asthma and conjunctivitis. Although multimorbid patients appear to have more severe symptoms related to each component of their allergic disease constellation, it is not yet known whether AIT is equally or more effective in these patients, compared to patients with no multimorbidity. This can be tested using existing databases, but a controlled trial will also offer useful evidence. In the conditions and authorization of a SLIT mite tablet (95), multimorbidity was recognized as an indication for mite SLIT.

#### **d. Children**

In children, AIT is effective as shown by RCTs (105) and may have long-term effects after it is stopped (106). A recent study of SLIT (107), a previous study of subcutaneous grass pollen immunotherapy in children (108) and a meta-analysis (109) have provided some evidence that AIT can delay or prevent the onset of asthma in children. However, (i) the meta-analysis showed a limited reduced short-term risk of developing asthma in those with AR with unclear benefit over the longer term (109), and (ii) costs cannot be supported by health care systems due to the very large number of patients who might be treated with uncertainty on cost-effectiveness.

Thus, AIT can be initiated in children with moderate/severe AR that is not controlled by pharmacotherapy. In such children without asthma, the possibility of preventing the onset of asthma should be taken into consideration, although more studies are needed for an unreserved indication (9).

The lower age for initiating AIT has not been clearly established. In many countries, products are licensed for children without a lower age limit. Prospective observational trials and/or registries can help confirm AIT safety and performance in the youngest recipients, perhaps down to the age of 3 years.

AIT is a paradigm for precision medicine, as it takes into account the multitude of sensitization and multimorbidity profile of each patient, both cross-sectionally and in relation to their natural history. Indirect yet important evidence provides clues about young patients who may benefit the most: (i) the severity of respiratory allergic disease is associated with its persistence (110); (ii) epitope spreading and development of new sensitizations suggest benefit with early intervention (111); (iii) the effects of AR on school performance and education (56) support focusing of treatment on developmental/career milestones. Therefore, the consideration of AIT at early time points, using risk in addition to severity as a key selection criterion, is expected to maximize impact on the natural history of the disease as well as on cost/burden.

Accepted Article

More studies are needed to characterize the long-term effects of AIT. Such studies cannot be randomized and even less blinded. Therefore, observational approaches, such as registry research, need to be used (112).

In addition, there are opportunities for disease prevention that have not been adequately explored, such as primary prevention. We need more evidence on whether AIT may play a role for the prevention of allergic sensitization, the first allergic disease (9). Support for such studies needs to come from governmental organizations/public sources, in order to identify optimal cost-efficacy strategies.

#### **e. Allergen immunotherapy in older age adults**

The immunologic and allergic characteristics of older allergic patients differ from those of young and middle-aged adults. Limited studies have found that AIT may be effective in this population (113, 114). More data are certainly required for a strong recommendation. At this point, and before making the decision to initiate AIT in older patients, physicians need to have strong indications for the role of specific allergens in these patients' AR or asthma and to take into account non-allergic comorbidities that may impact on the safety of AIT.

### **11. mHealth in the AIT precision medicine approach**

#### **a. Patient stratification**

It is recommended to stratify AR patients who are uncontrolled despite appropriate treatment and adherence to treatment (115). This can easily be achieved using electronic diaries obtained by cell phones as demonstrated in MASK-air<sup>®</sup> (2, 3, 47). Such diaries should include the full list of medications. After a single year of survey, physicians can assess whether SCUAD is present and could initiate AIT if (i) symptoms are associated with pollen season, (ii) adherence to pharmacologic treatment is achieved, (iii) the duration of uncontrolled symptoms was long enough and (iv) an impact on work or school productivity was observed. Moreover, asthma and eye symptoms can be recorded, as in MASK-air<sup>®</sup> (2) and other Apps allowing to evaluate the role of multimorbidity.

#### **b. Follow-up of patients under AIT**

The same approach can be proposed for the follow up of patients on AIT to assess its efficacy as suggested by a panel of international experts in an AIT Position Paper (18).

### c. Electronic clinical decision support system

The AR algorithm has been digitalized in tablets for health care professionals (46).

### Conclusions

AIT is an effective treatment for allergic diseases caused by inhaled allergens. Its use should, however, be restricted to carefully selected patients who are unresponsive to appropriate pharmacotherapy according to guidelines and for whom effective and cost-effective AIT is available. The present report reviews care pathways for the administration of AIT using evidence-based criteria. It is hoped that these recommendations are considered by healthcare professionals, so that the appropriate usage of AIT maximizes its impact on allergic diseases.

### References

1. Bousquet J, Hellings PW, Agache I, Amat F, Annesi-Maesano I, Ansotegui IJ, et al. ARIA Phase 4 (2018): Change management in allergic rhinitis and asthma multimorbidity using mobile technology. *J Allergy Clin Immunol*. 2018;pii: S0091-6749(18)31359-9. doi: 10.1016/j.jaci.2018.08.049.
2. Bousquet J, Arnavielhe S, Bedbrook A, Bewick M, Laune D, Mathieu-Dupas E, et al. MASK 2017: ARIA digitally-enabled, integrated, person-centred care for rhinitis and asthma multimorbidity using real-world-evidence. *Clin Transl Allergy*. 2018;8:45.
3. Bousquet J, Anto JM, Annesi-Maesano I, Dedeu T, Dupas E, Pepin JL, et al. POLLAR: Impact of air POLLution on Asthma and Rhinitis; a European Institute of Innovation and Technology Health (EIT Health) project. *Clin Transl Allergy*. 2018;8:36.
4. Bousquet J, Hellings P, Agache I, Amat F, Annesi-Maesano I, Ansotegui I, et al. ARIA Phase 4 (2018): Change management in allergic rhinitis and asthma multimorbidity using mobile technology. *J Allergy Clin Immunol* 2018;in press.
5. Campbell H, Hotchkiss R, Bradshaw N, Porteous M. Integrated care pathways. *BMJ*. 1998;316(7125):133-7.

- Accepted Article
6. Hujala A, Taskinen H, Rissanen S. In: Richardson E, van Ginneken E, editors. How to support integration to promote care for people with multimorbidity in Europe? European Observatory Policy Briefs. Copenhagen (Denmark)2017.
  7. Palmer K, Marengoni A, Forjaz MJ, Jureviciene E, Laatikainen T, Mammarella F, et al. Multimorbidity care model: Recommendations from the consensus meeting of the Joint Action on Chronic Diseases and Promoting Healthy Ageing across the Life Cycle (JA-CHRODIS). *Health Policy*. 2018;122(1):4-11.
  8. Brozek JL, Bousquet J, Baena-Cagnani CE, Bonini S, Canonica GW, Casale TB, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines: 2010 revision. *J Allergy Clin Immunol*. 2010;126(3):466-76.
  9. Halken S, Larenas-Linnemann D, Roberts G, Calderon MA, Angier E, Pfaar O, et al. EAACI guidelines on allergen immunotherapy: Prevention of allergy. *Pediatr Allergy Immunol*. 2017;28(8):728-45.
  10. Bonertz A, Roberts G, Slater JE, Bridgewater J, Rabin RL, Hoefnagel M, et al. Allergen manufacturing and quality aspects for allergen immunotherapy in Europe and the United States: An analysis from the EAACI AIT Guidelines Project. *Allergy*. 2018;73(4):816-26.
  11. Muraro A, Roberts G, Halken S, Agache I, Angier E, Fernandez-Rivas M, et al. EAACI guidelines on allergen immunotherapy: Executive statement. *Allergy*. 2018;73(4):739-43.
  12. Roberts G, Pfaar O, Akdis CA, Ansotegui IJ, Durham SR, Gerth van Wijk R, et al. EAACI Guidelines on Allergen Immunotherapy: Allergic rhinoconjunctivitis. *Allergy*. 2018;73(4):765-98.
  13. Ryan D, Gerth van Wijk R, Angier E, Kristiansen M, Zaman H, Sheikh A, et al. Challenges in the implementation of the EAACI AIT guidelines: A situational analysis of current provision of allergen immunotherapy. *Allergy*. 2018;73(4):827-36.
  14. Pfaar O, Bachert C, Bufe A, Buhl R, Ebner C, Eng P, et al. Guideline on allergen-specific immunotherapy in IgE-mediated allergic diseases: S2k Guideline of the German Society for Allergology and Clinical Immunology (DGAKI), the Society for Pediatric Allergy and Environmental Medicine (GPA), the Medical Association of German Allergologists (AeDA), the Austrian Society for Allergy and Immunology (OGAI), the Swiss Society for Allergy and Immunology (SGAI), the German Society of Dermatology (DDG), the German Society of Oto-Rhino-Laryngology, Head and Neck Surgery (DGHNO-KHC), the German Society of Pediatrics and Adolescent Medicine (DGKJ), the Society for Pediatric Pneumology (GPP), the German Respiratory Society (DGP), the German Association of ENT Surgeons (BV-HNO), the Professional Federation of Paediatricians and Youth Doctors (BVKJ), the Federal Association of Pulmonologists (BDP) and the German Dermatologists Association (BVDD). *Allergo J Int*. 2014;23(8):282-319.

15. Zielen S, Devillier P, Heinrich J, Richter H, Wahn U. Sublingual immunotherapy provides long-term relief in allergic rhinitis and reduces the risk of asthma: A retrospective, real-world database analysis. *Allergy*. 2018;73(1):165-77.
16. Bousquet J, Khaltaev N, Cruz AA, Denburg J, Fokkens WJ, Togias A, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy*. 2008;63 Suppl 86:8-160.
17. Larenas-Linnemann DES, Antolin-Amerigo D, Parisi C, Nakonechna A, Luna-Pech JA, Wedi B, et al. National clinical practice guidelines for allergen immunotherapy: An international assessment applying AGREE-II. *Allergy*. 2018;73(3):664-72.
18. Pfaar O, Demoly P, Gerth van Wijk R, Bonini S, Bousquet J, Canonica GW, et al. Recommendations for the standardization of clinical outcomes used in allergen immunotherapy trials for allergic rhinoconjunctivitis: an EAACI Position Paper. *Allergy*. 2014;69(7):854-67.
19. Bousquet J, Addis A, Adcock I, Agache I, Agustí A, Alonso A, et al. Integrated care pathways for airway diseases (AIRWAYS-ICPs). *Eur Respir J*. 2014;44(2):304-23.
20. Bousquet J, Barbara C, Bateman E, Bel E, Bewick M, Chavannes NH, et al. AIRWAYS-ICPs (European Innovation Partnership on Active and Healthy Ageing) from concept to implementation. *Eur Respir J*. 2016;47(4):1028-33.
21. Calderon MA, Demoly P, Casale T, Akdis CA, Bachert C, Bewick M, et al. Allergy immunotherapy across the life cycle to promote active and healthy ageing: from research to policies: An AIRWAYS Integrated Care Pathways (ICPs) programme item (Action Plan B3 of the European Innovation Partnership on active and healthy ageing) and the Global Alliance against Chronic Respiratory Diseases (GARD), a World Health Organization GARD research demonstration project. *Clin Transl Allergy*. 2016;6:41.
22. Bonertz A, Roberts GC, Hoefnagel M, Timon M, Slater JE, Rabin RL, et al. Challenges in the implementation of EAACI guidelines on allergen immunotherapy: A global perspective on the regulation of allergen products. *Allergy*. 2018;73(1):64-76.
23. Committee for medicinal products for human use (CPMP). Guideline on allergen products: production and quality issues. EMEA/CHMP/BWP/304831/2007. London, 20 November 2008. 2008.
24. Kowalski ML, Ansotegui I, Aberer W, Al-Ahmad M, Akdis M, Ballmer-Weber BK, et al. Risk and safety requirements for diagnostic and therapeutic procedures in allergology: World Allergy Organization Statement. *World Allergy Organ J*. 2016;9(1):33.
25. Bachert C, Larche M, Bonini S, Canonica GW, Kundig T, Larenas-Linnemann D, et al. Allergen immunotherapy on the way to product-based evaluation-a WAO statement. *World Allergy Organ J*. 2015;8(1):29.
26. Nelson HS, Ikle D, Buchmeier A. Studies of allergen extract stability: the effects of dilution and mixing. *J Allergy Clin Immunol*. 1996;98(2):382-8.

27. Reid MJ, Lockey RF, Turkeltaub PC, Platts-Mills TA. Survey of fatalities from skin testing and immunotherapy 1985-1989. *J Allergy Clin Immunol.* 1993;92(1 Pt 1):6-15.
28. Bousquet J, Lockey R, Malling HJ, Alvarez-Cuesta E, Canonica GW, Chapman MD, et al. Allergen immunotherapy: therapeutic vaccines for allergic diseases. World Health Organization. American academy of Allergy, Asthma and Immunology. *Ann Allergy Asthma Immunol.* 1998;81(5 Pt 1):401-5.
29. Didier A, Malling HJ, Worm M, Horak F, Jager S, Montagut A, et al. Optimal dose, efficacy, and safety of once-daily sublingual immunotherapy with a 5-grass pollen tablet for seasonal allergic rhinitis. *J Allergy Clin Immunol.* 2007;120(6):1338-45.
30. Nelson H, Blaiss M, Nolte H, Wurtz SO, Andersen JS, Durham SR. Efficacy and safety of the SQ-standardized grass allergy immunotherapy tablet in mono- and polysensitized subjects. *Allergy.* 2013;68(2):252-5.
31. Durham SR, Emminger W, Kapp A, de Monchy JG, Rak S, Scadding GK, et al. SQ-standardized sublingual grass immunotherapy: confirmation of disease modification 2 years after 3 years of treatment in a randomized trial. *J Allergy Clin Immunol.* 2012;129(3):717-25 e5.
32. Wheatley L, Wood R, Nadeau K, Liu A, Zoratti E, Bacharier L, et al. Mind the gaps: Clinical trial concepts to address unanswered questions in aeroallergen immunotherapy. An NIAID/AHRQ workshop. *J Allergy Clin Immunol.* 2019;in press.
33. Paving the way for personalized medicine, FDA's role in a new era of medical product development. <http://www.fda.gov/downloads/ScienceResearch/SpecialTopics/PersonalizedMedicine/UCM372421.pdf>. 2013.
34. Hellings PW, Fokkens WJ, Bachert C, Akdis CA, Bieber T, Agache I, et al. Positioning the principles of precision medicine in care pathways for allergic rhinitis and chronic rhinosinusitis - A EUFOREA-ARIA-EPOS-AIRWAYS ICP statement. *Allergy.* 2017;72(9):1297-305.
35. Muraro A, Fokkens WJ, Pietikainen S, Borrelli D, Agache I, Bousquet J, et al. European symposium on precision medicine in allergy and airways diseases: report of the European Union parliament symposium (October 14, 2015). *Rhinology.* 2015.
36. Canonica GW, Bachert C, Hellings P, Ryan D, Valovirta E, Wickman M, et al. Allergen Immunotherapy (AIT): a prototype of Precision Medicine. *World Allergy Organ J.* 2015;8(1):31.
37. Pfaar O, Bonini S, Cardona V, Demoly P, Jakob T, Jutel M, et al. Perspectives in allergen immunotherapy: 2017 and beyond. *Allergy.* 2018;73 Suppl 104:5-23.
38. Del-Rio Camacho G, Montes Arjona AM, Fernandez-Cantalejo Padias J, Rodriguez Catalan J. How molecular diagnosis may modify immunotherapy prescription in multi-sensitized pollen-allergic children. *Allergol Immunopathol (Madr).* 2018;46(6):552-6.
39. Sastre J, Sastre-Ibanez M. Molecular diagnosis and immunotherapy. *Curr Opin Allergy Clin Immunol.* 2016;16(6):565-70.



- Accepted Article
40. Frew AJ, Powell RJ, Corrigan CJ, Durham SR. Efficacy and safety of specific immunotherapy with SQ allergen extract in treatment-resistant seasonal allergic rhinoconjunctivitis. *J Allergy Clin Immunol*. 2006;117(2):319-25.
  41. Corrigan CJ, Kettner J, Doemer C, Cromwell O, Narkus A. Efficacy and safety of preseasonal-specific immunotherapy with an aluminium-adsorbed six-grass pollen allergoid. *Allergy*. 2005;60(6):801-7.
  42. Lee J, Kronborg C, O'Hehir RE, Hew M. Who's at risk of thunderstorm asthma? The ryegrass pollen trifecta and lessons learnt from the Melbourne thunderstorm epidemic. *Respir Med*. 2017;132:146-8.
  43. O'Hehir RE, Varese NP, Deckert K, Zubrinich CM, van Zelm MC, Rolland JM, et al. Epidemic Thunderstorm Asthma Protection with Five-Grass Pollen Tablet Sublingual Immunotherapy: A Clinical Trial. *Am J Respir Crit Care Med*. 2018;198(1):126-8.
  44. Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin Pharmacol Ther*. 2001;69(3):89-95.
  45. Shamji MH, Kappen JH, Akdis M, Jensen-Jarolim E, Knol EF, Kleine-Tebbe J, et al. Biomarkers for monitoring clinical efficacy of allergen immunotherapy for allergic rhinoconjunctivitis and allergic asthma: an EAACI Position Paper. *Allergy*. 2017;72(8):1156-73.
  46. Courbis AL, Murray RB, Arnavielhe S, Caimmi D, Bedbrook A, Van Eerd M, et al. Electronic Clinical Decision Support System for allergic rhinitis management: MASK e-CDSS. *Clin Exp Allergy*. 2018.
  47. Bousquet J, Arnavielhe S, Bedbrook A, Alexis-Alexandre G, Eerd Mv, Murray R, et al. Treatment of allergic rhinitis using mobile technology with real world data: The MASK observational pilot study. *Allergy*. 2018:sous presse.
  48. Bousquet J, Hellings PW, Agache I, Bedbrook A, Bachert C, Bergmann KC, et al. ARIA 2016: Care pathways implementing emerging technologies for predictive medicine in rhinitis and asthma across the life cycle. *Clin Transl Allergy*. 2016;6:47.
  49. Bousquet J, Rabe K, Humbert M, Chung KF, Berger W, Fox H, et al. Predicting and evaluating response to omalizumab in patients with severe allergic asthma. *Respir Med*. 2007.
  50. Diaz RA, Charles Z, George E, Adler AI. NICE guidance on omalizumab for severe asthma. *Lancet Respir Med*. 2013;1(3):189-90.
  51. Committee for medicinal products for human use. Guideline on the clinical development of products for specific immunotherapy for the treatment of allergic diseases. CHMP/EWP/18504/2006. London. European Medicines Agency. Pre-authorisation evaluation of medicines for human use. 20 November. 2008.
  52. Bousquet J, Lockey R, Malling H. WHO Position Paper. Allergen Immunotherapy: Therapeutic Vaccines for allergic diseases. *Allergy*. 1998;53, suppl 54.

53. Bousquet J, Demoly P, Michel FB. Specific immunotherapy in rhinitis and asthma. *Ann Allergy Asthma Immunol.* 2001;87(1 Suppl 1):38-42.
54. Vandenas O, Vinnikov D, Blanc PD, Agache I, Bachert C, Bewick M, et al. Impact of Rhinitis on Work Productivity: A Systematic Review. *J Allergy Clin Immunol Pract.* 2018;6(4):1274-86 e9.
55. Zuberbier T, Lotvall J, Simoens S, Subramanian SV, Church MK. Economic burden of inadequate management of allergic diseases in the European Union: a GA(2) LEN review. *Allergy.* 2014;69(10):1275-9.
56. Walker S, Khan-Wasti S, Fletcher M, Cullinan P, Harris J, Sheikh A. Seasonal allergic rhinitis is associated with a detrimental effect on examination performance in United Kingdom teenagers: case-control study. *J Allergy Clin Immunol.* 2007;120(2):381-7.
57. Bousquet J, Bachert C, Canonica GW, Casale TB, Cruz AA, Lockey RJ, et al. Unmet needs in severe chronic upper airway disease (SCUAD). *J Allergy Clin Immunol.* 2009;124(3):428-33.
58. Hellings PW, Fokkens WJ, Akdis C, Bachert C, Cingi C, Dietz de Loos D, et al. Uncontrolled allergic rhinitis and chronic rhinosinusitis: where do we stand today? *Allergy.* 2013;68(1):1-7.
59. Bousquet PJ, Demoly P, Devillier P, Mesbah K, Bousquet J. Impact of Allergic Rhinitis Symptoms on Quality of Life in Primary Care. *Int Arch Allergy Immunol.* 2013;160(4):393-400.
60. Meadows A, Kaambwa B, Novielli N, Huissoon A, Fry-Smith A, Meads C, et al. A systematic review and economic evaluation of subcutaneous and sublingual allergen immunotherapy in adults and children with seasonal allergic rhinitis. *Health Technol Assess.* 2013;17(27):vi, xi-xiv, 1-322.
61. Devlin N, Parkin D. Does NICE have a cost-effectiveness threshold and what other factors influence its decisions? A binary choice analysis. *Health Econ.* 2004;13(5):437-52.
62. Asaria M, Dhami S, van Ree R, Gerth van Wijk R, Muraro A, Roberts G, et al. Health economic analysis of allergen immunotherapy for the management of allergic rhinitis, asthma, food allergy and venom allergy: A systematic overview. *Allergy.* 2018;73(2):269-83.
63. Bruggenjurgan B, Reinhold T. Cost-effectiveness of grass pollen subcutaneous immunotherapy (SCIT) compared to sublingual immunotherapy (SLIT) and symptomatic treatment in Austria, Spain, and Switzerland. *J Med Econ.* 2018;21(4):374-81.
64. Faria R, McKenna C, Palmer S. Optimizing the position and use of omalizumab for severe persistent allergic asthma using cost-effectiveness analysis. *Value Health.* 2014;17(8):772-82.
65. Bermejo I, Stevenson M, Cooper K, Harnan S, Hamilton J, Clowes M, et al. Mepolizumab for Treating Severe Eosinophilic Asthma: An Evidence Review Group Perspective of a NICE Single Technology Appraisal. *Pharmacoeconomics.* 2018;36(2):131-44.
66. Cooper K, Frampton G, Harris P, Rose M, Chorozoglou M, Pickett K. Reslizumab for Treating Asthma with Elevated Blood Eosinophils Inadequately Controlled by Inhaled Corticosteroids:

An Evidence Review Group Perspective of a NICE Single Technology Appraisal.

Pharmacoeconomics. 2018;36(5):545-53.

67. Cox L, Larenas-Linnemann D, Lockey RF, Passalacqua G. Speaking the same language: The World Allergy Organization Subcutaneous Immunotherapy Systemic Reaction Grading System. *J Allergy Clin Immunol*. 2010;125(3):569-74, 74 e1-74 e7.
68. Passalacqua G, Baena-Cagnani CE, Bousquet J, Canonica GW, Casale TB, Cox L, et al. Grading local side effects of sublingual immunotherapy for respiratory allergy: speaking the same language. *J Allergy Clin Immunol*. 2013;132(1):93-8.
69. Baiardini I, Puggioni F, Menoni S, Boot JD, Diamant Z, Braido F, et al. Patient knowledge, perceptions, expectations and satisfaction on allergen-specific immunotherapy: a survey. *Respir Med*. 2013;107(3):361-7.
70. Nam YH, Lee SK. Physician's recommendation and explanation is important in the initiation and maintenance of allergen immunotherapy. *Patient Prefer Adherence*. 2017;11:381-7.
71. Chivato T, Alvarez-Calderon P, Panizo C, Abengozar R, Alias C, Al-Baech A, et al. Clinical management, expectations, and satisfaction of patients with moderate to severe allergic rhinoconjunctivitis treated with SQ-standardized grass-allergen tablet under routine clinical practice conditions in Spain. *Clin Mol Allergy*. 2017;15:1.
72. Skoner DP, Blaiss MS, Dykewicz MS, Smith N, Leatherman B, Bielory L, et al. The Allergies, Immunotherapy, and Rhinoconjunctivitis (AIRS) survey: patients' experience with allergen immunotherapy. *Allergy Asthma Proc*. 2014;35(3):219-26.
73. Pitsios C, Dietis N. Ways to increase adherence to allergen immunotherapy. *Curr Med Res Opin*. 2018:1-9.
74. Bender BG, Lockey RF. Solving the Problem of Nonadherence to Immunotherapy. *Immunol Allergy Clin North Am*. 2016;36(1):205-13.
75. Bachert C, Gräfin-von-Strachwitz-Helmstatt K. Zur Diskussion gestellt: Der Arzt und die Spezifische Immuntherapie im Spannungsfeld von Leitlinie, Wirtschaftlichkeit und Medizinrecht. *Allergologie*. 2016;39:381-8.
76. Kuehl BL, Abdunour S, O'Dell M, Kyle TK. Understanding the role of the healthcare professional in patient self-management of allergic rhinitis. *SAGE Open Med*. 2015;3:2050312115595822.
77. Bosnic-Anticevich S, Kritikos V, Carter V, Yan KY, Armour C, Ryan D, et al. Lack of asthma and rhinitis control in general practitioner-managed patients prescribed fixed-dose combination therapy in Australia. *J Asthma*. 2018;55(6):684-94.
78. Bosnic-Anticevich S, Costa E, Menditto E, Lourenco O, Novellino E, Bialek S, et al. ARIA pharmacy 2018 "Allergic rhinitis care pathways for community pharmacy". *Allergy*. 2018.
79. Jutel M, Papadopoulos NG, Gronlund H, Hoffman HJ, Bohle B, Hellings P, et al. Recommendations for the allergy management in the primary care. *Allergy*. 2014;69(6):708-18.

80. Finlay I, Egner W. Allergy--will we ever meet the unmet need? *J R Soc Med.* 2010;103(11):430-1.
81. Shehata Y, Ross M, Sheikh A. Undergraduate allergy teaching in a UK medical school: comparison of the described and delivered curriculum. *Prim Care Respir J.* 2007;16(1):16-21.
82. Ewan PW, Durham SR. NHS allergy services in the UK: proposals to improve allergy care. *Clin Med (Lond).* 2002;2(2):122-7.
83. Agache I, Ryan D, Rodriguez MR, Yusuf O, Angier E, Jutel M. Allergy management in primary care across European countries -- actual status. *Allergy.* 2013;68(7):836-43.
84. Ellis J, Rafi I, Smith H, Sheikh A. Identifying current training provision and future training needs in allergy available for UK general practice trainees: national cross-sectional survey of General Practitioner Specialist Training programme directors. *Prim Care Respir J.* 2013;22(1):19-22.
85. Gerth van Wijk R, Eguiluz-Gracia I, Gayraud J, Gutermuth J, Hamelmann E, Heffler E, et al. The roadmap for allergology in Europe: The subspecialty of allergology as "stop-over" on the way to a full specialty. An EAACI position statement. *Allergy.* 2018;73(3):540-8.
86. Levy ML, Fletcher M, Price DB, Hausen T, Halbert RJ, Yawn BP. International Primary Care Respiratory Group (IPCRG) Guidelines: diagnosis of respiratory diseases in primary care. *Prim Care Respir J.* 2006;15(1):20-34.
87. Flokstra-de Blok BMJ, Brakel TM, Wubs M, Skidmore B, Kocks JWH, Oude Elberink JNG, et al. The feasibility of an allergy management support system (AMSS) for IgE-mediated allergy in primary care. *Clin Transl Allergy.* 2018;8:18.
88. Jutel M, Angier L, Palkonen S, Ryan D, Sheikh A, Smith H, et al. Improving allergy management in the primary care network--a holistic approach. *Allergy.* 2013;68(11):1362-9.
89. Pinnock H, Thomas M, Tsiligianni I, Lisspers K, Ostrem A, Stallberg B, et al. The International Primary Care Respiratory Group (IPCRG) Research Needs Statement 2010. *Prim Care Respir J.* 2010;19 Suppl 1:S1-20.
90. Kiel MA, Roder E, Gerth van Wijk R, Al MJ, Hop WC, Rutten-van Molken MP. Real-life compliance and persistence among users of subcutaneous and sublingual allergen immunotherapy. *J Allergy Clin Immunol.* 2013;132(2):353-60 e2.
91. Alvarez-Cuesta E, Bousquet J, Canonica GW, Durham SR, Malling HJ, Valovirta E. Standards for practical allergen-specific immunotherapy. *Allergy.* 2006;61 Suppl 82:1-20.
92. Landi M, Meglio P, Praitano E, Lombardi C, Passalacqua G, Canonica GW. The perception of allergen-specific immunotherapy among pediatricians in the primary care setting. *Clin Mol Allergy.* 2015;13(1):15.
93. Zuberbier T, Bachert C, Bousquet PJ, Passalacqua G, Walter Canonica G, Merk H, et al. GA(2) LEN/EAACI pocket guide for allergen-specific immunotherapy for allergic rhinitis and asthma. *Allergy.* 2010;65(12):1525-30.

94. Stokes JR, Casale TB. Allergy immunotherapy for primary care physicians. *Am J Med.* 2006;119(10):820-3.
95. Summary of product characteristics. Acarizax 12 SQ-HDM oral lyophilisate. European Medicines Agency. [https://mriacts-mrpeu/Human/Downloads/DE\\_H\\_1947\\_001\\_FinalSPCpdf](https://mriacts-mrpeu/Human/Downloads/DE_H_1947_001_FinalSPCpdf). 2016.
96. Bousquet J, Schunemann HJ, Fonseca J, Samolinski B, Bachert C, Canonica GW, et al. MACVIA-ARIA Sentinel Network for allergic rhinitis (MASK-rhinitis): The new generation guideline implementation. *Allergy.* 2015.
97. Brozek JL, Bousquet J, Agache I, Agarwal A, Bachert C, Bosnic-Anticevich S, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) Guidelines - 2016 Revision. *J Allergy Clin Immunol.* 2017;140(4):950-8.
98. Dykewicz MS, Wallace DV, Baroody F, Bernstein J, Craig T, Finegold I, et al. Treatment of seasonal allergic rhinitis: An evidence-based focused 2017 guideline update. *Ann Allergy Asthma Immunol.* 2017;119(6):489-511 e41.
99. Church MK, Maurer M, Simons FE, Bindslev-Jensen C, van Cauwenberge P, Bousquet J, et al. Risk of first-generation H(1)-antihistamines: a GA(2)LEN position paper. *Allergy.* 2010;65(4):459-66.
100. Bousquet J, Schunemann HJ, Hellings PW, Arnavielhe S, Bachert C, Bedbrook A, et al. MACVIA clinical decision algorithm in adolescents and adults with allergic rhinitis. *J Allergy Clin Immunol.* 2016;138(2):367-74 e2.
101. Bousquet J, Meltzer EO, Couroux P, Koltun A, Kopietz F, Munzel U, et al. Onset of Action of the Fixed Combination Intranasal Azelastine-Fluticasone Propionate in an Allergen Exposure Chamber. *J Allergy Clin Immunol Pract.* 2018;6(5):1726-32.
102. Pitsios C, Demoly P, Bilo MB, Gerth van Wijk R, Pfaar O, Sturm GJ, et al. Clinical contraindications to allergen immunotherapy: an EAACI position paper. *Allergy.* 2015;70(8):897-909.
103. Global Strategy for Asthma Management and Prevention (2018 update). *wms-GINA-2018-report-V13-002pdf.* 2018.
104. Amaral R, Fonseca JA, Jacinto T, Pereira AM, Malinowski A, Janson C, et al. Having concomitant asthma phenotypes is common and independently relates to poor lung function in NHANES 2007-2012. *Clin Transl Allergy.* 2018;8:13.
105. Masuyama K, Okamoto Y, Okamiya K, Azuma R, Fujinami T, Riis B, et al. Efficacy and safety of SQ house dust mite sublingual immunotherapy-tablet in Japanese children. *Allergy.* 2018.
106. Penagos M, Eifan AO, Durham SR, Scadding GW. Duration of Allergen Immunotherapy for Long-Term Efficacy in Allergic Rhinoconjunctivitis. *Curr Treat Options Allergy.* 2018;5(3):275-90.

107. Valovirta E, Petersen TH, Piotrowska T, Laursen MK, Andersen JS, Sorensen HF, et al. Results from the 5-year SQ grass sublingual immunotherapy tablet asthma prevention (GAP) trial in children with grass pollen allergy. *J Allergy Clin Immunol.* 2018;141(2):529-38 e13.
108. Moller C, Dreborg S, Ferdousi HA, Halken S, Host A, Jacobsen L, et al. Pollen immunotherapy reduces the development of asthma in children with seasonal rhinoconjunctivitis (the PAT-study). *J Allergy Clin Immunol.* 2002;109(2):251-6.
109. Kristiansen M, Dhami S, Netuveli G, Halken S, Muraro A, Roberts G, et al. Allergen immunotherapy for the prevention of allergy: A systematic review and meta-analysis. *Pediatr Allergy Immunol.* 2017;28(1):18-29.
110. Tai A, Tran H, Roberts M, Clarke N, Gibson AM, Vidmar S, et al. Outcomes of childhood asthma to the age of 50 years. *J Allergy Clin Immunol.* 2014;133(6):1572-8 e3.
111. Lau S, Matricardi PM, Wahn U, Lee YA, Keil T. Allergy and atopy from infancy to adulthood: Messages from the German birth cohort MAS. *Ann Allergy Asthma Immunol.* 2018.
112. Price D, Bateman ED, Chisholm A, Papadopoulos NG, Bosnic-Anticevich S, Pizzichini E, et al. Complementing the randomized controlled trial evidence base. Evolution not revolution. *Ann Am Thorac Soc.* 2014;11 Suppl 2:S92-8.
113. Bozek A, Kolodziejczyk K, Kozłowska R, Canonica GW. Evidence of the efficacy and safety of house dust mite subcutaneous immunotherapy in elderly allergic rhinitis patients: a randomized, double-blind placebo-controlled trial. *Clin Transl Allergy.* 2017;7:43.
114. Bozek A, Kolodziejczyk K, Warkocka-Szoltyssek B, Jarzab J. Grass pollen sublingual immunotherapy: a double-blind, placebo-controlled study in elderly patients with seasonal allergic rhinitis. *Am J Rhinol Allergy.* 2014;28(5):423-7.
115. Bousquet J, Schunemann HJ, Fonseca J, Samolinski B, Bachert C, Canonica GW, et al. MACVIA-ARIA Sentinel Network for allergic rhinitis (MASK-rhinitis): the new generation guideline implementation. *Allergy.* 2015;70(11):1372-92.
116. Canonica GW, Ansotegui IJ, Pawankar R, Schmid-Grendelmeier P, van Hage M, Baena-Cagnani CE, et al. A WAO - ARIA - GA(2)LEN consensus document on molecular-based allergy diagnostics. *World Allergy Organ J.* 2013;6(1):17.
117. Hampel FC, Ratner PH, Van Bavel J, Amar NJ, Daftary P, Wheeler W, et al. Double-blind, placebo-controlled study of azelastine and fluticasone in a single nasal spray delivery device. *Ann Allergy Asthma Immunol.* 2010;105(2):168-73.
118. Carr W, Bernstein J, Lieberman P, Meltzer E, Bachert C, Price D, et al. A novel intranasal therapy of azelastine with fluticasone for the treatment of allergic rhinitis. *J Allergy Clin Immunol.* 2012;129(5):1282-9 e10.
119. Meltzer EO. Pharmacotherapeutic strategies for allergic rhinitis: matching treatment to symptoms, disease progression, and associated conditions. *Allergy Asthma Proc.* 2013;34(4):301-11.

120. Seidman MD, Gurgel RK, Lin SY, Schwartz SR, Baroody FM, Bonner JR, et al. Clinical practice guideline: allergic rhinitis executive summary. *Otolaryngol Head Neck Surg.* 2015;152(2):197-206.
121. Seidman MD, Gurgel RK, Lin SY, Schwartz SR, Baroody FM, Bonner JR, et al. Clinical practice guideline: Allergic rhinitis. *Otolaryngol Head Neck Surg.* 2015;152(1 Suppl):S1-43.
122. Bachert C, Bousquet J, Hellings P. Rapid onset of action and reduced nasal hyperreactivity: new targets in allergic rhinitis management. *Clin Transl Allergy.* 2018;8:25.

**Table 1: Gaps in AIT proposed by AIRWAYS ICs (modified from (21))**

- Better understand the role of AIT across the life cycle, particularly in preschool children (prevention and treatment) and in the elderly.
- Increase the awareness of the impact of AIT across the life cycle to promote active and healthy ageing.
- Stratify patients who benefit the most from AIT in all age groups.
- Launch a collaboration to develop care pathways for chronic respiratory allergic diseases integrating AIT in European countries and regions in the frame of AIRWAYS ICs.
- Follow and implement actions and plans suggested by this integrated collaboration.
- Provide evidence for regulatory decisions including cost-effectiveness.
- Follow and implement actions and plans suggested by this integrated collaboration, endorsed by national (or regional) health authorities.
- Encourage research strategies for novel approaches and biomarker discovery in AIT
- Encourage research strategies for 'responders/no-responders' in AIT

**Table 2: Precision Medicine in the indication of AIT (adapted from (16) and (36))**

1. Precise diagnosis with history, skin prick tests and/or specific IgE and, if needed, component-resolved *in vitro* diagnosis (CRD) (116). In some rare instances, provocation tests may be needed.
2. Proven indications: Allergic rhinitis, conjunctivitis and/or asthma.
3. Allergic symptoms predominantly induced by the relevant allergen exposure.
4. Patient stratification: Poor control of symptoms despite appropriate pharmacotherapy according to guidelines with adherence to treatment during the allergy season and/or the alteration of the natural history of allergy. Mobile technology may become of relevant importance to stratify patients (mHealth biomarker).
5. Demonstration of efficacy and safety for the product with relevant trials.
6. The patient (and caregiver)'s views represent an essential component.

**Table 3: Summary of recommendations for the treatment of allergic rhinitis and conjunctivitis used in the algorithm (adapted from (100))**

<p>Overall, GRADE-based AR guidelines agree on some important points (8, 97, 98, 100):</p> <ul style="list-style-type: none"><li>• Oral or intra-nasal H<sub>1</sub>-anti-histamines are less effective than intra-nasal corticosteroids (INCS) for the control of all rhinitis symptoms. H<sub>1</sub>-anti-histamines are however effective in many patients with mild disease and many patients prefer oral medications than intra-nasal ones.</li><li>• Consensus has not been reached as to the relative efficacy of oral versus intra-nasal H<sub>1</sub>-anti-histamines.</li><li>• In patients with severe rhinitis, INCS represent the first line treatment. However, they need a few days to be fully effective.</li><li>• The combination of oral H<sub>1</sub>-anti-histamines and INCS does not offer a better efficacy than INCS alone (97, 98).</li><li>• MPAzeFlu, the combined intranasal FP and Azelastine (Aze) in a single device, is more effective than monotherapy and indicated for those patients in whom monotherapy with intranasal glucocorticoid is insufficient (117-121) (117-121), patients with severe AR or those who want rapid symptom relief (97, 98, 122). An allergen chamber study has confirmed the speed of onset of the combination (101).</li></ul>
---

**Figure 1: Organizations supporting the meeting**

**Figure 2: Flow of precision medicine for AIT (adapted from (16) and (36))**

**Figure 3: Potential Biomarkers for AIT**

**Figure 4: Step-up algorithm in treated patients using visual analogue scale (adolescents and adults) (adapted from (100))**


*The proposed algorithm considers the treatment steps and patient's preference*

*VAS levels in ratio*

*In the case of remaining ocular symptoms add intra-ocular treatment*







\*: There are some exceptions

Fig 3




Fig 4

