



REPORT *RAPPORT*

MAS

Modelling, Analysis and Simulation



Modelling, Analysis and Simulation

Dependence of the transition from Townsend to glow discharge on secondary emission

Yu.P. Raizer, U.M. Ebert, D.D. Šijačić

REPORT MAS-E0412 AUGUST 2004

CWI is the National Research Institute for Mathematics and Computer Science. It is sponsored by the Netherlands Organization for Scientific Research (NWO).

CWI is a founding member of ERCIM, the European Research Consortium for Informatics and Mathematics.

CWI's research has a theme-oriented structure and is grouped into four clusters. Listed below are the names of the clusters and in parentheses their acronyms.

Probability, Networks and Algorithms (PNA)

Software Engineering (SEN)

Modelling, Analysis and Simulation (MAS)

Information Systems (INS)

Copyright © 2004, Stichting Centrum voor Wiskunde en Informatica

P.O. Box 94079, 1090 GB Amsterdam (NL)

Kruislaan 413, 1098 SJ Amsterdam (NL)

Telephone +31 20 592 9333

Telefax +31 20 592 4199

ISSN 1386-3703

Dependence of the transition from Townsend to glow discharge on secondary emission

ABSTRACT

In a recent paper Sijacic and Ebert have systematically studied the transition from Townsend to glow discharge, referring to older work from von Engel (1934) up to Raizer (1991), and they stated a strong dependence on secondary emission γ from the cathode. We here show that the earlier results of von Engel and Raizer on the small current expansion about the Townsend limit actually are the limit of small γ of the new expression; and that for larger γ the old and the new results vary by no more than a factor of 2. We discuss the γ -dependence of the transition which is rather strong for short gaps.

2000 Mathematics Subject Classification: 92F05, 34C99

Keywords and Phrases: current-voltage-characteristics, small current expansion, dependence on secondary emission γ , Townsend discharge, glow discharge

Dependence of the transition from Townsend to glow discharge on secondary emission

Yu.P. Raizer¹, Ute Ebert^{2,3} and D.D. Šijačić²

¹*Inst. for Problems in Mechanics of the Russian Academy of Sciences, Moscow, Russia,*

²*CWI, P.O.Box 94079, 1090 GB Amsterdam, The Netherlands, and*

³*Dept. Physics, Eindhoven Univ. Techn., The Netherlands.*

(Dated: February 24, 2004)

In a recent paper Šijačić and Ebert have systematically studied the transition from Townsend to glow discharge, referring to older work from von Engel (1934) up to Raizer (1991), and they stated a strong dependence on secondary emission γ from the cathode. We here show that the earlier results of von Engel and Raizer on the small current expansion about the Townsend limit actually are the limit of small γ of the new expression; and that for larger γ the old and the new results vary by no more than a factor of 2. We discuss the γ -dependence of the transition which is rather strong for short gaps.

In the recent article [1], the transition from Townsend to glow discharge was re-investigated with analytical and numerical means. On the analytical side, a systematic small current expansion about the Townsend limit was performed and it was stated:

“The result agrees qualitatively with the one given by Raizer [2] and Engel and Steenbeck [3]. In particular, the leading order correction is also of order $\alpha''(j/\mu)^2$. However, the explicit coefficient of j^2 differs: while the coefficient in [2, 3] does not depend on γ at all, we find that the dependence on γ is essential, as the plot of F in Fig. 1 (of [1]) clearly indicates. In fact, within the relevant range of $10^{-6} \leq \gamma \leq 10^0$, this coefficient varies by almost four orders of magnitude. We remark that it indeed would be quite a surprising mathematical result if the Townsend limit itself would depend on γ , but the small current expansion about it would not.”

Here we remark that while the new systematic calculation was correct, the interpretation and comparison to earlier work requires some correction.

To be precise, the model treated in [1–3] and by many other authors is a one-dimensional time independent Townsend or glow discharge characterized by the classical equations

$$\partial_x J_e = |J_e| \bar{\alpha}(|E|), \quad \partial_x J_+ = |J_+| \bar{\alpha}(|E|), \quad (1)$$

$$\partial_x E = \frac{e}{\epsilon_0} (n_+ - n_e), \quad (2)$$

for electron and ion particle current

$$J_e = -n_e \mu_e E, \quad J_+ = n_+ \mu_+ E, \quad (3)$$

and electric field E . Impact ionization in the bulk of the discharge is given by the Townsend approximation

$$\bar{\alpha}(|E|) = \alpha_0 e^{-E_0/|E|}. \quad (4)$$

(In [1], the generalized case $\bar{\alpha}(|E|) = \alpha_0 \exp(-E_0/|E|)^s$ was treated.) Since ions are generated by impact within the gap and drift towards the cathode, there are no ions at the anode at $x = 0$

$$n_+(0) = 0. \quad (5)$$

The ions impacting on the cathode liberate free electrons with rate γ , therefore secondary emission from the cathode is given by

$$|J_e(d)| = \gamma |J_+(d)|, \quad (6)$$

where $x = d$ is the position at the cathode. The electrical potential difference between the electrodes is

$$U = \int_0^d dx E(x), \quad (7)$$

and the total electric current is

$$J = e (n_+ \mu_+ + n_e \mu_e) E, \quad \partial_x J = 0. \quad (8)$$

It is useful to introduce dimensionless voltage and current

$$u = \frac{U}{E_0/\alpha_0}, \quad \bar{j} = \frac{J}{\epsilon_0 \alpha_0 E_0 \mu_+ E_0}, \quad (9)$$

where $\bar{j} = j/\mu$ with the definition of j from [1]. It should be noted that only bulk gas parameters have been used as units; therefore the dimensionless u and \bar{j} are independent of γ .

Further dimensional analysis yields that the current-voltage-characteristics $u = u(\bar{j})$ can depend on three parameters only, namely on the dimensionless gap length $L = \alpha_0 d$, on the coefficient γ of secondary emission and on the mobility ratio $\mu = \mu_+/\mu_e$. In practice, the dependence on the small parameter μ is almost negligibly weak [1], therefore $u = u(\bar{j}, L, \gamma)$. Here the dimensionless gap length L is related to pd through $L = A pd$ as long as the coefficient α_0 is related to pressure like $\alpha_0 = Ap$.

How strongly does the characteristics $u = u(\bar{j}, L, \gamma)$ depend on γ ? In [1], Šijačić and Ebert calculated the whole Townsend-to-glow regime numerically and derived by expanding systematically in powers of current \bar{j} about the Townsend limit

$$u = u_T - A_{SE} \bar{j}^2 + O(\bar{j}^3), \quad (10)$$

$$A_{SE} = \frac{\mathcal{E}_T \alpha''}{2 \alpha'} \frac{F(\gamma, \mu)}{(\alpha \mathcal{E}_T)^3}, \quad (11)$$

$$\alpha(\mathcal{E}_T) = e^{-1/|\mathcal{E}_T|}, \quad (12)$$

which gave an excellent fit to the numerical solutions. Here

$$F(\gamma, \mu) = \frac{L_\gamma^3}{12} + (1 + \mu) (2 - L_\gamma - 2e^{-L_\gamma} - L_\gamma e^{-L_\gamma}) \\ + (1 + \mu)^2 \left(\frac{1 - e^{-2L_\gamma}}{2} - \frac{(1 - e^{-L_\gamma})^2}{L_\gamma} \right), \\ L_\gamma = \ln \frac{1 + \gamma}{\gamma}, \quad (13)$$

and \mathcal{E}_T and u_T are field and potential in the Townsend limit of “vanishing” current, i.e., with breakdown values

$$\mathcal{E}_T = \frac{1}{\ln(L/L_\gamma)}, \quad u_T = \frac{L}{\ln(L/L_\gamma)}. \quad (14)$$

The minimal potential u_T is $L_\gamma e^1$, it is attained for gap length $L = L_\gamma e^1$ on the Paschen curve $u_T = u_T(L)$ [1–3].

In [1], it was argued that the coefficient A_{SE} in (10) strongly depends on γ due to the factor $F(\gamma, \mu)$ in (11). This factor strongly depends on γ , for small γ actually in leading order like $L_\gamma^3/12$. (Note that there is a discrepancy between equation (50) in [1] for $F(\gamma, \mu)$ which is reproduced as Eq. (13) in the present paper, and the plot in Fig. 1 of [1] for $10^{-1} < \gamma < 10^0$. Equation (50) in [1] is correct and the figure erroneous. $F(\gamma, \mu)$ actually varies by five orders of magnitude on $10^{-6} < \gamma < 10^0$, not only by four.)

At this point, the question how the remaining factor depends on γ was omitted. In fact, the denominator $(\alpha \mathcal{E}_T)^3$ in (10) has in leading order the same strong dependence on γ , since

$$\frac{1}{(\alpha \mathcal{E}_T)^3} = \left(\frac{L}{L_\gamma} \ln(L/L_\gamma) \right)^3, \quad (15)$$

according to the Townsend breakdown criterion $\alpha L = L_\gamma$, cf. (12)–(14). Therefore the leading order dependence on L_γ^3 of the coefficient of \bar{j}^2 in (10) is cancelled and replaced by a dependence on L^3 , while the term with α'' has the classical explicit form

$$\frac{\mathcal{E}_T \alpha''}{2\alpha'} = \frac{1 - 2\mathcal{E}_T}{2\mathcal{E}_T} = \frac{\ln(L/L_\gamma) - 2}{2}. \quad (16)$$

In [2, 3], another small current expansion was derived from (1)–(3), assuming $n_+ \gg n_e$ and $n_+(x) \approx \text{const}$. This approximation was criticised in [1], since it is in contradiction with the boundary condition (5) — however, for very small γ , it is a good approximation in a large part of the gap. The resulting equations (8.8) and (8.10) from [2] read in the notation of the present paper

$$U = U_T - \frac{U_T}{48} \frac{1 - 2\mathcal{E}_T}{2\mathcal{E}_T} \left(\frac{J}{J_L} \right)^2, \quad (17)$$

$$J_L = \frac{\epsilon_0 \mu_+ U_T^2}{2d^3}. \quad (18)$$

(Here a misprint in [2] was corrected, namely the missing factor U_T in the coefficient of J^2 in (17), and the factor $1/(8\pi)$ in (8.8) is substituted by $\epsilon_0/2$ in (18), since we here write the Poisson equation (2) in MKS units rather than in Gaussian units, cf. (8.6) in [2].)

In (17), the physical current density J is compared to J_L . J_L is the current density where deviations from the Townsend limit through space charges start to occur; it explicitly depends on γ through U_T (14).

Comparison of the results of Šijačić/Ebert (SE) (10) and Engel/Raizer (ER) (17) show that the coefficients $A_{SE,ER}$ in the expansion (10) are related like

$$A_{SE} = A_{ER} \frac{12 F(\gamma, \mu)}{L_\gamma^3}, \quad A_{ER} = \frac{1 - 2\mathcal{E}_T}{2\mathcal{E}_T} \frac{L^3}{12 \mathcal{E}_T^3}. \quad (19)$$

The coefficients A_{SE} and A_{ER} depend in the same way on L , and they are essentially independent of μ for realistic values of μ . Therefore the ratio A_{SE}/A_{ER} depends only on γ as shown in Fig. 1. For $\gamma \rightarrow 0$, the ratio tends to unity. For a large range of γ values, the deviation is not too large, approaching a factor 0.44 for $\gamma = 10^{-1}$.




FIG. 1: The ratio A_{SE}/A_{ER} of the small current expansions by Šijačić/Ebert and Engel/Raizer as a function of γ .

Fig. 2 shows that the factor A_{SE} indeed strongly depends on γ for the given L .

The strong dependence of A_{SE} or A_{ER} on γ for a given short gap length L means that we can obtain both negative and positive differential resistance dU/dJ close to the Townsend limit for the same gap length. Therefore the choice of γ is important since it can change the differential conductivity and therefore the stability of a Townsend discharge in a short gap.

Acknowledgments

Y.R. acknowledges hospitality of CWI Amsterdam and D.S. a Ph.D. grant of the Dutch physics funding agency FOM.




FIG. 2: The normalized coefficient $A = 24 A_{SE}/(L^3 \ln^4 L)$ as a function of γ for gap lengths $L = A pd = 15, 30, 60, 120, 240$ (dashed and solid lines with labels).

-
- [1] D.D. Šijačić, U. Ebert, Phys. Rev. E **66**, 066410 (2002).
 [2] Yu.P. Raizer, *Gas Discharge Physics* (Springer, Berlin, 1991).
 [3] A. von Engel and M. Steenbeck, **Elektrische Gasent-**

ladungen. Ihre Physik und Technik, Vol. II (Springer, Berlin 1934).