

**stichting
mathematisch
centrum**



AFDELING TOEGEPASTE WISKUNDE
(DEPARTMENT OF APPLIED MATHEMATICS)

TW 172/77

NOVEMBER

O. DIEKMANN & H.G. KAPER

ON THE BOUNDED SOLUTIONS OF A NONLINEAR
CONVOLUTION EQUATION

Preprint

2e boerhaavestraat 49 amsterdam

5778.821

BIBLIOTHEEK MATHEMATISCH CENTRUM
AMSTERDAM

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).

On the bounded solutions of a nonlinear convolution equation *

by

O. Diekmann & H.G. Kaper **

ABSTRACT

This investigation is concerned with the nonlinear convolution equation $u(x) - (g \circ u) * k(x) = 0$ on the real line \mathbb{R} . The kernel k is nonnegative and integrable on \mathbb{R} , with $\int_{\mathbb{R}} k(x) dx = 1$; the function g is real-valued and continuous on \mathbb{R} , $g(0) = 0$, and there exists a $p > 0$ such that $g(x) > x$ for $x \in (0, p)$ and $g(p) = p$. Sufficient conditions are given for the non-existence of bounded nontrivial solutions. Implications for the solution of the inhomogeneous equation $u(x) - (g \circ u) * k(x) = f(x)$, $x \in \mathbb{R}$, are discussed. Finally, uniqueness (modulo translation) is shown to hold. The results are applied to a problem of mathematical epidemiology.

KEY WORDS & PHRASES: *nonlinear convolution equation on the real line; non-existence; asymptotic behavior; uniqueness modulo translation; Tauberian methods; mathematical epidemiology*

* * Applied Mathematics Division, Argonne National Laboratory, Argonne, IL 60439, USA. Work performed under the auspices of the U.S. Department of Energy.

* This report will be submitted for publication elsewhere

1. INTRODUCTION

Certain deterministic models for the spatial spread of an epidemic or an advantageous gene among a population along a line can be analyzed in terms of the nonlinear convolution equation

$$(1.1) \quad u(x) = (g \circ u) * k(x) \quad x \in \mathbb{R},$$

where $g \circ u$ is the composite of g and u , $g \circ u(x) = g(u(x))$, and $*$ denotes convolution, $\phi * \psi(x) = \int_{\mathbb{R}} \phi(x-y)\psi(y)dy$. In the epidemic model, g typically has the form $g(x) = \alpha(1-e^{-x})$, α some constant ($\alpha > 1$), cf. DIEKMANN [6], while in the genetic model g is given by $g(x) = [\alpha x^2 + \beta x(1-x)] / [\alpha x^2 + 2\beta x(1-x) + \gamma(1-x)^2]$, α , β , and γ positive constants, cf. WEINBERGER [16]. In both cases, k is a nonnegative function, normalized such that $\int_{\mathbb{R}} k(x)dx = 1$.

We consider eq. (1.1) under the following hypotheses:

(H_g¹) $g: \mathbb{R} \rightarrow \mathbb{R}$ is continuous; $g(0) = 0$; there exists a $p > 0$ such that $g(x) > x$ for $0 < x < p$ and $g(p) = p$.

(H_k¹) $k: \mathbb{R} \rightarrow \mathbb{R}$ is nonnegative; $k \in L_1(\mathbb{R})$ with $\int_{\mathbb{R}} k(x)dx = 1$; $\int_{\mathbb{R}} |x|k(x)dx < \infty$.

Further hypotheses on g and k will be introduced as needed.

Equation (1.1) has at least two constant solutions, viz., $u_0: x \mapsto 0$ and $u_p: x \mapsto p$. These we call trivial solutions. By a nontrivial solution of (1.1) we mean a Lebesgue measurable function $u: \mathbb{R} \rightarrow \mathbb{R}$ which satisfies (1.1) and the inequalities $0 \leq u(x) \leq p$ for almost all $x \in \mathbb{R}$ and which is not a trivial solution. We are interested in nontrivial solutions of (1.1) and in the solution of the corresponding inhomogeneous equation

$$(1.2) \quad u(x) = (g \circ u) * k(x) + f(x) \quad x \in \mathbb{R},$$

where f is a given nonnegative function.

In section 2 we establish various general properties of the solutions of equation (1.1). In Sections 3 and 4 we prove some non-existence results, i.e., we establish conditions on g and k under which one can prove that the equation (1.1) does not have a nontrivial solution. These results are

complementary to some earlier results of WEINBERGER [16] and of one of the authors [6] concerning the existence of nontrivial solutions of (1.1). As we will show, the existence or non-existence of such solutions hinges upon the distribution of the mass of the kernel k . Formally, this is determined by the existence or non-existence of real zeros of the function $\lambda \mapsto 1 - g'(0)K(\lambda)$ ($\lambda \in \mathbb{R}$), where K is the two-sided Laplace transform of k . The problem has some Tauberian aspects and part of our arguments are based on Pitt's form of Wiener's general Tauberian theorem. This approach is related to the use of Tauberian methods in the study of linear convolution equations by KARLIN [9] and ESSÉN [7], cf. [14, Chapter 9].

In Section 5 we study the inhomogeneous equation (1.2). In Section 6 we turn our attention to the case when a nontrivial solution of (1.1) exists and investigate the question of its uniqueness (modulo translation, as eq. (1.1) is translation invariant). Here, our method of proof forces us to impose further conditions on g and k .

In the final Section 7 we apply our results to a problem in mathematical epidemiology, viz., the traveling wave problem for the Kermack-McKendrick-Kendall model for the spatial spread of an epidemic, cf. [1], [2], [3], [4], [6] [10], [11], [12].

NOTATION. \mathbb{N}_+ denotes the set of all positive integers; $\mathbb{R}_+ = \{x \in \mathbb{R} : x \geq 0\}$ and $\mathbb{R}_- = \{x \in \mathbb{R} : x \leq 0\}$. The letter C is used to denote a generic positive constant.

2. PRELIMINARIES

Starting from k we form a sequence of functions $\{k^{n*} : n \in \mathbb{N}_+\}$. We interpret k^{1*} as k , and define for $n \in \mathbb{N}_+$, $k^{(n+1)*} := k^{n*} * k$. Thus, $k^{n*} \in L_1(\mathbb{R})$ and $\int_{\mathbb{R}} k^{n*}(x) dx = 1$ for all n . Following FELLER [8, Sections V.2, V.4], we call x a point of increase of k if $\int_I k(y) dy > 0$ for every open interval I containing x , and we define $\Sigma := \bigcup_{n=1}^{\infty} \Sigma_n$, where Σ_n denotes the set formed by the points of increase of k^{n*} . We say that k is concentrated on a set I if $\int_{\mathbb{R} \setminus I} k(x) dx = 0$. The following result will be used several times.

LEMMA 2.1. *If k is concentrated on \mathbb{R}_+ (\mathbb{R}_-), then there exists an $a \geq 0$ such that \sum contains the interval $[a, \infty)$ ($(-\infty, -a]$). If k is not concentrated on either \mathbb{R}_+ or \mathbb{R}_- , then $\sum = \mathbb{R}$.*

PROOF. For an arbitrary real number $b > 0$ we define $\ell(x) := \min\{k(x), b\}$. Then $\ell \in L_\infty(\mathbb{R}) \cap L_1(\mathbb{R})$ and, furthermore,

$$k^{2*}(x) \geq \ell^{2*}(x) \geq \int_0^x \ell(x-y)\ell(y)dy =: L(x).$$

We claim that $L(x)$ is uniformly continuous. Indeed, $|L(x_1) - L(x_2)| \leq b^2|x_1 - x_2| + b \int_{\mathbb{R}} |\ell(y) - \ell_{x_1 - x_2}(y)| dy$, where $\ell_x(y) := \ell(y-x)$, and the translation map $x \mapsto \ell_x$ from \mathbb{R} into $L_1(\mathbb{R})$ is uniformly continuous, see [13, Section 1.1.5].

Suppose $\int_{\mathbb{R}_+} k(x)dx > 0$. Then L cannot be identically zero on \mathbb{R}_+ and, hence, there exist $x_1 \geq 0$ and $\delta > 0$ such that $k^{2*}(x) \geq L(x) > 0$ for $x \in (x_1, x_1 + \delta)$. From this we deduce that $k^{4*}(x) = (k^{2*} * k^{2*})(x) > 0$ for $x \in (2x_1, 2x_1 + 2\delta)$ and, by induction, $k^{2n*}(x) > 0$ for $x \in (nx_1, nx_1 + n\delta)$, $n \in \mathbb{N}_+$. For sufficiently large $n(n > \delta^{-1}x_1)$, successive intervals become overlapping and, consequently, \sum contains an interval $[a, \infty)$ for some $a \geq 0$. A similar argument shows that if $\int_{\mathbb{R}_-} k(x)dx > 0$, then \sum contains an interval $(-\infty, -a]$ for some $a \geq 0$.

Suppose now that both $\int_{\mathbb{R}_+} k(x)dx > 0$ and $\int_{\mathbb{R}_-} k(x)dx > 0$. Let $x_0 \in \mathbb{R}$ be arbitrary. For sufficiently large y we know that there exist positive integers m and n such that $k^{n*}(x) > 0$ for x in a neighborhood of $\frac{1}{2}x_0 + y$ and $k^{m*}(x) > 0$ for x in a neighborhood of $\frac{1}{2}x_0 - y$. Consequently, $k^{(n+m)*}(x) > 0$ for x in a neighborhood of x_0 and $x_0 \in \sum$. \square

The next lemma contains some basic information concerning nontrivial solutions of (1.1).

LEMMA 2.2. *Any nontrivial solution u of (1.1) is uniformly continuous, strictly positive, and such that $\inf\{u(x) : x \in \mathbb{R}\} = 0$.*

PROOF. Let u be a nontrivial solution of (1.1). The uniform continuity of u follows from the continuity of translation in $L_1(\mathbb{R})$ (cf. the proof of Lemma 2.1).

Next, we observe that $g \circ u(x) \geq u(x)$ for all $x \in \mathbb{R}$, so $u(x) \geq u * k(x)$ and, upon iteration, $u(x) \geq u * k^{n*}(x)$ for $n \in \mathbb{N}_+$. Hence, if $u(x_0) = 0$ for some $x_0 \in \mathbb{R}$, then $u * k^{n*}(x_0) = 0$ for $n \in \mathbb{N}_+$, which implies $u(x) = 0$ for all x such that $x_0 - x \in \sum$. On account of Lemma 2.1 this implies that $u(x) = 0$ for all $x \in \mathbb{R}$ if k is not concentrated on either \mathbb{R}_+ or \mathbb{R}_- . If, on the other hand, k is concentrated on a half-axis, \mathbb{R}_+ say, then Lemma 2.1 implies that $u(x) = 0$ for all $x \in (-\infty, x_0 - a]$ for some $a \in \mathbb{R}_+$ and, consequently, $u(x) = 0$ for all $x \in \mathbb{R}$, since the equation (1.1) has to be satisfied. In all cases we thus arrive at the conclusion that $u(x) = 0$ for all $x \in \mathbb{R}$, contrary to the assumption that u is nontrivial.

Let $\alpha := \inf\{u(x) : x \in \mathbb{R}\}$ and suppose $\alpha > 0$. Then $g \circ u(x) \geq \min\{g(y) : \alpha \leq y \leq p\}$. Since g is continuous, there exists a $\beta \in [\alpha, p]$ such that $g(\beta) = \min\{g(y) : \alpha \leq y \leq p\}$. So $u(x) = (g \circ u) * k(x) \geq g(\beta) = \alpha + (g(\beta) - \alpha) \geq \alpha + (g(\beta) - \beta) = \alpha + \varepsilon$, where $\varepsilon := g(\beta) - \beta > 0$. Hence, $\inf\{u(x) : x \in \mathbb{R}\} \geq \alpha + \varepsilon$, which is in contradiction with the definition of α . We conclude that $\alpha = 0$. \square

3. THE FIRST NON-EXISTENCE RESULT

In this section we give a non-existence result which can be proved by elementary arguments. Let L be defined by

$$(3.1) \quad L := \limsup_{x \downarrow 0} \frac{g(x)}{x}.$$

Note that $L \geq 1$; L may be infinite. If g is differentiable at the origin with derivative value $g'(0)$, then $L = g'(0)$.

LEMMA 3.1. *Suppose g is monotone nondecreasing on $[0, p]$. If $\int_{\mathbb{R}_+} k(x) dx > L^{-1}$, then any nontrivial solution u of (1.1) is such that $\liminf_{x \rightarrow \pm\infty} u(x) > 0$.*

PROOF. We prove the lemma only for the upper sign. Let u be a nontrivial solution of (1.1) and suppose that $\liminf_{x \rightarrow +\infty} u(x) = 0$. Choose a monotonically decreasing sequence $\{y_n : n \in \mathbb{N}_+\}$ such that $y_n \downarrow 0$ and $g(y_n)/y_n \rightarrow L$ as $n \rightarrow \infty$. Define x_n for $n = n_0, n_0+1, \dots$, where n_0 is such that $u(0) > y_{n_0}^{-1}$, by $x_n := \sup\{x \in \mathbb{R}_+ : u(y) \geq y_n \text{ for all } y \in [0, x]\}$. Then $u(x_n) = y_n$ and

$x_n \rightarrow \infty$ as $n \rightarrow \infty$. Furthermore, as g is monotone nondecreasing,

$$u(x_n) \geq \int_0^{x_n} g \circ u(x_n - y) k(y) dy \geq g \circ u(x_n) \int_0^{x_n} k(y) dy,$$

so $(g(y_n)/y_n) \int_0^{x_n} k(y) dy \leq 1$ for all n . Letting $n \rightarrow \infty$ we obtain the inequality $L \int_{\mathbb{R}_+} k(y) dy \leq 1$, a contradiction. We conclude that

$$\liminf_{x \rightarrow \infty} u(x) > 0. \quad \square$$

The following theorem is an immediate consequence of Lemmas 2.2 and 3.1.

THEOREM 3.2. *Suppose g is monotone nondecreasing on $[0, p]$. If $\int_{\mathbb{R}_+} k(x) dx > L^{-1}$ and $\int_{\mathbb{R}_-} k(x) dx > L^{-1}$, then there is no nontrivial solution of (1.1).*

PROOF. Let u be a nontrivial solution of (1.1). Then either $\liminf_{x \rightarrow +\infty} u(x) = 0$ or $\liminf_{x \rightarrow -\infty} u(x) = 0$, by Lemma 2.2. However, from Lemma 3.1 we infer that $\liminf_{x \rightarrow +\infty} u(x) > 0$ and $\liminf_{x \rightarrow -\infty} u(x) > 0$, and we arrive at a contradiction. We conclude that there is no nontrivial solution of (1.1).

REMARK 3.3. Up to this point we have not made use of the hypothesis that $\int_{\mathbb{R}} |x| k(x) dx < \infty$, so all foregoing results remain valid if this hypothesis is actually violated.

4. ANOTHER NON-EXISTENCE RESULT

In the present section we establish a non-existence result whose proof is rather involved. It is based on various reformulations of (1.1) as an inhomogeneous linear convolution equation. First we rewrite (1.1) in the form

$$(4.1) \quad u(x) - u * k(x) = \phi(x) \quad x \in \mathbb{R},$$

where

$$(4.2) \quad \phi(x) := (g \circ u - u) * k(x).$$

The function ϕ is nonnegative on \mathbb{R} and we will exploit this fact in the investigation of solutions of (4.1), cf. ESSÉN [7].

Let m denote the first moment of the kernel k ,

$$(4.3) \quad m = \int_{\mathbb{R}} xk(x)dx.$$

The first theorem requires barely more than a reference to the literature.

THEOREM 4.1. *If $m = 0$ then there is no nontrivial solution of (1.1).*

PROOF. It is known that the condition $m \neq 0$ is necessary and sufficient for the existence of a bounded continuous function u such that $u(x) - u*k(x) \geq 0$, with strict inequality in some point, cf. [7, Theorem 3.1]. \square

Having established this non-existence result for $m = 0$, we may henceforth assume $m \neq 0$.

Let

$$(4.4) \quad n(x) := \begin{cases} \int_x^{\infty} k(y)dy & x \geq 0, \\ -\int_{-\infty}^x k(y)dy & x < 0. \end{cases}$$

Note that $n(x) = H(x) - \int_{-\infty}^x k(y)dy$ and, consequently, $n' = \delta - k$ in the sense of distributions (here H denotes the Heaviside step function and δ the Dirac distribution). The condition $\int_{\mathbb{R}} |x|k(x)dx < \infty$ implies that $n \in L_1(\mathbb{R})$ with $\|n\|_1 = \int_{\mathbb{R}} |x|k(x)dx$, cf. [8, Section V.6.]. The Fourier transform \hat{n} of n , which is a continuous function, is related to the Fourier transform \hat{k} of k via the identity $\hat{n}(\lambda) = (1 - \hat{k}(\lambda))/(i\lambda)$ for $\lambda \neq 0$, while $\hat{n}(0) = m$. If $m \neq 0$, then $\hat{n}(\lambda) \neq 0$ for all $\lambda \in \mathbb{R}$.

LEMMA 4.2. *Suppose $m > 0$. If u is any nontrivial solution of (1.1), then $\lim_{x \rightarrow -\infty} u(x)$ and $\lim_{x \rightarrow \infty} u(x)$ exist and $u(-\infty) = 0$, $u(\infty) = p$.*

PROOF. Let u be a nontrivial solution of (1.1). Then u satisfies (4.1) with ϕ given by (4.2). We integrate both members of (4.1) from a to x ,

$$\int_a^x u(\xi) d\xi - \int_a^x \int_{-\infty}^{\infty} u(\xi-\eta)k(\eta)d\eta d\xi = \int_a^x \phi(\xi) d\xi.$$

Interchanging the order of the integrals in the second term and integrating by parts we obtain the identity

$$\int_a^x \int_{-\infty}^{\infty} u(\xi-\eta)k(\eta)d\eta d\xi = \int_a^x u(\xi) d\xi - \int_{-\infty}^{\infty} n(x-\xi)u(\xi) d\xi + \int_{-\infty}^{\infty} n(a-\xi)u(\xi) d\xi,$$

where n is defined by (4.4). Hence, u satisfies

$$(4.5) \quad u * n(x) - u * n(a) = \int_a^x \phi(\xi) d\xi \quad x \in \mathbb{R}.$$

Since u is bounded and $n \in L_1(\mathbb{R})$, $u * n$ is bounded. Also, $\phi(x)$ is nonnegative for all $x \in \mathbb{R}$, so the integral $\int_a^x \phi(\xi) d\xi$, besides being bounded, depends monotonically on x . It follows that $\lim_{x \rightarrow \infty} \int_a^x \phi(\xi) d\xi$ exists and hence, from (4.5), that $\lim_{x \rightarrow \infty} u * n(x)$ exists. In fact,

$$u * n(\infty) = u * n(a) + \int_a^{\infty} \phi(\xi) d\xi.$$

Now, n belongs to the Wiener class W (i.e., $n \in L_1(\mathbb{R})$ and $\{\lambda \in \mathbb{R} : \hat{n}(\lambda) = 0\} = \emptyset$), and u is bounded and uniformly continuous. It follows from Pitt's form of Wiener's fundamental Tauberian theorem, cf. [14, Section 9.7] or [17, Section V.10], that $\lim_{x \rightarrow \infty} u(x)$ exists and is given by

$$u(\infty) = m^{-1} \left(u * n(a) + \int_a^{\infty} \phi(x) dx \right).$$

Similarly, one shows that $\lim_{x \rightarrow -\infty} u(x)$ exists and is given by

$$u(-\infty) = m^{-1} \left(u * n(a) - \int_{-\infty}^a \phi(x) dx \right).$$

A simple argument shows that both $u(\infty)$ and $u(-\infty)$ satisfy the scalar equation $y = g(y)$, so $\{u(\infty), u(-\infty)\} \subset \{0, p\}$. Since $u(\infty) - u(-\infty) = \int_{-\infty}^{\infty} \phi(x) dx > 0$, we conclude that $u(-\infty) = 0$ and $u(\infty) = p$. \square

REMARK 4.3. The case $m < 0$ can be reduced to the case $m > 0$ by a change of variable $\tau: x \mapsto -x$.

The next step towards a non-existence result for $m \neq 0$ consists of an analysis of the rate of convergence of a nontrivial solution towards its limiting value 0 as $x \rightarrow -\infty$. The following lemma is needed in the proof of Lemma 4.5.

LEMMA 4.4. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be nonnegative, $f \in L_1(\mathbb{R})$. Define $f^{(0)}(x) := f(x)$ and, recursively, $f^{(k)}(x) := \int_{-\infty}^x f^{(k-1)}(y) dy$ for those values of $k \in \mathbb{N}_+$ for which $f^{(k-1)} \in L_1(\mathbb{R}_-)$. Then, for any $r \in \mathbb{R}$,

$$\int_{-\infty}^r (r-x)^k f(x) dx = k! \int_{-\infty}^r f^{(k)}(x) dx,$$

in the sense that if one side converges so does the other.

PROOF. The identity given in the lemma follows upon induction from the identity

$$\int_{-\infty}^r (r-x)^k f(x) dx = k \int_{-\infty}^r (r-x)^{k-1} f^{(1)}(x) dx.$$

We omit the proof of the latter identity, as it is similar to the proof of Lemma 1 in [8, Section V.6]. \square

Let ℓ be defined by

$$(4.6) \quad \ell := \liminf_{x \downarrow 0} \frac{g(x)}{x}.$$

Note that $1 \leq \ell \leq L$, cf. (3.1); ℓ may be infinite. If g is differentiable at the origin with derivative value $g'(0)$, then $\ell = L = g'(0)$.

LEMMA 4.5. Let u be a nontrivial solution of (1.1) such that $\lim_{x \rightarrow -\infty} u(x) = 0$. If $\ell > 1$, then there exists a $\delta > 0$ such that the integral $\int_{-\infty}^{\infty} e^{-\lambda x} u(x) dx$ converges for $\lambda \in S_\delta$, where $S_\delta := \{\lambda \in \mathbb{C}: 0 < \operatorname{Re} \lambda < \delta\}$.

PROOF. It suffices to prove the convergence of the integral $\int_{-\infty}^r e^{-\lambda x} u(x) dx$ for some $r < \infty$ and λ real, $0 < \lambda < \delta$.

Take $\alpha \in (1, \ell)$ and let c be such that $\int_{-c}^c k(x) dx = (\alpha+1)/(2\alpha)$. Define a new function k_c on \mathbb{R} by putting $k_c(x) := k(x)$ if $|x| \leq c$, $k_c(x) := 0$ if $|x| > c$. If u is a nontrivial solution of (1.1), then $u(x) = (g \circ u) * k_c(x) + \psi_1(x)$ for all $x \in \mathbb{R}$, with $\psi_1(x) := (g \circ u) * (k - k_c)(x) \geq 0$. We rewrite the latter identity in the form

$$(4.7) \quad u(x) - \frac{2\alpha}{\alpha+1} u * k_c(x) = \frac{\alpha-1}{\alpha+1} u(x) + \psi(x) \quad x \in \mathbb{R},$$

where $\psi(x) := \frac{2}{\alpha+1} [\psi_1(x) + \psi_2(x)]$, with $\psi_2(x) := (g \circ u - \alpha u) * k_c(x)$. Let

$$n_c(x) := \begin{cases} \frac{2\alpha}{\alpha+1} \int_x^\infty k_c(y) dy & x \geq 0, \\ -\frac{2\alpha}{\alpha+1} \int_{-\infty}^x k_c(y) dy & x < 0. \end{cases}$$

Then $\text{supp}(n_c) \subset [-c, c]$. We integrate both members of (4.7) from x to some $r < \infty$. Thus,

$$\int_x^r u(\xi) d\xi - \frac{2\alpha}{\alpha+1} \int_x^r \int_{-\infty}^\infty u(\xi-\eta) k_c(\eta) d\eta d\xi = \frac{\alpha-1}{\alpha+1} \int_x^r u(\xi) d\xi + \int_x^r \psi(\xi) d\xi.$$

Interchanging the order of the integrals in the second term and integrating by parts we obtain the identity

$$(4.8) \quad u * n_c(r) - u * n_c(x) = \frac{\alpha-1}{\alpha+1} \int_x^r u(\xi) d\xi + \int_x^r \psi(\xi) d\xi,$$

cf. (4.5). Because $\lim_{x \rightarrow -\infty} u(x) = 0$, there exists $x_\alpha \in \mathbb{R}$ such that $g \circ u(x) \geq \alpha u(x)$ for all $x \in (-\infty, x_\alpha]$ and, hence, $\psi_2(x) \geq 0$ for all $x \in (-\infty, x_\alpha - c]$. The same is then true of $\psi(x)$ and we conclude, as in the proof of Lemma 4.2, that $\lim_{x \rightarrow -\infty} \int_x^r u(\xi) d\xi$ and $\lim_{x \rightarrow -\infty} \int_x^r \psi(\xi) d\xi$ exist. Letting x tend to $-\infty$ we obtain the identity

$$\int_{-\infty}^r u(x) dx = \frac{\alpha+1}{\alpha-1} \left[u * n_c(r) - \int_{-\infty}^r \psi(x) dx \right].$$

If r is chosen such that $r \leq x_\alpha - c$ it follows that

$$\int_{-\infty}^r u(x) dx \leq \frac{\alpha+1}{\alpha-1} \max\{u(x) : x \in [r-c, r+c]\} \int_{\mathbb{R}} |n_c(x)| dx.$$

Next, we put $u^{(0)}(x) := u(x)$ and define $u^{(k)}(x) := \int_{-\infty}^x u^{(k-1)}(\xi) d\xi$, $x \in \mathbb{R}$, for those values of $k \in \mathbb{N}_+$ for which the integral exists. The functions $\psi^{(k)}$ are defined similarly. Integration of (4.7) from $-\infty$ to x yields an equation for $u^{(1)}$ similar to (4.7), with ψ replaced by $\psi^{(1)}$, from which one deduces the relation

$$u^{(1)} * n_c(r) - u^{(1)} * n_c(x) = \frac{\alpha-1}{\alpha+1} \int_x^r u^{(1)}(\xi) d\xi + \int_x^r \psi^{(1)}(\xi) d\xi,$$

cf. (4.8). The integrability of u on $(-\infty, r)$ and the continuity of u imply the boundedness of $u^{(1)}$ on $(-\infty, r+c]$. As before, it follows that $\lim_{x \rightarrow -\infty} \int_x^r u^{(1)}(\xi) d\xi$ and $\lim_{x \rightarrow -\infty} \int_x^r \psi^{(1)}(\xi) d\xi$ exist,

$$\int_{-\infty}^r u^{(1)}(x) dx = \frac{\alpha+1}{\alpha-1} \left[u^{(1)} * n_c(r) - \int_{-\infty}^r \psi^{(1)}(x) dx \right]$$

and, therefore,

$$\int_{-\infty}^r u^{(1)}(x) dx \leq \frac{\alpha+1}{\alpha-1} \max\{u^{(1)}(x) : x \in [r-c, r+c]\} \int_{\mathbb{R}} |n_c(x)| dx.$$

Continuing in this manner we obtain the sequence of inequalities,

$$\int_{-\infty}^r u^{(k)}(x) dx \leq \frac{\alpha+1}{\alpha-1} \max\{u^{(k)}(x) : x \in [r-c, r+c]\} \int_{\mathbb{R}} |n_c(x)| dx,$$

for $k \in \mathbb{N}_+$. Now, for each $k \in \mathbb{N}_+$, $u^{(k)}$ is a monotonically increasing function, so

$$\begin{aligned} \max\{u^{(k)}(x) : x \in [r-c, r+c]\} &= u^{(k)}(r+c) \\ &\leq \int_{-\infty}^r u^{(k-1)}(x) dx + cu^{(k-1)}(r+c) \\ &\leq \delta^{-1} u^{(k-1)}(r+c), \end{aligned}$$

where $\delta := [(\alpha+1)(\alpha-1)^{-1} \int_{\mathbb{R}} |n_c(x)| dx + c]^{-1}$. Hence, by induction,

$$\max\{u^{(k)}(x) : x \in [r-c, r+c]\} \leq \delta^{-k} \max\{u(x) : x \in [r-c, r+c]\} \leq p\delta^{-k}.$$

Thus, there exists a positive constant C such that

$$\int_{-\infty}^r u^{(k)}(x) dx \leq C\delta^{-k} \quad k \in \mathbb{N}_+.$$

Next, we apply Lemma 4.4 to the function u and conclude that each of the integrals $\int_{-\infty}^r (r-x)^k u(x) dx$ converges. Moreover,

$$\frac{1}{k!} \int_{-\infty}^r (r-x)^k u(x) dx \leq C\delta^{-k}.$$

This result, in turn, implies the convergence of the infinite sum $\sum_{k=0}^{\infty} (k!)^{-1} \lambda^k \int_{-\infty}^r (r-x)^k u(x) dx$ and therefore of the integral $\int_{-\infty}^r e^{\lambda(r-x)} u(x) dx$, at least for $\lambda < \delta$. \square

LEMMA 4.6. *Let u be a nontrivial solution of (1.1). If the integral $\int_{-\infty}^{\infty} e^{-\lambda x} u(x) dx$ converges in the strip S_{δ} (cf. Lemma 4.5) then so does the integral $\int_{-\infty}^{\infty} e^{-\lambda x} k(x) dx$.*

PROOF. It suffices to prove the convergence of the integral $\int_{-\infty}^{\infty} e^{-\lambda x} k(x) dx$ for λ real, $0 < \lambda < \delta$.

Since u is a nontrivial solution of (1.1) we have $u * k(x) \leq u(x)$ for all $x \in \mathbb{R}$, so $\int_{-\infty}^{\infty} e^{-\lambda x} u * k(x) dx \leq \int_{-\infty}^{\infty} e^{-\lambda x} u(x) dx < \infty$ for any $\lambda \in (0, \delta)$. On the other hand, $\int_{-\infty}^{\infty} e^{-\lambda x} u * k(x) dx = (\int_{-\infty}^{\infty} e^{-\lambda x} k(x) dx) (\int_{-\infty}^{\infty} e^{-\lambda x} u(x) dx)$, in the sense that if the integral on the left hand side exists then so do the integrals on the right hand side. Hence, $\int_{-\infty}^{\infty} e^{-\lambda x} k(x) dx$ exists for $\lambda \in (0, \delta)$ and satisfies the inequality $\int_{-\infty}^{\infty} e^{-\lambda x} k(x) dx \leq 1$ there. \square

We remark that Lemmas 4.5 and 4.6 together imply that, if $\ell > 1$ then a necessary condition for the existence of nontrivial solutions of (1.1) is that $\int_{-\infty}^{\infty} e^{-\lambda x} k(x) dx$ converges in S_{δ} for some $\delta > 0$.

The above results lead us to define a real number Λ_k ,

$$(4.9) \quad \Lambda_k := \sup \left\{ \lambda \in \mathbb{R} : \int_{\mathbb{R}} e^{-\lambda x} k(x) dx < \infty \right\}$$

and to associate with each nontrivial solution u of (1.1) a number Λ_u thus,

$$(4.10) \quad \Lambda_u := \sup \left\{ \lambda \in \mathbb{R} : \int_{\mathbb{R}} e^{-\lambda x} u(x) dx < \infty \right\}.$$

It follows from Lemma 4.6 that $\Lambda_u \leq \Lambda_k$. Furthermore, let

$$K(\lambda) := \hat{k}(-i\lambda) = \int_{\mathbb{R}} e^{-\lambda x} k(x) dx,$$

$$U(\lambda) := \hat{u}(-i\lambda) = \int_{\mathbb{R}} e^{-\lambda x} u(x) dx,$$

whenever the defining integrals converge. It is a standard result from Laplace transform theory that if $\Lambda_u > 0$ then U is analytic in the strip $S_u := \{\lambda \in \mathbb{C} : 0 < \operatorname{Re} \lambda < \Lambda_u\}$, cf. [17, Section VI.4]. The following result is a consequence of the positivity of $u(x)$.

LEMMA 4.7. *If u is a nontrivial solution of (1.1) and $0 < \Lambda_u < \infty$, then $U(\lambda)$ is singular at $\lambda = \Lambda_u$.*

PROOF. Cf. [17, Section II.5]. \square

At this point we make two further hypotheses concerning the functions g and k :

(H_g²) the function g is differentiable at 0 with derivative value $g'(0)$ and $g(x) = g'(0)x + o(x^{1+\varepsilon})$ as $x \downarrow 0$ for some $\varepsilon > 0$;

(H_k²) the kernel k is exponentially small at $-\infty$, i.e., there exists a $\delta > 0$ such that $e^{-\lambda x} k(x)$ is bounded for $\lambda \in (0, \delta)$.

Under these additional hypotheses we can prove the following result.

LEMMA 4.8. *Let u be a nontrivial solution of (1.1) with $\lim_{x \rightarrow -\infty} u(x) = 0$. If $0 < \Lambda_u < \Lambda_k$, then $1 - g'(0)K(\Lambda_u) = 0$.*

PROOF. We rewrite (1.1) as a linear inhomogeneous integral equation,

$$u(x) - g'(0)u * k(x) = r(x) \quad x \in \mathbb{R},$$

where $r(x) := (g \circ u - g'(0)u) * k(x)$. A two-sided Laplace transform yields the equation

$$(4.11) \quad (1 - g'(0)K(\lambda))U(\lambda) = R(\lambda),$$

which is valid for all $\lambda \in S_{\Lambda_u}$. In fact, the right abscissa of convergence of R lies to the right of Λ_u . This can be shown in the following manner. Because of the hypothesis (H_k^2) , $u(x)e^{-\lambda x} = (g \circ u) * k(x)e^{-\lambda x} \leq C \int_{\mathbb{R}} g \circ u(y)e^{-\lambda y} dy$. The integral is bounded for $\text{Re} \lambda$ sufficiently small, since $g(x) < Cx$ for x near zero ($x > 0$). Hence, using the hypothesis (H_g^2) we obtain for $\lambda \in (0, \Lambda_u)$ and ν sufficiently small ($\nu > 0$),

$$\begin{aligned} & \left| \int_{-\infty}^{\infty} e^{-(\lambda+\nu)x} (g \circ u - g'(0)u) * k(x) dx \right| \\ &= K(\lambda+\nu) \int_{-\infty}^{\infty} e^{-(\lambda+\nu)x} |(g \circ u - g'(0)u)(x)| dx \\ &\leq C K(\lambda+\nu) \int_{-\infty}^{\infty} e^{-\lambda x} u(x) (e^{-(\nu/\varepsilon)x} u(x))^\varepsilon dx \\ &\leq C K(\lambda+\nu) U(\lambda) (\sup\{e^{-(\nu/\varepsilon)x} u(x) : x \in \mathbb{R}\})^\varepsilon. \end{aligned}$$

This result shows that the abscissa of convergence of R lies to the right of Λ_u . Consequently, $R(\lambda)$ is analytic in the neighborhood of $\lambda = \Lambda_u$. Since $U(\lambda)$ is singular at $\lambda = \Lambda_u$ (Lemma 4.7) we conclude from (4.11) that $1 - g'(0)K(\lambda) = 0$ at $\lambda = \Lambda_u$. \square

The preceding lemmas lead to the following non-existence theorem.

THEOREM 4.9. *Suppose $m > 0$, $g'(0) > 1$ and $1 - g'(0)K(\lambda) \neq 0$ for all $\lambda \in [0, \Lambda_k)$. If there exists a $\lambda < \Lambda_k$ such that $\int_{-\infty}^0 e^{-\lambda x} k(x) dx > (g'(0))^{-1}$, then there is no nontrivial solution of (1.1).*

PROOF. Suppose (1.1) has a nontrivial solution u . Then $\lim_{x \rightarrow -\infty} u(x) = 0$ by Lemma 4.2, and $\Lambda_u > 0$ by Lemma 4.5 and the definition (4.10) of Λ_u . For any $\alpha \in (1, g'(0))$ there exists an x_α such that $g \circ u(x) \geq \alpha u(x)$ for all $x \leq x_\alpha$. Take any $c > 0$. Then $u(x) \geq \alpha \int_{-c}^{\infty} u(x-y)k(y)dy$ whenever $x \leq x_\alpha - c$ and, consequently,

$$u(x)e^{-\lambda x} \geq \alpha \int_{-c}^{\infty} e^{-\lambda(x-y)} u(x-y)e^{-\lambda y} k(y) dy$$

for real λ . Now, $\liminf_{x \rightarrow -\infty} u(x)e^{-\lambda x} = 0$ for any $\lambda \in [0, \Lambda_u)$, so if we take a monotonically decreasing sequence $\{y_n : n \in \mathbb{N}_+\}$ such that $y_n \downarrow 0$ as $n \rightarrow \infty$, and define x_n for $n = n_0, n_0+1, \dots$, where n_0 is such that $u(0) > y_{n_0}$, by $x_n := \inf\{x \in \mathbb{R}_- : u(y)e^{-\lambda y} \geq y_n \text{ for all } y \in [x, 0]\}$, then $u(x_n)e^{-\lambda x_n} = y_n$ and $x_n \rightarrow -\infty$ as $n \rightarrow \infty$. Now $[-c, 0] \subset [x_n, 0]$ for n sufficiently large, so $u(x_n - y)e^{-\lambda(x_n - y)} \geq y_n$ for all $y \in [-c, 0]$ and

$$\begin{aligned} y_n = u(x_n)e^{-\lambda x_n} &\geq \alpha \int_{-c}^{\infty} e^{-\lambda(x_n - y)} u(x_n - y)e^{-\lambda y} k(y) dy \\ &\geq \alpha \int_{-c}^0 e^{-\lambda(x_n - y)} u(x_n - y)e^{-\lambda y} k(y) dy \\ &\geq \alpha y_n \int_{-c}^0 e^{-\lambda y} k(y) dy. \end{aligned}$$

Thus, we obtain the inequality

$$(4.12) \quad \alpha \int_{-c}^0 e^{-\lambda y} k(y) dy \leq 1,$$

which is valid for any $\lambda \in [0, \Lambda_u)$. If Λ_u were strictly less than Λ_k , then the function $\lambda \mapsto 1 - g'(0)K(\lambda)$ would have a zero at Λ_u by Lemma 4.8, i.e., inside the interval $[0, \Lambda_k)$, contrary to the hypothesis of the theorem. Hence, $\Lambda_u = \Lambda_k$ and the inequality (4.12) holds for all $\lambda \in [0, \Lambda_k)$. Since α and c were chosen arbitrarily subject to the constraints $1 < \alpha < g'(0)$ and $c > 0$ we conclude that the assumption of the existence of a nontrivial solution of (1.1) leads to the validity of the inequality $g'(0) \int_{-\infty}^0 e^{-\lambda y} k(y) dy \leq 1$ for all $\lambda \in [0, \Lambda_k)$. This proves the theorem. \square

If $\Lambda_k = \infty$, or if $K(\lambda) \rightarrow \infty$ for $\lambda \uparrow \Lambda_k$, then there certainly exists a $\lambda < \Lambda_k$ such that $\int_{-\infty}^0 e^{-\lambda x} k(x) dx > (g'(0))^{-1}$. If, on the other hand, $K(\lambda)$ approaches a finite limit as $\lambda \uparrow \Lambda_k$, then Theorem 4.9 is not necessarily applicable. In that case, one can conclude from a Tauberian theorem for Laplace transforms, cf. [17, Section V.4], that the function $u_\lambda: x \mapsto u(x)e^{-\lambda x}$ belongs to $L_1(\mathbb{R})$ for $\lambda = \Lambda_k$. This implies that, under the hypothesis that $k_\lambda: x \mapsto k(x)e^{-\lambda x}$ is bounded for $\lambda = \Lambda_k$, u_λ is bounded for $\lambda = \Lambda_k$. Then one can repeat the analysis of the present section, starting from the identity $u_\lambda(x) = \int_{\mathbb{R}} e^{-\lambda y} (g \circ u)(y) k_\lambda(x-y) dy$ for $\lambda = \Lambda_k$, to obtain a non-existence result. We do not elaborate this idea any further.

REMARK 4.10. The condition that k has a finite first absolute moment ($\int_{\mathbb{R}} |x|k(x) dx < \infty$) has been used only once, in Lemma 4.2, to prove that any nontrivial solution of (1.1) has limits as $x \rightarrow \pm \infty$. The proofs of Lemmas 4.5 through 4.8 and Theorem 4.9 proceeded without an explicit reference to this condition. It remains an open question whether, if k does not have a finite first absolute moment, there exist nontrivial solutions of (1.1) which do not have limits as $x \rightarrow \pm \infty$.

5. THE INHOMOGENEOUS EQUATION

As the greater part of the proofs in the foregoing section were based on inequalities, rather than equalities, the same or similar arguments can be used in the analysis of the inhomogeneous equation

$$(5.1) \quad u(x) = (g \circ u) * k(x) + f(x) \quad x \in \mathbb{R}.$$

In fact, let f satisfy the following hypothesis:

(H_f) $f: \mathbb{R} \rightarrow \mathbb{R}$ is uniformly continuous, nonnegative and not equal to the zero function.

We then have the following result.

THEOREM 5.1. *Suppose $g(x) \geq p$ for $x \geq p$. The same hypotheses on g and k which lead to the non-existence results of Theorem 4.1 and Theorem 4.9 guarantee that any bounded nonnegative solution u of (5.1) necessarily*

satisfies the inequality $u(x) \geq p$ for all $x \in \mathbb{R}$.

PROOF. Let u satisfy (5.1). We introduce the function w by the definition $w(x) := \min\{u(x), p\}$, $x \in \mathbb{R}$. Then $g \circ u(x) \geq w(x)$ for all $x \in \mathbb{R}$, so $u(x) \geq (g \circ u) * k(x) \geq w * k(x)$. But $w(x) \leq p$ for all $x \in \mathbb{R}$, so $w * k(x) \leq p \int_{\mathbb{R}} k(x) dx = p$. Combining these two results we see that w satisfies the inequality $w * k(x) \leq \min\{u(x), p\} = w(x)$ for all $x \in \mathbb{R}$. Under the hypotheses of Theorem 4.1, this inequality leads to the conclusion that w is constant, $w(x) = c$ say. The case $0 \leq c < p$ is excluded, because then $u(x) = c$ and (5.1) cannot be satisfied. Hence, $w(x) = p$ and $u(x) \geq p$ for all $x \in \mathbb{R}$. Under the hypotheses of Theorem 4.9 it follows that $w(-\infty) := \lim_{x \rightarrow -\infty} w(x)$ and $w(\infty) := \lim_{x \rightarrow \infty} w(x)$ exist and are such that $w(\infty) - w(-\infty) \geq 0$ and $\{w(-\infty), w(\infty)\} \subset \{0, p\}$, cf. Lemma 4.2. One shows as in Lemma 2.2 that $w(x) > 0$ for all $x \in \mathbb{R}$, and that either $w(x) = p$ for all $x \in \mathbb{R}$ or $\inf\{w(x) : x \in \mathbb{R}\} = 0$. It follows that either $w(x) = p$ for all $x \in \mathbb{R}$ or $w(-\infty) = 0$. In the latter case, the chain of arguments leading from Lemma 4.4 to Theorem 4.9 can be repeated mutatis mutandis. Under the conditions of Theorem 4.9 the assumption that there exists a nontrivial w such that $\lim_{x \rightarrow -\infty} w(x) = 0$ leads to a contradiction, so necessarily $w(x) = p$, i.e., $u(x) \geq p$, for all $x \in \mathbb{R}$. \square

REMARK 5.2. A close examination of the proof of Theorem 5.1 shows that the condition (H_f) can be weakened somewhat if there exists a monotonically nondecreasing function g_0 on $[0, p]$ such that $g(x) \geq g_0(x) \geq x$ for all $x \in [0, p]$. In fact, let $f: \mathbb{R} \rightarrow \mathbb{R}$ be such that $f(x) \geq f_0(x)$ for all $x \in \mathbb{R}$, where $f_0: \mathbb{R} \rightarrow \mathbb{R}$ satisfies (H_f) . Suppose u is a solution of (5.1). Consider the equation

$$(5.2) \quad v(x) = (g_0 \circ v) * k(x) + f_0(x) \quad x \in \mathbb{R}.$$

The minimal solution of (5.2) can be constructed via the method of successive iterations, $v^{(0)}(x) := f_0(x)$, $v^{(n)}(x) := (g_0 \circ v^{(n-1)}) * k(x) + f_0(x)$ ($n \in \mathbb{N}_+$). Any of these iterates is majorized by the solution u of (5.1). The monotonicity of g_0 implies that the sequence $\{v^{(n)}(x) : n \in \mathbb{N}_+\}$ is monotonically nondecreasing for each $x \in \mathbb{R}$, so the sequence converges to a limit function, v say, which satisfies the equation (5.2) and the inequality

$v(x) \leq u(x)$ for all $x \in \mathbb{R}$. Suppose now that the function g_0 is such that any bounded nonnegative solution v of (5.2) necessarily satisfies the inequality $v(x) \geq p$ for all x , according to the theorem, then the solution u of (5.1) must also satisfy the inequality $u(x) \geq p$ for all $x \in \mathbb{R}$.

6. ASYMPTOTIC BEHAVIOR AND UNIQUENESS CRITERIA

The key assumption leading to the non-existence result of Theorem 4.9 is that the function $\lambda \mapsto 1 - g'(0)K(\lambda)$ does not vanish in the interval $[0, \Lambda_k)$. In the present section we will drop this assumption, i.e., we assume that the function $\lambda \mapsto 1 - g'(0)K(\lambda)$ has a real positive zero. The non-negativity of k implies that K is a convex function on $[0, \Lambda_k)$, so $\lambda \mapsto 1 - g'(0)K(\lambda)$ has at most two real positive zeros. Let σ denote the smallest positive zero; we assume that it is simple.

In Section 4 we established various conditions which guarantee that any nontrivial solution of (1.1) has the property that $\lim_{x \rightarrow -\infty} u(x) = 0$. We also analyzed the rate of convergence of $u(x)$ to its limiting value 0 in terms of the Laplace transform of u . Our first objective in the present section is to establish conditions which guarantee that any nontrivial solution of (1.1) has the much stronger property that $u(x) \sim e^{\sigma x}$ as $x \rightarrow -\infty$. We will then use this knowledge of the asymptotic behavior to establish a uniqueness result for the nontrivial solutions of (1.1).

The following lemma is a modified version of Ikehara's Theorem and will be needed in the proof of Theorem 6.2.

LEMMA 6.1. *If the real-valued function ϕ is nonnegative and nonincreasing on \mathbb{R}_+ , and there exists a $\tau > 0$ such that the integral $f(\lambda) = \int_0^\infty e^{-\lambda x} \phi(x) dx$ converges for $\operatorname{Re} \lambda > -\tau$, and if, furthermore, for some constant A and some function $g: \mathbb{R} \rightarrow \mathbb{R}$,*

$$\lim_{x \downarrow -\tau} [f(x+iy) - A(\tau+x+iy)^{-1}] = g(y),$$

uniformly on compact subsets of \mathbb{R} , then

$$\lim_{x \rightarrow \infty} \phi(x) e^{\tau x} = A.$$

PROOF. Analogous to the proof of Ikehara's Theorem, see e.g. [17, Section V.17]. \square

THEOREM 6.2. *Suppose $m > 0$, $g'(0) > 1$ and $g(x) \leq g'(0)x$ for $x \in [0, p]$. If u is a monotone nondecreasing nontrivial solution of (1.1), then there exists a positive constant A such that $\lim_{x \rightarrow \infty} u(x)e^{-\sigma x} = A$.*

PROOF. Let u be a monotone nondecreasing nontrivial solution of (1.1). It follows from Lemma 4.5 and the definition (4.10) of Λ_u that $U(\lambda)$ is defined for $\lambda \in [0, \Lambda_u)$ for some $\Lambda_u > 0$. Moreover, $[1 - g'(0)K(\lambda)]U(\lambda) = R(\lambda)$, cf. (4.11). The continuity of g , together with the properties $g(0) = 0$, $g'(0) > 1$, and $g(0) = p$, imply that the inequality \leq in the condition $g(x) \leq g'(0)x$ for $x \in [0, p]$ in the statement of the theorem, is a strict inequality, $g(x) < g'(0)x$, at least on some subinterval of $[0, p]$. Hence, $g \circ u(x) < g'(0)u(x)$ on a set of positive measure on \mathbb{R} and, consequently, the function $R: \lambda \mapsto \int_{\mathbb{R}} e^{-\lambda x} (g \circ u - g'(0)u) * k(x) dx$ does not vanish at $\lambda = \sigma$. Thus,

$$U(\lambda) = \frac{R(\lambda)}{1 - g'(0)K(\lambda)} \sim \frac{R(\sigma)}{g'(0)K'(\sigma)} (\sigma - \lambda)^{-1} \quad \text{as } \lambda \uparrow \sigma.$$

The function R is regular in a neighborhood of the line $\operatorname{Re} \lambda = \sigma$. The zeros of the function $\lambda \mapsto 1 - g'(0)K(\lambda)$, which lie in a vertical strip left-adjacent to the line $\operatorname{Re} \lambda = \sigma$ lie, in fact, in a rectangle according to the Lemma of Riemann-Lebesgue [15, Section 1.8]. The analyticity of K then implies that there are only finitely many inside this rectangle. Hence, if we choose the width of the strip sufficiently small, then there are no other zeros of $\lambda \mapsto 1 - g'(0)K(\lambda)$ in this strip, besides the real zero $\lambda = \sigma$, and we can apply Lemma 6.1 (with the obvious modifications) to obtain the statement of the theorem. \square

The use of an Ikehara-type lemma makes the proof of the asymptotic behavior rather simple, but has the disadvantage that the conclusion holds only for monotone solutions. Our next objective is to give an alternate proof of the asymptotic behavior, under slightly different assumptions on k , without presupposing that the solution be monotone.

THEOREM 6.3. Suppose $m > 0$, $g'(0) > 1$, and $g(x) \leq g'(0)x$ for $x \in [0, p]$. Let $\varepsilon > 0$ be such that $g(x) = g'(0)x + o(x^{1+\varepsilon})$, cf. (H_g^2) . Suppose that $K(-\delta) < \infty$ for some $\delta > 0$, and that there exists a $\beta \in (\sigma, \sigma(1+\varepsilon))$ such that

- (i) $g'(0)K(\beta) < 1$, and
- (ii) $k(x)e^{-(\beta/(1+\varepsilon))x}$ is bounded.

If u is a nontrivial solution of (1.1), then there exists a positive constant A such that $\lim_{x \rightarrow -\infty} u(x)e^{-\sigma x} = A$.

PROOF. In this proof we use for any given function $\phi: \mathbb{R} \rightarrow \mathbb{R}$ the symbol ϕ_λ to denote the mapping $x \mapsto \phi(x)e^{-\lambda x}$, $x \in \mathbb{R}$. Let u be a nontrivial solution of (1.1). We know that $u_{\beta/(1+\varepsilon)} \in L_1(\mathbb{R})$ and $k_{\beta/(1+\varepsilon)} \in L_\infty(\mathbb{R})$. Hence, $u_{\beta/(1+\varepsilon)}(x) \leq g'(0)u_{\beta/(1+\varepsilon)} * k_{\beta/(1+\varepsilon)}(x) \leq C$, or, in other words, $u(x) \leq Ce^{-(\beta/(1+\varepsilon))x}$ for all $x \in \mathbb{R}$. We rewrite (1.1) as an inhomogeneous linear convolution equation,

$$(6.1) \quad u(x) = g'(0)u * k(x) + r(x) \quad x \in \mathbb{R},$$

where $r(x) := (g \circ u - g'(0)u) * k(x)$. There exists a positive constant C such that $0 \geq r(x) \geq -Cu^{1+\varepsilon} * k(x) \geq -Ce^{-\beta x}$ for all $x \in \mathbb{R}$. Multiplying both sides of (6.1) with $e^{-\beta x}$ we obtain

$$(6.2) \quad u_\beta(x) = g'(0)u_\beta * k_\beta(x) + r_\beta(x) \quad x \in \mathbb{R}.$$

The function r_β is nonpositive and bounded. Consider the iterative scheme $v^{(0)}(x) := r_\beta(x)$, $v^{(n)}(x) := g'(0)v^{(n-1)} * k_\beta(x) + r_\beta(x)$, $n \in \mathbb{N}_+$. For each x , the sequence $\{v^{(n)}(x): n \in \mathbb{N}_+\}$ is monotone nonincreasing. Moreover, $v^{(n)}(x) \geq \inf\{r_\beta(x): x \in \mathbb{R}\} (1 + \alpha + \alpha^2 + \dots + \alpha^n)$, where $\alpha = g'(0)K(\beta) < 1$. So the sequence $\{v^{(n)}: n \in \mathbb{N}_+\}$ being bounded and monotone, converges to a limit function, v say, which satisfies the equation $v(x) = g'(0)v * k_\beta(x) + r_\beta(x)$ for all $x \in \mathbb{R}$. The same equation is satisfied by the function u_β , cf. (6.2). Since v is nonpositive and u_β is positive, the difference $u_\beta - v$ is not the zero function. Let the function w be defined by $w_\beta(x) := u_\beta(x) - v(x)$. Then w satisfies the homogeneous linear convolution equation

$$(6.3) \quad w(x) = g'(0)w^*k(x) \quad x \in \mathbb{R},$$

and the estimates $0 \leq w(x) \leq C(1+e^{\beta x})$ for all $x \in \mathbb{R}$. We claim that this implies that $w(x) = Ae^{\sigma x}$ for some $A > 0$.

In order to support this claim we first collect some information concerning the roots of the characteristic equation $g'(0)K(\lambda) = 1$ in the strip $S_\beta = \{\lambda \in \mathbb{C}: 0 \leq \operatorname{Re}\lambda \leq \beta\}$. In this strip, σ is the only real root; all other roots occur in complex conjugate pairs. The nonnegativity of k implies that the real part of any root other than σ is strictly less than σ . As in the proof of Theorem 6.2 it follows that there are only finitely many roots in S_β .

The function w , being a solution of (6.3) which satisfies the estimates $0 \leq w(x) \leq C(1+e^{\beta x})$, is a linear combination of exponential functions, possibly multiplied by polynomial functions, where the exponents are found as the roots of the characteristic equation in S_β , cf. [15, Section 11.2]. (It is in this step that the assumption $K(-\delta) < \infty$ is used.) Next, observe that any contribution from a pair of complex conjugate roots (with real parts less than σ) leads to an oscillatory behavior of w which is asymptotically dominant over $e^{\sigma x}$ as $x \rightarrow -\infty$. Hence, if there were a contribution to w from any complex conjugate pair of roots of the characteristic equation in the strip S_β , $w(x)$ would certainly become negative for some x sufficiently large negative, which is impossible because $w(x)$ is nonnegative, as we have seen. The conclusion is, therefore, that the only nonzero coefficient in the finite expansion for w is the one corresponding to σ . In other words, $w(x) = Ae^{\sigma x}$ for some $A > 0$, as claimed.

Finally, $u(x) = w(x) + v(x)e^{\beta x}$ and so $\lim_{x \rightarrow -\infty} u(x)e^{-\sigma x} = \lim_{x \rightarrow -\infty} w(x)e^{-\sigma x} = A$. \square

Our final results concern the uniqueness of nontrivial solutions of (1.1). We observe that the homogeneous equation (1.1) is invariant under translation, so uniqueness is to be understood in the sense of uniqueness modulo translation.

THEOREM 6.4. *Suppose that the conditions of Theorem 6.3 are satisfied and that, in addition, g is such that $|g(x)-g(y)| \leq g'(0)|x-y|$ for all pairs*

$x, y \in [0, p]$. Then there is at most one nontrivial solution (modulo translation) of (1.1).

PROOF. Let u_1^* and u_2 be two nontrivial solutions of (1.1). There exist positive constants C^* and C such that $\lim_{x \rightarrow -\infty} u_1^*(x)e^{-\sigma x} = C^*$ and $\lim_{x \rightarrow -\infty} u_2(x)e^{-\sigma x} = C$, according to Theorem 6.3. Define u_1 by putting $u_1(x) := u_1^*(x-\tau)$ for all $x \in \mathbb{R}$, where $\tau = \sigma^{-1} \ln(C^*/C)$. Then u_1 is also a nontrivial solution of (1.1) and $\lim_{x \rightarrow -\infty} u_1(x)e^{-\sigma x} = C$. Thus, if we consider the function v defined by $v(x) := (u_1 - u_2)(x)e^{-\sigma x}$ for all $x \in \mathbb{R}$, then v is continuous on \mathbb{R} , $\lim_{x \rightarrow \pm\infty} v(x) =: v(\pm\infty)$ exist, and $v(-\infty) = v(\infty) = 0$. Also, if $z := \sup\{|v(x)| : x \in \mathbb{R}\}$, then $z = v(x_0)$ for some $x_0 \in \mathbb{R}$. Now, $v(x) = (u_1 - u_2)(x)e^{-\sigma x} = (g \circ u_1 - g \circ u_2) * k(x)e^{-\sigma x}$, so since the Lipschitz constant of g on $[0, p]$ is at most equal to $g'(0)$ we have the inequality $|v(x)| \leq g'(0) |v| * k_\sigma(x)$, where $k_\sigma(x) := k(x)e^{-\sigma x}$, which is valid for all $x \in \mathbb{R}$. From this point on, the arguments parallel some of the arguments used in the proof of Lemma 2.2, cf. also FELLER [8, Section XI.2]. Upon induction we obtain the estimate

$$(6.4) \quad |v(x)| \leq (g'(0))^n |v| * k_\sigma^{n*}(x) \quad x \in \mathbb{R},$$

for any $n \in \mathbb{N}_+$. Now, the expression in the right member of this inequality is at most equal to $z(g'(0))^n \int_{\mathbb{R}} k_\sigma^{n*}(x) dx$ which, in turn, is equal to $z(g'(0) \int_{\mathbb{R}} k_\sigma(x) dx)^n = z(g'(0)K(\sigma))^n = z$. Hence, at $x = x_0$, the inequality (6.4) must be an equality, $|v(x_0)| = (g'(0))^n |v| * k_\sigma^{n*}(x_0)$ for all $n \in \mathbb{N}_+$. But this is only possible if $v(x_0 - y) = v(x_0)$ for all $y \in \Sigma$. (We recall that $\Sigma = \bigcup_{n=1}^{\infty} \Sigma_n$, Σ_n being the set consisting of the points of increase of k^{n*} .) It follows that $v(x) = v(x_0)$ for all $x \in \mathbb{R}$ if k is not concentrated on either \mathbb{R}_+ or \mathbb{R}_- , cf. Lemma 2.1. If k is concentrated on \mathbb{R}_+ , say, then it follows that $v(x) = v(x_0)$ for all $x \in (-\infty, x_0 - a]$ for some $a \in \mathbb{R}_+$, but as v satisfies the inequality (6.4) one may conclude again that $v(x) = v(x_0)$ for all $x \in \mathbb{R}$. Hence, v is a constant function on \mathbb{R} . Since its limiting values at $\pm\infty$ are zero, it follows that v is the zero function, i.e., $u_1(x) = u_2(x)$ for all $x \in \mathbb{R}$. \square

The proof of the following uniqueness theorem is analogous to the

proof of Theorem 6.4.

THEOREM 6.5. *Suppose that the conditions of Theorem 6.2 are satisfied and that, in addition, g is such that $|g(x)-g(y)| \leq g'(0)|x-y|$ for all pairs $x, y \in [0, p]$. Then there is at most one monotone nondecreasing nontrivial solution (modulo translation) of (1.1).*

REMARK 6.6. If g is monotone nondecreasing and sublinear (i.e., $g(\alpha x) \geq \alpha g(x)$ for $\alpha \in [0, 1]$, $x \in [0, p]$), then the Lipschitz constant of g on $[0, p]$ is at most $g'(0)$. The former properties were used by one of the authors in [5] to prove a uniqueness result. The fact that one can use the less restrictive condition involving the Lipschitz constant was observed by BARBOUR [3] for a function g which was explicitly given by an expression of the form $g(x) = \alpha(1-e^{-x})$.

REMARK 6.7. We observe that a combination of the existence result for monotone solutions (cf. [6], [16]) and the uniqueness result of Theorem 6.4 leads, in a very indirect way, to the conclusion that monotone functions g lead to monotone nontrivial solutions of (1.1) only. It would be of interest to have a direct proof of this fact.

7. APPLICATION TO A DETERMINISTIC MODEL FROM MATHEMATICAL EPIDEMIOLOGY

The spatio-temporal development of an epidemic among a closed population, with habitat \mathbb{R}^n , can be described by an integral equation of the form

$$(7.1) \quad u(t, x) = \int_0^{\infty} H(\tau) \int_{\mathbb{R}^n} g \circ u(t-\tau, y) V(x-y) dy d\tau$$

$-\infty < t < \infty$, $x \in \mathbb{R}^n$, cf. [6]. Here u is a measure for the density of susceptible individuals. The equation is based on various assumptions, the most important being

- (i) the members of the population can be categorized as either susceptible to or infected by the disease;
- (ii) the infectivity of an infected individual as a function of time

elapsed since exposure and position relative to the individual's own position is given by $H: \mathbb{R}_+ \rightarrow \mathbb{R}$ and $V: \mathbb{R}^n \rightarrow \mathbb{R}$, respectively; (iii) the disease induces permanent immunity, so an individual can pass from the class of susceptibles to the class of infectives, but not vice versa.

Both H and V are nonnegative, and normalized to have integral one. The assumption that the habitat \mathbb{R}^n is homogeneous and isotropic is reflected in the fact that V is a radial function. The nonlinearity of the equation comes about through the function g which has the form $g(x) = \alpha(1 - e^{-x})$. The parameter α has a threshold value 1, cf. [1], [6]; here, we restrict our attention to the more interesting case $\alpha > 1$. In the so-called traveling (plane) wave problem one looks for solutions of (7.1) of the form $u(t, x) = w(x \cdot v + ct)$, where v is a fixed unit vector. If one chooses a basis in \mathbb{R}^n such that $v = (1, 0, \dots, 0)$, then the function w must satisfy the following convolution equation on the line,

$$(7.2) \quad w(x) = (g \circ w) * V_c(x) \quad x \in \mathbb{R},$$

where $V_c(x) := \int_0^\infty H(\tau) \tilde{V}(x - c\tau) d\tau$, $x \in \mathbb{R}$, with $\tilde{V}(x_1) := \int_{\mathbb{R}^{n-1}} V(x_1, x_2, \dots, x_n) dx_2 \dots dx_n$, $x_1 \in \mathbb{R}$. If $c = 0$ the function V_c is symmetric. One can then apply Theorem 4.1, from which it follows that (7.1) does not admit any standing wave solution. Next, we consider the case $c \neq 0$. Because of the symmetry of V we can restrict our attention to the case $c > 0$.

As c increases, the mass of V_c shifts to the right. This can be seen by inspecting the qualitative behavior of the characteristic function L_c , associated with the linearized equation, as a function of the wave velocity c . By definition,

$$\begin{aligned} L_c(\lambda) &:= g'(0) \int_{\mathbb{R}} e^{-\lambda y} V_c(y) dy \\ &= g'(0) \int_{\mathbb{R}_+} e^{-\lambda c \tau} H(\tau) d\tau \int_{\mathbb{R}} e^{-\lambda y} \tilde{V}(y) dy. \end{aligned}$$

The nonnegativity of H and V implies that, for c fixed, L_c is a convex function of λ ($\lambda \in \mathbb{R}$) and, for λ fixed ($\lambda > 0$), it is a monotonically decreasing function of c . Furthermore, $L_c(0) = g'(0) = \alpha > 1$. Hence, the number $c_0 := \inf\{c > 0: \text{there exists } \lambda > 0 \text{ such that } L_c(\lambda) = 1\}$ is well-defined if H and V satisfy the appropriate hypotheses [6].

Various authors have given constructive proofs for the existence of (monotone) nontrivial solution of (7.2) in the case $c \geq c_0$, cf. [2], [4], [6], [16]. From the results of our investigation it follows that there exists exactly one such solution (modulo translation) at each speed $c > c_0$, at least if $\tilde{V}_\lambda: x \mapsto \tilde{V}(x)e^{-\lambda x}$ is bounded for the value of λ specified in Theorem 6.4. (If, for instance, V has compact support then this condition is certainly fulfilled.) A similar conclusion has been reached at by BARBOUR [3] who used probabilistic arguments.

Furthermore, it follows from the results of our investigation that no traveling waves exist with speed $c < c_0$. The same result has been obtained for the special case $H(t) = ae^{-\mu t}$ (a, μ nonnegative constants) by ATKINSON and REUTER [2] and by ARONSON [1], and for a discrete time model in population genetics by WEINBERGER [16]. Both Aronson and Weinberger based their proofs on the construction of a subsolution. This approach made it actually possible for them to conclude that c_0 is the asymptotic speed of propagation of disturbances from a rest state of the associated initial value problem (see also MOLLISON [12]).

In our opinion, the main advantage of the approach presented in this paper is that the same chain of purely analytical arguments leads to the nonexistence, as well as the uniqueness results.

ACKNOWLEDGMENT

Part of this investigation was carried out while the second author was on foreign assignment to the Mathematisch Centrum, Amsterdam (Neth.). The final draft of this paper was written while the first author was visiting the Applied Mathematics Division at Argonne National Laboratory. Both authors wish to thank their respective host institutions for their hospitality and financial support.

REFERENCES

- [1] ARONSON, D.G., *The asymptotic speed of propagation of a simple epidemic*, to appear in: *Nonlinear Diffusion*, W.E. Fitzgibbon & H.F. Walker, eds., *Research Notes in Mathematics*, Pitman Publishing Co., 1977.
- [2] ATKINSON, C. & G.E.H. REUTER, *Deterministic epidemic waves*, *Math. Proc. Camb. Phil. Soc.* 80 (1976) 315-330.
- [3] BARBOUR, A.D., *The uniqueness of Atkinson and Reuter's epidemic waves*, *Math. Proc. Camb. Phil. Soc.* 82 (1977) 127-130.
- [4] BROWN, K.J. & J. CARR, *Deterministic epidemic waves of critical velocity*, *Math. Proc. Camb. Phil. Soc.* 81 (1977) 431-435.
- [5] DIEKMANN, O., *Limiting behaviour in an epidemic model*, *Nonlinear Analysis, Theory, Methods & Applications* 1 (1977) 459-470.
- [6] DIEKMANN, O., *Thresholds and travelling waves for the geographical spread of infection*, *Mathematical Centre Report TW 166*, Amsterdam, 1977.
- [7] ESSÉN, M., *Studies on a convolution inequality*, *Ark. Mat.* 5 (1963) 113-152.
- [8] FELLER, W., *An Introduction to Probability Theory and Its Applications*, Vol. II, Wiley, New York, 1966.
- [9] KARLIN, S., *On the renewal equation*, *Pacific J. Math.* 5 (1955) 229-257.
- [10] KENDALL, D.G., *Mathematical models of the spread of infection*, in: *Mathematics and Computer Science in Biology and Medicine*, Medical Research Council, London, 1965, 213-224.
- [11] KERMACK, W.O. & A.G. MCKENDRICK, *A contribution to the mathematical theory of epidemics*, *Proc. Roy. Soc. A* 115 (1927) 700-721.
- [12] MOLLISON, D., *The rate of spatial propagation of simple epidemics*, in: *Proc. Sixth Berkeley Symposium on Mathematical Statistics and Probability*, Vol. III, L.M. le Cam, J. Neyman & E.L. Scott, eds., Univ. of California Press, Berkeley, 1972, 579-614.

- [13] RUDIN, W., *Fourier Analysis on Groups*, Interscience, New York, 1962.
- [14] RUDIN, W., *Functional Analysis*, McGraw-Hill, New York, 1973.
- [15] TITCHMARSCH, E.C., *Introduction to the Theory of Fourier Integrals*, Clarendon Press, Oxford, 1937.
- [16] WEINBERGER, H.F., *Asymptotic behavior of a model in population genetics*, to appear in: *Indiana University Seminar in Applied Mathematics*, J. Chadam, ed., Springer Lecture Notes.
- [17] WIDDER, D.V., *The Laplace Transform*, University Press, Princeton, 1946.