



Centrum voor Wiskunde en Informatica
REPORT*RAPPORT*

A Predicate Transformer for Unification

L. Colussi, E. Marchiori

Computer Science/Department of Software Technology

CS-R9227 1992

A Predicate Transformer for Unification

Livio Colussi

Dipartimento di Matematica Pura ed Applicata
Università di Padova, Via Belzoni 7, 35131 Padova, Italy

Elena Marchiori

CWI

P.O. Box 4079, 1009 AB Amsterdam, The Netherlands

and

Dipartimento di Matematica Pura ed Applicata
Università di Padova, Via Belzoni 7, 35131 Padova, Italy

Abstract

In this paper we study unification as predicate transformer. Given a unification problem expressed as a set of sets of terms \mathcal{U} and a predicate P , we are interested in the strongest predicate R (w.r.t. the implication) s.t. if P holds before the unification of \mathcal{U} then R holds when the unification is performed. We introduce a Dijkstra-style calculus that given P and \mathcal{U} computes R . We prove the soundness, completeness and termination of the calculus. The predicate language considered contains monotonic predicates together with some non-monotonic predicates like *var*, \neg *ground*, *share* and \neg *share*. This allows to use the calculus for the static analysis of run-time properties of Prolog programs.

1985 Mathematics Subject Classification: 68Q40, 68Q60, 68T15.

CR Categories: F.3.1., F.4.1, I.2.3.

Keywords and Phrases: Unification, strongest postcondition, Prolog programs, forward semantics.

Note: With minor variations, this paper will appear in the Proceedings of the 1992 Joint International Conference and Symposium on Logic Programming, Washington, D.C., U.S.A., The MIT Press.

1 Introduction

The standard view of logic programming is declarative, i.e. a program describes some predicate or function without referring to the way it will be computed. Nevertheless computational aspects become fundamental for the study of run-time properties of Prolog programs, like the actual form of the arguments of a goal before and after its call. In Prolog unification is the main computational mechanism since it produces the value of the variables during the execution of a goal in a program. To study its effect on the values of variables we study unification by means of predicate transformers. The use of predicate transformers for semantic analysis has been studied in the setting of imperative programming: it was advocated by Floyd [5] and by Dijkstra [3] for program verification. The use of predicate transformers in the framework of logic programming is new. Given a unification problem expressed by a set of sets of terms \mathcal{U} , we introduce the predicate transformer $sp.\mathcal{U}$ such that $sp.\mathcal{U}.P$ is semantically equivalent to the

strongest predicate R (w.r.t. implication) s.t. if P holds before the unification of \mathcal{U} , then R holds when the unification is performed. We show that $sp\mathcal{U}.P$ could be computed in one step if P were a monotonic predicate. Since our aim is to infer run-time properties of Prolog programs, then the predicate language considered contains also non-monotonic predicates like *var* or *share*. For this reason a careful analysis of some intermediate steps of the unification process is necessary. This yields to a non-trivial system of syntactic rules to compute $sp\mathcal{U}.P$. The soundness, completeness and termination of the system is proved. The calculus can be used to infer run-time properties of logic programs. In Cousot and Cousot's original paper on abstract interpretation of imperative programs [2] everything was couched in terms of predicate transformers. Predicate transformers were used to define deductive semantics. Deductive semantics was used to design approximate program analysis frameworks. To propose a similar approach for logic programs we need the correspondent of program point for a logic program. In [7] Nilsson introduced a scheme for inferring run-time properties of logic programs based on a semantic description of logic programs that uses the concept of program point. We will show that the predicate transformer sp can be easily cast in such a theory.

The rest of the paper is organized as follows. The next section contains some preliminaries and introduces the predicate transformer $sp\mathcal{U}$. Section 3 introduces the transformation rules to compute $sp\mathcal{U}.P$. In section 4 the soundness, completeness and termination of the calculus are proved. In section 5 we illustrate the use of the calculus for defining a forward semantics of Prolog programs.

2 Unification as Predicate Transformer

The computational meaning of unification in Prolog relies on the concept of substitution. A *substitution* is a mapping from variables to terms such that $dom(\vartheta) \stackrel{\text{def}}{=} \{v \mid v\vartheta \neq v\}$ is finite. The notion of unification can be given w.r.t. a set of sets of terms [4] or w.r.t. a set of equations [6]. We choose the first approach. Let \mathcal{U} be a finite set of sets of terms. A *unifier* for \mathcal{U} is a substitution ϑ such that every set in \mathcal{U} , under the application of ϑ , becomes a singleton, i.e. $\forall S \in \mathcal{U} \forall t, t' \in S (t\vartheta = t'\vartheta)$. A *most general unifier* for \mathcal{U} is a unifier ϑ such that for every unifier σ there exists a substitution γ such that $\vartheta\gamma = \sigma$. The *set of idempotent most general unifiers* for \mathcal{U} will be denoted by $mgu(\mathcal{U})$. The operational meaning of \mathcal{U} can be described as the partial function $\lambda\alpha.\alpha\mu$, where α is a substitution and μ is a fixed mgu in $mgu(\mathcal{U}\alpha)$; clearly $\lambda\alpha.\alpha\mu$ is undefined if $mgu(\mathcal{U}\alpha) = \emptyset$. We study unification by means of the predicate transformer $sp\mathcal{U}$ (where sp stands for strongest postcondition [5]) with the following operational meaning.

Definition 2.1 $sp\mathcal{U}.P$ is true in precisely those substitutions $\alpha\mu$ such that $P\alpha$ is true and $\mu \in mgu(\mathcal{U}\alpha)$.

The choice to represent the unification process as set of sets of terms is motivated by the following observations:

$$mgu(\{\{f(t_1, \dots, t_n), f(s_1, \dots, s_n)\}\}) = mgu(\{\{t_1, s_1\}, \dots, \{t_n, s_n\}\}) \text{ and} \\ mgu(\{S_1, \dots, S_n\}) = mgu(\{S_1 \cup S_2, S_3, \dots, S_n\}) \text{ if } S_1 \cap S_2 \neq \emptyset.$$

These two equalities will be used in our calculus for $sp\mathcal{U}$ and they clearly lead to consider sets of sets of terms. For sake of clarity, we use double square brackets to enclose sets of terms $S = \llbracket t_1, \dots, t_m \rrbracket$ and braces to enclose sets of sets of terms $\mathcal{U} = \{S_1, \dots, S_n\}$.

We call a predicate P *monotonic* if it is (semantically) invariant under instantiation, that is for all substitutions α, β if $P\alpha$ is true then $P\alpha\beta$ is true. Now let \mathcal{U} be

$\{\llbracket t_1^1, \dots, t_{n_1}^1 \rrbracket, \dots, \llbracket t_1^m, \dots, t_{n_m}^m \rrbracket\}$: we denote by U the predicate $((t_1^1 = \dots = t_{n_1}^1) \wedge \dots \wedge (t_1^m = \dots = t_{n_m}^m))$. Then the following lemma holds.

Lemma 2.2 *Let P be a monotonic predicate. Then $P \wedge U$ is equivalent to $sp\mathcal{U}.P$.*

Proof. Let α be s.t. $P\alpha$ is true and let $\mu \in mgu(\mathcal{U}\alpha)$. Then $U\alpha\mu$ is true and from P monotonic it follows that $P\alpha\mu$ is true.

Viceversa let α be s.t. $(P \wedge U)\alpha$ is true. Then $P\alpha$ is true and $\epsilon \in mgu(\mathcal{U}\alpha)$. So by Definition 2.1 $(sp\mathcal{U}.P)\alpha$ is true. \square

Lemma 2.2 allows to compute $sp\mathcal{U}.P$ when P is a monotonic predicate.

2.1 The Language

However we are interested also in properties that describe the structure of terms, like *var* or \neg *ground*, since we want to use the predicate transformer to infer run-time properties of logic programs. Thus we introduce the language \mathcal{A} defined on the alphabet containing the following classes of symbols:

- a countable set VAR of *variables*;
- a set FUN of *functions*;
- a set $PRED = Pred \cup \{free, var, \neg ground, share, \neg share, inst\}$ of *predicate* symbols where $Pred$ is a finite set of monotonic predicate symbols s.t. $=, ground, \neg var, \prec, \preceq, invar$ are in $Pred$;
- the *connectives* \wedge and \vee ;
- the *existential quantifier* \exists ;
- (and) as punctuation symbols.

Variables will be normally denoted by the letters u, v, w, x, y, z (possibly subscripted or superscripted) and functions will be normally denoted by the letters f, g, h (possibly subscripted). Let $TERM$ be the set of *terms* built on FUN and VAR . Terms will be normally denoted by the letters r, s, t (possibly subscripted or superscripted). Given a term t , the set $vars(t) \subseteq VAR$ denotes the set of variables that occur in t . We call *structured term* a term of the form $f(t_1, \dots, t_m)$, where $m \geq 1$; we call *proper subterm* of t every subterm of t but t . We assume that sequences are contained in \mathcal{A} . We denote by \underline{t} a sequence t_1, \dots, t_k and we write $\underline{t}_{(k)}$ or $\langle t_1, \dots, t_k \rangle$ if respectively the size or the elements of the sequence are relevant. Moreover we indicate with $\underline{x}\rho$ the sequence of terms obtained applying the substitution ρ to every element of the sequence \underline{x} . We call *atom* a predicate of the form $p(t_1, \dots, t_n)$ where p is a predicate symbol of arity n and t_1, \dots, t_n are terms. When ambiguity does not arise we write $r(t_1, \dots, t_m)$ as a shorthand for the predicate $r(t_1) \wedge \dots \wedge r(t_m)$, where r is a predicate symbol of arity 1.

The *truth value* of a predicate $P \in \mathcal{A}$ w.r.t. a substitution α s.t. $vars(P) \subseteq dom(\alpha)$ is defined inductively on the structure of P , and the meaning of an atom is specified as follows:

- $\neg var(t)\alpha$ is true iff $t\alpha \notin VAR$;
- $ground(t)\alpha$ is true iff $vars(t\alpha) = \emptyset$;
- $(t_1 = t_2)\alpha$ is true iff $t_1\alpha = t_2\alpha$ syntactically;
- $(s \preceq t)\alpha$ is true iff $s\alpha$ is a subterm of $t\alpha$;
- $(s \prec t)\alpha$ is true iff $s\alpha$ is a proper subterm of $t\alpha$;
- $invar(s, t)\alpha$ is true iff $vars(s\alpha) \subseteq vars(t\alpha)$;
- $free(x)\alpha$ is true iff $x\alpha \in VAR$ and $x\alpha \notin vars(y\alpha)$ for all $y \in dom(\alpha)$ s.t. $y \neq x$;

- $var(x)\alpha$ is true iff $x\alpha \in VAR$;
- $\neg ground(t)\alpha$ is true iff $vars(t\alpha) \neq \emptyset$;
- $share(s, t)\alpha$ is true iff $vars(s\alpha) \cap vars(t\alpha) \neq \emptyset$;
- $\neg share(s, t)\alpha$ is true iff $vars(s\alpha) \cap vars(t\alpha) = \emptyset$;
- $inst(x, r_1, r_2, y)\alpha$ is true iff $r_1\alpha$ is the sequence $\langle x_1, \dots, x_m \rangle$, with $x_i \in vars(x\alpha)$ and $x_i \notin vars(y\alpha)$ for $i \in [1, m]$, $r_2\alpha$ is the sequence $\langle t_1, \dots, t_m \rangle$ and $\{x_1/t_1, \dots, x_m/t_m\} \in mgu(\{\llbracket x\alpha, y\alpha \rrbracket\})$.

Notice that x and y in $inst(x, r_1, r_2, y)$ represent two terms the second of which is an instance of the first. Thus the predicate $inst$ expresses a special case of the unification.

Given two predicates P and Q , we write $P \equiv Q$ to indicate that P and Q are semantically equivalent. We can assume that the predicates $TRUE$ (the predicate true w.r.t. all substitutions) and $FALSE$ (the predicate false w.r.t. all substitutions) are in \mathcal{A} , since $TRUE \equiv (var(x) \vee \neg var(x))$ and $FALSE \equiv (var(x) \wedge \neg var(x))$.

Predicates in \mathcal{A} are not in general monotonic, since all atoms built on predicate symbols not in $Pred$ are non-monotonic by definition. So Lemma 2.2 is not sufficient to characterize $sp\mathcal{U}$: consider for instance the unification $\{\llbracket x, a \rrbracket\}$ and the predicate $var(x)$. Thus a careful analysis of the effect of the unification process on non-monotonic predicates is necessary. The fact that the connective \neg is not in our language guarantees that atoms built on predicate symbols not in $Pred$ are the only non-monotonic atoms of the language; this allows a case analysis of the effect of unification on non-monotonic predicates.

2.2 Some Useful Assumptions

The following assumptions will be used to (relatively) simplify the form of the rules for $sp\mathcal{U}$ that will be introduced in the next section.

Assumption 1 Predicates are of the form $\exists \underline{x}P$ where P doesn't contain any quantifier, it is in disjunctive normal form (i.e. it is a disjunction of conjunctions of atoms) and the equalities that occur in each conjunct are denoted by a set of equations in solved form.

This assumption is not restrictive, since it is well known that a predicate can be transformed in an equivalent one satisfying the first two conditions, while the third condition can be obtained applying for instance the algorithm in [6] to compute the solved form of a set of equations, otherwise the predicate is equivalent to $FALSE$.

Assumption 2 Atoms with predicate symbol $free, var, \neg var, ground, \neg ground, share, \neg share, invar$ have variables as arguments.

This assumption is not restrictive, since we can transform a predicate of \mathcal{A} in an equivalent one satisfying this property, by repeatedly applying the following equivalences:

- $free(f(\underline{t}_{(m)})) \equiv FALSE$,
- $var(f(\underline{t}_{(m)})) \equiv FALSE$,
- $\neg var(f(\underline{t}_{(m)})) \equiv TRUE$,
- $ground(f(\underline{t}_{(m)})) \equiv (ground(t_1) \wedge \dots \wedge ground(t_m))$,
- $\neg ground(f(\underline{t}_{(m)})) \equiv (\neg ground(t_1) \vee \dots \vee \neg ground(t_m))$,
- $share(f(\underline{t}_{(m)}), t) \equiv (share(t_1, t) \vee \dots \vee share(t_m, t))$,
 $share(x, f(\underline{t}_{(m)})) \equiv (share(x, t_1) \vee \dots \vee share(x, t_m))$,
- $\neg share(f(\underline{t}_{(m)}), t) \equiv (\neg share(t_1, t) \wedge \dots \wedge \neg share(t_m, t))$,
 $\neg share(x, f(\underline{t}_{(m)})) \equiv (\neg share(x, t_1) \wedge \dots \wedge \neg share(x, t_m))$,

$$\begin{aligned}
- \text{invar}(f(\underline{t}_{(m)}), t) &\equiv (\text{invar}(t_1, t) \wedge \dots \wedge \text{invar}(t_m, t)), \\
\text{invar}(x, f(\underline{t}_{(m)})) &\equiv (\text{invar}(x, t_1) \vee \dots \vee \text{invar}(x, t_m)).
\end{aligned}$$

Assumption 3 For any formula $sp\mathcal{U}.P$ the predicate P does not contain (existential) quantifiers.

This assumption is not restrictive as the following lemma shows.

Lemma 2.3 *If the variable x does not occur in \mathcal{U} then $sp\mathcal{U}.\exists xP$ is equivalent to $\exists x(sp\mathcal{U}.P)$ w.r.t. Definition 2.1.*

Proof. Since x doesn't occur in \mathcal{U} then the truth value of $\exists x(sp\mathcal{U}.P)\beta$ and of $(sp\mathcal{U}.\exists xP)\beta$ does not depend on $x\beta$. Thus we can assume without loss of generality $x \notin \text{dom}(\beta)$. Then $(sp\mathcal{U}.\exists xP)\beta$ is true iff there exist α and μ s.t. $x \notin \text{dom}(\alpha)$, $x \notin \text{dom}(\mu)$, $\mu \in \text{mgu}(\mathcal{U}\alpha)$, $(\exists xP)\alpha$ is true and $\beta = \alpha\mu$ iff there exist α , μ and t s.t. $x \notin \text{dom}(\alpha)$, $x \notin \text{dom}(\mu)$, $\mu \in \text{mgu}(\mathcal{U}\alpha)$, $P(\alpha \cup \{x/t\})$ is true and $\beta = \alpha\mu$ iff there exist α , μ and t s.t. $\mu \in \text{mgu}(\mathcal{U}(\alpha \cup \{x/t\}))$, $P(\alpha \cup \{x/t\})$ is true and $(\beta \cup \{x/t\}) = (\alpha \cup \{x/t\})\mu$ iff $(sp\mathcal{U}.P)(\beta \cup \{x/t\})$ is true iff $(\exists xsp\mathcal{U}.P)\beta$ is true. \square

3 A Calculus for $sp\mathcal{U}$

The following conditions on P and \mathcal{U} characterize the types of formulas which will specify the scope of applicability of the rules for $sp\mathcal{U}.P$.

- (i) P is a conjunction of atoms.
- (ii) For each equation $x = t$ in P , x does not occur in \mathcal{U} .
- (iii) For every x occurring in \mathcal{U} either $\text{var}(x)$ or $\neg\text{var}(x)$ occurs in P .
- (iv) For all distinct variables x occurring in \mathcal{U} and y occurring in P either $\text{share}(x, y)$ or $\neg\text{share}(x, y)$ occurs in P .
- (v) $\mathcal{U} = \{S_1, \dots, S_n\}$ contains disjoint sets, i.e. $S_i \cap S_j = \emptyset$ for $i \neq j$.
- (vi) Each set in \mathcal{U} contains more than one element.
- (vii) Each set in \mathcal{U} contains at most one structured element $f(v_1, \dots, v_m)$ and in such a case $\text{free}(v_1), \dots, \text{free}(v_m)$ occur in P .
- (viii) Every element x of a set $S \in \mathcal{U}$ is s.t. $\text{free}(x)$ occurs in P if x occurs in the structured element of another set in \mathcal{U} and $\neg\text{var}(x)$ occurs in P otherwise. Moreover, each set that contains a structured element also contains an element y s.t. $\text{free}(y)$ occurs in P . (Hence y occurs in the structured element of another set).

We introduce 3 types of formulas $sp\mathcal{U}.P$ as follows.

type 1: those which satisfy conditions (i)–(iii).

type 2: those which satisfy conditions (i)–(vii).

type 3: those which satisfy conditions (i)–(viii).

Each type of formula characterizes a simpler form of P and \mathcal{U} . The final form will be a disjunction of formulas in the so called *reduced form*.

A formula $sp\mathcal{U}.P$ is in **reduced form** if P is a conjunction of atoms, for each equation $x = t$ in P x does not occur in \mathcal{U} , \mathcal{U} contains only disjoint sets of two or more variables, for all x occurring in \mathcal{U} both $\neg\text{var}(x)$ and $\neg\text{ground}(x)$ occur in P and for all x occurring in \mathcal{U} and y occurring in P either $\text{share}(x, y)$ or $\neg\text{share}(x, y)$ occurs in P .

We are now ready to present the rules for $sp.\mathcal{U}.P$. The notation E_t^x will be used to indicate the formula obtained by replacing the occurrences of x in E with t .

- If $P = P_1 \vee \dots \vee P_n$ then

$$sp.\mathcal{U}.P \equiv sp.\mathcal{U}.P_1 \vee \dots \vee sp.\mathcal{U}.P_n \quad \text{OR}$$

- If x occurs in \mathcal{U} and neither $var(x)$ nor $\neg var(x)$ occurs in P then

$$sp.\mathcal{U}.P \equiv sp.\mathcal{U}.(P \wedge var(x)) \vee sp.\mathcal{U}.(P \wedge \neg var(x)) \quad \text{VAR1}$$

- If P is a conjunction of atoms and $x = t$ occurs in P then:

$$sp.\mathcal{U}.P \equiv sp.\mathcal{U}_t^x.P \quad \text{EQ}$$

- $sp.\mathcal{U}.FALSE \equiv FALSE$

F

The following eight rules may be applied only to **type 1** formulas.

- If x occurs in \mathcal{U} and y occurs in P and neither $share(x, y)$ nor $\neg share(x, y)$ occurs in P then

$$sp.\mathcal{U}.P \equiv sp.\mathcal{U}.(P \wedge share(x, y)) \vee sp.\mathcal{U}.(P \wedge \neg share(x, y)) \quad \text{SH1}$$

- If $\mathcal{U} = \{\llbracket f_1(\underline{s}), f_2(\underline{t}), \dots \rrbracket, S_2, \dots, S_n\}$ and $f_1 \neq f_2$, then

$$sp.\mathcal{U}.P \equiv FALSE \quad \text{MIS1}$$

- If $\mathcal{U} = \{\llbracket x, s, \underline{t} \rrbracket, S_2, \dots, S_n\}$ and either $x \in vars(s)$ or the conjunct $x \prec s$ occurs in P then

$$sp.\mathcal{U}.P \equiv FALSE \quad \text{MIS2}$$

- If $\mathcal{U} = \{\llbracket f(\underline{s}_{(k)}^1), \dots, f(\underline{s}_{(k)}^m) \rrbracket, S_2, \dots, S_n\}$ then

$$sp.\mathcal{U}.P \equiv sp.\mathcal{U}'.P \quad \text{STR1}$$

where $\mathcal{U}' = \{\llbracket \underline{s}_j^{(m)} \rrbracket_{j \in [1, k]}, S_2, \dots, S_n\}$

- If $\mathcal{U} = \{\llbracket f(\underline{s}_{(k)}^1), \dots, f(\underline{s}_{(k)}^i), x_{i+1}, \dots, x_m \rrbracket, S_2, \dots, S_n\}$ with $i < m$ and either $i \geq 2$ or at least one s_j^i is not a variable or at least one s_j^i is a variable such that $\neg var(s_j^i)$ occurs in P , then

$$sp.\mathcal{U}.P \equiv \exists \underline{y}_{(k)} (sp.\mathcal{U}'.P') \quad \text{STR2}$$

where $\mathcal{U}' = \{\llbracket f(\underline{y}_{(k)}), x_{i+1}, \dots, x_m \rrbracket, \llbracket y_j, \underline{s}_j^{(i)} \rrbracket_{j \in [1, k]}, S_2, \dots, S_n\}$,

$P' = P \wedge free(\underline{y}_{(k)})$ and $\underline{y}_{(k)}$ are fresh variables.

- If $\mathcal{U} = \{\llbracket t, \underline{t}_{(m)}, \underline{s}_{(m')} \rrbracket, S_3, \dots, S_n\}$ then

$$sp.\mathcal{U}.P \equiv sp.\mathcal{U}'.P \quad \text{SH2}$$

where $\mathcal{U}' = \{\llbracket t, \underline{t}_{(m)}, \underline{s}_{(m')} \rrbracket, S_3, \dots, S_n\}$

- If $\mathcal{U} = \{\llbracket t \rrbracket, S_2, \dots, S_n\}$ then

$$sp.\mathcal{U}.P \equiv sp.\mathcal{U}'.P \quad \text{SI}$$

where $\mathcal{U}' = \{S_2, \dots, S_n\}$

The following two rules may be applied only to **type 2** formulas.

- If $\mathcal{U} = \{\llbracket t, \underline{x}_{(m)} \rrbracket, S_2, \dots, S_n\}$ where x_m does not occur in the structured term of any set of \mathcal{U} , $\text{var}(x_m)$ and $\neg \text{share}(x_m, y)$ occurs in P for all $y \in \text{vars}(t)$, then

$$\text{sp.}\mathcal{U}.P \equiv \exists \underline{z}' \text{ sp.}\mathcal{U}'.R \quad \text{VAR2}$$

where $\mathcal{U}' = \{\llbracket t, \underline{x}_{(m-1)} \rrbracket, S_2, \dots, S_n\}$,

$R = (\bigwedge_{z \in \underline{z}} \text{inst}(z\rho, \langle x_m\rho \rangle, \langle t \rangle, z) \wedge P' \wedge x_m = t)$,

$\underline{z} = \langle z \in \text{vars}(P) \mid P \Rightarrow \text{share}(z, x_m) \rangle$, $\underline{z}' = \underline{z}\rho$ is a variant of \underline{z} disjoint from P and $P' = P_{\underline{z}'}^{\underline{z}}$.

- If $\mathcal{U} = \{\llbracket f(\underline{s}_{(k)}), \underline{x}_{(m)} \rrbracket, S_2, \dots, S_n\}$ and $\neg \text{var}(x_1), \dots, \neg \text{var}(x_m)$ occur in P then

$$\text{sp.}\mathcal{U}.P \equiv \exists \underline{y} \text{ sp.}\mathcal{U}'.(P \wedge x_1 = f(\underline{y}_{(k)}^1) \wedge \dots \wedge x_m = f(\underline{y}_{(k)}^m)) \quad \text{VAR3}$$

where $\mathcal{U}' = \{\llbracket s_i, \underline{y}_i^{(m)} \rrbracket_{i \in [1, k]}, S_2, \dots, S_n\}$ and \underline{y} is the sequence $\underline{y}_{(k)}^1, \dots, \underline{y}_{(k)}^m$ of fresh variables.

The following three rules may be applied only to **type 3** formulas.

- If there is a set $S \in \mathcal{U}$ that contains a structured term then

$$\text{sp.}\mathcal{U}.P \equiv \text{FALSE} \quad \text{MIS3}$$

- If x occurs in \mathcal{U} and neither $\text{ground}(x)$ nor $\neg \text{ground}(x)$ occurs in P then

$$\text{sp.}\mathcal{U}.P \equiv \text{sp.}\mathcal{U}.(P \wedge \text{ground}(x)) \vee \text{sp.}\mathcal{U}.(P \wedge \neg \text{ground}(x)) \quad \text{GR1}$$

- If $\mathcal{U} = \{\llbracket \underline{x}_{(m)} \rrbracket, S_2, \dots, S_n\}$ and $\text{ground}(x_m)$ occurs in P then

$$\text{sp.}\mathcal{U}.P \equiv \exists \underline{x}', z_{\underline{x}}, y_{\underline{x}} \text{ sp.}\mathcal{U}'.R \quad \text{GR2}$$

where $\mathcal{U}' = \{S_2, \dots, S_n\}$,

$\underline{x} = \langle x \in \text{vars}(P) \mid P \Rightarrow \text{share}(x, x_i) \text{ for some } i \in [1, m-1] \rangle$, $\underline{x}' = \underline{x}\rho$ is a variant of \underline{x} disjoint from P , $z_{\underline{x}}$ and $y_{\underline{x}}$ are the sequences of fresh variables z_x and y_x with $x \in \underline{x}$, $P' = P_{\underline{x}'}^{\underline{x}}$ and R is the predicate

$$\bigwedge_{x \in \underline{x}} (\text{inst}(x\rho, z_x, y_x, x) \wedge \bigwedge_{i \mid P \Rightarrow \text{share}(x, x_i)} \text{invar}(z_x, z_{x_i}) \wedge \bigwedge_{y \in \text{vars}(P)} \neg \text{share}(z_x, y)) \wedge P' \wedge x_1 = \dots = x_m.$$

To a formula in **reduced form** we can apply the following rule.

- If $\text{sp.}\mathcal{U}.P$ is in reduced form, where $\mathcal{U} = \{\llbracket \underline{x}_{(m_1)}^1 \rrbracket, \dots, \llbracket \underline{x}_{(m_n)}^n \rrbracket\}$, then

$$\text{sp.}\mathcal{U}.P \equiv \exists \underline{x}', z_{\underline{x}}, y_{\underline{x}} (R \wedge U) \quad \text{RF}$$

where U is the predicate $(x_1^1 = \dots = x_{m_1}^1) \wedge \dots \wedge (x_1^n = \dots = x_{m_n}^n)$,

$\underline{x} = \langle x \in \text{vars}(P) \mid P \Rightarrow \text{share}(x, x_i^j) \text{ for some } i \in [1, m_j], j \in [1, n] \rangle$, $\underline{x}' = \underline{x}\rho$ is a variant of \underline{x} disjoint from P , $z_{\underline{x}}$ and $y_{\underline{x}}$ are the sequences of fresh variables z_x and y_x with $x \in \underline{x}$, $P' = P_{\underline{x}'}^{\underline{x}}$ and $R = (\bigwedge_{x \in \underline{x}} \text{inst}(x\rho, z_x, y_x, x) \wedge P')$.

The previous rules are natural abstractions of the relative unification step except rules **MIS3**, **VAR2**, **GR2** and **RF**. Rule **MIS3** relies on the condition that the formula is of type 3 and \mathcal{U} contains at least a set with a structured element. In this case it can be proven that \mathcal{U} has no unifier.

Rules **VAR2**, **GR2** and **RF** take into account how sharing among variables can propagate the bindings produced by the considered transformation and how the transformations affect the truth of the non-monotonic atoms. To keep track of the way the predicate is modified suitable variables are renamed with fresh variables existentially quantified and suitable predicates are introduced to specify the link among the original variables and the renamed ones.

All the rules are syntactic. Thus the set of rules provides a (nondeterministic) algorithm. We will see in the following section that this algorithm terminates and computes $sp\mathcal{U}.P$. We conclude this section with some examples.

Example 3.1 Let $P = free(x, y)$ and $\mathcal{U} = \{\llbracket f(x), y \rrbracket, \llbracket g(y), x \rrbracket\}$. Since $sp\mathcal{U}.P$ is of **type 3**, then by rule **MIS3** it is equivalent to $FALSE$. In fact an occur check does occur.

Example 3.2 Let $P = (free(x, y) \wedge \neg share(x, y))$ and $\mathcal{U} = \{\llbracket f(y), x \rrbracket\}$. Since $sp\mathcal{U}.P$ is of **type 2**, then we can apply rule **VAR2**. We obtain

$$\exists x' (sp.\{\llbracket f(y) \rrbracket\}.(P_x^x \wedge inst(x', \langle x' \rangle, \langle f(y) \rangle, x) \wedge x = f(y))).$$

By rules **SI** and **RF** we obtain

$$\exists x' (P_x^x \wedge inst(x', \langle x' \rangle, \langle f(y) \rangle, x) \wedge x = f(y)),$$

which is equivalent to $(free(y) \wedge x = f(y))$.

Example 3.3 Let $P = (ground(y) \wedge \neg var(x) \wedge \neg ground(x) \wedge \neg share(x, y))$ and $\mathcal{U} = \{\llbracket x, y \rrbracket\}$. Since $sp\mathcal{U}.P$ is of **type 3**, then we can apply rule **GR2**. We obtain

$$\exists x', z_x, y_x (sp.\{\}.(P_{x'}^x \wedge inst(x', z_x, y_x, x) \wedge invar(z_x, z_x) \wedge \neg share(z_x, x) \wedge \neg share(z_x, y) \wedge x = y)).$$

By rule **RF** we obtain

$$\exists x', z_x, y_x (P_x^x \wedge inst(x', z_x, y_x, x) \wedge invar(z_x, z_x) \wedge \neg share(z_x, x) \wedge \neg share(z_x, y) \wedge x = y)$$

which is equivalent to $(ground(y) \wedge \neg var(y) \wedge y = x)$.

Example 3.4 Let $P = (\neg var(x, y) \wedge \neg ground(x, y) \wedge share(x, y))$ and $\mathcal{U} = \{\llbracket x, y \rrbracket\}$. Since $sp\mathcal{U}.P$ is in **reduced form**, then we can apply rule **RF**. We obtain

$$\exists x', y', z_x, y_x, z_y, y_y (P_{x', y'}^{x, y} \wedge inst(x', z_x, y_x, x) \wedge inst(x', z_x, y_x, y) \wedge inst(y', z_y, y_y, y) \wedge x = y),$$

which is equivalent to $(x = y \wedge \neg var(x, y))$, if CON contains at least a function of arity greater than one and a constant; otherwise it is equivalent to $(x = y \wedge \neg var(x, y) \wedge \neg ground(x, y))$.

4 Soundness and Completeness of the Calculus

We indicate by \mathcal{H}_{sp} the set of rules but **RF**. We first show that all the rules are equivalences. Then we show that a formula $sp\mathcal{U}.P$ can be reduced in a finite number of steps to a disjunction of formulas in reduced form, by applying rules from \mathcal{H}_{sp} . Finally rule **RF** applied to each disjunct will give the desired predicate (of \mathcal{A}) relative to $sp\mathcal{U}.P$.

The following properties of most general unifiers are useful:

- 1) Let $\mathcal{U} = \{\llbracket t_{(m)} \rrbracket, S_2, \dots, S_n\}$. If $\beta \in mgu(\{\llbracket t_{(i)} \rrbracket\})$ and $\mu \in mgu(\mathcal{U}\beta)$ then $\mu \cup \mu' \in mgu(\mathcal{U})$, where $\mu' = (\beta\mu)_{dom(\beta)}$.
- 2) $mgu(\{\llbracket t \rrbracket, S_2, \dots, S_n\}) = mgu(\{S_2, \dots, S_n\})$.

Theorem 4.1 *All rules are equivalences (with respect to Definition 2.1)*

Proof.

OR Because $P\alpha$ is true if and only if $P_j\alpha$ is true for at least one $j = 1, \dots, n$.

VAR1 Because $P\alpha$ is true if and only if either $(P \wedge \text{var}(x))\alpha$ is true or $(P \wedge \neg \text{var}(x))\alpha$ is true.

EQ Because, if $x = t$ occurs in P , then $x\alpha = t\alpha$ and so $\text{mgu}(\mathcal{U}_t^x\alpha) = \text{mgu}(\mathcal{U}\alpha)$.

F There is no α that satisfy *FALSE*.

SH1 Because $P\alpha$ is true if and only if either $(P \wedge \text{share}(x, y))\alpha$ is true or $(P \wedge \neg \text{share}(x, y))\alpha$ is true.

MIS1, MIS2 There is no μ in $\text{mgu}(\mathcal{U}\alpha)$, for all α .

STR1 Because for all α ,

$$\text{mgu}(\{\llbracket f(\underline{s}_{(k)}^1), \dots, f(\underline{s}_{(k)}^m) \rrbracket, S_2, \dots, S_n\}\alpha)$$

is equal to

$$\text{mgu}(\{\llbracket \underline{s}_j^{(m)} \rrbracket_{j=1, \dots, k}, S_2, \dots, S_n\}\alpha).$$

STR2 Let

$$P' = P \wedge \text{free}(y_1) \wedge \dots \wedge \text{free}(y_k),$$

$$\mathcal{U} = \{\llbracket f(\underline{s}_{(k)}^1), \dots, f(\underline{s}_{(k)}^i), x_{i+1}, \dots, x_m \rrbracket, S_2, \dots, S_n\},$$

$$\mathcal{U}' = \{\llbracket f(\underline{y}_{(k)}), x_{i+1}, \dots, x_m \rrbracket, \llbracket y_j, \underline{s}_j^{(i)} \rrbracket_{j=1, \dots, k}, S_2, \dots, S_n\}.$$

Let α be such that $P\alpha$ is true and let $\mu \in \text{mgu}(\mathcal{U}\alpha)$. Let u_1, \dots, u_k be fresh variables and let α', μ' be such that

$$x\alpha' = \begin{cases} u_j & \text{if } x = y_j \text{ for a } j \in [1, k], \\ x\alpha & \text{otherwise;} \end{cases}$$

$$v\mu' = \begin{cases} s_j^i\alpha\mu & \text{if } v = u_j \text{ for a } j \in [1, k], \\ v\mu & \text{otherwise.} \end{cases}$$

Then $P'\alpha'$ is true and $\mu' \in \text{mgu}(\mathcal{U}'\alpha')$. Thus $(\text{sp}\mathcal{U}'.P')\alpha'\mu'$ is true and, since $x\alpha'\mu' = x\alpha\mu$ for all x in $\text{vars}(P)$, then $(\exists y_1, \dots, y_k \text{sp}\mathcal{U}'.P')\alpha\mu$ is true.

Viceversa, let α' be such that $P'\alpha'$ is true and let $\mu' \in \text{mgu}(\mathcal{U}'\alpha')$. Then $y_i\alpha'$ is a free variable, say u_i , for all $i \in [1, k]$. Let α be equal to α' with the domain restricted to $\text{dom}(\alpha') \setminus \{\underline{y}_{(k)}\}$ and let μ be equal to μ' with the domain restricted to $\text{dom}(\mu') \setminus \{\underline{u}_{(k)}\}$. Then $P\alpha$ is true and $\mu \in \text{mgu}(\mathcal{U}\alpha)$. Thus $(\text{sp}\mathcal{U}.P)\alpha\mu$ is true and, since $x\alpha\mu = x\alpha'\mu'$ for all x in $\text{vars}(P)$, then $(\text{sp}\mathcal{U}.P)\alpha'\mu'$ is true.

SH2 Because for all α

$$\text{mgu}(\{\llbracket t, \underline{t}_{(m)}, \underline{s}_{(m')} \rrbracket, S_3, \dots, S_n\}\alpha)$$

is equal to

$$\text{mgu}(\{\llbracket t, \underline{t}_{(m)} \rrbracket, \llbracket t, \underline{s}_{(m')} \rrbracket, S_3, \dots, S_n\}\alpha).$$

SI Because of property 2).

VAR2 Let α be such that $P\alpha$ is true and let $\mu \in mgu(\mathcal{U}\alpha)$. Let α' be such that

$$x\alpha' = \begin{cases} t\alpha & \text{if } x = x_m, \\ z\alpha & \text{if } x = z\rho, \\ (x\alpha)_{t\alpha}^{x_m\alpha} & \text{otherwise.} \end{cases}$$

Let A' be an atom in P' . Then $A' = A_{\underline{z}}$, with A atom in P . If A' is monotonic then $A'\alpha'$ is an instance of $A\alpha$. Otherwise $A'\alpha' = A\alpha$. Thus in both cases $A'\alpha'$ is true. From $t\alpha' = t\alpha$ it follows that $inst(z\rho, \langle x_m\rho, \langle t, z \rangle)\alpha'$ and $(x_m = t)\alpha'$ are both true. Then $R\alpha'$ is true. Now let μ' be s.t. $\mu = \mu' \cup \{x_m\alpha/t\alpha\mu'\}$. Then from $\mathcal{U}'\alpha' = (\mathcal{U}\alpha)_{t\alpha}^{x_m\alpha}$ it follows by property 1) that $\mu' \in mgu(\mathcal{U}'\alpha')$. Thus $(sp.\mathcal{U}'.R)\alpha'\mu'$ is true and, since $x\alpha\mu = x\alpha'\mu'$ for all x in P , then $(\exists \underline{z}' sp.\mathcal{U}'.R)\alpha\mu$ is true. Viceversa let α' be such that $R\alpha'$ is true and let $\mu' \in mgu(\mathcal{U}'\alpha')$. Let α be such that

$$x\alpha = \begin{cases} x\rho\alpha' & \text{if } x \text{ in } \underline{z}, \\ x\alpha' & \text{otherwise.} \end{cases}$$

Then $P\alpha = P'\alpha'$ is true. Let $\mu = \mu' \cup \{x_m\alpha/t\alpha\mu'\}$. By $inst(z\rho, \langle x_m\rho, \langle t, z \rangle)\alpha'$ true for all z in \underline{z} it follows that $\mathcal{U}'\alpha' = (\mathcal{U}\alpha)_{t\alpha}^{x_m\alpha}$. Then by property 1) $\mu \in mgu(\mathcal{U}\alpha)$. Thus $(sp.\mathcal{U}.P)\alpha\mu$ is true and, since $x_m\alpha\mu' = t\alpha'\mu' = t\alpha\mu' = x_m\alpha\mu$, then $(sp.\mathcal{U}.P)\alpha'\mu'$ is true.

VAR3 Let

$$\begin{aligned} \mathcal{U} &= \{[f(\underline{s}_{(k)}), x_2, \dots, x_m], S_2, \dots, S_n\}, \\ \mathcal{U}' &= \{[s_i, y_i^2, \dots, y_i^m]_{i \in [1, k]}, S_2, \dots, S_n\}, \\ P' &= P \wedge x_2 = f(\underline{y}_{(k)}^2) \wedge \dots \wedge x_m = f(\underline{y}_{(k)}^m). \end{aligned}$$

Let α be such that $P\alpha$ is true and let $\mu \in mgu(\mathcal{U}\alpha)$. From $P \Rightarrow \neg var(x_j)$ it follows that $x_j\alpha$ is a structured term of the form $f(\underline{t}_{(k)}^j)$ for suitable terms t_1^j, \dots, t_k^j (otherwise $mgu(\mathcal{U}\alpha)$ would be empty).

Let α' be such that

$$x\alpha' = \begin{cases} t_i^j & \text{if } x = y_i^j \text{ for } j \in [2, m] \text{ and } i \in [1, k], \\ x\alpha & \text{otherwise} \end{cases}$$

Then $P'\alpha'$ is true and $\mu \in mgu(\mathcal{U}'\alpha')$ by construction. Thus $(sp.\mathcal{U}'.P')\alpha'\mu$ is true and, since $x\alpha' = x\alpha$ for all x occurring in P , then $(\exists \mathbf{y} sp.\mathcal{U}'.P')\alpha\mu$ is true.

Viceversa, let α' be such that $P'\alpha'$ is true and let $\mu' \in mgu(\mathcal{U}'\alpha')$. Then $s_i\alpha'\mu' = y_i^2\alpha'\mu' = \dots = y_i^m\alpha'\mu'$ for $i \in [1, k]$.

Then $P\alpha'$ is true and since $x_j\alpha'\mu' = f(y_1, \dots, y_k)\alpha'\mu'$ for all $j \in [2, m]$, then $\mu' \in mgu(\mathcal{U}\alpha)$. So $(sp.\mathcal{U}.P)\alpha'\mu'$ is true.

MIS3 By hypothesis the formula is of type 3 and \mathcal{U} contains at least a set with a structured element. Then by condition (vii) each set that contains a structured element $f(y_1, \dots, y_k)$ also contains at least a variable x that occurs in the structured element of another set. In such a situation we can eventually extract from \mathcal{U} a subset $\{S_1, \dots, S_t\}$ of sets such that

$$\begin{aligned} S_1 &= [f_1(\dots, x_t, \dots), x_1, \dots] \\ S_2 &= [f_2(\dots, x_1, \dots), x_2, \dots] \\ &\dots \\ S_t &= [f_t(\dots, x_{t-1}, \dots), x_t, \dots]. \end{aligned}$$

Clearly $\{S_1, \dots, S_t\}\alpha$ has no unifier.

GR1 Immediate since $P\alpha$ is true if and only if $(P \wedge \text{ground}(x))\alpha$ is true or $(P \wedge \neg\text{ground}(x))\alpha$ is true.

GR2 Let α and μ be such that $P\alpha$ is true and $\mu \in \text{mgu}(\mathcal{U}\alpha)$, let $\mu_i = \mu|_{\text{vars}(x_i\alpha)}$. From $\text{ground}(x_m)\alpha$ true it follows that $x_i\alpha\mu = x_m\alpha$ for $i \in [1, m-1]$. Let α' be s.t.

$$w\alpha' = \begin{cases} x\alpha & \text{if } w = x\rho \text{ with } x \text{ in } \underline{x}, \\ x_m\alpha & \text{if } w = x_i \text{ for } i \in [1, m-1], \\ \underline{y}^1 \dots \underline{y}^{m-1} & \text{if } w = z_x \text{ with } x \text{ in } \underline{x}, \\ (\underline{y}^1 \dots \underline{y}^{m-1})\mu & \text{if } w = y_x \text{ with } x \text{ in } \underline{x}, \\ (w\alpha)\mu_1 \dots \mu_{m-1} & \text{otherwise.} \end{cases}$$

where \underline{y}^i is the sequence of variables in $\text{vars}(x\alpha) \cap \text{vars}(x_i\alpha)$ for $i \in [1, m-1]$.

Let A' be an atom of P' . Then $A' = A \frac{x}{\underline{x}}$, with A atom in P . If A' is monotonic then $A'\alpha'$ is an instance of $A\alpha$. If A' is non-monotonic then $A'\alpha' = A\alpha$. In both cases $A'\alpha'$ is true. Moreover $(x_1 = \dots = x_m)\alpha'$ is true because $x_m\alpha' = x_m\alpha = x_i\alpha'$ for all $i \in [1, m-1]$, $\neg\text{share}(z_x, x)\alpha'$ is true because all variables in $z_x\alpha'$ occur in $x_i\alpha$ for some $i \in [1, m-1]$ and $x\alpha'$ is obtained replacing the variables in all $x_i\alpha$ with ground terms, $\text{inst}(x\rho, z_x, y_x, x)\alpha'$ is true because $x\rho\alpha' = x\alpha$, $x\alpha' = (x\alpha)_{y_x\alpha'}$ and $\neg\text{share}(z_x, x)\alpha'$ true imply $\{z_x\alpha'/y_x\alpha'\} \in \text{mgu}(\{\llbracket x\rho\alpha', x\alpha' \rrbracket\})$; finally $\text{invar}(z_x, \langle z_{x_1}, \dots, z_{x_{m-1}} \rangle)\alpha'$ is true by construction. Thus $R\alpha'$ is true. Now let $\mu' = \mu|_{\text{vars}(\mathcal{U}'\alpha')}$. We have that $\mu_1 \dots \mu_{m-1}$ is in $\text{mgu}(\{\llbracket x_1, \dots, x_m \rrbracket\}\alpha)$, $\text{range}(\mu_1 \dots \mu_{m-1}) = \emptyset$ because $x_m\alpha$ is ground, $\mathcal{U}'\alpha' = \mathcal{U}'\alpha\mu_1 \dots \mu_{m-1}$. Then $\mu = \mu' \cup \mu_1 \dots \mu_{m-1}$ and by properties 1) and 2) μ' is in $\text{mgu}(\mathcal{U}'\alpha')$. Then $(\text{sp}.\mathcal{U}'.R)\alpha'\mu'$ is true and, since $x\alpha'\mu' = x\alpha\mu$ for all x occurring in P , then $(\exists \underline{x}, z_{\underline{x}}, \underline{y}_{\underline{x}} \text{sp}.\mathcal{U}.R)\alpha\mu$ is true.

Viceversa let α' be such that $R\alpha'$ is true and let $\mu' \in \text{mgu}(\mathcal{U}'\alpha')$. Let α be s.t.

$$x\alpha = \begin{cases} x\rho\alpha' & \text{if } x \text{ in } \underline{x}, \\ x\alpha' & \text{otherwise.} \end{cases}$$

Then $P\alpha = P'\alpha'$ is true. Let $\mu = \mu' \cup \beta$ with $\beta = \{(z_{x_i}\alpha'/y_{x_i}\alpha')_{i \in [1, m-1]}\}$. From $\text{inst}(x\rho, z_x, y_x, x)\alpha'$, $\neg\text{share}(z_{x_j}, x)\alpha'$ and $\text{invar}(z_x, \langle z_{x_1}, \dots, z_{x_{m-1}} \rangle)\alpha'$ true it follows that $x\alpha' = x\alpha\beta$ for all $x \in \underline{x}$. If $x \notin \underline{x}$ then from $\neg\text{share}(z_y, x)\alpha'$ true for all y it follows that $x\alpha' = (x\alpha')\beta = x\alpha\beta$. Then $x\alpha' = x\alpha\beta$ for all x occurring in P . Then $\mathcal{U}'\alpha' = \mathcal{U}'\alpha\beta$. From $x_1\alpha' = \dots = x_m\alpha'$ true, $x_m\alpha'$ ground and $\text{inst}(x_i\rho, z_{x_i}, y_{x_i}, x_i)\alpha'$ true for all $i \in [1, m-1]$ it follows that $\beta \in \text{mgu}(\{\llbracket x_1, \dots, x_m \rrbracket\}\alpha)$. Then by properties 1) and 2) it follows that $\mu \in \text{mgu}(\mathcal{U}\alpha)$. Thus $(\text{sp}.\mathcal{U}.P)\alpha\mu$ is true and, since $x\alpha'\mu' = (x\alpha)\beta\mu' = x\alpha\mu$ for all x occurring in P , then $(\text{sp}.\mathcal{U}.P)\alpha'\mu'$ is true.

RF Let α and μ be such that $P\alpha$ is true, $\mu \in \text{mgu}(\mathcal{U}\alpha)$. Let α' be s.t.

$$w\alpha' = \begin{cases} x\alpha & \text{if } w = x\rho \text{ with } x \text{ in } \underline{x}, \\ w\alpha\mu & \text{if } w \text{ occurs in } P, \\ \underline{y} & \text{if } w = z_x \text{ with } x \text{ in } \underline{x}, \\ \underline{y}\mu & \text{if } w = y_x \text{ with } x \text{ in } \underline{x}. \end{cases}$$

where \underline{y} is the sequence of variables occurring in $\text{dom}(\mu|_{\text{vars}(x\alpha)})$. Now $U\alpha'$ is true because $x_i^j\alpha' = x_i^j\alpha\mu$ for every $i \in [1, m_j]$, $j \in [1, n]$. Let A' be an atom of P' . Then $A' = A \frac{x}{\underline{x}}$, with A atom in P . If A' is monotonic then $A'\alpha'$ is an instance of $A\alpha$. If A' is non-monotonic then $A'\alpha' = A\alpha$. In both cases $A'\alpha'$ is true. Moreover $\text{inst}(x\rho, z_x, y_x, x)\alpha'$ is true because $x\rho\alpha' = x\alpha$, $x\alpha' = x\alpha\mu$ and the substitution relative to the two sequences $z_x\alpha'$ and $y_x\alpha'$ is equal to $\mu|_{\text{vars}(x\alpha)}$.

Since μ is idempotent by hypothesis, then $\mu|_{vars(x\alpha)} \in mgu(\{\llbracket x\alpha, x\alpha\mu \rrbracket\})$. Then $(R \wedge U)\alpha'$ is true and, since $x\alpha' = x\alpha\mu$ for every x occurring in P , then $(\exists \underline{x}', z_{\underline{x}}, y_{\underline{x}}(R \wedge U))\alpha\mu$ is true. Viceversa let α' be s.t. $(R \wedge U)\alpha'$ is true. Let α be s.t.

$$x\alpha = \begin{cases} x\rho\alpha' & \text{if } x \text{ in } \underline{x}, \\ x\alpha' & \text{otherwise.} \end{cases}$$

Then $P\alpha = P'\alpha'$ is true. Let μ be the substitution relative to the sequences $z_{x_i^j}\alpha', y_{x_i^j}\alpha'$ for all $i \in [1, m_j], j \in [1, n]$. Then $\mu \in mgu(\mathcal{U}\alpha)$. Thus $(sp.\mathcal{U}.P)\alpha\mu$ is true and, since $x\alpha\mu = x\rho\alpha'\mu = x\alpha'$ for every x that occurs in P , then $(sp.\mathcal{U}.P)\alpha'$ is true. \square

Theorem 4.2 *The system \mathcal{H}_{sp} is terminating.*

Proof.

We show that no proof tree built using \mathcal{H}_{sp} has an infinite branch. Rules **F**, **MIS1**, **MIS2** and **MIS3** have a predicate as right hand side, so they cannot belong to an infinite branch. To prove that only finitely many applications of the remaining rules are allowed, consider the tuple

$$\tau = (leq, comp, funct, elem, disj, unvar, unshare, unground)$$

of natural numbers with the lexicographic order. A structured term $f(t_1, \dots, t_n)$ will be called *compound* if either some t_i is not a variable or the variables t_1, \dots, t_n are not distinct. Then

leq denotes the number of variables in \mathcal{U} that occur as left hand side of an equation in P .
 $comp$ denotes the number of occurrences of compound subterms of terms in \mathcal{U} . Thus

$$comp = \sum_{j=1}^n \sum_{i=1}^{m_j} comp(t_i^j)$$

$$comp(t_i^j) = \begin{cases} 0 & \text{if } t_i^j \text{ not compound;} \\ 1 + \sum_{k=1}^m comp(s_k) & \text{if } t_i^j = f(s_1, \dots, s_m) \text{ compound} \end{cases}$$

$funct$ denotes the number of occurrences of function symbols in \mathcal{U} . Thus

$$funct = \sum_{j=1}^n \sum_{i=1}^{m_j} funct(t_i^j)$$

$$funct(t_i^j) = \begin{cases} 0 & \text{if } t_i^j \in VAR; \\ 1 + \sum_{k=1}^m funct(s_k) & \text{if } t_i^j = f(s_1, \dots, s_m) \end{cases}$$

$elem$ denotes the total number of elements in the sets of \mathcal{U} . Thus

$$elem = \sum_{j=1}^n |S_j|$$

where $|S_j|$ indicates the cardinality of the set S_j .

$disj$ denotes the number of disjuncts in the disjunctive normal form of the precondition P .

$unvar$ denotes the number of variables $x \in vars(P)$ such that neither $P \Rightarrow var(x)$ nor $P \Rightarrow \neg var(x)$.

$unshare$ denotes the number of variables $x \in vars(P)$ such that neither $P \Rightarrow share(x, y)$ nor $P \Rightarrow \neg share(x, y)$ for some distinct variable $y \in vars(\mathcal{U})$.

$unground$ denotes the number of variables $x \in vars(P)$ such that neither $P \Rightarrow ground(x)$ nor $P \Rightarrow \neg ground(x)$.

Let us list the remaina rules to show that their application decreases the value of τ .

EQ decreases leq .

OR decreases $disj$, does not change $leq, comp, funct$ and $elem$.

SH1 decreases *unshare*, does not change *unvar*, *leq*, *elem*, *funct*, *comp* and does not increase *disj* (*disj* may decrease if a disjunct contains $\neg share(x, y)$ or $share(x, y)$).

VAR1 decreases *unvar*, does not change *leq*, *elem*, *funct*, *comp* and does not increase *disj* (*disj* may decrease if a disjunct contains $\neg var(x)$ or $var(x)$).

STR1 decreases *funct* and does not increase *leq*, *comp* (*comp* may decrease if some s_i is a structured term).

STR2 decreases *comp* or does not change *leq*, *comp* and decreases *funct*.

SH2 decreases *elem*, and does not increase *leq*, *comp* and *funct* (*funct* may decrease if t is a structured term, *comp* may decrease if t is a compound term).

SI decrease *elem*, and does not increase *leq*, *funct* (*funct* may decrease if t is a structured term).

VAR2 decreases *elem* and does not change *leq*, *funct* and *comp*.

VAR3 decreases *funct* and does not change *leq* and *comp*.

GR1 decreases *unground*, does not change *leq*, *comp*, *funct*, *elem*, *unshare*, *unvar* and does not increase *disj* (*disj* may decrease if a disjunct contains $\neg ground(x)$ or $ground(x)$).

GR2 decreases *elem* and does not change *leq*, *comp*, *funct*. \square

Theorem 4.3 *Rules of \mathcal{H}_{sp} transform $sp.U.P$ in a (semantically unique) disjunction of formulas in reduced form.*

Proof.

By Theorem 4.1 all transformations are equivalencies (w.r.t. Definition 2.1). By Theorem 4.2 there is a final form. Thus the final form is semantically unique. By contraposition suppose that the final form is not a disjunction of formulas in reduced form. Then if condition (i) is not satisfied then rule **OR** can be applied; if condition (ii) is not satisfied then rule **EQ** can be applied; if condition (iii) is not satisfied then rule **VAR1** can be applied; if condition (iv) is not satisfied then rule **SH1** can be applied; if condition (v) is not satisfied then rule **SH2** can be applied; if condition (vi) is not satisfied then rule **SI** can be applied; if one set contains a structured term then either rule **STR1** or rule **STR2** can be applied or the set contains only one structured term that is a flat term: in such a case if the formula is not of type 3 then rule **VAR2** or rule **VAR3** can be applied, otherwise rule **MIS3** can be applied. If for a variable x occurring in \mathcal{U} $var(x)$ occurs in P then rule **MIS2** or **VAR2** can be applied. Finally if, for a variable x occurring in \mathcal{U} , $ground(x)$ occurs in P then rule **GR1** or **GR2** can be applied. \square

Corollary 4.4 *sp is a sound and complete calculus for $sp.U$.*

5 Applications

5.1 A Forward Semantics for Logic Programs

Predicate transformers are related to the core of abstract interpretation of imperative programs. In [2] predicate transformers are used to define deductive semantics. Deductive semantics is used to design approximate program analysis frameworks. To propose a similar approach in the setting of logic programming we need the correspondent of program point for a logic program. In [7] Nilsson introduced a scheme for inferring run-time properties of logic programs based on a semantic description of logic programs that uses the concept of program point. The predicate transformer sp can be easily cast in such a theory. A clause of a logic program \mathcal{P} is interpreted as a sequence of procedure calls. To each call A there corresponds a *calling point* $\bullet A$ and a *success*

point A_\bullet . The leftmost and rightmost points in the body of a clause C are called respectively *entry-* and *exit points* of the clause and are indicated respectively by $\bullet C$ and C_\bullet . Goals are represented as elements of the set $\mathcal{Cgoals} := (\mathcal{Points} \times \mathcal{Env})^*$, where \mathcal{Points} denotes the set of program points of \mathcal{P} and \mathcal{Env} is the set of predicates \mathcal{A} . A transition system for \mathcal{P} can be defined through two state transition schemes that transform elements of \mathcal{Cgoals} as follows.

$$\begin{aligned} \langle C_\bullet; R \rangle &:: y \models y, \\ \langle \bullet A; R \rangle &:: y \models (\langle \bullet C\sigma; TRUE \rangle :: \langle A_\bullet; R \rangle :: y)T, \end{aligned}$$

where A is a body atom, $C\sigma$ is a variant of a clause C of \mathcal{P} s.t. $vars(\langle \bullet A; R \rangle :: y) \cap vars(C\sigma) = \emptyset$, $T \equiv sp.\{\llbracket A, head(C\sigma) \rrbracket\} . (R \wedge free(vars(C\sigma))) \not\equiv FALSE$. We assume that the definition of \mathcal{U} in sp is generalized in the obvious way to atoms or terms. The application of a predicate R to a \mathcal{C} -goal is defined as follows:

$$\begin{aligned} (nil)R &= R, \\ (\langle x; T \rangle :: y)R &= \langle x; T \bullet R \rangle :: yR, \end{aligned}$$

where $T \bullet R$ is (equivalent to) $T' \wedge R$, with T' the strongest assertion (w.r.t. implication) s.t. $T \rightarrow T'$ and $(T' \wedge R) \not\equiv FALSE$. Notice that $T \bullet R$ is defined when R is consistent. For instance if $T = (x = y \wedge var(x))$ and $R = ground(y)$ then $T \bullet R \equiv (x = y \wedge ground(y))$.

The previous transitions schemes are obtained from those in [7] by taking as environment \mathcal{Env} predicates instead of substitutions, by using the predicate transformer sp instead of the mgu as operation in the transition and the operation \bullet to model the application of a predicate to a \mathcal{C} -goal. From [7], to each program point i is associated a set Θ_i of states which specifies when the program point becomes current. The set of states is defined as $\mathcal{Cgoals} \times \mathcal{Cgoals}$, where the first component describes the \mathcal{C} -goal that invoked the clause containing point i and the second component is the \mathcal{C} -goal when the point is current. The semantics of \mathcal{P} is defined as the least fixpoint of the system of equations relative to its program points. Every program point is either the entry point of a clause or the success point of a body atom. Then it is sufficient to define the meaning of entry- and success points. Let $A \rightsquigarrow C$ indicate that A and a variant of the head of the clause C can be unified. Then:

$$\begin{aligned} \Theta_{\bullet C} &= \bigcup_{A \rightsquigarrow C} \{ \langle G_i; G_{i+1} \rangle \mid \exists G (\langle G; G_i \rangle \in \Theta_{\bullet A} \wedge G_i \models^{C_1} G_{i+1}) \}, \\ \Theta_{A_\bullet} &= \bigcup_{A \rightsquigarrow C} \{ \langle G; tail(G_j) \rangle \mid \exists G_i (\langle G; G_i \rangle \in \Theta_{\bullet A} \wedge \langle G_i; G_j \rangle \in \Theta_{C_\bullet}) \}, \end{aligned}$$

Example 5.1 Consider the following simple case of concatenation of two lists:

$$\begin{aligned} C_0 &: \leftarrow_1 \text{append}([a], [], z)_2. \\ C_1 &: \text{append}([H|L1], L2, [H|L3]) \leftarrow_3 \text{append}(L1, L2, L3)_4. \\ C_2 &: \text{append}([], L, L) \leftarrow_5. \end{aligned}$$

Here the program points are explicitly labelled by integers. The meaning of this program, when $\text{append}([a], [], z)$ is called with z *free* variable, can be given as least fixpoint of the following set of equations, where we use the notation of [7].

$$\begin{aligned} \Theta_1 &= \{ \langle nil; \langle \bullet C_0; free(z) \rangle :: nil \rangle \}, \\ \Theta_2 &= \{ \langle G; tail(G_j) \rangle \mid \exists G_i (\langle G; G_i \rangle \in \Theta_1 \wedge (\langle G_i; G_j \rangle \in \Theta_4 \vee \langle G_i; G_j \rangle \in \Theta_5)) \}, \\ \Theta_3 &= \{ \langle G_i; G_{i+1} \rangle \mid \exists G ((\langle G; G_i \rangle \in \Theta_1 \vee \langle G; G_i \rangle \in \Theta_3) \wedge G_i \models^{C_1} G_{i+1}) \}, \\ \Theta_4 &= \{ \langle G; tail(G_j) \rangle \mid \exists G_i (\langle G; G_i \rangle \in \Theta_3 \wedge (\langle G_i; G_j \rangle \in \Theta_4 \vee \langle G_i; G_j \rangle \in \Theta_5)) \}, \\ \Theta_5 &= \{ \langle G_i; G_{i+1} \rangle \mid \exists G ((\langle G; G_i \rangle \in \Theta_1 \vee \langle G; G_i \rangle \in \Theta_3) \wedge G_i \models^{C_2} G_{i+1}) \}. \end{aligned}$$

Notice that in this case the fixpoint can be computed in finite time since the program terminates. We first calculate Θ_3 . We need to compute

$sp.\{\llbracket append([a], [], z), append([H|L1], L2, [H|L3]) \rrbracket\}.(free(z, H, L1, L2, L3)).$

By rule **STR1**, rule **VAR2** applied to $L2$ and z , rules **SI**, **STR1** and rule **VAR2** applied to H and $L1$ we obtain the predicate

$$T \equiv (H = a \wedge L1 = [] \wedge L2 = [] \wedge z = [a|L3] \wedge free(L3)).$$

Since $(free(z) \bullet T) = T$ then $\langle \bullet C_0; free(z) \rangle :: nil \models^{C_1} \langle \bullet C_1; T \rangle :: \langle C_{0\bullet}; T \rangle :: nil.$

By rules **STR1** and **MIS1**

$sp.\{\llbracket append(L1, L2, L3), append([H'|L1'], L2', [H'|L3']) \rrbracket\}.(free(H', L1', L2', L3') \wedge T)$
is equivalent to $FALSE$. Hence

$$\Theta_3 = \{\langle \bullet C_0; free(z) \rangle :: nil; \langle \bullet C_1; T \rangle :: \langle C_{0\bullet}; T \rangle :: nil\}.$$

Consider now Θ_5 . We need to compute

$$sp.\{\llbracket append([], L, L), append(L1, L2, L3) \rrbracket\}.(free(L) \wedge T).$$

By rule **STR1**, rule **EQ** applied to $L1$ and $L2$, rule **SI**, rule **SH2** applied to L and rule **VAR2** applied to L and $L3$ we obtain the predicate

$$R \equiv ((H = a \wedge L1 = L2 = L3 = L = [] \wedge z = [a]).$$

Since $T \bullet R = R$ then

$$\langle \bullet C_1; T \rangle :: \langle C_{0\bullet}; T \rangle :: nil \models^{C_2} \langle \bullet C_2; R \rangle :: \langle C_{1\bullet}; R \rangle :: \langle C_{0\bullet}; R \rangle :: nil.$$

By rules **STR1** and **MIS1**

$sp.\{\llbracket append([a], [], z), append([], L, L) \rrbracket\}.(free(L) \wedge T)$
is equivalent to $FALSE$. Hence

$$\Theta_5 = \{\langle \bullet C_1; T \rangle :: \langle C_{0\bullet}; T \rangle :: nil; \langle \bullet C_2; R \rangle :: \langle C_{1\bullet}; R \rangle :: \langle C_{0\bullet}; R \rangle :: nil\}.$$

Finally Θ_2 and Θ_4 can be easily calculated.

$$\Theta_2 = \{\langle nil; \langle C_{0\bullet}; R \rangle :: nil \rangle\};$$

$$\Theta_4 = \{\langle \bullet C_0; free(z) \rangle :: nil; \langle C_{1\bullet}; R \rangle :: \langle C_{0\bullet}; R \rangle :: nil\}.$$

Every set Θ_i describes the states associated to the program point i . Thus for instance Θ_3 specifies that the program point 3 becomes current only when the goal $append([a], [], z)$ invokes C_1 with z free variable and in such a case H becomes equal to a , $L1$ and $L2$ become equal to the empty list $[]$ and $L3$ remains a free variable.

6 Conclusions

In this paper we proposed a formal characterization of unification by means of the predicate transformer $sp\mathcal{U}$. We introduced a calculus for $sp\mathcal{U}$ and proved its soundness, completeness and termination. The properties considered include some non-monotonic predicates on the structure of terms, like var , $share$ and $\neg ground$. Non-monotonic predicates increase the complexity of the calculus that otherwise, i.e. in case of monotonic predicates, would consist of one simple rule.

The predicate transformer allows to define a forward semantics of logic programs that can be used to design approximate program analysis framework in the style of those proposed by Cousot and Cousot for imperative programs ([2]).

References

- [1] P. Cousot, R. Cousot. Abstract Interpretation : a Unified Lattice Model for Static Analysis of Programs by Construction or Approximation of Fixpoints. *Proceedings of the 4th ACM Symposium on Principles of Programming Languages*, 238–251, 1977.
- [2] P. Cousot, R. Cousot. Systematic Design of Program Analysis Frameworks. *Proceedings of the 6th ACM Symposium on Principles of Programming Languages*, 269–282, 1979.
- [3] E. W. Dijkstra. A Discipline of Programming. *Prentice-Hall*, 1976.
- [4] E. Eder. Properties of Substitutions and Unifications. *Journal of Symbolic Computation*, 1: 31–46, 1985.
- [5] R.W. Floyd. Assigning Meanings to Programs. *Proc. Symp. Appl. Math.*, American Math. Society, Providence, Rhode Island, 19: 15–32, 1967.
- [6] J-L. Lassez, M.J. Maher, K. Marriott. Unification revisited. *Foundations of Logic and Functional Programming, LNCS 306*, 1987.
- [7] Ulf Nilsson. Systematic Semantics Approximations of Logic Programs. *Proceedings of PLILP '90, Springer-Verlag*, 1990.