

STICHTING
MATHEMATISCH CENTRUM
2e BOERHAAVESTRAAT 49
AMSTERDAM
AFDELING ZUIVERE WISKUNDE

ZW 1967-003

On convex sublattices of
distributive lattices

by

J.W. de Bakker



May 1967

The Mathematical Centre at Amsterdam, founded the 11th of February, 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications, and is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.) and the Central Organization for Applied Scientific Research in the Netherlands (T.N.O.), by the Municipality of Amsterdam and by several industries.

1. Introduction

In this paper we study some properties of convex sublattices of distributive lattices.

The family of all convex sublattices of a lattice L will be denoted by $\mathcal{C}(L)$.

Section 2 contains some definitions and preliminary lemma's.

The first result of section 3 is the following: Let L be a distributive lattice and let $A, B \in \mathcal{C}(L)$ with $A \subset B$, $A \neq \emptyset$. In theorem 1 we prove that the family of all elements of $\mathcal{C}(L)$ which have the intersection A with B has a largest element, by means of an explicit construction of this element from A and B . The next theorems are concerned with congruence relations. Let $C \in \mathcal{C}(L)$. We construct the smallest congruence relation such that C is one of its congruence classes, and the largest congruence relation such that all elements of C are incongruent with respect to this congruence relation. Next, these results are related to the construction of theorem 1.

In section 4 we consider the lattice $(\mathcal{C}(L), \subset)$, i.e. the family of all convex sublattices of L , partially ordered by inclusion. We prove that in a distributive relatively complemented lattice L , all intervals $[\emptyset, A]$ of $(\mathcal{C}(L), \subset)$ are complemented. A necessary and sufficient condition that $(\mathcal{C}(L), \subset)$ be relatively complemented, is that L is also discrete (i.e., all intervals of L have finite length).

In section 5 we introduce an ordering \leq on $\overline{\mathcal{C}}(L)$ (i.e., the family of all non-empty convex sublattices of L), which is a variant of the ordering by inclusion. We prove that $(\overline{\mathcal{C}}(L), \leq)$ is a distributive lattice, if L is distributive. Next we consider the lattices $\overline{\mathcal{C}}^2(L) = \overline{\mathcal{C}}(\overline{\mathcal{C}}(L))$, ..., $\overline{\mathcal{C}}^i(L)$. We prove that $\overline{\mathcal{C}}^i(B_j)$, where B_j is the Boolean algebra with 2^j elements, is isomorphic with the direct union of j factors F_i , where F_i is the free distributive lattice with i generators, with an extra zero and unit element adjoined.

Section 6 is concerned with a ternary function which can be used to characterize convex sublattices of distributive relatively complemented

lattices. Finally, we exhibit a set of axioms for distributive relatively complemented lattices in terms of this ternary function.

I am indebted to A.B. Paalman-de Miranda for several helpful suggestions.

2. Definitions

Definition 1. Let X be a subset of a lattice L . The sets X_l, X_r are defined as follows:

$$\begin{aligned} X_l &= \{ a \in L \mid \exists x \in X \text{ such that } a \leq x \}, \\ X_r &= \{ a \in L \mid \exists x \in X \text{ such that } a \geq x \}. \end{aligned}$$

It is easily seen that:

1. l and r are closure-operators, i.e. for all $X, Y \subset L$ we have:
 $X \subset X_l, X_l = X_{l.l}, (X \cup Y)_l = X_l \cup Y_l$, and similarly for r .
2. $X_{lr} = X_{rl} = L$.
3. If X is closed with respect to $\vee(\wedge)$ then $X_l(X_r)$ is a \vee -ideal (\wedge -ideal).
4. If X is a \vee -ideal (\wedge -ideal) then $X = X_l$ ($X = X_r$).

Definition 2. Let X, Y be non-empty subsets of a lattice L . The sets $X \wedge Y, X \vee Y$ are defined as follows:

$$\begin{aligned} X \wedge Y &= \{ x \wedge y \mid x \in X \text{ and } y \in Y \}, \\ X \vee Y &= \{ x \vee y \mid x \in X \text{ and } y \in Y \}. \end{aligned}$$

Clearly, for all $X, Y \subset L$ we have $(X \wedge Y)_l = X_l \wedge Y_l$, and $(X \vee Y)_r = X_r \vee Y_r$. It is also easy to prove that $(X \vee Y)_l = X_l \vee Y_l$, for all $X, Y \subset L$, if and only if L is distributive (and dually).

Definition 3. A subset of a lattice L is called convex if and only if $X_l \cap X_r \subset X$.

In this paper we are only interested in convex sublattices. The family of all convex sublattices of L will be denoted by $\mathcal{C}(L)$. The family of all \vee -ideals (\wedge -ideals) of L will be denoted by $I(L)$ ($Y(L)$). Some of the simplest properties of $\mathcal{C}(L)$ are:

1. $I(L) \subset \mathcal{C}(L)$ and $J(L) \subset \mathcal{C}(L)$.
2. The intersection of a family of convex sublattices is a convex sublattice.
3. If A is closed with respect to \vee and B is closed with respect to \wedge then $A_{\perp} \cap B_r \in \mathcal{C}(L)$.
4. A subset C of L is a convex sublattice of L if and only if it has the following property: For all $c_1, c_2 \in C$ and all $x \in L$ we have: $c_1 \wedge (x \vee c_2) \in C$ and $c_1 \vee (x \wedge c_2) \in C$.

Clearly, if $C \in \mathcal{C}(L)$ then $C = C_{\perp} \cap C_r$. Hence, each convex sublattice can be written as the intersection of a \vee -ideal and a \wedge -ideal. The following lemma proves that this "decomposition" is unique:

Lemma 1. Let $C \in \mathcal{C}(L)$, $C \neq \emptyset$, and suppose that $C = I \cap J$, where I is a \vee -ideal and J is a \wedge -ideal. Then $I = C_{\perp}$ and $J = C_r$.

Proof. $C = I \cap J \subset I$, hence $C_{\perp} \subset I_{\perp} = I$. Also, $I \vee (I \cap J) = C$; hence, $I = I_{\perp} \subset \{I \vee (I \cap J)\}_{\perp} = C_{\perp}$. Thus $I = C_{\perp}$. Similarly, $J = C_r$.

From this lemma it follows that if $C, D \in \mathcal{C}(L)$, and $C \cap D \neq \emptyset$, then $(C \cap D)_{\perp} = C_{\perp} \cap D_{\perp}$, and $(C \cap D)_r = C_r \cap D_r$.

Definition 4. Let $C, D \in \mathcal{C}(L)$. The smallest convex sublattice of L that contains C and D is denoted by $C \sqcup D$.

From this definition it follows that if $C \neq \emptyset$, $D \neq \emptyset$, then $C \sqcup D = (C \wedge D)_r \cap (C \vee D)_{\perp}$.

The following two lemma's state some properties of convex sublattices that will be used later.

Lemma 2. Let $A, B, C \in \mathcal{C}(L)$, with $A \cap B \neq \emptyset$, $B \cap C \neq \emptyset$, and $C \cap A \neq \emptyset$. Then $A \cap B \cap C \neq \emptyset$.

Proof. Let $x \in A \cap B$, $y \in B \cap C$ and $z \in C \cap A$. Since $x, y \in B$ and $x, z \in A$, we have $x \wedge (y \vee z) \in A \cap B$.

Now consider the element $\{x \wedge (y \vee z)\} \vee (y \wedge z)$.

We have: $x \wedge (y \vee z)$ and y are elements of B ,
 $x \wedge (y \vee z)$ and z are elements of A ,
 $y \vee z$ and $y \wedge z$ are elements of C .

Therefore, $\{x \wedge (y \vee z)\} \vee (y \wedge z) \in A \cap B \cap C$.

Two consequences of this lemma are:

1. If $A_i \in \mathcal{C}(L)$, $1 \leq i \leq n$, and $\bigcap_{i=1}^n A_i = \emptyset$, then $A_i \cap A_j = \emptyset$, for some i, j with $1 \leq i, j \leq n$.
2. If $A_i \in \mathcal{C}(L)$, $1 \leq i \leq n$, and $\bigcap_{\substack{i=1 \\ i \neq j}}^n A_i \neq \emptyset$ for three values of j ($1 \leq j \leq n$), then $\bigcap_{i=1}^n A_i \neq \emptyset$.

Lemma 3. Let L be a distributive lattice and let $C, D \in \mathcal{C}(L)$ with $C \cap D = \emptyset$. Then there exist $C', D' \in \mathcal{C}(L)$, such that $C \subset C'$, $D \subset D'$, $C' \cap D' = \emptyset$ and $C' \cup D' = L$. Moreover, either C' is a \vee -ideal and D' is a \wedge -ideal or conversely.

Proof. (This proof is due to P.C. Baayen).

1. Either $C_1 \cap D_r = \emptyset$, or $C_r \cap D_1 = \emptyset$. For, suppose that there exist $c_1, c_2 \in C$ and $d_1, d_2 \in D$ with $c_1 \leq d_1$ and $c_2 \geq d_2$. Then $c_1 \leq d_1 \wedge (c_1 \vee d_2) \leq c_1 \vee c_2$; hence, $d_1 \wedge (c_1 \vee d_2) \in C \cap D$, a contradiction.
2. Suppose $C_1 \cap D_r = \emptyset$. We can then apply Stone's theorem [10] to the \wedge -ideal D_r .

3. Congruence relations

Theorem 1. Let L be a distributive lattice, let $A, B \in \mathcal{C}(L)$ with $A \subset B$, $A \neq \emptyset$. Let C be defined as:

$$C = (A_r \setminus (B \setminus A_1)_r)_1 \cap (A_1 \setminus (B \setminus A_r)_1)_r.$$

Then:

1. $C \in \mathcal{C}(L)$.
2. $B \cap C = A$.
3. $D \in \mathcal{C}(L)$ and $D \cap B = A$ imply $D \subset C$.

Proof.

1. In order to prove that $C \in \mathcal{C}(L)$ it is sufficient to prove that $A_r \setminus (B \setminus A_1)_r$ is closed with respect to \vee . Let $a'_1, a'_2 \in A_r \setminus (B \setminus A_1)_r$. Clearly, $a'_1 \vee a'_2 \in A_r$. Suppose that $a'_1 \vee a'_2 \geq b$ for some $b \in B \setminus A_1$. Since $a'_1 \in A_r$, there exists $a_1 \in A$ such that $a'_1 \geq a_1$. Then $b \geq b \wedge a'_1 \geq b \wedge a_1$. Since $A \subset B$ and B is a convex sublattice, we have $b \wedge a_1 \in B$, and $b \wedge a'_1 \in B$. Since $a'_1 \geq b \wedge a_1$, and $a'_1 \notin (B \setminus A_1)_r$, we see that $b \wedge a'_1 \in A_1$, so that there exists $a_3 \in A$, with $a_3 \geq b \wedge a'_1$. Similarly, there exists $a_4 \in A$ such that $a_4 \geq b \wedge a'_2$. Thus, $a_3 \vee a_4 \geq (b \wedge a'_1) \vee (b \wedge a'_2) = b \wedge (a'_1 \vee a'_2) = b$, which contradicts $b \in B \setminus A_1$. We conclude therefore that $a'_1 \vee a'_2 \in A_r \setminus (B \setminus A_1)_r$, where $A_r \setminus (B \setminus A_1)_r$ is closed with respect to \vee (it is easy to prove that $A_r \setminus (B \setminus A_1)_r$ is even a convex sublattice).
- 2.1. In order to prove that $A \subset B \cap C$ it is sufficient to prove that $A \subset A_r \setminus (B \setminus A_1)_r$. It is clear that $A \subset A_r$. Also, it is impossible that there exists $a \in A$ such that $a \geq b$ for some $b \in B \setminus A_1$.
- 2.2. Let $b \in B \cap C$. Then there exist $a' \in A_r \setminus (B \setminus A_1)_r$ and $a'' \in A_1 \setminus (B \setminus A_r)_1$ such that $a'' \leq b \leq a'$. From $a' \geq b$ and $a' \notin (B \setminus A_1)_r$ we see that $b \in A_1$. Similarly, from $a'' \leq b$ we infer that $b \in A_r$. Hence $b \in A_1 \cap A_r = A$, from which we conclude that $B \cap C \subset A$.
3. Suppose $D \in \mathcal{C}(L)$ and $D \cap B = A$. We have to prove that $D \subset C$. It is sufficient to show that for each $d \in D$ and $a \in A$: $d \vee a \in A_r \setminus (B \setminus A_1)_r$. Clearly, $d \vee a \in A_r$. Suppose that $d \vee a \geq b$ for some $b \in B \setminus A_1$. Then $b = b \wedge (d \vee a) \leq (b \wedge d) \vee a$. Since $A \subset B$ and $A \subset D$, we have $(b \wedge d) \vee a \in B$ and $(b \wedge d) \vee a \in D$; hence, $(b \wedge d) \vee a \in A$. This contradicts $b \in B \setminus A_1$.

Corollary. Let L be a distributive lattice, let $A, B \in \mathcal{C}(L)$ with $A \subset B$, $A \neq \emptyset$, and let $C(A, B)$ be the largest element of $\mathcal{C}(L)$ which has the intersection A with B . Then $C(A, B) = C(A, C(A, C(A, B)))$.

Proof. Since $C(A, C(A, B))$ is the largest convex sublattice which has the intersection A with $C(A, B)$, and since $B \cap C(A, B) = A$, we have $B \subset C(A, C(A, B))$. Thus, $B \cap C(A, C(A, C(A, B))) = B \cap C(A, C(A, B)) \cap C(A, C(A, C(A, B))) = B \cap A = A$.

Since $C(A, B)$ is the largest convex sublattice which has the intersection A with B , we have

$$(1) \quad C(A, C(A, C(A, B))) \subset C(A, B).$$

Since $C(A, B) \cap C(A, C(A, B)) = A$, and since $C(A, C(A, C(A, B)))$ is the largest convex sublattice which has the intersection A with $C(A, C(A, B))$, we have

$$(2) \quad C(A, B) \subset C(A, C(A, C(A, B))).$$

From (1) and (2) the assertion follows.

Remarks:

1. From this corollary it follows that $C(A, C(A, B))$ is the largest element of the family of all elements $B' \in \mathcal{C}(L)$ such that $C(A, B) = C(A, B')$:
 - a. If $B' = C(A, C(A, B))$ then $C(A, B') = C(A, C(A, C(A, B))) = C(A, B)$.
 - b. If $C(A, B) = C(A, B')$, then $B' \subset C(A, C(A, B')) = C(A, C(A, B))$.
2. In section 4 we shall derive a sufficient condition for L in order that for each $A, B \in \mathcal{C}(L)$ with $A \subset B$, $A \neq \emptyset$, we have $B = C(A, C(A, B))$.
3. Clearly, the corollary can be formulated more generally as a statement on sets instead of on lattices.

The next theorems are concerned with congruence relations.

In theorems 2 and 3 we investigate some general properties of congruence relations in distributive lattices, and in theorem 5 we relate these properties to the construction of theorem 1.

Theorem 2. Let L be a distributive lattice and let K be a sublattice of L . Let the relation R_K be defined as follows:

$$xR_K y \text{ if and only if there exist } k_1, k_2 \in K \text{ such that } k_1 \wedge x = k_1 \wedge y \\ \text{and } k_2 \vee x = k_2 \vee y.$$

Then R_K is the smallest congruence relation that contains K in one of its congruence classes.

Proof. It is clear that $xR_K x$ and that $xR_K y$ implies $yR_K x$. Now suppose that $xR_K y$ and $yR_K z$ hold. This means that there exist k_1, k_2, k_3, k_4 such that $k_1 \wedge x = k_1 \wedge y$, $k_2 \vee x = k_2 \vee y$, $k_3 \wedge y = k_3 \wedge z$ and $k_4 \vee y = k_4 \vee z$. Hence, $k_1 \wedge k_3 \wedge x = k_1 \wedge k_3 \wedge z$ and $k_2 \vee k_4 \vee x = k_2 \vee k_4 \vee z$. Since K is a sublattice, we see that $xR_K z$. It is easy to verify that if $xR_K y$ and

$t \in L$ than $x \wedge t R_K y \wedge t$ and $x \vee t R_K y \vee t$. Clearly, all elements of K are congruent with respect to R_K . There remains the proof that R_K is the smallest congruence relation with this property. Suppose S is a congruence relation such that for all $k_1, k_2 \in K : k_1 S k_2$.

We prove that $R_K \leq S$, i.e., $x R_K y$ implies $x S y$. From $x R_K y$ we see that there exist k_1, k_2 such that $k_1 \wedge x = k_1 \wedge y$ and $k_2 \vee x = k_2 \vee y$. From $k_1 S k_2$ it follows that $x \wedge k_1 S x \wedge k_2$; hence,

$$y = y \vee (y \wedge k_1) = y \vee (x \wedge k_1) S y \vee (x \wedge k_2).$$

also, $y \wedge k_1 S y \wedge k_2$; hence,

$$x = x \vee (x \wedge k_1) = x \vee (y \wedge k_1) S x \vee (y \wedge k_2).$$

$$y S y \vee (x \wedge k_2) = (y \vee x) \wedge (y \vee k_2) = (y \vee x) \wedge (x \vee k_2) = x \vee (y \wedge k_2) S x.$$

Corollary. 1. Let a, b be two elements of a distributive lattice L , with $a \leq b$. The smallest congruence relation $R_{[a,b]}$ with the property that $[a,b]$ is one of its congruence classes, can be defined as follows:

$$x R_{[a,b]} y \text{ if and only if } a \wedge x = a \wedge y \text{ and } b \vee x = b \vee y.$$

2. Let I be a \vee -ideal of the distributive lattice L . The smallest congruence relation R_I which has I as one of its congruence classes can be defined as follows:

$$x R_I y \text{ if and only if there exists } i \in I \text{ such that } x \vee i = y \vee i.$$

Proof.

1. $a \wedge x = a \wedge y$ and $b \vee x = b \vee y$ is equivalent to the existence of two elements $c_1, c_2 \in [a,b]$ with $c_1 \wedge x = c_1 \wedge y$ and $c_2 \vee x = c_2 \vee y$. It can be verified directly that $[a,b]$ is a congruence class of $R_{[a,b]}$.

2. It is only necessary to prove that there exists $i_1 \in I$ with $i_1 \wedge x = i_1 \wedge y$. However, for each $i \in I$ we have $(i \wedge x \wedge y) \wedge x = (i \wedge x \wedge y) \wedge y$, and $i \wedge x \wedge y \in I$.

Remark: Grätzer and Schmidt [4] have given another definition of $R_{[a,b]}$, which requires a more complicated proof. Corollary 2 also occurs in [4], again with a more elaborate proof.

Theorem 3. Let L be a distributive lattice and let K be a sublattice of L . We define the relation θ_K as follows:

$$x \theta_K y \text{ if and only if for all } k_1, k_2 \in K : k_1 \wedge (x \vee k_2) = k_1 \wedge (y \vee k_2).$$

Then θ_K is a congruence relation such that different elements of K belong to different congruence classes of θ_K . If K is also convex, then θ_K is the largest congruence relation with this property.

Proof. It can be verified directly that θ_K is a congruence relation. Suppose $k_1 \theta_K k_2$ for some $k_1, k_2 \in K$. Then by the definition of θ_K :

$$k_1 \wedge (k_1 \vee k_2) = k_1 \wedge (k_2 \vee k_2) \text{ and}$$

$$k_2 \wedge (k_1 \vee k_1) = k_2 \wedge (k_2 \vee k_1).$$

Hence, $k_1 \leq k_2$ and $k_2 \leq k_1$. Thus, $k_1 = k_2$.

Suppose that K is also convex, and let θ^* be a congruence relation such that all elements of K belong to different congruence classes of θ^* . We prove that $\theta^* \leq \theta_K$. Let $x \theta^* y$. Then for all $k_1, k_2 \in K$:

$$k_1 \wedge (x \vee k_2) \theta^* k_1 \wedge (y \vee k_2).$$

Since $k_1 \wedge (x \vee k_2) \in K$ and $k_1 \wedge (y \vee k_2) \in K$ we have $k_1 \wedge (x \vee k_2) = k_1 \wedge (y \vee k_2)$, by the definition of θ^* .

This means that $x \theta_K y$.

Definition 5. Let L be a lattice. The zero element of the lattice of all congruence relations of L will be denoted by Ω , the unit element of this lattice will be denoted by \mathbf{U} .

Corollary. Let K be a sublattice of a distributive lattice. Let R_K and θ_K be defined as in theorems 2 and 3. Then $R_K \wedge \theta_K = \Omega$.

Proof. Suppose $x R_K \wedge \theta_K y$, i.e., $x R_K y$ and $x \theta_K y$ both hold.

From $x R_K y$ it follows that there exist $k_1, k_2 \in K$ such that $k_1 \wedge x = k_1 \wedge y$ and $k_2 \vee x = k_2 \vee y$. However, from $x \theta_K y$ we see that $k_2 \wedge (x \vee k_1) = k_2 \wedge (y \vee k_1)$. Also, $k_2 \vee (x \vee k_1) = k_2 \vee (y \vee k_1)$. Since L is distributive, we have $x \vee k_1 = y \vee k_1$. Together with $x \wedge k_1 = y \wedge k_1$, this yields $x = y$.

For the proof of theorem 5 we need a theorem of J. Hashimoto.

Definition 6. A lattice is called discrete if and only if all its intervals have finite length.

Theorem 4. The lattice of all congruence relations of a lattice L is a Boolean algebra if and only if L is distributive and discrete.

Proof. See [6], theorem 8.4.

Theorem 5. Let L be a distributive lattice and let $C \in \mathcal{C}(L)$, $C \neq \emptyset$. For $c \in C$, let C_c be the largest convex sublattice of L which has the intersection $\{c\}$ with C . Let the relation Γ_C be defined as follows:

$$x \Gamma_C y \text{ if and only if there exists } c \in C \text{ such that } x \in C_c \text{ and } y \in C_c.$$

Then:

1. $C_{c_1} \cap C_{c_2} = \emptyset$, if $c_1 \neq c_2$.
2. If $x \in C_{c_1}$ and $y \in C_{c_2}$, then $x \wedge y \in C_{c_1 \wedge c_2}$ and $x \vee y \in C_{c_1 \vee c_2}$.
3. If C is an interval then Γ_C is a congruence relation.
4. If Γ_C is a congruence relation, then Γ_C is equal to the congruence relation θ_C as introduced in theorem 3.
5. If L is also relatively complemented then the following two assertions are equivalent:
 - a) L is discrete.
 - b) Γ_C is a congruence relation for each $C \in \mathcal{C}(L)$.

Proof.

1. Since $C \cap C_{c_1} \cap C_{c_2} = \{c_1\} \cap \{c_2\} = \emptyset$, and since $C \cap C_{c_1} = \{c_1\}$, $C \cap C_{c_2} = \{c_2\}$, we conclude that $C_{c_1} \cap C_{c_2} = \emptyset$, by lemma 2.
2. Let $x \in C_{c_1}$, $y \in C_{c_2}$. We only prove that $x \wedge y \in C_{c_1 \wedge c_2}$. By theorem 1, there exist $s \in \{c_1\}_r \setminus (C \setminus \{c_1\}_1)_r$, and $t \in \{c_2\}_r \setminus (C \setminus \{c_2\}_1)_r$, such that $x \leq s$ and $y \leq t$. We show that $s \wedge t \in \{c_1 \wedge c_2\}_r \setminus (C \setminus \{c_1 \wedge c_2\}_1)_r$. Since $s \geq c_1$ and $t \geq c_2$, we have $s \wedge t \geq c_1 \wedge c_2$. Suppose $s \wedge t \in (C \setminus \{c_1 \wedge c_2\}_1)_r$. This means that there exists $\bar{c} \in C$ such that $s \wedge t \geq \bar{c}$, but $\bar{c} \notin \{c_1 \wedge c_2\}_1$. As in the proof of theorem 1, we have: $c_1 \geq \bar{c} \wedge s$ and $c_2 \geq \bar{c} \wedge t$; hence, $c_1 \wedge c_2 \geq \bar{c} \wedge s \wedge t = \bar{c}$, a contradiction. Thus, $s \wedge t \in \{c_1 \wedge c_2\}_r \setminus (C \setminus \{c_1 \wedge c_2\}_1)_r$. Since $x \leq s$ and $y \leq t$ we have $x \wedge y \leq s \wedge t$, whence $x \wedge y \in (\{c_1 \wedge c_2\}_r \setminus (C \setminus \{c_1 \wedge c_2\}_1)_r)_1$. Similarly, it can be shown that $x \wedge y \in (\{c_1 \wedge c_2\}_1 \setminus (C \setminus \{c_1 \wedge c_2\}_r)_1)_r$. We conclude that $x \wedge y \in C_{c_1 \wedge c_2}$.

3. Let C be an interval, say $C = \{x \in L \mid a \leq x \leq b\}$. By 1 and 2, in order to prove that Γ_C is a congruence relation, we only have to show that $\bigcup_{C \in \mathcal{C}} C = L$. By the maximality of the sets C_c , it is sufficient to show that for each $z \in L$ there exists a convex sublattice containing z , the intersection of which with C contains precisely one element. Let $D = \{y \in L \mid b \wedge z \leq y \leq a \vee z\}$. Then D has the required property: if $t \in C \wedge D$, then $a \leq t \leq b$ and $b \wedge z \leq t \leq a \vee z$; hence, $a \vee (b \wedge z) \leq t \leq b \wedge (a \vee z)$. Since L is distributive, we have $a \vee (b \wedge z) = t = b \wedge (a \vee z)$.

4. Let Γ_C be a congruence relation. Clearly, all elements of C belong to different congruence classes of Γ_C . By theorem 3, $\Gamma_C \leq \theta_C$. We prove that also $\theta_C \leq \Gamma_C$. Suppose $x \theta_C y$, and let $x \in C_{c_1}$, $y \in C_{c_2}$.

Since $c_1 \wedge (x \vee c_2) = c_1 \wedge (y \vee c_2)$, we have $C_{c_1 \wedge (c_1 \vee c_2)} = C_{c_1 \wedge (c_2 \vee c_2)}$, i.e., $c_1 \wedge (c_1 \vee c_2) = c_1 \wedge (c_2 \vee c_2)$ or $c_1 \leq c_2$.

Similarly, $c_2 \leq c_1$, from which $x \Gamma_C y$ follows.

5. Let L be distributive and relatively complemented.

a. Suppose L is discrete. We prove that Γ_C is a congruence relation for each $C \in \mathcal{C}(L)$. As in 3, it is sufficient to show that for each $x \in L$ there exists a convex sublattice C^* , containing x , which meets C in precisely one point. Let $x \in L$. Consider the congruence relation R_C . By theorem 4, R_C has a complement R_C^* . Let c be an arbitrary element of C . Then $x R_C \vee R_C^* c$. Since L is relatively complemented, we have $x R_C^* R_C c$, i.e., there exists $t \in L$ with $x R_C^* t$ and $t R_C c$. Let C^* be the congruence class of R_C^* which contains both x and t . It follows that $C \cap C^* = \{t\}$; hence, C^* has the desired property.

(We see that $R_C^* = \Gamma_C$; this can be shown as follows:

α . By the corollary of theorem 3, we have $R_C \wedge \Gamma_C = \Omega = R_C \wedge R_C^*$.

β . By 4, $R_C^* \leq \Gamma_C$. Since $R_C \vee R_C^* = \mathbf{1}$, we have $R_C \vee \Gamma_C = \mathbf{1} = R_C \vee R_C^*$.

γ . Since the lattice of all congruence relations of a lattice is distributive, we conclude that $\Gamma_C = R_C^*$).

b. Suppose that L is distributive and relatively complemented and that Γ_C is a congruence relation for each $C \in \mathcal{C}(L)$. We prove that L is discrete. By theorem 4, it is sufficient to prove that each

congruence relation of L has a complement. Let R be a congruence relation of L and let C be one of its congruence classes. Since in a distributive relatively complemented lattice each convex sublattice is congruence class of precisely one congruence relation [4] we have $R = R_C$. We show that Γ_C is the complement of R_C . $\Gamma_C \wedge R_C = \Omega$ was proved already. Let x, y be two arbitrary elements of L , and suppose $x \in C_{c_1}$, $y \in C_{c_2}$, with $c_1, c_2 \in C$. Then $x \Gamma_C c_1$, $c_1 R_C c_2$ and $c_2 \Gamma_C y$; hence, $x R_C \vee \Gamma_C y$, from which we conclude that $R_C \vee \Gamma_C = \sqcup$, i.e., Γ_C is the complement of $R_C = R$.

4. The lattice $(\mathcal{C}(L), \subset)$

Let L be a lattice. In this section we study some properties of the lattice $(\mathcal{C}(L), \subset)$ i.e., the lattice of all convex sublattices of L , partially ordered by inclusion. The join operation in $(\mathcal{C}(L), \subset)$ is denoted by \sqcup (definition 4.).

Lemma 4. Let L be a distributive lattice, and let $A, B, C \in \mathcal{C}(L)$.

Then:

1. If $A \cap B \neq \emptyset$ and $A \cap C \neq \emptyset$, then $A \cap (B \sqcup C) = (A \cap B) \sqcup (A \cap C)$.
2. If $B \cap C \neq \emptyset$, then $A \sqcup (B \cap C) = (A \sqcup B) \cap (A \sqcup C)$.

Proof.

1. Clearly, $A \cap (B \sqcup C) \supset (A \cap B) \sqcup (A \cap C)$. In order to prove that $A \cap (B \sqcup C) \subset (A \cap B) \sqcup (A \cap C)$, assume that $a \in A$ and $a \in B \sqcup C$. This means that there exist $b_1, b_2 \in B$ and $c_1, c_2 \in C$ such that $b_1 \wedge c_1 \leq a \leq b_2 \vee c_2$. Let $s \in A \cap B$ and $t \in A \cap C$. Then: $a \leq a \vee s \vee t = (a \wedge b_2) \vee (a \wedge c_2) \vee s \vee t$. However, $(a \wedge b_2) \vee s \in A \cap B$ and $(a \wedge c_2) \vee t \in A \cap C$. Thus, $a \in \{(A \cap B) \vee (A \cap C)\}_1$. Similarly, $a \in \{(A \cap B) \wedge (A \cap C)\}_r$. This proves that $a \in (A \cap B) \sqcup (A \cap C)$.
2. Similar to part 1.

Theorem 6. Let L be a distributive relatively complemented lattice. Let $C, D \in \mathcal{C}(L)$ with $C \subset D$. There exists $C' \in \mathcal{C}(L)$ such that $C \cap C' = \emptyset$, $C \sqcup C' = D$.

Proof.

1. First we prove that for each $C \in \mathcal{C}(L)$ there exists C' such that $C \cap C' = \emptyset$ and $C \sqcup C' = L$. If $C = L$ then $C' = \emptyset$. Otherwise, let $x \in L \setminus C$. Application of lemma 3 to the disjoint convex sublattices C and $\{x\}$ yields a prime ideal I , say a \vee -ideal, such that $C \cap I = \emptyset$. Since L is relatively complemented, I is maximal. We prove that $C \sqcup I = L$. $I \subset C \sqcup I \subset (C \sqcup I)_r$; hence, $I_r = L \subset (C \sqcup I)_r$. Thus, $(C \sqcup I)_r = L$, i.e. $C \sqcup I = (C \sqcup I)_1$. Since $I \subset (C \sqcup I)_1$ and since I is maximal, we have $(C \sqcup I)_1 = I$ or $(C \sqcup I)_1 = L$. $(C \sqcup I)_1 = I$ contradicts $C \cap I = \emptyset$. We conclude therefore that $(C \sqcup I)_1 = C \sqcup I = L$.

2. Let $C, D \in \mathcal{C}(L)$ with $C \subset D$. Since D is a convex sublattice, D is a relatively complemented (and distributive) lattice. We can therefore apply part 1, which yields a set C' such that:

- a. $C \cap C' = \emptyset$.
- b. The smallest convex sublattice of D that contains C and C' is D .
- c. C' is a \vee -ideal of D .

From b. it follows that $C \sqcup C' = D$ (since each convex sublattice of L which is contained in D is a convex sublattice of D). Also, C' is a convex sublattice of L : It is clear that C' is a sublattice.

Suppose that $c'_1 \leq x \leq c'_2$, for some $c'_1, c'_2 \in C'$, $x \in L$. Since $c'_1, c'_2 \in C \subset D$, we have $x \in D$. Together with the fact that C' is a \vee -ideal of D and $x \leq c'_2$, this gives $x \in C'$; hence, C' is convex.

Theorem 6 asserts that if L is distributive relatively complemented lattice then each interval $[\emptyset, C]$ of $(\mathcal{C}(L), \subset)$ is complemented.

Theorem 8 shows that an extra condition is necessary (and sufficient) in order that each interval $[C, D]$ of $(\mathcal{C}(L), \subset)$ be complemented (i.e., in order that $(\mathcal{C}(L), \subset)$ be relatively complemented).

For the proof of theorem 8 we need the following theorem of J. Hashimoto:

Theorem 7. The lattice of all \vee -ideals (\wedge -ideals) of a lattice L is distributive and relatively complemented if and only if L is distributive, relatively complemented and discrete.

Proof. See [6], theorem 4.3.

Theorem 8.

1. Let L be a distributive lattice $(\mathcal{C}(L), \subset)$ is relatively complemented if and only if L is relatively complemented and discrete.
2. Let L be a distributive lattice. Let $A, B, C \in \mathcal{C}(L)$ with $A \subset B \subset C$; $A \neq \emptyset$. Then: B has at most one complement in $[A, C]$.

Proof.

- 1.1. Suppose L is distributive, relatively complemented and discrete.

Let A, B, C be elements of $\mathcal{C}(L)$ with $A \subset B \subset C$. We prove that there exists $B^* \in \mathcal{C}(L)$ such that $B \cap B^* = A$ and $B \cup B^* = C$. We may assume that $A \neq \emptyset$, since the case that $A = \emptyset$ was already treated in theorem 6. $A \subset B \subset C$ implies $A_1 \subset B_1 \subset C_1$ and $A_r \subset B_r \subset C_r$. Let $\mathfrak{V}(L)$ be the family of all \vee -ideals of L and $\mathfrak{J}(L)$ the family of all \wedge -ideals. By theorem 7, $(\mathfrak{V}(L), \subset)$ and $(\mathfrak{J}(L), \subset)$ are relatively complemented.

Therefore, there exists $B_1^* \in \mathfrak{V}(L)$ such that $B_1 \cap B_1^* = A_1$ and such that C_1 is the smallest \vee -ideal that contains B_1 and B_1^* . Since L is distributive, this means that $B_1 \vee B_1^* = C_1$. Similarly, there exists $B_r^* \in \mathfrak{J}(L)$ such that $B_r \cap B_r^* = A_r$ and $B_r \wedge B_r^* = C_r$. We prove that $B_1^* \cap B_r^*$ is the relative complement of B in the interval $[A, C]$.

Clearly, $B_1 \cap B_r \cap B_1^* \cap B_r^* = A_1 \cap A_r = A$. Also, $B \cup (B_1^* \cap B_r^*) = \{B \wedge (B_1^* \cap B_r^*)\}_r \cup \{B \vee (B_1^* \cap B_r^*)\}_1 = (B_r \wedge (B_1^* \cap B_r^*)) \cap (B_1 \vee (B_1^* \cap B_r^*)) = (B_r \wedge B_r^*) \cap (B_1 \vee B_1^*) = C_r \cap C_1 = C$.

- 1.2. Let L be distributive and suppose that $(\mathcal{C}(L), \subset)$ is relatively complemented. We show that then $\mathfrak{J}(L)$ is also relatively complemented. Theorem 7 then gives the desired result. Let $I_1 \subset I_2 \subset I_3$ be three elements of $\mathfrak{J}(L)$. There exists $C \in \mathcal{C}(L)$ such that $C \cap I_2 = I_1$, $C \cup I_2 = I_3$. Since $I_1 \subset C$ we have $I_1 \subset C_r$; hence, $I_{12} = L \subset C_r$. This means that $C = C_1$; i.e., C is a \vee -ideal, from which we conclude that $\mathfrak{J}(L)$ is relatively complemented.

2. Let L be distributive, let $A \subset B \subset C \in \mathcal{G}(L)$ with $A \neq \emptyset$, and suppose that B has two relative complements B_1^* and B_2^* in $[A, C]$. As above, it follows that B_{11}^* and B_{21}^* are two relative complements (in $\mathcal{M}(L)$) of B_1 in the interval $[A_1, C_1]$. Since $\mathcal{M}(L)$ is distributive, we have $B_{11}^* = B_{21}^*$. Similarly, $B_{1r}^* = B_{2r}^*$, whence $B_1^* = B_2^*$.

Remark: In the assertion that complementation in each interval $[A, C]$ of $\mathcal{G}(L)$ is unique (for L distributive), we may not omit the condition that $A \neq \emptyset$. This can be seen as follows: Suppose that complementation in the whole of $\mathcal{G}(L)$ is unique, for L distributive. If L is also relatively complemented and discrete, we would have: $(\mathcal{G}(L), \subset)$ is a lattice in which complements always exist and unique. Together with the atomicity of $(\mathcal{G}(L), \subset)$ this would give the result that $(\mathcal{G}(L), \subset)$ is distributive ([7], p. 57), which is clearly not the case.

Corollary 1. Let L be a distributive lattice, let $A, B \in \mathcal{G}(L)$ with $A \subset B$, $A \neq \emptyset$, and let $C(A, B)$ be the largest element of $\mathcal{G}(L)$ that has the intersection A with B (theorem 1). Then $B \sqcup C(A, B) = L$ for all A, B , if and only if L is relatively complemented and discrete.

Proof.

1. Let L be distributive relatively complemented and discrete. Let $A \subset B$, $A \neq \emptyset$ ($A, B \in \mathcal{G}(L)$), and let B^* be the complement of B in the interval $[A, L]$. Then $B \cap B^* = A$. By the definition of $C(A, B)$: $B^* \subset C(A, B)$; hence, $B \sqcup C(A, B) \supset B \cup B^* = L$.
Thus, $B \sqcup C(A, B) = L$.
2. Let L be distributive and suppose that for each $A, B \in \mathcal{G}(L)$ with $A \subset B$, $A \neq \emptyset$, we have $B \sqcup C(A, B) = L$. In particular, if I and H are two \vee -ideals of L with $I \subset H$, we have $I \sqcup C(I, H) = L$. By theorem 1, $C(I, H) = (I_1 \setminus (H \setminus I_r)_1)_r \cap (I_r \setminus (H \setminus I_1)_r)_1 = (I \setminus (H \setminus L)_1)_r \cap (L \setminus (H \setminus I)_r)_1 = L \cap (L \setminus (H \setminus I)_r)_1 = (L \setminus (H \setminus I)_r)_1$. Thus, $C(I, H)$ is a \vee -ideal and we see that each interval $[I, L]$ of $\mathcal{M}(L)$ is complemented. Since (L) and $\mathcal{M}(L)$ is distributive, $\mathcal{M}(L)$ is relatively complemented. By theorem 7, L is then relatively complemented and discrete.

Corollary 2. Let L be a distributive relatively complemented and discrete lattice, let $A, B \in \overline{\mathcal{C}}(L)$ with $A \subset B$, $A \neq \emptyset$. Let $C(A, B)$ be defined as in corollary 1. Then we have: $C(A, C(A, B)) = B$.

Proof. By corollary 1, we have

$$B \cap C(A, B) = A \quad \text{and} \quad B \cup C(A, B) = L,$$

$$C(A, C(A, B)) \cap C(A, B) = A \quad \text{and} \quad C(A, C(A, B)) \cup C(A, B) = L.$$

Uniqueness of complementation in $[A, L]$ yields $B = C(A, C(A, B))$.

5. The lattice $(\overline{\mathcal{C}}(L), \leq)$

In this section we study a partial ordering on $\overline{\mathcal{C}}(L)$ which is a variant of the ordering by inclusion. $\overline{\mathcal{C}}(L)$ is used to denote the family of all non-empty convex sublattices of L .

Definition 7. Let L be a lattice and let $C, D \in \overline{\mathcal{C}}(L)$. We define the partial ordering \leq as follows:

$$C \leq D \text{ if and only if } C \subset D_l \text{ and } D \subset C_r.$$

Lemma 5. \leq is a partial ordering on $\overline{\mathcal{C}}(L)$.

Proof. We prove only anti-symmetry. Let $C, D \in \overline{\mathcal{C}}(L)$, with $C \leq D$ and $D \leq C$. Then $C \subset D_l$, $D \subset C_r$, $D \subset C_l$ and $C \subset D_r$. Hence, $C \subset D_l \cap D_r = D$ and $D \subset C_l \cap C_r = C$, which gives $C = D$.

Lemma 6. Let $C \in \overline{\mathcal{C}}(L)$. C is a \vee -ideal (\wedge -ideal) of L if and only if $C \leq L$ ($C \geq L$).

Proof.

1. From $C \leq L$ we see that $L \subset C_r$, whence $C = C_l \cap L = C_l$.
2. Let I be a \vee -ideal. Clearly, $I \subset L_l = L$. Also, $L \subset I_r$, since $L = I_l \cap I_r = I_r$.

Lemma 7. Let $C, D \in \overline{\mathcal{C}}(L)$. Then $C \leq D$ if and only if $C \wedge D = C$ ($C \vee D = D$).

Proof. Follows directly from the definitions.

Theorem 9. Let L be a distributive lattice. Then $(\bar{C}(L), \leq)$ is a distributive lattice.

Proof.

1. $C, D \in \bar{C}(L)$ implies $C \wedge D \in \bar{C}(L)$:

a. Clearly, $(c_1 \wedge d_1) \wedge (c_2 \wedge d_2) \in C \wedge D$.

b. $(c_1 \wedge d_1) \vee (c_2 \wedge d_2) = \{c_1 \vee (c_2 \wedge d_2)\} \wedge \{d_1 \vee (c_2 \wedge d_2)\} \in C \wedge D$.

c. Suppose $c_1 \wedge d_1 \leq x \leq c_2 \wedge d_2$, for some $x \in L$. Then:

$c_1 \leq x \vee c_1 \leq c_1 \vee (c_2 \wedge d_2)$; hence, $x \vee c_1 \in C$. Also, $x \vee d_1 \in D$,
whence $x = x \vee (c_1 \wedge d_1) = (x \vee c_1) \wedge (x \vee d_1) \in C \wedge D$.

2. Similarly, $C \vee D \in \bar{C}(L)$.

3. The commutative, associative and absorption laws follow directly.

4. Distributivity is proved by showing that, for $C, D, E \in \bar{C}(L)$:

$$C \wedge (D \vee E) = (C \wedge D) \vee (C \wedge E).$$

a. It is clear that $C \wedge (D \vee E) \subset (C \wedge D) \vee (C \wedge E)$.

b. Let $(c_1 \wedge d) \vee (c_2 \wedge e) \in (C \wedge D) \vee (C \wedge E)$. Then:

$$\begin{aligned} (c_1 \wedge d) \vee (c_2 \wedge e) &= \{(c_1 \wedge d) \vee c_2\} \wedge \{(c_1 \wedge d) \vee e\} = \\ c_3 \wedge \{(c_1 \wedge d) \vee e\} &= c_3 \wedge \{(c_1 \vee e) \wedge (d \vee e)\} = c_4 \wedge (d \vee e) \in C \wedge (D \vee E). \end{aligned}$$

Corollary. Let L be a distributive lattice. Then $(\mathcal{M}(L), \leq)$ is a \vee -ideal of $(\bar{C}(L), \leq)$ and $(\mathcal{M}(L), \leq)$ is a \wedge -ideal of $(\bar{C}(L), \leq)$.

Proof. Follows from lemma 6 and theorem 9.

In the remainder of this section we shall omit indication of the partial ordering \leq on $\bar{C}(L)$, i.e., when we write $\bar{C}(L)$, we mean $(\bar{C}(L), \leq)$.

Theorem 10. Let F_i ($i \geq 0$) be the free distributive lattice with i generators, with an (extra) zero and unit element adjoined. Let B_j ($j \geq 1$) be the Boolean algebra with 2^j elements. For L distributive, we define $\bar{C}^0(L) = L$ and $\bar{C}^i(L) = \bar{C}(\bar{C}^{i-1}(L))$, ($i \geq 0$). Then we have: $\bar{C}^i(B_j)$ is isomorphic with the direct union of j factors F_i (cf. [1], chapter IX, section 10).

Proof. We use induction on i .

1. $\bar{C}^0(B_j)$ is clearly isomorphic with the direct union of j factors F_0 ,

- since $F_0 \cong B_1$ 1)
2. Suppose $\bar{C}^i(B_j) \cong F_i^j$ (The direct union of two lattices L_1, L_2 is denoted by $L_1 \times L_2$; the direct union of j factors L is denoted by L^j). In order to prove that $\bar{C}^{i+1}(B_j) \cong F_{i+1}^{j+1}$, we have to prove that $\bar{C}(F_i^j) \cong F_{i+1}^j$. However, it is easy to verify that for two finite distributive lattices L_1, L_2 we have $\bar{C}(L_1 \times L_2) \cong \bar{C}(L_1) \times \bar{C}(L_2)$. Therefore, there remains the proof of $\bar{C}(F_i) \cong F_{i+1}$. Let $C = \{f_i \in F_i \mid a \leq f_i \leq b\}$ be an element of $\bar{C}(F_i)$, where a and b are finite joins of meets of the generators, say x_1, x_2, \dots, x_i , of F_i . (Verification of the following argument in the case that a or b is the zero or unit element of F_i is straight forward and is therefore omitted). We define the isomorphism $\psi: \bar{C}(F_i) \rightarrow F_{i+1}$ as follows: We introduce $y (\neq x_1, x_2, \dots, x_i)$ as the $i+1$ -th generator of F_{i+1} . Consider the element $(b \wedge y) \vee a$ of F_{i+1} . It may be possible to "reduce" this element: E.g., let $b = x_1 \vee x_2$, and $a = x_1$. Then $(b \wedge y) \vee a = ((x_1 \vee x_2) \wedge y) \vee x_1$ can be reduced to $(x_2 \wedge y) \vee x_1$. Clearly, however, each element $(b \wedge y) \vee a$ has an "irreducible" form. From now on we assume that all elements of F_{i+1} are in reduced form. We then define $\psi(C)$ as $(b \wedge y) \vee a$. We prove that ψ is an isomorphism:
- Let $C_1 = \{f_i \in F_i \mid a \leq f_i \leq b\}$ and $C_2 = \{f_i \in F_i \mid c \leq f_i \leq d\}$. Then:
- $$C_1 \wedge C_2 = \{f_i \in F_i \mid a \wedge c \leq f_i \leq b \wedge d\}, \text{ and}$$
- $$C_1 \vee C_2 = \{f_i \in F_i \mid a \vee c \leq f_i \leq b \vee d\}.$$
- $$\psi(C_1) \wedge \psi(C_2) = \{(b \wedge y) \vee a\} \wedge \{(d \wedge y) \vee c\} =$$
- $$(b \wedge d \wedge y) \vee (a \wedge d \wedge y) \vee (b \wedge c \wedge y) \vee (a \wedge c) = (b \wedge d \wedge y) \vee (a \wedge c) = \psi(C_1 \wedge C_2)$$
- $$\psi(C_1) \vee \psi(C_2) = \{(b \wedge y) \vee a\} \vee \{(d \wedge y) \vee c\} = \{(b \vee d) \wedge y\} \vee (a \vee c) = \psi(C_1 \vee C_2)$$
- Suppose $\psi(C_1) = \psi(C_2)$. This means that $(b \wedge y) \vee a = (d \wedge y) \vee c$. From the irreducibility of $(b \wedge y) \vee a$ and $(d \wedge y) \vee c$, it follows that $a = c$ and $b = d$. Hence, ψ is 1-1.
- Also, ψ is onto: Each element of F_{i+1} can be written as $(a \wedge y) \vee b$, with $a, b \in F_i$. However, $(a \wedge y) \vee b$ is the image of the convex sublattice $\{f_i \in F_i \mid b \leq f_i \leq a \vee b\}$ of F_i . This follows from $\{(a \vee b) \wedge y\} \vee b = (a \wedge y) \vee b$.
- This completes the proof of theorem 10.

1)

\cong is used to denote isomorphism.

Remark: Let K_n be the chain of n elements. We state without proof the following formulae:

Let $\gamma_n^{(i)}$ be the number of elements of $\bar{C}^{(i)}(K_n)$, $i=1,2,3$.

$$\text{Then } \gamma_n^{(1)} = \sum_{i=1}^n (n-i+1) = \frac{1}{2}n(n+1),$$

$$\gamma_n^{(2)} = \sum_{i=1}^n (n-i+1)i^2 = \frac{1}{12}n(n+1)^2(n+2)$$

$$\gamma_n^{(3)} = \frac{1}{120} \sum_{i=1}^n (n-i+1)(8i^6 + 24i^5 + 35i^4 + 30i^3 + 17i^2 + 6i).$$

6. A ternary function in distributive relatively complemented lattices

Theorem 11. Let L be a distributive relatively complemented lattice.

Let $f: L^3 \rightarrow L$ be defined as follows: $f(a,b,c)$ is the relative complement of a in the interval $[a \wedge b, a \vee c]$. Then we have:

A subset C of L is an element of $\bar{C}(L)$ if and only if $f(L,C,C) \subset C$.

Proof.

1. Suppose that $f(L,C,C) \subset C$. Clearly, $f(c_1, c_2, c_1) = c_1 \wedge c_2 \in C$, and $f(c_1, c_1, c_2) = c_1 \vee c_2 \in C$; hence, C is a sublattice. Also, if $c_1 \leq x \leq c_2$, then $x = f(x, c_2, c_1) \in C$.

2. Suppose $C \in \bar{C}(L)$. Let $f(x, c_1, c_2)$ be an element of $f(L,C,C)$.

We prove that $f(x, c_1, c_2)$ (abbreviated to x^*) is an element of C .

We have $x \wedge x^* = x \wedge c_1$ and $x \vee x^* = x \vee c_2$.

Hence,

$$\begin{aligned} x^* &= x^* \vee (x \wedge c_1) = (x^* \vee x) \wedge (x^* \vee c_1) = (x \vee c_2) \wedge (x^* \vee c_1) = \\ &= \{x \wedge (x^* \vee c_1)\} \vee \{c_2 \wedge (x^* \vee c_1)\} = \{(x \wedge x^*) \vee (x \wedge c_1)\} \vee c_3 = \\ &= (x \wedge c_1) \vee c_3 \in C. \end{aligned}$$

Theorem 12. A set L is a distributive relatively complemented lattice if and only if there exists a function $f: L^3 \rightarrow L$ with the following properties: For all $a,b,c,d,e \in L$:

P1. $f(a,a,a) = a$.

P2.1. $f(a,b,a) = f(b,a,b)$

P2.2. $f(a,a,b) = f(b,b,a)$.

$$P3. \quad f(a, f(a, b, c), f(a, d, e)) = f(a, b, e),$$

$$P4.1. \quad f(a, f(b, b, c), a) = f(f(a, b, a), f(a, b, a), f(a, c, a)).$$

$$P4.2. \quad f(a, a, f(b, b, c)) = f(f(a, a, b), f(a, a, b), f(a, a, c)).$$

Proof.

1. The condition is sufficient.

$$\text{We define } a \wedge b = f(a, b, a) \text{ and } a \vee b = f(a, a, b).$$

1.1. The commutativity of \wedge and \vee follows from P2.

$$\begin{aligned} 1.2. \quad a \wedge (a \vee b) &= f(a, f(a, a, b), a) = f(a, f(a, a, b), f(a, a, a)) = \\ &= f(a, a, a) = a, \text{ by P1, P3 and P1.} \end{aligned}$$

$$\text{Similarly, } a \vee (a \wedge b) = a.$$

1.3. In order to prove that $a \wedge (b \wedge c) = (a \wedge b) \wedge c$, we have to show that:

$$f(a, f(b, c, b), a) = f(f(a, b, a), c, f(a, b, a)).$$

$$\text{Let } A = f(a, f(b, c, b), a) \text{ and } B = f(f(a, b, a), c, f(a, b, a)).$$

First we prove that $a \wedge A = a \wedge B$ and $a \vee A = a \vee B$:

$$f(a, A, a) = f(a, f(a, f(b, c, b), a), a) = f(a, f(b, c, b), a) \text{ by P3 and P1.}$$

$$\begin{aligned} f(a, B, a) &= f(a, f(a, b, a), c, f(a, b, a)), a) \\ &= f(f(a, f(a, b, a), a), f(a, c, a), f(a, f(a, b, a), a)) \\ &= f(f(a, b, a), f(a, c, a), f(a, b, a)) \\ &= f(a, f(b, c, b), a) \text{ by P4.1, P1 and P3, and P4.1.} \end{aligned}$$

$$f(a, a, A) = f(a, a, f(a, f(b, c, b), a)) = a, \text{ by P1 and P3.}$$

$$\begin{aligned} f(a, a, B) &= f(a, a, f(f(a, b, a), c, f(a, b, a))) \\ &= f(f(a, a, f(a, b, a)), f(a, a, c), f(a, a, f(a, b, a))) \\ &= f(a, f(a, a, c), a) = a, \text{ by P4.2, P1 and P3.} \end{aligned}$$

The rest of the proof that $A = B$ is standard:

$$\begin{aligned} A &= A \vee (A \wedge a) = A \vee (B \wedge a) = (A \vee B) \wedge (A \vee a) \\ &= (A \vee B) \wedge (B \vee a) = B \vee (a \wedge A) = B \vee (a \wedge B) = B, \end{aligned}$$

by application of P2 and P4.2.

For the proof of $(a \vee b) \vee c = a \vee (b \vee c)$ we need the dual equalities of P4.1 and P4.2, i.e:

$$(1) \quad f(a, f(b, b, c), a) = f(f(a, b, a), f(a, b, a), f(a, c, a)) \text{ and}$$

$$(2) \quad f(a, a, f(b, b, c)) = f(f(a, a, b), f(a, a, b), f(a, a, c)).$$

(1) is established as usual:

$$\begin{aligned} (a \wedge b) \vee (a \wedge c) &= \{(a \wedge b) \vee a\} \wedge \{(a \wedge b) \vee c\} \\ &= a \wedge \{(a \wedge b) \vee c\} = a \wedge \{(a \vee c) \wedge (b \vee c)\} \\ &= \{a \wedge (a \vee c)\} \wedge (b \vee c) = a \wedge (b \vee c), \end{aligned}$$

by the associativity of \wedge and P4.2.

To prove (2), we consider:

$$\begin{aligned} \{a \vee (b \vee c)\} \wedge \{(a \vee b) \vee (a \vee c)\} &= \\ [\{a \wedge \{(a \vee b) \vee (a \vee c)\}\}] \vee [(b \vee c) \wedge \{(a \vee b) \vee (a \vee c)\}] &= \\ [\{a \wedge (a \vee b)\} \vee \{a \wedge (a \vee c)\}] \vee [(b \wedge \{(a \vee b) \vee (a \vee c)\}) \vee \\ (c \wedge \{(a \vee b) \vee (a \vee c)\})] &= a \vee [\{(b \wedge (a \vee b)) \vee \{b \wedge (a \vee c)\}\} \\ \vee \{(c \wedge (a \vee b)) \vee \{c \wedge (a \vee c)\}\}] &= a \vee [(b \vee \{b \wedge (a \vee c)\}) \vee \\ (\{c \wedge (a \vee b)\} \vee c)] &= a \vee (b \vee c) \end{aligned}$$

and:

$$\begin{aligned} \{a \vee (b \vee c)\} \wedge \{(a \vee b) \vee (a \vee c)\} &= \\ [\{a \vee (b \vee c)\} \wedge (a \vee b)] \vee [\{a \vee (b \vee c)\} \wedge (a \vee c)] &= \\ [a \vee \{(b \vee c) \wedge b\}] \vee [a \vee \{(b \vee c) \wedge c\}] &= \\ (a \vee b) \vee (a \vee c). \end{aligned}$$

Hence, $a \vee (b \vee c) = (a \vee b) \vee (a \vee c)$.

Finally, the proof of the associativity of \vee is now dual to the proof of the associativity of \wedge .

1.4. The distributivity of L follows from P4.2 and (1).

1.5. Let $a \leq c \leq b$. Then:

$$c \wedge f(c, a, b) = f(c, f(c, a, b), c) = f(c, a, c) = c \wedge a = a \text{ and}$$

$$c \vee f(c, a, b) = f(c, c, f(c, a, b)) = f(c, c, b) = c \vee b = b.$$

Thus, $f(c, a, b)$ is the relative complement of c in the interval $[a, b]$.

2. The condition is necessary.

Let L be a distributive relatively complemented lattice. Let $f(a, b, c)$ be the relative complement of a in the interval $[a \wedge b, a \vee c]$.

Then $f(a, b, c)$ has the properties P1 to P4.

We prove only P2.1 and P3.

Clearly, the relative complement of a in the interval $[a \wedge b, a]$ is $a \wedge b$. Thus, $f(a, b, a) = a \wedge b = b \wedge a = f(b, a, b)$.

Furthermore, by the definition of f ,

$$a \wedge f(a, f(a,b,c), f(a,d,e)) = a \wedge f(a,b,c) = a \wedge b \text{ and}$$

$$a \wedge f(a,b,e) = a \wedge b.$$

$$a \vee f(a, f(a,b,c), f(a,d,e)) = a \vee f(a,d,e) = a \vee e \text{ and}$$

$$a \vee f(a,b,e) = a \vee e.$$

$$\text{Hence, } f(a, f(a,b,c), f(a,d,e)) = f(a,b,e).$$

This completes the proof of theorem 12.

Finally we mention some properties of the function f that can be verified directly from its definition:

$$P5. \quad f(b,a,a) = a.$$

$$P6. \quad f(a,b,f(a,b,c)) = f(a, f(a,b,c), c) = f(f(f(a,b,c),b,c),b,c) = f(a,b,c).$$

$$P7. \quad f(f(a,b,c),b,c) = (a \wedge b) \vee (b \wedge c) \vee (c \wedge a).$$

$$P8. \quad f(a, f(b,c,d), f(b,e,g)) = f(b, f(a,c,e), f(a,d,g)).$$

Remarks:

1. The function f has been used to define Boolean algebra's and distributive relatively complemented lattices with zero in [3].
2. From P7 we see that the function f is related to the well-known ternary function $(a \wedge b) \vee (b \wedge c) \vee (c \wedge a)$, which has been used for the axiomatics of distributive lattices by several authors (these investigations started with [5]; for recent results see [9]).
3. From P3, P5 and P8 we see that f is one of the "selection functions" as studied in [2]. In particular, if L is the Boolean algebra with two elements, then f coincides with the "conditional Boolean expression" if a then b else c , as used in the programming language ALGOL 60 [8].

References

1. G. Birkhoff Lattice Theory.
American Math. Society, New York, 1948.
2. A. Caraccialo di N-ary selection functions and formal selec-
Forino tive systems. Part I.
Calcolo, 1964, vol. 1, pp 49-81.
3. R.M. Dicker A set of independent postulates for Boolean
algebra.
Proc. London Math. Soc. (3), 1963, vol. 3,
pp 20-30.
4. G. Grätzer and Ideals and congruence relations in lattices.
E.T. Schmidt Acta Math. Acad. Sci. Hung., 1958, vol. 9,
pp 137-175.
5. A.A. Grau Ternary Boolean algebra.
Bull. Amer. Math. Soc. 1947, vol. 53,
pp 567-572.
6. J. Hashimoto Ideal Theory for Lattices.
Math. Japonicae, 1952, vol. 2, pp 149-185.
7. H. Hermes Einführung in die Verbandstheorie.
Springer-Verlag, Berlin, 1955.
8. P. Naur (Ed.) Revised Report on the Algorithmic Language
ALGOL 60.
Regnecentralen, Copenhagen, 1962.
9. B. Sobocinski Six new sets of independent axioms for
distributive lattices with 0 and I.
Notre Dame J. of Formal logic, 1962, vol. 3,
pp 187-192
10. M.H. Stone Topological representations of distributive
lattices and Brouwerian logics.
Casopis Pest. Mat. Fyz. 1937-'38. vol. 67,
pp 1-25.