
**stichting
mathematisch
centrum**



AFDELING ZUIVERE WISKUNDE
(DEPARTMENT OF PURE MATHEMATICS)

ZW 135/80

JANUARI

A.M. COHEN & H.A. WILBRINK

THE STABILIZER OF DYE'S SPREAD ON A HYPERBOLIC QUADRIC
IN $PG(4n-1,2)$ WITHIN THE ORTHOGONAL GROUP

Preprint

2e boerhaavestraat 49 amsterdam

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).

The stabilizer of Dye's spread on a hyperbolic quadric in $PG(4n-1,2)$ within the orthogonal group^{*)}

by

A.M. Cohen & H.A. Wilbrink

ABSTRACT

Recently, R.H. DYE [3] constructed spreads as indicated in the title. He determined their stabilizers within the relevant orthogonal group in the cases $n = 2, 3$. The present note deals with all $n \geq 3$. Use is made of Holt's characterisation of certain triply transitive permutation groups of degree $2^{2n-1} + 1$.

KEY WORDS & PHRASES: *finite classical geometry, spreads, permutation groups*

^{*)}

This report will be submitted for publication elsewhere.

1. INTRODUCTION

The projective space $PG(4n-1,2)$ is viewed in the usual way as the incidence structure of 1- and 2-dimensional subspaces of the vector space \mathbb{F}_2^{4n} . The hyperbolic quadric Ω will be fixed as the set of projective points X in $PG(4n-1,2)$ whose homogenous coordinates $(X_1, X_2, \dots, X_{4n})$ satisfy

$$q(X) = X_1X_2 + X_3X_4 + \dots + X_{4n-1}X_{4n} = 0.$$

The hyperbolic quadratic form q on $PG(4n-1,2)$ admits a symplectic polarity that we shall denote by B . A spread on the quadric Ω is defined to be a partitioning $S = \{S_1, \dots, S_{2^{2n-1}+1}\}$ of Ω into $2^{2n-1}+1$ projective $(2n-1)$ -dimensional totally isotropic subspaces of $(PG(4n-1,2), q)$.

2. CONSTRUCTION OF THE SPREAD

The following construction of a spread on Ω is to be found in [3]. Fix a nonisotropic point P and an isotropic point Q of $(PG(4n-1,2), q)$ such that $B(P, Q) \neq 0$. Then the projective space H underlying $P^\perp \cap Q^\perp$ is a $PG(4n-3,2)$ with symplectic polarity B_0 induced by B . By means of scalar restriction from the Galois field $\mathbb{F}_{2^{2n-1}}$ to \mathbb{F}_2 , the projective line $PG(1, 2^{2n-1})$ with nondegenerate symplectic polarity B_1 can be regarded as a $PG(4n-3,2)$ with nondegenerate symplectic polarity $\text{trace}_{\mathbb{F}_{2^{2n-1}}|\mathbb{F}_2} \circ B_1$. Thus (H, B_0) can be identified with $(PG(1, 2^{2n-1}), \text{trace}_{\mathbb{F}_{2^{2n-1}}|\mathbb{F}_2} \circ B_1)$ whenever the latter is viewed as a projective space over \mathbb{F}_2 . Under this identification, the points of $PG(1, 2^{2n-1})$ correspond to totally isotropic $(2n-2)$ -dimensional subspaces of (H, B_0) partitioning H . Next, H is mapped bijectively onto $P^\perp \cap \Omega$ by means of projection from P . Note that totally isotropic subspaces of (H, B_0) map into totally isotropic subspaces of $(P^\perp, q|_{P^\perp})$ inside Ω , so that the partitioning of (H, B_0) maps onto a partitioning of $P^\perp \cap \Omega$ into totally isotropic subspaces. In order to obtain a spread, note that each of these $(2n-2)$ -dimensional subspaces should be extended to a maximal totally isotropic subspace of $(PG(4n-1,2), q)$. It follows from [2] that this can be done in precisely two different

ways such that no two subspaces intersect. The two resulting spreads on Ω are mapped into one another by the symmetry with center P . Moreover, the subspaces belonging to one of these two spreads are all in the same $\Omega_{4n}^+(2)$ -orbit, where $\Omega_{4n}^+(2)$ stands for the commutator subgroup of the orthogonal group $O_{4n}^+(2)$ with respect to q . Hence, the spread is uniquely determined by the requirement that its elements are maximal totally isotropic subspaces from a fixed $\Omega_{4n}^+(2)$ -orbit. The spread thus constructed will be denoted \mathcal{P} .

3. THE STABILIZER OF THE SPREAD

Let G denote the stabilizer of the spread \mathcal{P} within $O_{4n}^+(2)$ and let G_R for R a point of $PG(4n-1, 2)$ stand for the subgroup of G fixing R . Since $P\Gamma\ell_2(2^{2n-1})$ is in a canonical way a group of automorphisms of $(PG(1, 2^{2n-1}), \text{trace } \mathbb{F}_{2^{2n-1}} | \mathbb{F}_2 \circ B_1)$ and thus of (H, B_0) , it can be embedded uniquely into G_P . This implies that G_P contains a subgroup K isomorphic to $P\Gamma\ell_2(2^{2n-1})$. The following lemma summarizes what is known about G from [3].

LEMMA. (Let q, \mathcal{P}, K and G be as above)

- (i) K acts on \mathcal{P} as $P\Gamma\ell_2(2^{2n-1})$ acts on $PG(1, 2^{2n-1})$;
- (ii) $G_P = K \cong P\Gamma\ell_2(2^{2n-1})$; G_P has three orbits on the set of nonisotropic points of $(PG(4n-1, 2), q)$ with cardinalities $1, 2^{4n-2}-1, 2^{2n-1}(2^{2n-1}-1)$;
- (iii) If $n = 2$, then $G \cong \text{Alt}(9)$;
- (iv) if $n = 3$, then $G = G_P \cong P\Gamma\ell_2(2^5)$.

The proof of (ii) can be found on page 191 in [3] in an argument that is valid in the present situation (though not explicitly stated).

Statement (iv) is demonstrated by use of specific knowledge of the subgroups of $Sp_6(2)$.

The theorem which we aim to prove, shows that (iv) is representative for what happens for $n \geq 3$.

THEOREM. Let $n \geq 3$. Suppose P is a nonisotropic point and Q an isotropic point of a nondegenerate hyperbolic space $(PG(4n-1, 2), q)$ such that $P + Q$ is a hyperbolic line. Let \mathcal{P} be the spread constructed in 2 departing from P and Q , and let G be as defined in 3. Then $G = G_P \cong P\Gamma\ell_2(2^{2n-1})$.

4. PROOF OF THE THEOREM

We proceed in four steps.

(4.1) *G does not possess a normal subgroup which is regular on the set of nonisotropic points of $(PG(4n-1,2),q)$.*

PROOF. Suppose N is a counterexample. Then G_p acts on N by conjugation as it does on the nonisotropic points. In particular N has two G_p -orbits distinct from $\{1\}$. Let p and q denote the orders of representatives from these two orbits. Then by Cauchy's lemma N has order $p^a q^b$ for $a, b \in \mathbb{N}$; moreover p and q are prime numbers. On the other hand, the regularity of N implies that its order is $2^{2n-1}(2^{2n}-1)$. The comparison of these two expressions for $|N|$ yields that $2^{2n}-1$ is a prime power, which is absurd. \square

(4.2) *If N is a nontrivial normal subgroup of G , then $[G:N] = [G_p:N_p]$ is a divisor of $2n-1$.*

PROOF. If $G = G_p$, the statement concerns $G \cong P\Gamma\ell_2(2^{2n-1})$ and is known to hold. So we may assume $G > G_p$ for the rest of the proof. In view of the orbit structure of G_p described in (ii) of the lemma, this means that G is primitive on the set of nonisotropic points. So any nontrivial normal subgroup N of G is transitive on these $2^{2n-1}(2^{2n}-1)$ points, so $[G:N] = [G_p:N_p]$. Moreover N_p is normal in $G_p \cong P\Gamma\ell_2(2^{2n-1})$, whence $N_p = 1$ or we are through. The former possibility, however, is excluded by (4.1) \square

(4.3) *The permutation representation of G on \mathcal{P} is faithful.*

PROOF. Let N be the kernel of this representation. If N is nontrivial, then $[G:N] = [G_p:N_p]$ by (4.2); but (i) of the lemma states that $N_p = 1$, whence $[G:N] = |G_p|$, contradicting (4.2). The conclusion is that N is trivial. \square

(4.4) *If $n \geq 3$, then $G = G_p$.*

PROOF. By (4.3) the group G can be regarded as a triply transitive permutation group of degree $2^{2n-1}+1$. Application of a theorem by Holt [4] yields that G contains a normal subgroup N isomorphic to either

$\text{Sym}(2^{2n-1} + 1)$, $\text{Alt}(2^{2n-1} + 1)$ or $\text{PSL}_2(2^{2n-1})$. Comparing orders with $|G|$, we obtain that N is an isomorph of $\text{PSL}_2(2^{2n-1})$. From (4.2) it follows that $G = G_p \quad \square$.

5. REMARKS

For $n = 2$, the arguments of the proof are equally valid. They result in: $G \cong \text{P}\Gamma\mathcal{L}_2(2^{2n-1})$ or $G \cong \text{Alt}(9)$. Together with the observation that all spreads are in a single $O_{4n}^+(2)$ -orbit, this reestablishes (iii) of the lemma.

De Clerck, Dye and Thas [1] have shown that any spread leads to a partial geometry with parameters $(s, t, \alpha) = (2^{2n-1}-1, 2^{2n-1}, 2^{2n-2})$ on the nonisotropic points of $\text{PG}(4n-1, q)$. Using the above theorem, it is not hard to see that G is the part of the automorphism group of the partial geometry derived from \mathcal{P} that is contained in $O_{4n}^+(2)$.

6. REFERENCES

- [1] DE CLERCK; DYE, R.H.; THAS, J.; preprint
- [2] DIEUDONNÉ J; *La géométrie des groupes classiques*,
- [3] DYE, R.H.; *Partitions and their stabilizers for line complexes and quadrics*, *Annali di Matematica pura ed applicata* (4) 114 (1977), 173-194.
- [4] HOLT, D.F.; *Triply-transitive permutation groups in which an involution central in a Sylow 2-subgroup fixes a unique point*, *J. London Math. Soc* (2) 15 (1977), 55-65.

ONTVANGEN 3 2 JAN. 1980