



Centrum voor Wiskunde en Informatica

**REPORTRAPPORT**

On periodic cohort solutions of a size-structured population model

W. Huyer

Department of Analysis, Algebra and Geometry

**AM-R9504 1995**

Report AM-R9504  
ISSN 0924-2953

CWI  
P.O. Box 94079  
1090 GB Amsterdam  
The Netherlands

CWI is the National Research Institute for Mathematics and Computer Science. CWI is part of the Stichting Mathematisch Centrum (SMC), the Dutch foundation for promotion of mathematics and computer science and their applications.

SMC is sponsored by the Netherlands Organization for Scientific Research (NWO). CWI is a member of ERCIM, the European Research Consortium for Informatics and Mathematics.

Copyright © Stichting Mathematisch Centrum  
P.O. Box 94079, 1090 GB Amsterdam (NL)  
Kruislaan 413, 1098 SJ Amsterdam (NL)  
Telephone +31 20 592 9333  
Telefax +31 20 592 4199

# On Periodic Cohort Solutions of a Size-Structured Population Model

Waltraud Huyer

*Institut für Mathematik, Universität Graz, Heinrichstraße 36, A-8010 Graz, Austria*

## Abstract

We consider a size-structured population model with discontinuous reproduction and feedback through the environmental variable “substrate”. The model admits solutions with finitely many cohorts and in that case the problem is described by a system of ODEs involving a bifurcation parameter  $\beta$ . Existence of nontrivial periodic  $n$ -cohort solutions is investigated. Moreover, we discuss the question whether  $n$  cohorts ( $n \geq 2$ ) with small size differences will tend to a periodic one-cohort solution as  $t \rightarrow \infty$ .

*AMS Subject Classification (1991):* 92D25

*Keywords & Phrases:* size-structured populations, discontinuous reproduction, variable environment, periodic cohort solutions, discrete time dynamical systems, stability, implicit function theorem, local bifurcation

## 1. INTRODUCTION

Reproduction is an event in the life of individuals. When there is quite some variation in the individual states at which such events occur, a formal appeal to the law of large numbers allows one to model reproduction by rates at the population level. When reproduction occurs upon reaching a specified individual state, one can still work with rates at the population level provided the distribution of the population with respect to individual state is smooth (i.e. absolutely continuous). However, when “giving birth” is restricted to specific individual states and, in addition, the population is concentrated in some (moving) individual states, reproduction is an event even at the population level. The aim of this paper is to initiate the study of models incorporating these two features.

We consider a size-structured population model of this type, i.e. the state of an individual is characterised by its size. The individuals are born at size  $x_\alpha$ , and when they reach size  $x_\omega$  ( $x_\omega > x_\alpha > 0$ ), they produce  $k$  offspring of size  $x_\alpha$  and die (or, equivalently, split into  $k$  parts). There is feedback through the environment, i.e. the growth rate  $g$  depends on the food density  $S$ . The mortality rate  $\mu$  is assumed to be a positive constant.

We are interested in the special case that the population consists of finitely many cohorts. A cohort refers to a group of individuals of exactly the same state (i.e. size in the present context) and is mathematically described by a measure concentrated in a point, i.e. a Dirac measure. The full problem is described by an integral equation with a measure on  $[x_\alpha, x_\omega)$  as initial condition, but in the case of finitely many cohorts it reduces to a system of coupled ordinary differential equations. If we start with  $n$  cohorts, we shall have  $n$  cohorts all the time. Let  $N_i$  denote the population number of cohort  $i$  and  $x_i$  the corresponding size. In the case of  $n$  cohorts the model equations read

$$\left\{ \begin{array}{l} \frac{dN_i}{dt} = -\mu N_i, \quad i = 1, \dots, n, \\ \frac{dx_i}{dt} = g(S, x_i), \quad i = 1, \dots, n, \\ \frac{dS}{dt} = h(S, \beta) - \sum_{i=1}^n \gamma(S, x_i) N_i \\ + \text{initial conditions and jumps in } N_i \text{ and } x_i \text{ at times } t \text{ such that } x_i(t) = x_\omega \text{ for some } i. \end{array} \right. \quad (1.1)$$

The function  $h$  describes the dynamics of the unstructured variable  $S$  in the absence of a consumer, and  $\gamma$  is the consumption rate. We want to study the asymptotic behaviour and, in particular, the existence and stability of periodic  $n$ -cohort solutions. This behaviour might change if certain parameters in the model equations are varied. We assume that the function  $h$  also depends on a scalar parameter  $\beta$  that will be the bifurcation parameter for our problem. The following assumptions on the functions  $g$ ,  $h$  and  $\gamma$  are used throughout this paper:

$$\begin{aligned} &g \in C([0, \infty) \times [x_\alpha, x_\omega]; \mathbb{R}), \text{ for each } x \in [x_\alpha, x_\omega] \text{ there exists an } S_x \geq 0 \\ &\text{such that } g(S, x) > 0 \text{ for } S > S_x \text{ and } g(S, x) = 0 \text{ for } 0 \leq S \leq S_x, \\ &g \text{ is decreasing in } x, \text{ strictly increasing in } S \text{ on } [S_x, \infty) \text{ for all } x \in [x_\alpha, x_\omega] \\ &\text{and twice continuously differentiable at those } (S, x) \text{ for which } g(S, x) > 0. \end{aligned} \quad (\text{H}_g)$$

$$\begin{aligned} &h \in C^1([0, \infty) \times (0, \infty); \mathbb{R}), h \text{ is strictly increasing in } \beta, \text{ for each } \beta > 0 \\ &\text{there exists a unique } \tilde{S}_\beta > 0 \text{ such that } h(\tilde{S}_\beta, \beta) = 0 \text{ and } \frac{\partial h}{\partial S}(\tilde{S}_\beta, \beta) =: -d_\beta < 0. \end{aligned} \quad (\text{H}_h)$$

$$\begin{aligned} &\gamma \in C^1([0, \infty) \times [x_\alpha, x_\omega]; \mathbb{R}), \gamma(0, x) = 0 \text{ for } x \in [x_\alpha, x_\omega] \text{ and } \gamma > 0 \text{ elsewhere,} \\ &\gamma \text{ is increasing in } S \text{ and } x. \end{aligned} \quad (\text{H}_\gamma)$$

Moreover, we consider only values of  $\beta$  for which

$$\tilde{S}_\beta > S_{x_\omega}. \quad (\text{H}_\beta)$$

The typical growth rates we have in mind are the so-called growth rates of von Bertalanffy type (cf. [2, p. 22] and [1], for example)

$$g(S, x) = c(f(S) - x)^+, \quad (1.2)$$

where  $c > 0$ ,  $z^+ := \max(0, z)$ ,  $f$  is a continuously differentiable, strictly increasing function of  $S$  and  $x$  is of dimension ‘‘length’’. The condition  $(\text{H}_h)$  implies that  $h(S, \beta) > 0$  for  $0 < S < \tilde{S}_\beta$ ,  $h(S, \beta) < 0$  for  $S > \tilde{S}_\beta$  and  $\tilde{S}_\beta$  is strictly increasing in  $\beta$ . Typical examples are  $h(S, \beta) = \beta - S$  (constant inflow of fresh, nonreproducing food particles combined with a constant food deterioration) and  $h(S, \beta) = S(1 - S/\beta)$  (logistic growth). The condition  $(\text{H}_\beta)$  ensures that size  $x_\omega$  is reached under the food conditions  $S \equiv \tilde{S}_\beta$ ; otherwise size  $x_\omega$  would never be reached for initial data  $S(0) \leq \tilde{S}_\beta$  and the population would become extinct.

We consider  $n$  cohorts (that we assume to be arranged according to size; without loss of generality we can also assume that the first cohort is a cohort of newborns) and associate a mapping (discrete time dynamical system) with the ODE system. Let  $n \in \mathbb{N}$ ,  $\beta > 0$  with  $(\text{H}_\beta)$ ,  $x_\alpha = x_{1,0} \leq x_{2,0} \leq \dots \leq x_{n,0} \leq x_\omega$ ,  $0 < S_0 \leq \tilde{S}_\beta$ ,  $N_{i,0} \geq 0$  for  $i = 1, \dots, n$ , and solve the equations

$$\begin{cases} \frac{dx_i}{dt} = g(S, x_i), & x_i(0) = x_{i,0}, & i = 1, \dots, n, \\ \frac{dS}{dt} = h(S, \beta) - e^{-\mu t} \sum_{i=1}^n \gamma(S, x_i) N_{i,0}, & S(0) = S_0. \end{cases} \quad (1.3)$$

**Remark 1.1.** a)  $(\text{H}_h)$  implies that it is necessary to choose  $S_0 \in (0, \tilde{S}_\beta]$  in order to possibly obtain a periodic solution.

b) We have to extend the mapping artificially to the case that the initial sizes can become  $x_\omega$  in order to be able to deal with the degenerate case that  $x_{i,0} = x_{i+1,0}$  for some  $i$ . In this case, as well as in the case where  $N_{i,0} = 0$  for some  $i$ , we actually have less than  $n$  cohorts.

We define a mapping  $F_n : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ ,  $(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0}) \mapsto (S'_0, N'_{1,0}, \dots, N'_{n,0}, x'_{2,0}, \dots, x'_{n,0})$  as follows. Let  $T = T(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0}) \geq 0$  be such that  $x_n(T) = x_\omega$ ,  $x_n(t) < x_\omega$  for  $0 \leq t < T$  (if  $x_{n,0} = x_\omega$ , we have  $T = 0$ ), and set

$$\begin{cases} S'_0 := S(T), \\ N'_{1,0} := kN_{n,0}e^{-\mu T}, \\ N'_{i,0} := N_{i-1,0}e^{-\mu T}, \quad i = 2, \dots, n, \\ x'_{i,0} := x_{i-1}(T), \quad i = 2, \dots, n. \end{cases} \quad (1.4)$$

$T$  is well-defined because of assumptions  $(H_\beta)$ ,  $(H_h)$  and  $(H_\gamma)$ , and we have  $S'_0 > 0$ ,  $S'_0 \in [S_{x_\omega}, \tilde{S}_\beta]$ ,  $N_{i,0} \geq 0$  for  $i = 1, \dots, n$  and  $x_\alpha \leq x'_{2,0} \leq \dots \leq x'_{n,0} \leq x_\omega$ .

**Remark 1.2.** Since the size growth is not necessarily strictly monotonic, the mappings  $F_n$  might have discontinuities if  $S'_0 = S_{x_\omega} > 0$ ,  $\dot{S}(T-) < 0$ . (If  $S(T) = S_{x_\omega}$ ,  $\dot{S}(T-) = 0$ , then  $\dot{S}(T-) = \mu e^{-\mu T} \sum_{i=1}^n \gamma(S_{x_\omega}, x_i) N_{i,0} > 0$ , i.e.  $S$  would have a local minimum at  $t = T$  if there were no jumps, and this case is all right.)

In Section 2 the mapping  $F_1$  is investigated. We assume that the trivial fixed point  $(\tilde{S}_\beta, 0)$  becomes unstable at  $\beta = 1$ , and existence of a stable nontrivial fixed point of  $F_1$  (resp. a periodic one-cohort solution of (1.1)) is proved for  $1 < \beta < \beta_1$  for some  $\beta_1 > 1$ . In Section 3 a criterion is derived to determine whether for  $1 < \beta < \beta_1$ ,  $\beta$  sufficiently close to 1, the iterations of  $F_2$  will converge to the fixed point  $(\tilde{S}_0(\beta), \tilde{N}_0(\beta))$  of  $F_1$  for  $(S_0, N_{1,0} + N_{2,0}, x_{2,0})$  sufficiently close to  $(\tilde{S}_0(\beta), \tilde{N}_0(\beta), x_\alpha)$ , i.e. whether the nontrivial fixed point of  $F_1$  is stable with respect to splitting into two cohorts. The same criterion applies to splitting into  $n$  cohorts for all  $n \geq 2$ . Moreover, local existence of nontrivial fixed points of  $F_n$  is proved for  $n \geq 2$ . In Section 4 we present some numerical examples that show that both stability and instability of the fixed point of  $F_1$  with respect to cohort splitting can occur in the case of linear growth rates (1.2).

## 2. THE ONE-COHORT PROBLEM

We consider now the special case  $n = 1$ ,  $F_1 =: F$ , and we drop the indices. Let  $\beta > 0$  be such that  $(H_\beta)$  is satisfied. We have to solve the coupled ODE system

$$\begin{cases} \frac{dx}{dt} = g(S, x), & x(0) = x_\alpha, \\ \frac{dS}{dt} = h(S, \beta) - N_0 e^{-\mu t} \gamma(S, x), & S(0) = S_0. \end{cases} \quad (2.1)$$

If  $T = T(S_0, N_0) > 0$  is such that  $x(T) = x_\omega$ ,  $x(t) < x_\omega$  for  $0 \leq t < T$ , the mapping  $F : (S_0, N_0) \mapsto (S'_0, N'_0)$  is given by

$$\begin{cases} S'_0 := S(T), \\ N'_0 := kN_0 e^{-\mu T}. \end{cases} \quad (2.2)$$

**Remark 2.1.** a) We have  $T(S_0, N_0) > T(S_0, 0)$  for all  $S_0 \in (0, \tilde{S}_\beta]$ ,  $N_0 > 0$ , and  $S_0 \mapsto T(S_0, 0)$  is strictly decreasing on  $(0, \tilde{S}_\beta]$ . If

$$\tilde{T}_\beta := T(\tilde{S}_\beta, 0) \geq \frac{1}{\mu} \ln k,$$

we have  $\lim_{t \rightarrow \infty} N(t) = 0$  and  $\lim_{t \rightarrow \infty} S(t) = \tilde{S}_\beta$  for all choices of  $(S_0, N_0)$ , and  $F$  has only the trivial fixed point  $(S_0, N_0) = (\tilde{S}_\beta, 0)$ . Note that  $\tilde{T}_\beta$  is strictly decreasing in  $\beta$ .

b) Remark 1.2 implies that discontinuities might occur if  $S(T) = S'_0 = S_{x_\omega} > 0$ ,  $\dot{S}(T^-) < 0$ , i.e.

$$h(S_{x_\omega}, \beta) - N_0 e^{-\mu T} \gamma(S_{x_\omega}, x_\omega) < 0$$

or

$$N_0 > \frac{e^{\mu T}}{\gamma(S_{x_\omega}, x_\omega)} h(S_{x_\omega}, \beta).$$

Since  $T > \tilde{T}_\beta$  for  $N_0 > 0$ , the mapping is in particular continuously differentiable (differentiability of ODE solutions with respect to initial conditions and parameters) in the neighbourhood of  $(S_0, N_0)$  with  $S_0 \in (0, \tilde{S}_\beta]$  and

$$0 \leq N_0 \leq \frac{e^{\mu \tilde{T}_\beta}}{\gamma(S_{x_\omega}, x_\omega)} h(S_{x_\omega}, \beta).$$

At the points where  $F$  is differentiable we have

$$\begin{cases} \frac{\partial S'_0}{\partial S_0} = \frac{\partial S}{\partial S_0}(T) + \dot{S}(T) \frac{\partial T}{\partial S_0}, \\ \frac{\partial S'_0}{\partial N_0} = \frac{\partial S}{\partial N_0}(T) + \dot{S}(T) \frac{\partial T}{\partial N_0}, \\ \frac{\partial N'_0}{\partial S_0} = -\mu \frac{\partial T}{\partial S_0} k e^{-\mu T} N_0, \\ \frac{\partial N'_0}{\partial N_0} = -\mu \frac{\partial T}{\partial N_0} k e^{-\mu T} N_0 + k e^{-\mu T}. \end{cases}$$

We have to compute the eigenvalues of the Jacobi matrix of  $F$  in order to determine the stability of the trivial fixed point  $(0, \tilde{S}_\beta)$ . To this end we first evaluate the derivatives of  $x$  and  $S$  with respect to initial data and parameters at  $(S_0, N_0) = (\tilde{S}_\beta, 0)$ .

**Lemma 2.2.** *Let  $(x, S)$  be the solution of (2.1), and let  $\bar{x}$  be the solution of*

$$\frac{d\bar{x}}{dt} = g(\bar{x}, \tilde{S}_\beta), \quad \bar{x}(0) = x_\alpha.$$

Then we have at  $(S_0, N_0) = (\tilde{S}_\beta, 0)$

$$\frac{\partial S}{\partial S_0}(t) = e^{-d_\beta t}, \tag{2.3}$$

$$\frac{\partial x}{\partial S_0}(t) = \int_0^t \exp\left(\int_\tau^t \frac{\partial g}{\partial x}(\tilde{S}_\beta, \bar{x}(\sigma)) d\sigma\right) \frac{\partial g}{\partial S}(\tilde{S}_\beta, \bar{x}(\tau)) e^{-d_\beta \tau} d\tau, \tag{2.4}$$

$$\frac{\partial S}{\partial N_0}(t) = -e^{-d_\beta t} \int_0^t e^{(d_\beta - \mu)\tau} \gamma(\tilde{S}_\beta, \bar{x}(\tau)) d\tau, \tag{2.5}$$

$$\frac{\partial x}{\partial N_0}(t) = \int_0^t \exp\left(\int_\tau^t \frac{\partial g}{\partial x}(\tilde{S}_\beta, \bar{x}(\sigma)) d\sigma\right) \frac{\partial g}{\partial S}(\tilde{S}_\beta, \bar{x}(\tau)) \frac{\partial S}{\partial N_0}(\tau) d\tau, \tag{2.6}$$

$$\frac{\partial S}{\partial \beta}(t) = \frac{1}{d_\beta} \frac{\partial h}{\partial \beta}(\tilde{S}_\beta, \beta) (1 - e^{-d_\beta t}), \tag{2.7}$$

$$\frac{\partial x}{\partial \beta}(t) = \int_0^t \exp\left(\int_\tau^t \frac{\partial g}{\partial x}(\tilde{S}_\beta, \bar{x}(\sigma)) d\sigma\right) \frac{\partial g}{\partial S}(\tilde{S}_\beta, \bar{x}(\tau)) \frac{\partial S}{\partial \beta}(\tau) d\tau. \tag{2.8}$$

Moreover we have at  $(S_0, N_0) = (\tilde{S}_\beta, 0)$

$$\frac{\partial T}{\partial \alpha} = -g(\tilde{S}_\beta, x_\omega)^{-1} \frac{\partial x}{\partial \alpha}(\tilde{T}_\beta) \tag{2.9}$$

for  $\alpha = S_0, N_0$  and  $\beta$ .

*Proof.* Application of theorems about differentiability of ODE solutions with respect to initial data and parameters yields formulas (2.3)–(2.8), and by differentiating the relationship  $x(T) = x_\omega$  we obtain (2.9).  $\square$

The following lemma shows that the stability of the trivial fixed point of  $F$  is as we might expect from Remark 2.1, a).

**Lemma 2.3.** *The Jacobi matrix of  $F$  at the trivial fixed point  $(S_0, N_0) = (\tilde{S}_\beta, 0)$  reads*

$$(DF)(\tilde{S}_\beta, 0) = \begin{pmatrix} e^{-d_\beta \tilde{T}_\beta} & \frac{\partial S}{\partial N_0}(\tilde{T}_\beta) \\ 0 & ke^{-\mu \tilde{T}_\beta} \end{pmatrix}$$

and has the two real positive eigenvalues  $\lambda_1 = e^{-d_\beta \tilde{T}_\beta} < 1$  and  $\lambda_2 = ke^{-\mu \tilde{T}_\beta}$ . The trivial fixed point is stable if  $\lambda_2 < 1$  or

$$\tilde{T}_\beta > \frac{1}{\mu} \ln k$$

and unstable if  $\lambda_2 > 1$  or

$$\tilde{T}_\beta < \frac{1}{\mu} \ln k.$$

We want to discuss the existence of nontrivial fixed points of  $F$  for  $\tilde{T}_\beta < \frac{1}{\mu} \ln k$ . The mapping  $F$  can be extended to  $S_0 > \tilde{S}_\beta$  and  $N_0 < 0$ , but the case  $N_0 < 0$  has no biological meaning. If  $\beta_0 \geq 0$  is such that  $(H_\beta)$  is satisfied for  $\beta > \beta_0$ , we can define a mapping  $G : (0, \infty) \times (-\infty, \infty) \times (\beta_0, \infty) \rightarrow \mathbb{R}^2$  by

$$\begin{cases} G_1(S_0, N_0, \beta) := S(T) - S_0, \\ G_2(S_0, N_0, \beta) := ke^{-\mu T} - 1, \end{cases} \quad (2.10)$$

where  $S$  is the solution of the second equation of (2.1) and  $T = T(S_0, N_0, \beta)$  is defined as before. By (2.2) we must have  $G(S_0, N_0, \beta) = 0$  if  $(S_0, N_0)$  is a nontrivial fixed point of  $F$  pertaining to the parameter value  $\beta$ , and in particular

$$T(S_0, N_0, \beta) = \bar{T} := \frac{1}{\mu} \ln k. \quad (2.11)$$

Without loss of generality we assume that the trivial fixed point becomes unstable at  $\beta = 1$ , i.e.

$$\tilde{T}_1 = \frac{1}{\mu} \ln k. \quad (2.12)$$

Moreover, we assume henceforth

$$d_1 = 1, \quad (2.13)$$

which can be achieved by scaling the time variable and serves to simplify notation.

**Theorem 2.4.** *Assume that (2.12) and (2.13) are satisfied. Then there exist a neighbourhood  $U$  of 1 in  $(0, \infty)$  and a neighbourhood  $V$  of  $(\tilde{S}_1, 0)$  in  $\mathbb{R}^2$  such that for each  $\beta \in U$  there exists a unique  $(S_0, N_0) \in V$  with  $G(S_0, N_0, \beta) = 0$ . Moreover, for  $\beta \in U$  with  $\beta < 1$  we have  $S_0 > \tilde{S}_\beta$ ,  $N_0 < 0$ , and for  $\beta \in U$  with  $\beta > 1$  we have  $S_0 < \tilde{S}_\beta$ ,  $N_0 > 0$ , i.e. only the second case yields a biologically meaningful nontrivial fixed point of  $F$  resp. a periodic one-cohort solution of (1.1).*

*Proof.* By (2.12) we have  $T(\tilde{S}_1, 0, 1) = \bar{T}$  and  $G(\tilde{S}_1, 0, 1) = 0$ , and there exists a neighbourhood  $W$  of  $(\tilde{S}_1, 0, 1)$  such that  $G$  is continuously differentiable on  $W$ . Let  $y := (S_0, N_0)$ . Then we have

$$(D_y G)(y, \beta) = \begin{pmatrix} \frac{\partial S}{\partial S_0}(T) + \dot{S}(T) \frac{\partial T}{\partial S_0} - 1 & \frac{\partial S}{\partial N_0}(T) + \dot{S}(T) \frac{\partial T}{\partial N_0} \\ -\mu k e^{-\mu T} \frac{\partial T}{\partial S_0} & -\mu k e^{-\mu T} \frac{\partial T}{\partial N_0} \end{pmatrix},$$

and in particular

$$(D_y G)(\tilde{S}_1, 0, 1) = \begin{pmatrix} k^{-\frac{1}{\mu}} - 1 & \frac{\partial S}{\partial N_0}(\bar{T}) \\ -\mu \frac{\partial T}{\partial S_0} & -\mu \frac{\partial T}{\partial N_0} \end{pmatrix}.$$

The determinant of this matrix is

$$\Delta := \det(D_y G)(\tilde{S}_1, 0, 1) = \mu \frac{\partial T}{\partial N_0} (1 - k^{-\frac{1}{\mu}}) + \mu \frac{\partial T}{\partial S_0} \frac{\partial S}{\partial N_0}(\bar{T}), \quad (2.14)$$

which is positive by Lemma 2.2 and assumptions  $(H_g)$  and  $(H_\gamma)$ . The Implicit Function Theorem implies that there exist neighbourhoods  $U$  of 1 and  $V$  of  $(\tilde{S}_1, 0)$  and a  $C^1$ -function  $H : U \rightarrow V$  such that  $V \times U \subset W$  and  $G(H(\beta), \beta) = 0$  for all  $\beta \in U$ . Moreover, for  $\beta \in U$  the equation  $G(y, \beta) = 0$  possesses one and only one solution  $y = H(\beta)$  in  $V$ .

Only three kinds of choices for  $(S_0, N_0)$  are possible such that  $S$  might return to its initial value: a)  $S_0 > \tilde{S}_\beta$ ,  $N_0 < 0$ ; b)  $S_0 = \tilde{S}_\beta$ ,  $N_0 = 0$ ; c)  $S_0 < \tilde{S}_\beta$ ,  $N_0 > 0$ . Let  $\beta > 1$ . Then  $T(S_0, N_0) \leq \bar{T}_\beta < \frac{1}{\mu} \ln k$  for  $S_0 \geq \tilde{S}_\beta$ ,  $N_0 \leq 0$ , i.e. we must have case c). Analogously, case a) applies to  $\beta < 1$ .  $\square$

The principle of exchange of stability (cf. [3, p. 93, Theorem 5.1], for example) implies

**Theorem 2.5.** *Let the assumptions of Theorem 2.4 be satisfied. If  $U$  is chosen sufficiently small, the biologically meaningful nontrivial fixed point of  $F$  for  $\beta \in U \cap (1, \infty)$  is stable.*

We assume in the following that  $U$  is chosen sufficiently small such that the conclusion of Theorem 2.5 holds.

### 3. THE $n$ -COHORT PROBLEM FOR $n \geq 2$

In this section we address the following question: If we start with  $n \geq 2$  cohorts with small size differences, will these differences be larger after all  $n$  cohorts have reproduced, or will they be smaller and eventually tend to zero? Will the solution of the  $n$ -cohort problem converge to a periodic solution of the one-cohort problem? This phenomenon cannot occur in the case of a constant food density or if the growth rate is independent of  $S$  since then the size differences always stay the same. First we consider the case  $n = 2$ , i.e. for  $(S_0, N_1, N_2, x_0) \in (0, \tilde{S}_\beta) \times [0, \infty) \times [0, \infty) \times [x_\alpha, x_\omega]$ ,  $\beta > 0$ , we solve

$$\begin{cases} \frac{dx_1}{dt} = g(S, x_1), & x_1(0) = x_\alpha, \\ \frac{dx_2}{dt} = g(S, x_2), & x_2(0) = x_0, \\ \frac{dS}{dt} = h(S, \beta) - e^{-\mu t} (N_1 \gamma(S, x_1) + N_2 \gamma(S, x_2)), & S(0) = S_0, \end{cases} \quad (3.1)$$

where we again assume that  $(H_\beta)$ , (2.12) and (2.13) are satisfied. The mapping  $F_2 : (S_0, N_1, N_2, x_0) \mapsto (S'_0, N'_1, N'_2, x'_0)$  is defined by

$$\begin{cases} S'_0 := S(T), \\ N'_1 := k N_2 e^{-\mu T}, \\ N'_2 := N_1 e^{-\mu T}, \\ x'_0 := x_1(T), \end{cases} \quad (3.2)$$



where  $T = T(S_0, N_1, N_2, x_0) \geq 0$  is such that  $x_2(T) = x_\omega$ ,  $x_2(t) < x_\omega$  for  $0 \leq t < T$ .

For  $\beta \in U$  we use the abbreviations

$$\begin{cases} \bar{S}_0(\beta) := H_1(\beta), \\ \bar{N}_0(\beta) := H_2(\beta), \end{cases} \quad (3.3)$$

where  $H$  is defined by Theorem 2.4 (see the proof). The nontrivial fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta))$  of  $F_1$  for  $\beta \in U \cap (1, \infty)$  can be viewed as a periodic point  $(S_0, N_1, N_2, x_0)$  of period two of  $F_2$  with  $S_0 = \bar{S}_0(\beta)$ ,  $N_1 + N_2 = \bar{N}_0(\beta)$ ,  $x_0 = x_\alpha$  (or  $N_1 + kN_2 = \bar{N}_0(\beta)$ ,  $x_0 = x_\omega$ ), i.e. the artificial splitting of one cohort into two cohorts is not unique and instead of a fixed point of  $F_1$  we obtain a family of periodic points of period two of  $F_2$ . In order to investigate the ‘‘stability’’ of these periodic points of period two, we have to consider the second iterate  $F_2^2 =: K : (S_0, N_1, N_2, x_0) \mapsto (S_0'', N_1'', N_2'', x_0'')$ . If  $T_1 := T(S_0, N_1, N_2, x_0)$  and  $T_2 := T(F_2(S_0, N_1, N_2, x_0))$ , we have  $N_1'' = kN_1 e^{-\mu(T_1+T_2)}$  and  $N_2'' = kN_2 e^{-\mu(T_1+T_2)}$ , i.e. the quotient  $\frac{N_1}{N_2}$  or  $\frac{N_2}{N_1}$  remains invariant under  $K$  if the denominator is not zero. This allows us to restrict  $K$  to any set  $N_2 = \eta N_1$ ,  $\eta$  a positive constant, and thus to reduce the problem by one dimension. Let  $K_\eta : (S_0, N_0, x_0) \mapsto (S_0'', N_0'', x_0'')$  be defined by

$$(S_0'', \frac{1}{1+\eta}N_0'', \frac{\eta}{1+\eta}N_0'', x_0'') := K(S_0, \frac{1}{1+\eta}N_0, \frac{\eta}{1+\eta}N_0, x_0).$$

The nontrivial fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta))$  of  $F_1$  for  $\beta \in U \cap (1, \infty)$  corresponds to the fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha)$  (or  $(\bar{S}_0(\beta), \frac{1+\eta}{1+k\eta}\bar{N}_0(\beta), x_\omega)$ ) of  $K_\eta$ . The Jacobi matrix of  $K_\eta$  at the point  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha)$  is

$$(DK_\eta)(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha) = \begin{pmatrix} A & * \\ 0 & 0 \end{pmatrix}, \quad (3.4)$$

where

$$A := (DF_1)(\bar{S}_0(\beta), \bar{N}_0(\beta)) \quad (3.5)$$

is the  $(2 \times 2)$ -Jacobi matrix of  $F_1$  evaluated at its stable fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta))$  and ‘‘\*’’ denotes a not so interesting number. Note that  $x_0''(S_0, N_0, x_\alpha) = x_\alpha$  for all  $S_0, N_0$ , which accounts for the two zeros in the third row of the matrix (3.4). Thus the eigenvalues of the Jacobi matrix (3.4) are given by the eigenvalues of  $A$  and

$$\lambda := \frac{\partial x_0''}{\partial x_0}(\bar{S}_0(\beta), \frac{1}{1+\eta}\bar{N}_0(\beta), \frac{\eta}{1+\eta}\bar{N}_0(\beta), x_\alpha), \quad (3.6)$$

where  $x_0''$  now again denotes the fourth component of the mapping  $K$ . The former are of absolute value  $< 1$  by assumption, and the stability of the fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha)$  of  $K_\eta$  depends on the absolute value of  $\lambda$ .

**Lemma 3.1.** *The eigenvalue  $\lambda$  defined by (3.6) is independent of  $\eta$  and given by*

$$\lambda(\beta) = \frac{g(\bar{S}_0(\beta), x_\alpha)}{g(\bar{S}_0(\beta), x_\omega)} \exp\left(\int_0^T \frac{\partial g}{\partial x}(\bar{S}_\beta(t), \bar{x}_\beta(t)) dt\right), \quad (3.7)$$

where  $(\bar{x}_\beta, \bar{S}_\beta)$  is the solution of (2.1) corresponding to  $S_0 = \bar{S}_0(\beta)$ ,  $N_0 = \bar{N}_0(\beta)$ ,  $\beta \in U$ , and  $\bar{T}$  is defined by (2.11). In particular, for a growth rate of the form (1.2)  $\lambda$  reads

$$\lambda(\beta) = \frac{f(\bar{S}_0(\beta)) - x_\alpha}{f(\bar{S}_0(\beta)) - x_\omega} e^{-c\bar{T}} = \frac{f(\bar{S}_0(\beta)) - x_\alpha}{f(\bar{S}_0(\beta)) - x_\omega} k^{-\frac{c}{\mu}}. \quad (3.8)$$

*Proof.* In order to compute  $\lambda$ , we first have to solve the following ODE system:

$$\begin{cases} \frac{dx_1}{dt} = g(S, x_1), & x_1(0) = x_\alpha, \\ \frac{dx_2}{dt} = g(S, x_2), & x_2(0) = x_0, \\ \frac{dS}{dt} = h(S, \beta) - \frac{\bar{N}_0(\beta)}{1 + \eta} e^{-\mu t} (\gamma(S, x_1) + \eta\gamma(S, x_2)), & S(0) = \bar{S}_0(\beta). \end{cases} \quad (3.9)$$

Let  $T_1 \geq 0$  be such that  $x_2(T_1) = x_\omega$ ,  $x_2(t) < x_\omega$  for  $0 \leq t < T_1$ . We set  $S'_0 := S(T_1)$ ,  $x'_0 := x_1(T_1)$ ,  $N'_0 := e^{-\mu T_1} \bar{N}_0(\beta)$  and solve

$$\begin{cases} \frac{d\tilde{x}_1}{dt} = g(\tilde{S}, \tilde{x}_1), & \tilde{x}_1(0) = x_\alpha, \\ \frac{d\tilde{x}_2}{dt} = g(\tilde{S}, \tilde{x}_2), & \tilde{x}_2(0) = x'_0, \\ \frac{d\tilde{S}}{dt} = h(\tilde{S}, \beta) - \frac{N'_0}{1 + \eta} e^{-\mu t} (k\eta\gamma(\tilde{S}, \tilde{x}_1) + \gamma(\tilde{S}, \tilde{x}_2)), & \tilde{S}(0) = S'_0. \end{cases}$$

Let  $T_2 \geq 0$  be such that  $\tilde{x}_2(T_2) = x_\omega$ ,  $\tilde{x}_2(t) < x_\omega$  for  $0 \leq t < T_2$ , and define  $x''_0 := \tilde{x}_1(T_2)$ . Then

$$\frac{dx''_0}{dx_0} = \frac{\partial \tilde{x}_1}{\partial x'_0}(T_2) \frac{dx'_0}{dx_0} + \frac{\partial \tilde{x}_1}{\partial S'_0}(T_2) \frac{dS'_0}{dx_0} + \frac{\partial \tilde{x}_1}{\partial N'_0}(T_2) \frac{dN'_0}{dx_0} + \dot{\tilde{x}}_1(T_2) \frac{dT_2}{dx_0},$$

where all derivatives exist for  $0 \leq x_0 - x_\alpha < \varepsilon$ ,  $\varepsilon > 0$  suitably chosen, and the derivatives with respect to  $x_0$  are replaced by the right-hand derivatives at  $x_0 = x_\alpha$ . For  $x_0 = x_\alpha$  we have  $T_2 = 0$ ,  $S'_0 = \bar{S}_0(\beta)$  and

$$\frac{\partial \tilde{x}_1}{\partial x'_0}(0) = \frac{\partial \tilde{x}_1}{\partial S'_0}(0) = \frac{\partial \tilde{x}_1}{\partial N'_0}(0) = 0, \quad \dot{\tilde{x}}_1(0) = g(\bar{S}_0(\beta), x_\alpha).$$

Differentiation of the relationship  $\tilde{x}_2(T_2) = x_\omega$  with respect to  $x_0$  yields

$$\dot{\tilde{x}}_2(T_2) \frac{dT_2}{dx_0} + \frac{\partial \tilde{x}_2}{\partial x'_0}(T_2) \frac{dx'_0}{dx_0} + \frac{\partial \tilde{x}_2}{\partial S'_0}(T_2) \frac{dS'_0}{dx_0} + \frac{\partial \tilde{x}_2}{\partial N'_0}(T_2) \frac{dN'_0}{dx_0} = 0,$$

and for  $x_0 = x_\alpha$  we obtain

$$\frac{dT_2}{dx_0}(x_\alpha) = -\frac{1}{g(\bar{S}_0(\beta), x_\omega)} \frac{dx'_0}{dx_0}(x_\alpha)$$

and

$$\lambda = \frac{dx''_0}{dx_0}(x_\alpha) = -\frac{g(\bar{S}_0(\beta), x_\alpha)}{g(\bar{S}_0(\beta), x_\omega)} \frac{dx'_0}{dx_0}(x_\alpha).$$

We have

$$\frac{dx'_0}{dx_0} = \frac{\partial x_1}{\partial x_0}(T_1) + \dot{x}_1(T_1) \frac{dT_1}{dx_0},$$

and for  $x_0 = x_\alpha$  we obtain similarly as above, using the definition of  $T_1$  and the fact that  $T_1(x_\alpha) = \bar{T}$ ,

$$\frac{dx'_0}{dx_0}(x_\alpha) = \frac{\partial x_1}{\partial x_0}(\bar{T}) - \frac{\partial x_2}{\partial x_0}(\bar{T}).$$

If we differentiate the solutions of system (3.9) with respect to the initial condition  $x_0$  at  $x_0 = x_\alpha$ , we obtain the differential equations

$$\begin{cases} \frac{d}{dt} \frac{\partial x_1}{\partial x_0} = \frac{\partial g}{\partial x}(\bar{S}_\beta, \bar{x}_\beta) \frac{\partial x_1}{\partial x_0} + \frac{\partial g}{\partial S}(\bar{S}_\beta, \bar{x}_\beta) \frac{\partial S}{\partial x_0}, & \frac{\partial x_1}{\partial x_0}(0) = 0, \\ \frac{d}{dt} \frac{\partial x_2}{\partial x_0} = \frac{\partial g}{\partial x}(\bar{S}_\beta, \bar{x}_\beta) \frac{\partial x_2}{\partial x_0} + \frac{\partial g}{\partial S}(\bar{S}_\beta, \bar{x}_\beta) \frac{\partial S}{\partial x_0}, & \frac{\partial x_2}{\partial x_0}(0) = 1, \end{cases}$$

which implies that  $z(t) := \frac{\partial x_2}{\partial x_0}(t) - \frac{\partial x_1}{\partial x_0}(t)$  is the solution of the equation

$$\frac{dz}{dt} = \frac{\partial g}{\partial x}(\bar{S}_\beta, \bar{x}_\beta)z, \quad z(0) = 1.$$

From this we infer

$$\frac{\partial x_2}{\partial x_0}(t) - \frac{\partial x_1}{\partial x_0}(t) = \exp\left(\int_0^t \frac{\partial g}{\partial x}(\bar{S}_\beta(\tau), \bar{x}_\beta(\tau)) d\tau\right),$$

and formulas (3.7) and (3.8) follow.  $\square$

The equations  $\frac{d\bar{x}_\beta}{dt} = g(\bar{S}_\beta, \bar{x}_\beta)$ ,  $\bar{x}_\beta(0) = x_\alpha$  and  $\bar{x}_\beta(\bar{T}) = x_\omega$  yield

$$\frac{g(\bar{S}_0(\beta), x_\alpha)}{g(\bar{S}_0(\beta), x_\omega)} = \exp\left(-\int_0^{\bar{T}} \frac{d}{dt} \ln g(\bar{S}_0(\beta), \bar{x}_\beta(t)) dt\right) = \exp\left(-\int_0^{\bar{T}} \frac{\partial g}{\partial x}(\bar{S}_0(\beta), \bar{x}_\beta(t)) \frac{g(\bar{S}_\beta(t), \bar{x}_\beta(t))}{g(\bar{S}_0(\beta), \bar{x}_\beta(t))} dt\right),$$

and we obtain the useful representation

$$\lambda(\beta) = \exp\left(\int_0^{\bar{T}} \left(\frac{\partial g}{\partial x}(\bar{S}_\beta(t), \bar{x}_\beta(t)) - \frac{\partial g}{\partial x}(\bar{S}_0(\beta), \bar{x}_\beta(t)) \frac{g(\bar{S}_\beta(t), \bar{x}_\beta(t))}{g(\bar{S}_0(\beta), \bar{x}_\beta(t))}\right) dt\right). \quad (3.10)$$

For  $\beta = 1$  we have by assumption (2.12)  $\bar{S}_1(t) = \bar{S}_0(1) = \tilde{S}_1$ , and (3.10) immediately implies

$$\lambda(1) = 1.$$

In particular we obtain for a linear growth rate from (3.8)

$$\frac{f(\tilde{S}_1) - x_\alpha k^{-\frac{1}{\mu}}}{f(\tilde{S}_1) - x_\omega} = 1. \quad (3.11)$$

**Corollary 3.2.** a) *If in addition to (H<sub>g</sub>) we have*

$$S \mapsto \frac{\partial}{\partial x}(\ln g(S, x)) \text{ is strictly increasing in } S \text{ for } S > S_x, x \in [x_\alpha, x_\omega], \quad (3.12)$$

and

$$\bar{S}_0(\beta) = \max_{t \geq 0} \bar{S}_\beta(t) \quad (3.13)$$

for a  $\beta \in U \cap (1, \infty)$ , then the nontrivial fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha)$  of  $K_\eta$  is stable for all  $\eta > 0$ .

b) *In case of a linear growth rate (1.2) the nontrivial fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha)$  of  $K_\eta$  is stable for all  $\eta > 0$  for a  $\beta \in U \cap (1, \infty)$  if and only if  $\bar{S}_0(\beta) > \tilde{S}_1$ .*

*Proof.* The assertion a) follows from the representation (3.10) and the assumptions (H<sub>g</sub>), (3.12) and (3.13).

In the case of a linear growth rate (1.2), eqs. (3.8) and (3.11) and the strict monotonicity of  $f$  imply that  $\lambda(\beta) < 1$  if and only if  $\bar{S}_0(\beta) > \tilde{S}_1$ , which proves statement b).  $\square$

**Remark 3.3.** In numerical computations of nontrivial periodic one-cohort solutions for linear growth rates (1.2), linear  $h$ , consumption rates as in Section 4 and conservation of mass (cf. Section 4)  $S$  did attain its maximum at the instants of reproduction. However, we do not know if the property (3.13) was a just a peculiarity of our examples or if it may also occur for growth rates that satisfy (3.12) but are not of type (1.2).

We have  $\text{sign}(\lambda(\beta) - 1) = \text{sign} \lambda'(1)$  for  $\beta > 1$  sufficiently close to 1, and the stability of the fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha)$  of  $K_\eta$  depends on the sign of  $\lambda'(1)$ . In order to compute  $\lambda'(1)$ , we need the following lemma.

**Lemma 3.4.** *Let  $(x, S)$  be the solution of (2.1), let  $T$  be defined as in Section 2, and let  $\bar{S}_0(\beta)$  and  $\bar{N}_0(\beta)$  be defined by (3.3). Then the derivatives with respect to  $\beta$  at  $\beta = 1$  are given by*

$$\frac{d}{d\beta} \bar{S}_0(1) = \frac{\mu}{\Delta} \left( (1 - k^{-\frac{1}{\mu}}) \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) \frac{\partial T}{\partial N_0} - \frac{\partial T}{\partial \beta} \frac{\partial S}{\partial N_0}(\bar{T}) \right) \quad (3.14)$$

and

$$\frac{d}{d\beta} \bar{N}_0(1) = -\frac{\mu}{\Delta} (1 - k^{-\frac{1}{\mu}}) \left( \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) \frac{\partial T}{\partial S_0} + \frac{\partial T}{\partial \beta} \right), \quad (3.15)$$

where  $\Delta$  is defined by (2.14),  $\bar{T}$  is defined by (2.11) and all partial derivatives are evaluated at  $(S_0, N_0, \beta) = (\tilde{S}_1, 0, 1)$ .

*Proof.* If  $G$  is given by (2.10) and  $H$  and  $U$  are defined as in the proof of Theorem 2.4, differentiation of the relationship  $G(H(\beta), \beta) = 0$ ,  $\beta \in U$ , with respect to  $\beta$  yields

$$(D_y G)(H(\beta), \beta) \frac{dH}{d\beta}(\beta) + \frac{\partial G}{\partial \beta}(H(\beta), \beta) = 0, \quad \beta \in U.$$

Since the matrix  $(D_y G)(\tilde{S}_1, 0, 1)$  is invertible, we obtain in particular

$$\begin{aligned} \frac{dH}{d\beta}(1) &= \begin{pmatrix} \frac{d}{d\beta} \bar{S}_0(1) \\ \frac{d}{d\beta} \bar{N}_0(1) \end{pmatrix} = -((D_y G)(\tilde{S}_1, 0, 1))^{-1} \frac{\partial G}{\partial \beta}(H(1), 1) \\ &= -\frac{1}{\Delta} \begin{pmatrix} -\mu \frac{\partial T}{\partial N_0} & -\frac{\partial S}{\partial N_0}(\bar{T}) \\ \mu \frac{\partial T}{\partial S_0} & k^{-\frac{1}{\mu}} - 1 \end{pmatrix} \begin{pmatrix} \frac{\partial S}{\partial \beta}(\bar{T}) \\ -\mu \frac{\partial T}{\partial \beta} \end{pmatrix}, \end{aligned}$$

where all derivatives are evaluated at  $(S_0, N_0, \beta) = (\tilde{S}_1, 0, 1)$ . From (2.7) we obtain  $\frac{\partial S}{\partial \beta}(\bar{T}) = \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1)(1 - k^{-\frac{1}{\mu}})$ , which finishes the proof.  $\square$

**Lemma 3.5.**  *$\lambda'(1)$  is given by*

$$\begin{aligned} \lambda'(1) &= -\frac{\mu}{\Delta} \left( \frac{\partial T}{\partial \beta} + \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) \frac{\partial T}{\partial S_0} \right) \left( \frac{\partial S}{\partial N_0}(\bar{T}) \int_0^T \frac{\frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t))}{g(\tilde{S}_1, \bar{x}_1(t))} e^{-t} dt \right. \\ &\quad \left. + (1 - k^{-\frac{1}{\mu}}) \int_0^T \frac{\frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t))}{g(\tilde{S}_1, \bar{x}_1(t))} \left( \frac{\partial S}{\partial N_0}(t) + e^{-\mu t} \gamma(\tilde{S}_1, \bar{x}_1(t)) \right) dt \right) \end{aligned} \quad (3.16)$$

and

$$\begin{aligned} \text{sign } \lambda'(1) &= \text{sign} \left( \frac{\partial S}{\partial N_0}(\bar{T}) \int_0^T \frac{\frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t))}{g(\tilde{S}_1, \bar{x}_1(t))} e^{-t} dt \right. \\ &\quad \left. + (1 - k^{-\frac{1}{\mu}}) \int_0^T \frac{\frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t))}{g(\tilde{S}_1, \bar{x}_1(t))} \left( \frac{\partial S}{\partial N_0}(t) + e^{-\mu t} \gamma(\tilde{S}_1, \bar{x}_1(t)) \right) dt \right), \end{aligned} \quad (3.17)$$

where all partial derivatives are evaluated at  $(S_0, N_0, \beta) = (\tilde{S}_1, 0, 1)$ .

*Proof.* We differentiate (3.10) at  $\beta = 1$ , and by integration by parts we obtain

$$\begin{aligned}\lambda'(1) &= \int_0^{\bar{T}} \left( \frac{\partial^2 g}{\partial S \partial x}(\tilde{S}_1, \bar{x}_1(t)) - \frac{\partial g}{\partial x}(\tilde{S}_1, \bar{x}_1(t)) \frac{\frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t))}{g(\tilde{S}_1, \bar{x}_1(t))} \right) \left( \frac{d\bar{S}_\beta}{d\beta} \Big|_{\beta=1}(t) - \frac{d}{d\beta} \bar{S}_0(1) \right) dt \\ &= \int_0^{\bar{T}} \frac{d}{dt} \left( \frac{\frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t))}{g(\tilde{S}_1, \bar{x}_1(t))} \right) \left( \frac{d\bar{S}_\beta}{d\beta} \Big|_{\beta=1}(t) - \frac{d}{d\beta} \bar{S}_0(1) \right) dt \\ &= - \int_0^{\bar{T}} \frac{\frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t))}{g(\tilde{S}_1, \bar{x}_1(t))} \frac{d}{dt} \left( \frac{d\bar{S}_\beta}{d\beta} \Big|_{\beta=1}(t) \right) dt,\end{aligned}\tag{3.18}$$

where we have used the fact that  $\lambda(1) = 1$ . We have

$$\frac{d\bar{S}_\beta}{d\beta} \Big|_{\beta=1}(t) = \frac{\partial S}{\partial \beta}(t) + \frac{\partial S}{\partial S_0}(t) \frac{d}{d\beta} \bar{S}_0(1) + \frac{\partial S}{\partial N_0}(t) \frac{d}{d\beta} \bar{N}_0(1),$$

where all partial derivatives are evaluated at  $(S_0, N_0, \beta) = (\tilde{S}_1, 0, 1)$ , and (2.3), (2.5) and (2.7) yield

$$\frac{d}{dt} \left( \frac{d\bar{S}_\beta}{d\beta}(t) \right) = \left( \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) - \frac{d}{d\beta} \bar{S}_0(1) \right) e^{-t} - \frac{d}{d\beta} \bar{N}_0(1) \left( \frac{\partial S}{\partial N_0}(t) + e^{-\mu t} \gamma(\tilde{S}_1, \bar{x}_1(t)) \right).\tag{3.19}$$

Eq. (3.14) and the definition of  $\Delta$  (cf. (2.14)) imply

$$\frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) - \frac{d}{d\beta} \bar{S}_0(1) = \frac{\mu}{\Delta} \left( \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) \frac{\partial T}{\partial S_0} + \frac{\partial T}{\partial \beta} \right) \frac{\partial S}{\partial N_0}(\bar{T}),\tag{3.20}$$

and by using (3.19), (3.20) and (3.15) in (3.18) we obtain formula (3.16).

Eqs. (2.4) and (2.7)–(2.9) and assumptions  $(H_g)$  and  $(H_h)$  yield

$$\frac{\partial T}{\partial \beta} + \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) \frac{\partial T}{\partial S_0} = -g(\tilde{S}_1, x_\omega)^{-1} \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) \int_0^{\bar{T}} \exp\left(\int_t^{\bar{T}} \frac{\partial g}{\partial x}(\tilde{S}_1, \bar{x}_1(\tau)) d\tau\right) \frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_1(t)) dt < 0,$$

from which assertion (3.17) follows.  $\square$

**Remark 3.6.** The sign of  $\lambda'(1)$  does not depend on the particular shape of  $h$  but only on the value of its positive zero at  $\beta = 1$ .

**Lemma 3.7.** *In the special case of a linear growth rate (1.2) we have  $\text{sign } \lambda'(1) = -\text{sign } \frac{d}{d\beta} \bar{S}_0(1)$  and*

$$\begin{aligned}\frac{d}{d\beta} \bar{S}_0(1) &= \frac{\mu}{\Delta} (f(\tilde{S}_1) - x_\omega)^{-1} f'(\tilde{S}_1) \frac{\partial h}{\partial \beta}(\tilde{S}_1, 1) k^{-\frac{c}{\mu}} \left( (1 - k^{-\frac{1}{\mu}}) \int_0^{\bar{T}} e^{(c-1)t} \int_0^t e^{(1-\mu)\tau} \gamma(\tilde{S}_1, \bar{x}_1(\tau)) d\tau dt \right. \\ &\quad \left. - k^{-\frac{1}{\mu}} \int_0^{\bar{T}} e^{ct} (1 - e^{-t}) dt \int_0^{\bar{T}} e^{(1-\mu)t} \gamma(\tilde{S}_1, \bar{x}_1(t)) dt \right),\end{aligned}$$

and consequently

$$\begin{aligned}\text{sign } \lambda'(1) &= \text{sign} \left( k^{-\frac{1}{\mu}} \int_0^{\bar{T}} e^{ct} (1 - e^{-t}) dt \int_0^{\bar{T}} e^{(1-\mu)t} \gamma(\tilde{S}_1, \bar{x}_1(t)) dt \right. \\ &\quad \left. - (1 - k^{-\frac{1}{\mu}}) \int_0^{\bar{T}} e^{(c-1)t} \int_0^t e^{(1-\mu)\tau} \gamma(\tilde{S}_1, \bar{x}_1(\tau)) d\tau dt \right).\end{aligned}$$

*Proof.* Eq. (3.8) and the monotonicity of  $f$  imply  $\text{sign } \lambda'(1) = -\text{sign } \frac{d}{d\beta} \bar{S}_0(1)$ , and we have  $\frac{\partial g}{\partial x}(S, x) = -c$  and  $\frac{\partial g}{\partial S}(S, x) = cf'(S)$  at those  $(S, x)$  for which  $f(S) > x$ . From (2.5)–(2.9) we obtain at  $(S_0, N_0, \beta) = (\bar{S}_1, 0, 1)$

$$\frac{\partial T}{\partial N_0} = \frac{f'(\bar{S}_1)}{f(\bar{S}_1) - x_\omega} e^{-cT} \int_0^T e^{(c-1)t} \int_0^t e^{(1-\mu)\tau} \gamma(\bar{S}_1, \bar{x}_1(\tau)) d\tau dt$$

and

$$\frac{\partial T}{\partial \beta} = -\frac{\partial h}{\partial \beta}(\bar{S}_1, 1) \frac{f'(\bar{S}_1)}{f(\bar{S}_1) - x_\omega} e^{-cT} \int_0^T e^{ct} (1 - e^{-t}) dt,$$

and (3.14), (2.5) and the assumptions on  $f$  and  $h$  yield the assertions.  $\square$

We consider now the case  $n \geq 3$ . The nontrivial fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta))$  of  $F_1$  for  $\beta \in U \cap (1, \infty)$  corresponds to a family of periodic points  $(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0})$ ,  $S_0 = \bar{S}_0(\beta)$ ,  $\sum_{i=1}^n N_{i,0} = \bar{N}_0(\beta)$ ,  $x_{i,0} = x_\alpha$  for  $i = 2, \dots, n$ , of period  $n$  of  $F_n$ . Let  $F_n^n : (S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0}) \mapsto (S_0^{(n)}, N_{1,0}^{(n)}, \dots, N_{n,0}^{(n)}, x_{2,0}^{(n)}, \dots, x_{n,0}^{(n)})$  and  $T_i := T(F_n^{i-1}(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0}))$ ,  $i = 1, \dots, n$ . Then  $N_{i,0}^{(n)} = k N_{i,0} e^{-\mu \sum_{j=1}^n T_j}$ ,  $i = 1, \dots, n$ , so the quotients  $N_{i,0}/N_{j,0}$  remain invariant under  $F_n^n$  if the denominators are not zero. For a vector  $\eta = (\eta_2, \dots, \eta_n)$ ,  $\eta_i > 0$  for  $i = 2, \dots, n$ , we define  $K_\eta : (S_0, N_0, x_{2,0}, \dots, x_{n,0}) \mapsto (S_0^{(n)}, N_0^{(n)}, x_{2,0}^{(n)}, \dots, x_{n,0}^{(n)})$  by

$$(S_0^{(n)}, \frac{1}{\zeta} N_0^{(n)}, \frac{\eta_2}{\zeta} N_0^{(n)}, \dots, \frac{\eta_n}{\zeta} N_0^{(n)}, x_{2,0}^{(n)}, \dots, x_{n,0}^{(n)}) := F_n^n(S_0, \frac{1}{\zeta} N_0, \frac{\eta_2}{\zeta} N_0, \dots, \frac{\eta_n}{\zeta} N_0, x_{2,0}, \dots, x_{n,0}),$$

where  $\zeta := 1 + \sum_{i=2}^n \eta_i$ . The Jacobi matrix of  $K_\eta$  at the nontrivial fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha, \dots, x_\alpha)$  is given by

$$(DK_\eta)(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha, \dots, x_\alpha) = \begin{pmatrix} A & * & \cdots & \cdots & \cdots & * \\ & * & \cdots & \cdots & \cdots & * \\ 0 & 0 & \frac{\partial x_{2,0}^{(n)}}{\partial x_{2,0}} & 0 & \cdots & 0 \\ \vdots & \vdots & 0 & \frac{\partial x_{3,0}^{(n)}}{\partial x_{3,0}} & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ \vdots & \vdots & \vdots & & \ddots & 0 \\ 0 & 0 & 0 & \cdots & \cdots & 0 \end{pmatrix} \frac{\partial x_{n,0}^{(n)}}{\partial x_{n,0}}, \quad (3.21)$$

where the  $(2 \times 2)$ -matrix  $A$  is again defined by (3.5) and the zeros in the last  $(n-1)$  rows are due to the fact that  $x_{i,0}^{(n)} = x_\alpha$  for  $x_{i,0} = x_\alpha$  independent of  $S_0, N_0$  and  $x_{j,0}$ ,  $j \neq i$ , for  $i = 2, \dots, n$ .

The following lemma is a generalisation of Lemma 3.1 to the case  $n \geq 3$ .

**Lemma 3.8.** *We have*

$$\frac{\partial x_{i,0}^{(n)}}{\partial x_{i,0}}(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha, \dots, x_\alpha) = \lambda(\beta), \quad i = 2, \dots, n,$$

where  $\lambda$  is given by (3.7).

*Proof.* Let  $n \geq 3$ . The quantities  $x_{i,0}^{(n)}$  for  $S_0 = \bar{S}_0(\beta)$ ,  $N_0 = \bar{N}_0(\beta)$  are obtained by solving the following recursive ODE systems. First solve the system

$$\begin{cases} \frac{dx_1^{(1)}}{dt} = g(S^{(1)}, x_1^{(1)}), & x_1^{(1)}(0) = x_\alpha, \\ \frac{dx_i^{(1)}}{dt} = g(S^{(1)}, x_i^{(1)}), & x_i^{(1)}(0) = x_{i,0}, \quad i = 2, \dots, n, \\ \frac{dS^{(1)}}{dt} = h(S^{(1)}, \beta) - \bar{N}_0(\beta)\zeta^{-1}e^{-\mu t} \sum_{i=1}^n \eta_i \gamma(S^{(1)}, x_i^{(1)}), & S^{(1)}(0) = \bar{S}_0(\beta), \end{cases}$$

where  $\eta_1 := 1$  and  $\zeta := \sum_{i=1}^n \eta_i$  as before. Let  $T_1 \geq 0$  be such that  $x_n^{(1)}(T_1) = x_\omega$ ,  $x_n^{(1)}(t) < x_\omega$  for  $0 \leq t < T_1$ , and set  $S_0^{(1)} := S^{(1)}(T_1)$ ,  $N_0^{(1)} := \bar{N}_0(\beta)e^{-\mu T_1}$  and  $x_{i,0}^{(1)} := x_{i-1}^{(1)}(T_1)$ ,  $i = 2, \dots, n$ .

For  $j = 2, \dots, n$  we solve recursively the ODE systems

$$\begin{cases} \frac{dx_1^{(j)}}{dt} = g(S^{(j)}, x_1^{(j)}), & x_1^{(j)}(0) = x_\alpha, \\ \frac{dx_i^{(j)}}{dt} = g(S^{(j)}, x_i^{(j)}), & x_i^{(j)}(0) = x_{i,0}^{(j-1)}, \quad i = 2, \dots, n, \\ \frac{dS^{(j)}}{dt} = h(S^{(j)}, \beta) - N_0^{(j-1)}\zeta^{-1}e^{-\mu t} \left( k \sum_{i=1}^{j-1} \eta_{n+i-j+1} \gamma(S^{(j)}, x_i^{(j)}) + \sum_{i=j}^n \eta_{i-j+1} \gamma(S^{(j)}, x_i^{(j)}) \right), \\ S^{(j)}(0) = S_0^{(j-1)}. \end{cases}$$

Let  $T_j \geq 0$  be such that  $x_n^{(j)}(T_j) = x_\omega$ ,  $x_n^{(j)}(t) < x_\omega$  for  $0 \leq t < T_j$ , and define  $S_0^{(j)} := S^{(j)}(T_j)$ ,  $N_0^{(j)} := N_0^{(j-1)}e^{-\mu T_j}$ ,  $x_{i,0}^{(j)} := x_{i-1}^{(j)}(T_j)$ ,  $i = 2, \dots, n$ .

For  $(x_{2,0}, \dots, x_{n,0}) = (x_\alpha, \dots, x_\alpha)$  we have  $T_1 = \bar{T}$  and  $T_j = 0$  for  $j = 2, \dots, n$ . Moreover, for  $j = 1, \dots, n$  we have  $x_{i,0}^{(j)} = x_\alpha$ ,  $i = 2, \dots, j$ , and  $x_{i,0}^{(j)} = x_\omega$ ,  $i = j+1, \dots, n$ . We shall in the following often use the fact that

$$\frac{\partial x_i^{(j)}}{\partial x_{i,0}^{(j-1)}}(0) = 1, \quad i = 2, \dots, n, \quad (3.22)$$

and

$$\frac{\partial x_i^{(j)}}{\partial N_0^{(j-1)}}(0) = \frac{\partial x_i^{(j)}}{\partial S_0^{(j-1)}}(0) = \frac{\partial x_i^{(j)}}{\partial x_{k,0}^{(j-1)}}(0) = 0, \quad k \neq i, \quad i = 1, \dots, n, \quad (3.23)$$

for  $j = 2, \dots, n$ .

For the remainder of the proof we assume that all partial derivatives are taken at  $(x_{2,0}, \dots, x_{n,0}) = (x_\alpha, \dots, x_\alpha)$ . By differentiating the equations  $x_n^{(j)}(T_j) = x_\omega$ ,  $j = 1, \dots, n$ , and by using (3.22) and (3.23) we obtain for  $i = 2, \dots, n$

$$\frac{\partial T_1}{\partial x_{i,0}} = -\frac{1}{g(\bar{S}_0(\beta), x_\omega)} \frac{\partial x_n^{(1)}}{\partial x_{i,0}}(\bar{T})$$

and

$$\frac{\partial T_j}{\partial x_{i,0}} = -\frac{1}{g(\bar{S}_0(\beta), x_\omega)} \frac{\partial x_{n,0}^{(j-1)}}{\partial x_{i,0}}, \quad j = 2, \dots, n.$$

For  $j = 2, \dots, n$ ,  $i = 2, \dots, n$  we have

$$\frac{\partial x_{2,0}^{(j)}}{\partial x_{i,0}} = \dot{x}_1^{(j)}(0) \frac{\partial T_j}{\partial x_{i,0}} = -\frac{g(\bar{S}_0(\beta), x_\alpha)}{g(\bar{S}_0(\beta), x_\omega)} \frac{\partial x_{n,0}^{(j-1)}}{\partial x_{i,0}} \quad (3.24)$$

and

$$\begin{aligned} \frac{\partial x_{k,0}^{(j)}}{\partial x_{i,0}} &= \dot{x}_{k-1}^{(j)}(0) \frac{\partial T_j}{\partial x_{i,0}} + \frac{\partial x_{k-1,0}^{(j-1)}}{\partial x_{i,0}} = \frac{\partial x_{k-1,0}^{(j-1)}}{\partial x_{i,0}} - \frac{g(\bar{S}_0(\beta), x_{k-1,0}^{(j-1)})}{g(\bar{S}_0(\beta), x_\omega)} \frac{\partial x_{n,0}^{(j-1)}}{\partial x_{i,0}} \\ &= \begin{cases} \frac{\partial x_{k-1,0}^{(j-1)}}{\partial x_{i,0}} - \frac{g(\bar{S}_0(\beta), x_\alpha)}{g(\bar{S}_0(\beta), x_\omega)} \frac{\partial x_{n,0}^{(j-1)}}{\partial x_{i,0}}, & k = 3, \dots, j, \\ \frac{\partial x_{k-1,0}^{(j-1)}}{\partial x_{i,0}} - \frac{\partial x_{n,0}^{(j-1)}}{\partial x_{i,0}}, & k = j+1, \dots, n. \end{cases} \end{aligned} \quad (3.25)$$

The derivatives of  $x_{k,0}^{(1)}$  are given by

$$\frac{\partial x_{k,0}^{(1)}}{\partial x_{i,0}} = \dot{x}_{k-1}^{(1)}(\bar{T}) \frac{\partial T_1}{\partial x_{i,0}} + \frac{\partial x_{k-1,0}^{(1)}}{\partial x_{i,0}}(\bar{T}) = \frac{\partial x_{k-1,0}^{(1)}}{\partial x_{i,0}}(\bar{T}) - \frac{\partial x_n^{(1)}}{\partial x_{i,0}}(\bar{T}), \quad k, i = 2, \dots, n. \quad (3.26)$$

From (3.24)–(3.26) we obtain recursively

$$\frac{\partial x_{i,0}^{(n)}}{\partial x_{i,0}} = -\frac{g(\bar{S}_0(\beta), x_\alpha)}{g(\bar{S}_0(\beta), x_\omega)} \left( \frac{\partial x_1^{(1)}}{\partial x_{i,0}}(\bar{T}) - \frac{\partial x_i^{(1)}}{\partial x_{i,0}}(\bar{T}) \right), \quad i = 2, \dots, n.$$

Similarly as in the proof of Lemma 3.1 we have

$$\frac{\partial x_i^{(1)}}{\partial x_{i,0}}(\bar{T}) - \frac{\partial x_1^{(1)}}{\partial x_{i,0}}(\bar{T}) = \exp\left(-\int_0^{\bar{T}} \frac{\partial g}{\partial x}(\bar{S}_\beta(t), \bar{x}_\beta(t)) dt\right), \quad i = 2, \dots, n,$$

which finishes the proof.  $\square$

Thus the eigenvalues of the Jacobi matrix (3.21) are the eigenvalues of the matrix  $A$  and the  $(n-1)$ -fold eigenvalue  $\lambda$ , and we obtain the following result.

**Theorem 3.9.** *Let  $\beta \in U \cap (1, \infty)$ . Then the iterations of  $F_n$  converge to the stable fixed point  $(\bar{S}_0(\beta), \bar{N}_0(\beta))$  of  $F_1$  for initial values  $(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0})$  such that  $(S_0, \sum_{i=1}^n N_{i,0}, x_{2,0}, \dots, x_{n,0})$  is sufficiently close to  $(\bar{S}_0(\beta), \bar{N}_0(\beta), x_\alpha, \dots, x_\alpha)$  for all  $n \geq 2$  if and only if this is the case for  $n = 2$ .*

We summarise our results in

**Theorem 3.10.** *Let  $\lambda'(1) < 0$ . Then there exists a  $\beta_1 > 1$  such that for each  $\beta \in (1, \beta_1)$  there exist a neighbourhood  $V_\beta$  of  $(\bar{S}_0(\beta), \bar{N}_0(\beta))$  and an  $\varepsilon_\beta > 0$  such that  $\lim_{k \rightarrow \infty} F_n^{nk}(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0}) = (\bar{S}_0(\beta), \frac{N_{1,0}\bar{N}_0(\beta)}{\sum_{i=1}^n N_{i,0}}, \dots, \frac{N_{n,0}\bar{N}_0(\beta)}{\sum_{i=1}^n N_{i,0}}, x_\alpha, \dots, x_\alpha)$  for  $(S_0, \sum_{i=1}^n N_{i,0}) \in V_\beta$ ,  $0 \leq x_{i,0} - x_\alpha < \varepsilon_\beta$  for  $i = 2, \dots, n$  and all  $n \geq 2$ .*

No general statement can be made about sign  $\lambda'(1)$ ; cf. Section 4.

We conclude this section with an investigation of fixed points of  $F_n$  for  $n \geq 2$ . For a fixed point  $(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0})$  of  $F_n$  we must have  $N_{1,0} = ke^{-\mu T} N_{n,0}$  and  $N_{i,0} = e^{-\mu T} N_{i-1,0}$ ,  $i = 2, \dots, n$ , which implies that either  $N_{i,0} = 0$  for  $i = 1, \dots, n$  or

$$T = \frac{1}{n\mu} \ln k \quad (3.27)$$



and

$$N_{i,0} = k^{-(i-1)/n} N_{1,0}, \quad i = 2, \dots, n. \quad (3.28)$$

For values of  $\beta$  such that  $(H_\beta)$  is satisfied we define the function  $P_\beta : [x_\alpha, x_\omega] \rightarrow \mathbb{R}$  by

$$P_\beta(x) := \int_{x_\alpha}^x \frac{d\sigma}{g(\tilde{S}_\beta, \sigma)}, \quad x \in [x_\alpha, x_\omega].$$

Then  $P_\beta$  is strictly increasing on  $[x_\alpha, x_\omega]$ ,  $P_\beta(x_\alpha) = 0$ ,  $P_\beta(x_\omega) = \tilde{T}_\beta$ , which implies that  $P_\beta^{-1} : [0, \tilde{T}_\beta] \rightarrow [x_\alpha, x_\omega]$  exists. The point  $(\tilde{S}_\beta, 0, \dots, 0, x_{2,0}, \dots, x_{n,0})$  is a periodic point of period  $\leq n$  for all choices of  $x_{i,0}$ ,  $i = 2, \dots, n$ . It is a fixed point of  $F_n$  if and only if

$$x_{i,0} = P_\beta^{-1}\left(\frac{i-1}{n} \tilde{T}_\beta\right), \quad i = 2, \dots, n,$$

and in that case we have  $T = \frac{1}{n} \tilde{T}_\beta$ . We define the function  $G : W \subset \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ ,  $W$  an appropriately chosen neighbourhood of  $(\tilde{S}_1, 0, \bar{x}_{2,0}, \dots, \bar{x}_{n,0}, 1)$ , where

$$\bar{x}_{i,0} := P_1^{-1}\left(\frac{i-1}{n\mu} \ln k\right), \quad i = 2, \dots, n, \quad (3.29)$$

as follows. For  $(S_0, N_0, x_{2,0}, \dots, x_{n,0}, \beta) \in W$  we solve the ODE system

$$\begin{cases} \frac{dx_1}{dt} = g(S, x_1), & x_1(0) = x_\alpha, \\ \frac{dx_i}{dt} = g(S, x_i), & x_i(0) = x_{i,0}, \quad i = 2, \dots, n, \\ \frac{dS}{dt} = h(S, \beta) - N_0 e^{-\mu t} \sum_{i=1}^n k^{-(i-1)/n} \gamma(S, x_i), & S(0) = S_0. \end{cases} \quad (3.30)$$

Let  $T \geq 0$  be such that  $x_n(T) = x_\omega$ ,  $x_n(t) < x_\omega$  for  $0 \leq t < T$ , and define

$$\begin{cases} G_1(S_0, N_0, x_{2,0}, \dots, x_{n,0}, \beta) := S(T) - S_0, \\ G_2(S_0, N_0, x_{2,0}, \dots, x_{n,0}, \beta) := k e^{-n\mu T} - 1, \\ G_i(S_0, N_0, x_{2,0}, \dots, x_{n,0}, \beta) := x_{i-2}(T) - x_{i-1,0}, \quad i = 3, \dots, n. \end{cases} \quad (3.31)$$

If  $W$  is chosen sufficiently small,  $G$  is continuously differentiable in  $W$ , and (2.12) and (3.29) imply  $G(\tilde{S}_1, 0, \bar{x}_{2,0}, \dots, \bar{x}_{n,0}, 1) = 0$ . In view of (3.27) and (3.28)  $(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0})$  is a non-trivial fixed point of  $F_n$  if and only if  $N_{i,0} = k^{-(i-1)/n} N_0$ ,  $i = 1, \dots, n$ , and  $G(S_0, N_0, x_{2,0}, \dots, x_{n,0}, \beta) = 0$ .

We want to apply the Implicit Function Theorem to the mapping  $G$ . To this end we first compute the derivatives of the solutions of the ODE system (3.30) with respect to initial data and parameters.

**Lemma 3.11.** *Let  $(x_1, \dots, x_n, S)$  be the solution of (3.30), let  $T$  be defined as above and let*

$$\bar{x}_i(t) := P_1^{-1}\left(t + \frac{i-1}{n\mu} \ln k\right), \quad 0 \leq t \leq \frac{n-i+1}{n\mu} \ln k, \quad i = 1, \dots, n.$$

*Then the partial derivatives at  $(S_0, N_0, x_{2,0}, \dots, x_{n,0}, \beta) = (\tilde{S}_1, 0, \bar{x}_{2,0}, \dots, \bar{x}_{n,0}, 1)$  are given by*

$$\frac{\partial S}{\partial S_0}(t) = e^{-t}, \quad (3.32)$$

$$\frac{\partial x_i}{\partial S_0}(t) = \int_0^t \exp\left(\int_\tau^t \frac{\partial g}{\partial x}(\tilde{S}_1, \bar{x}_i(\sigma)) d\sigma\right) \frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_i(\tau)) e^{-\tau} d\tau, \quad i = 1, \dots, n, \quad (3.33)$$

$$\frac{\partial S}{\partial x_{j,0}}(t) = 0, \quad j = 2, \dots, n, \quad (3.34)$$

$$\frac{\partial x_i}{\partial x_{j,0}}(t) = 0, \quad i = 1, \dots, n, \quad j = 2, \dots, n, \quad j \neq i, \quad (3.35)$$

$$\frac{\partial x_i}{\partial x_{i,0}}(t) = \exp\left(\int_0^t \frac{\partial g}{\partial x}(\tilde{S}_1, \bar{x}_i(\tau)) d\tau\right), \quad i = 2, \dots, n, \quad (3.36)$$

$$\frac{\partial S}{\partial N_0}(t) = -e^{-t} \sum_{i=1}^n k^{-(i-1)/n} \int_0^t e^{(1-\mu)\tau} \gamma(\tilde{S}_1, \bar{x}_i(\tau)) d\tau, \quad (3.37)$$

$$\frac{\partial x_i}{\partial N_0}(t) = \int_0^t \exp\left(\int_\tau^t \frac{\partial g}{\partial x}(\tilde{S}_1, \bar{x}_i(\sigma)) d\sigma\right) \frac{\partial g}{\partial S}(\tilde{S}_1, \bar{x}_i(\tau)) \frac{\partial S}{\partial N_0}(\tau) d\tau, \quad i = 1, \dots, n, \quad (3.38)$$

$$\frac{\partial T}{\partial \alpha} = -g(\tilde{S}_1, x_\omega)^{-1} \frac{\partial x_n}{\partial \alpha}(T) \text{ for } \alpha = S_0, N_0 \text{ and } x_{i,0}, \quad i = 2, \dots, n, \quad (3.39)$$

and in particular

$$\frac{\partial T}{\partial x_{j,0}} = 0, \quad j = 2, \dots, n-1. \quad (3.40)$$

The following theorem is an analogue to Theorem 2.4 for  $F_n$ ,  $n \geq 2$ .

**Theorem 3.12.** *Let  $n \geq 2$ . There exists a  $\beta_1 > 1$  such that for each  $\beta \in (1, \beta_1)$  there exists a fixed point  $(S_0, N_{1,0}, \dots, N_{n,0}, x_{2,0}, \dots, x_{n,0})$ ,  $N_{i,0} = k^{-(i-1)/n} N_0$ ,  $i = 1, \dots, n$ , of  $F_n$  with  $N_0 > 0$ .*

*Proof.* We apply the Implicit Function Theorem to the function  $G$  defined by (3.31). Let  $y := (S_0, N_0, x_{2,0}, \dots, x_{n,0})$  and  $\bar{y} := (\tilde{S}_1, 0, \bar{x}_{2,0}, \dots, \bar{x}_{n,0})$ . Then we have

$$(D_y G)(\bar{y}, 1) = \begin{pmatrix} \frac{\partial G_1}{\partial S_0} & \frac{\partial G_1}{\partial N_0} & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & 0 \\ \frac{\partial G_2}{\partial S_0} & \frac{\partial G_2}{\partial N_0} & 0 & \cdots & \cdots & \cdots & \cdots & 0 & \frac{\partial G_2}{\partial x_{n,0}} \\ \frac{\partial G_3}{\partial S_0} & \frac{\partial G_3}{\partial N_0} & -1 & 0 & \cdots & \cdots & \cdots & 0 & \frac{\partial G_3}{\partial x_{n,0}} \\ \frac{\partial G_4}{\partial S_0} & \frac{\partial G_4}{\partial N_0} & \frac{\partial G_4}{\partial x_{2,0}} & -1 & \ddots & & & \vdots & \vdots \\ \frac{\partial G_5}{\partial S_0} & \frac{\partial G_5}{\partial N_0} & 0 & \frac{\partial G_5}{\partial x_{3,0}} & \ddots & \ddots & & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \vdots & & \ddots & \ddots & \ddots & 0 & \frac{\partial G_{n-1}}{\partial x_{n,0}} \\ \vdots & \vdots & \vdots & & & \ddots & \ddots & -1 & \frac{\partial G_n}{\partial x_{n,0}} \\ \frac{\partial G_{n+1}}{\partial S_0} & \frac{\partial G_{n+1}}{\partial N_0} & 0 & \cdots & \cdots & \cdots & 0 & \frac{\partial G_{n+1}}{\partial x_{n-1,0}} & \frac{\partial G_{n+1}}{\partial x_{n,0}} \end{pmatrix}$$

since (3.31), (3.34), (3.35) and (3.40) imply  $\frac{\partial G_1}{\partial x_{i,0}} = 0$ ,  $i = 2, \dots, n$ ,  $\frac{\partial G_2}{\partial x_{i,0}} = 0$ ,  $i = 2, \dots, n-1$ ,  $\frac{\partial G_{i+1}}{\partial x_{i,0}} = -1$ ,  $i = 2, \dots, n-1$ , and  $\frac{\partial G_i}{\partial x_{j,0}} = 0$ ,  $i = 3, \dots, n$ ,  $j \neq i-2, i-1, n$  at  $(y, \beta) = (\bar{y}, 1)$ . After a tedious but elementary computation we obtain

$$\begin{aligned} & \det(D_y G)(\bar{y}, 1) \\ &= (-1)^n \sum_{i=3}^{n+1} \left( \frac{\partial G_1}{\partial S_0} \left( \frac{\partial G_2}{\partial N_0} \frac{\partial G_i}{\partial x_{n,0}} - \frac{\partial G_2}{\partial x_{n,0}} \frac{\partial G_i}{\partial N_0} \right) - \frac{\partial G_1}{\partial N_0} \left( \frac{\partial G_2}{\partial S_0} \frac{\partial G_i}{\partial x_{n,0}} - \frac{\partial G_2}{\partial x_{n,0}} \frac{\partial G_i}{\partial S_0} \right) \right) \prod_{j=i-1}^{n-1} \frac{\partial G_{j+2}}{\partial x_{j,0}}. \end{aligned}$$

Differentiation of (3.31) at  $(y, \beta) = (\bar{y}, 1)$  yields

$$\frac{\partial G_2}{\partial N_0} \frac{\partial G_i}{\partial x_{n,0}} - \frac{\partial G_2}{\partial x_{n,0}} \frac{\partial G_i}{\partial N_0} = n\mu \left( \frac{\partial T}{\partial N_0} \delta_{i,n+1} + \frac{\partial T}{\partial x_{n,0}} \frac{\partial x_{i-2}}{\partial N_0}(\bar{T}_n) \right)$$

and

$$\frac{\partial G_2}{\partial S_0} \frac{\partial G_i}{\partial x_{n,0}} - \frac{\partial G_2}{\partial x_{n,0}} \frac{\partial G_i}{\partial S_0} = n\mu \left( \frac{\partial T}{\partial S_0} \delta_{i,n+1} + \frac{\partial T}{\partial x_{n,0}} \frac{\partial x_{i-2}}{\partial S_0}(\bar{T}_n) \right)$$

for  $i = 3, \dots, n+1$ , where we have used (2.12),  $\bar{T}_n := \frac{1}{n\mu} \ln k$  and  $\delta_{ij}$  denotes the Kronecker delta, and further

$$\begin{aligned} \det(D_y G)(\bar{y}, 1) &= (-1)^n n\mu \left( - \left( 1 - \frac{\partial S}{\partial S_0}(\bar{T}_n) \right) \frac{\partial T}{\partial N_0} - \frac{\partial S}{\partial N_0}(\bar{T}_n) \frac{\partial T}{\partial S_0} \right. \\ &\quad \left. + \sum_{i=3}^{n+1} \frac{\partial T}{\partial x_{n,0}} \left( - \left( 1 - \frac{\partial S}{\partial S_0}(\bar{T}_n) \right) \frac{\partial x_{i-2}}{\partial N_0}(\bar{T}_n) - \frac{\partial S}{\partial N_0}(\bar{T}_n) \frac{\partial x_{i-2}}{\partial S_0}(\bar{T}_n) \right) \prod_{j=i-1}^{n-1} \frac{\partial x_j}{\partial x_{j,0}}(\bar{T}_n) \right). \end{aligned}$$

From (3.32), (3.33) and (3.36)–(3.39) and assumptions  $(H_g)$  and  $(H_\gamma)$  we infer  $\frac{\partial S}{\partial S_0}(\bar{T}_n) < 1$ ,  $\frac{\partial S}{\partial N_0}(\bar{T}_n) < 0$ ,  $\frac{\partial T}{\partial S_0} < 0$ ,  $\frac{\partial T}{\partial N_0} > 0$ ,  $\frac{\partial T}{\partial x_{n,0}} < 0$ ,  $\frac{\partial x_{i-2}}{\partial S_0}(\bar{T}_n) > 0$ ,  $\frac{\partial x_{i-2}}{\partial N_0}(\bar{T}_n) < 0$ ,  $i = 3, \dots, n+1$ , and  $\frac{\partial x_j}{\partial x_{j,0}}(\bar{T}_n) > 0$ ,  $j = 2, \dots, n$ , which implies  $\det(D_y G)(\bar{y}, 1) \neq 0$  and  $\text{sign}(\det(D_y G)(\bar{y}, 1)) = (-1)^{n-1}$ . Thus we can apply the Implicit Function Theorem to  $G$  and obtain a nontrivial biological meaningful fixed point of  $F_n$  for  $\beta \in (1, \beta_1)$ ,  $\beta_1 > 1$  suitably chosen.  $\square$

#### 4. NUMERICAL RESULTS

In the first example we investigate  $\text{sign } \lambda'(1)$  numerically for the case of a linear growth rate (1.2) and a particular kind of consumption rate  $\gamma$ .

**Example 4.1.** We consider a linear growth rate (1.2) and scale the size variable such that  $x_\alpha = 1$ . Let the consumption rate be given by (cf. [2, p. 21])

$$\gamma(S, x) = f_1(S)x^2, \quad (4.1)$$

where  $f_1$  is a strictly increasing function of  $S$  with  $f_1(0) = 0$ , and assume that the conditions (2.12) and (2.13) are satisfied, i.e.

$$\frac{1}{\mu} \ln k = \frac{1}{c} \ln \frac{f(\tilde{S}_1) - 1}{f(\tilde{S}_1) - x_\omega}$$

or

$$f(\tilde{S}_1) = \frac{k^{\frac{c}{\mu}} x_\omega - 1}{k^{\frac{c}{\mu}} - 1}.$$

Then we have

$$\bar{x}_1(t) = \frac{k^{\frac{c}{\mu}} x_\omega - 1}{k^{\frac{c}{\mu}} - 1} - e^{-ct} \frac{k^{\frac{c}{\mu}} (x_\omega - 1)}{k^{\frac{c}{\mu}} - 1}.$$

By Lemma 3.7 the stability of the nontrivial fixed point of the one-cohort problem within the set of  $n$ -cohort solutions depends on the sign of the function

$$\begin{aligned} q(c, \mu, k, x_\omega) &:= k^{-\frac{1}{\mu} \ln k} \int_0^{\frac{1}{\mu} \ln k} e^{ct} (1 - e^{-t}) dt \int_0^{\frac{1}{\mu} \ln k} e^{(1-\mu)t} \left( \frac{k^{\frac{c}{\mu}} x_\omega - 1}{k^{\frac{c}{\mu}} - 1} - e^{-ct} \frac{k^{\frac{c}{\mu}} (x_\omega - 1)}{k^{\frac{c}{\mu}} - 1} \right)^2 dt \\ &\quad - (1 - k^{-\frac{1}{\mu}}) \int_0^{\frac{1}{\mu} \ln k} e^{(c-1)t} \int_0^t e^{(1-\mu)\tau} \left( \frac{k^{\frac{c}{\mu}} x_\omega - 1}{k^{\frac{c}{\mu}} - 1} - e^{-c\tau} \frac{k^{\frac{c}{\mu}} (x_\omega - 1)}{k^{\frac{c}{\mu}} - 1} \right)^2 d\tau dt, \end{aligned}$$

where  $c > 0$ ,  $\mu > 0$ ,  $k > 1$  and  $x_\omega > 1$  and these parameters are in general not connected by any condition and cannot be normalised by any additional scaling.

First we considered the case  $kx_\alpha^3 = k = x_\omega^3$  (conservation of mass). The function  $q$  was computed numerically for a grid of  $(c, \mu, x_\omega) \in [0.001, 100] \times [0.001, 100] \times [1.25, 4.65]$  (i.e. the cases  $k = 2$  up to  $k = 100$  are included in this set) and the function  $q$  was always negative, which means that the fixed point of the one-cohort problem is stable for  $F_n$ ,  $n \geq 2$ .

Next we let  $k = 2$  fixed (division into two parts) and computed  $q$  for a grid of  $(c, \mu, x_\omega) \in [0.001, 100] \times [0.001, 100] \times [\sqrt[3]{2}, 10]$  (i.e.  $kx_\alpha^3 \leq x_\omega^3$ ). There exists an increasing function  $\mu_1 : [0.001, 100] \rightarrow [0.001, 10]$  such that for  $(c, \mu) \in [0.001, 100] \times [0.001, 100]$  with  $\mu \geq \mu_1(c)$  there exists an  $x_1(c, \mu) \in [\sqrt[3]{2}, 10]$  such that  $q(c, \mu, x_\omega) \geq 0$  for  $x_1(c, \mu) \leq x_\omega \leq 10$  and  $q$  negative elsewhere on  $[0.001, 100] \times [0.001, 100] \times [\sqrt[3]{2}, 10]$ . The function  $x_1$  is decreasing in  $\mu$  and increasing in  $c$ . For other values of  $k$  we obtained similar results, i.e. positive values of  $q$  could only occur for  $kx_\alpha^3 < x_\omega^3$ .

The mapping  $F_n$ ,  $n \leq 10$ , was iterated numerically for linear growth rates (1.2) and consumption rates (4.1) with

$$f(S) = \frac{c_1 S}{c_2 S + 1}, \quad f_1(S) = \frac{c_3 S}{c_2 S + 1}, \quad (4.2)$$

where  $c_1$ ,  $c_2$  and  $c_3$  are positive constants (Holling type II functional response, cf. [2]), and linear or logistic  $h$  for  $kx_\alpha^3 = x_\omega^3$  (i.e. we considered cases where  $\lambda'(1) < 1$  according to Example 4.1),  $\beta > 1$  and different values of the initial data and the parameters. In all of our examples the iterations converged to a nontrivial fixed point of  $F_1$  even if the initial data were not chosen close to this fixed point.

In the following example we investigate the asymptotic behaviour numerically for a case where the fixed point of the one-cohort problem is not stable for  $F_n$ ,  $n \geq 2$ .

**Example 4.2.** We consider a linear growth rate (1.2) and a consumption rate (4.1) with (4.2), where  $c = 1.5$ ,  $c_1 = c_3 = 1$ ,  $c_2 = 0.4$ , and let  $\mu = 0.5$ ,  $k = 2$ ,  $x_\alpha = 1$ ,  $x_\omega = 2$  and  $h(S, \beta) = 15\beta - S$ . The trivial fixed point of  $F_1$  becomes unstable at  $\beta = 1$  and we have  $q(1.5, 0.5, 2, 2) > 0$  (i.e. we chose the parameters such that  $q$  is positive). The nontrivial fixed point of  $F_1$  was computed for  $\beta = \frac{16}{15}$ ,  $\frac{17}{15}$ ,  $\frac{6}{5}$ ,  $\frac{19}{15}$  and  $\frac{4}{3}$  (cf. Table 1). The function  $\bar{S}_0$  is strictly decreasing on  $[1, \frac{4}{3}]$ .

$\beta$	$\bar{S}_0(\beta)$	$\bar{N}_0(\beta)$
$\frac{16}{15}$	14.97	0.2392
$\frac{17}{15}$	14.94	0.4782
$\frac{6}{5}$	14.92	0.7170
$\frac{19}{15}$	14.89	0.9556
$\frac{4}{3}$	14.86	1.1939

Table 1

For  $\beta = \frac{19}{15}$  we investigated the asymptotic behaviour of  $F_n$  numerically for  $n = 2, 3, 4$ . Note that the quotients  $N_{i,0}/N_{j,0}$  remain invariant under  $F_n^n$ . For  $n = 2$  we have after one application of  $F_2$   $N'_{1,0}/N'_{2,0} = kN_{2,0}/N_{1,0}$ . We iterated  $F_2$  for different start values and the iterations converged to different periodic points of period two, depending on the start value of  $N_{1,0}/N_{2,0}$ . In the case that we started with  $N_{1,0}/N_{2,0} = \sqrt{k}$  the periodic point of period two degenerated to the fixed point (14.95, 0.4847, 0.3427, 1.739) of  $F_2$ . In Table 2 we show some periodic points of period two of  $F_2$ . Similarly periodic points of period  $n$  were found by iterating  $F_n$  for  $n = 3, 4$ , and these periodic points

can again degenerate to fixed points. By replacing  $h$  by  $h(S, \beta) = S(1 - \frac{S}{15\beta})$  we obtained similar results.

$S_0$	$N_{1,0}$	$N_{2,0}$	$x_{2,0}$
15.09	0.0814	0.8145	1.222
14.89	0.8748	0.0437	1.966
15.07	0.2960	0.5921	1.330
14.88	0.6632	0.1658	1.942
15.06	0.3534	0.5301	1.383
14.88	0.6074	0.2025	1.928
15.03	0.4345	0.4345	1.515
14.89	0.5304	0.2652	1.883

Table 2

Example 4.2 suggests that in the “unstable” case ( $\lambda'(1) > 1$ ) there exist families of periodic points of period  $n$  of  $F_n$ ,  $n \geq 2$ , for  $1 < \beta < \beta_1$  that can be “degenerate” (i.e. periodic points of period  $m$ ,  $1 \leq m < n$ ,  $m$  a divisor of  $n$ ) and belong to the domain of attraction of  $F_n$ .

## 5. CONCLUDING REMARKS

This simple population model exhibits the feature that cohorts “synchronise” in the course of time at least for particular choices of the functions, parameters and initial values. This behaviour is well-documented by numerical results. It still remains to determine the domains of attraction in the “unstable” case ( $\lambda'(1) > 0$ ). We conjecture that in that case there are families of periodic points of period  $n$  (that can be “degenerate”, i.e. periodic points of period  $m$ ,  $m$  a divisor of  $n$ ) of  $F_n$  for all  $n$  belonging to the domain of attraction of  $F_n$ . Moreover, the results we obtained are only local in nature. Since the mappings  $F_n$  might have discontinuities and it is not easy to quantify where they occur, it is difficult to make global statements about the existence and stability of periodic points. However, if  $S_x = 0$  for all  $x \in [x_\alpha, x_\omega]$  in the condition  $(H_g)$ , there are no discontinuities.

## ACKNOWLEDGEMENT

Part of this work was done during a ten-month stay at the CWI supported by NUFFIC. I would like to thank the CWI for its hospitality. Moreover, I am grateful to Odo Diekmann for stimulating discussions and valuable suggestions, which considerably helped to improve an earlier version of this paper.

## REFERENCES

1. S.A.L.M. Kooijman, *Dynamic Energy Budgets in Biological Systems: Theory and Applications in Ecotoxicology*, Cambridge University Press, Cambridge, 1993.
2. J.A.J. Metz and O. Diekmann (eds.), *The Dynamics of Physiologically Structured Populations*, Lecture Notes in Biomathematics **68**, Springer, Berlin, 1986.
3. N.M. Temme (ed.), *Nonlinear Analysis Vol. I*, MC Syllabus, Mathematisch Centrum, Amsterdam, 1976.