



Centrum voor Wiskunde en Informatica
REPORTRAPPORT

Non-existence and Uniqueness Results for the Extended Fisher-Kolmogorov
Equation

M. A. Peletier

Modelling, Analysis and Simulation (MAS)

MAS-R9903 March 1999

Report MAS-R9903
ISSN 1386-3703

CWI
P.O. Box 94079
1090 GB Amsterdam
The Netherlands

CWI is the National Research Institute for Mathematics and Computer Science. CWI is part of the Stichting Mathematisch Centrum (SMC), the Dutch foundation for promotion of mathematics and computer science and their applications.

SMC is sponsored by the Netherlands Organization for Scientific Research (NWO). CWI is a member of ERCIM, the European Research Consortium for Informatics and Mathematics.

Copyright © Stichting Mathematisch Centrum
P.O. Box 94079, 1090 GB Amsterdam (NL)
Kruislaan 413, 1098 SJ Amsterdam (NL)
Telephone +31 20 592 9333
Telefax +31 20 592 4199

Non-existence and Uniqueness Results for the Extended Fisher-Kolmogorov Equation

Mark A Peletier

CWI

P.O. Box 94079, 1090 GB Amsterdam, The Netherlands

ABSTRACT

We give a classification of all bounded solutions of the equation

$$u'''' + pu'' + F'(u) = 0, \quad -\infty < t < \infty,$$

in which F is a general quartic polynomial and p is restricted to various subsets of $(-\infty, 0]$. These results are obtained by combining an a priori estimate with geometric arguments in the (u, u'') -plane.

1991 Mathematics Subject Classification: 58F05, 34A26, 34A34, 34A46

Keywords and Phrases: Hamiltonian systems, Lorentz-Lagrangian systems, phase plane analysis

Note: This work was partially carried out under CWI Theme 'Modelling, Analysis, and Simulation'

1. INTRODUCTION

Starting with the work of Toland [1] and Hofer & Toland [2] a technique has been developed to exploit the special structure of certain Hamiltonian systems with an indefinite kinetic energy [3, 4, 5]. This technique provides a geometrical tool akin to the phase plane analysis that has proven so powerful in the case of two-dimensional dynamical systems. Geometrical properties of a pseudo-phase space of reduced dimension are used to prove existence, uniqueness, and many qualitative properties of solutions. The technique and the monotonicity property on which it is based are briefly explained in Section 4.

In this paper we combine this technique with an a priori estimate to study the solution set of the equation

$$u'''' + pu'' + F'(u) = 0, \quad -\infty < t < \infty, \quad p \leq 0. \quad (1.1)$$

While the methods used in this paper fall inside the pseudo-phase plane setting mentioned above, equation (1.1) has also been studied using several other techniques, such as variational methods [6, 7, 8, 9], and a topological shooting method [10, 11, 12, 13, 14]. The novelty that this paper brings to the pseudo-phase plane concept is the extension of the existing uniqueness and non-existence results, which typically are proven under

certain restrictions on the solutions, to the set of all bounded solutions. A particularly interesting application is the change in bifurcation diagram that is discussed in Section 2.

In equation (1.1) we are interested in potentials F that have one or two wells, and for simplicity we assume that F is an exact quartic. This is not essential, and for any double-well potential of similar type analogous results should hold. By exploiting the various symmetries present in (1.1) we can reduce the case of a general quartic to

$$F(u) = a + \frac{1}{2}u^2 - \frac{1}{3}u^3 + \frac{b}{4}u^4, \quad b > 0,$$

without losing generality. Note that one well is at $u = 0$, that $F''(0) = 1$, and that the second well, if it exists (which is the case for $0 < b < 1/4$) has $u > 0$. When $b = 2/9$, equation (1.1) is the Extended Fisher-Kolmogorov equation with wells of equal depth (at $u = 0$ and $u = 3$).

For a solution of (1.1), the Hamiltonian,

$$H(u) = u'u''' - \frac{1}{2}u''^2 + \frac{p}{2}u'^2 + F(u), \quad (1.2)$$

is constant in time. By incorporating this constant into the definition of F (in the zero-order term a) we can restrict ourselves to solutions with zero Hamiltonian, again without any loss of generality. In order to prevent needless repetition, we state here that *all solutions are assumed to satisfy $H = 0$* .

Equation (1.1) has a constant solution $u \equiv \alpha$ if and only if $F'(\alpha) = 0$. If $b \geq 1/4$, then F has only one well, and the only bounded solutions are constants:

Theorem 1 (Single-well case). *If $b \geq 1/4$ and $p \leq 0$, then equation (1.1) has no bounded solutions other than (possibly) the unique constant solution.*

Note that for $b = 1/4$ the function F has two stationary points: $F'(0) = F'(2) = 0$, and therefore there are two stationary solutions. However, since $F(0) \neq F(2)$, these occur for different values of a , so that the constant solutions are indeed unique (as stated in Theorem 1) under the condition $H = 0$.

More interesting are the double-well potentials. In Figure 1 we depict eight different classes of functions F , where the distinction is made on the basis of the distribution of signs and zeros of F , and their position with respect to the wells. When one well is at level zero, as in cases (b) and (f), we assume for definiteness that this is the well at $u = 0$.

In the theorem below, the main result of this paper, we state uniqueness and non-existence results for bounded solutions. These hold for values of p in intervals of the type $(-\infty, \bar{p}]$ or $(-\infty, \bar{p})$, where $-\infty < \bar{p} \leq 0$ is determined by the function F . It is of course possible to specify these limits exactly in terms of the values of a and b , but here we only do that in isolated cases.




Figure 1. Eight different cases for F , classified by the character of the roots.

- Theorem 2 (Double-well case).**
1. In cases (a) and (d) there are no bounded solutions for any $p \leq p_1$, $p_1 \leq 0$.
 2. In case (b) the only bounded solution for $p \leq p_2$, $p_2 > -2$, is the stationary solution $u \equiv 0$.
 3. In case (c) the only bounded solutions for $p \leq -2$ are the constants $u \equiv 0$ and $u \equiv 3$ and two heteroclinic solutions connecting 0 and 3.
 4. In case (e) there are no bounded solutions for any $p \leq 0$.
 5. In case (f) the only bounded solutions for $p \leq p_3$, $p_3 \leq -2$, are the stationary solution $u \equiv 0$ and a unique homoclinic orbit (homoclinic to zero).
 6. In case (g) the only bounded solution for $p \leq 0$ is the unique stationary solution.
 7. In case (h) the only bounded solution for $p \leq p_4$, $p_4 \leq -2$, is a unique periodic orbit.

Remark 1. Some of the bounds on p given above are sharp. For $p > -2$ (but not too large) in case (f) there exists a large collection of homoclinic orbits [4, 5]. In case (c), corresponding to the EFK equation with equal wells, there are many heteroclinic connections for $p > -2$ [10, 8, 9], as well as periodic solutions [12] and non-periodic bounded solutions [11]. In case (b) homoclinics have been found numerically (see [15, 16]) for values of p larger than -2 .

Remark 2. The fact that F is assumed to be an exact quartic is not essential to most of the proof; however, it simplifies many of the arguments by creating a phase plane of simple geometry. For instance, it is used a number of times that the regions C and C_{\pm} are strictly convex. As another example, the fact that $p_2 > -2$ is the result of an explicit calculation for this function; for a different nonlinearity the result may also be different. In Section 8 we discuss a generalization of these results to a different potential function.

Theorem 1 is proved in Section 3; the subsequent sections discuss the proof of Theorem 2.

2. EXAMPLE: HOMOCLINICS FOR A RESTIFFENING STRUT

In the context of structural mechanics, equation (1.1) arises as a model for a thin elastic strut on a nonlinear elastic foundation. Such struts are known experimentally to buckle in a localized fashion when loaded axially (in this model the parameter p represents the axial load). These localized profiles correspond to solutions of (1.1) that converge to zero at plus and minus infinity—homoclinics. An interesting class of solutions of (1.1) is found as minimizers of the elastic strain energy

$$\frac{1}{2} \int u'^2 + \int F(u)$$

under the condition of fixed total shortening of the strut,

$$\frac{1}{2} \int u'^2 = \lambda. \tag{2.1}$$

For a description of the derivation of this model, see [17].

The function F' models the stress-strain relationship of the foundation, $F'(u)$ being the stress associated with a strain u . The localization property, which corresponds to the existence of homoclinic solutions of this minimization problem, is related to the decrease of F'' as u increases from zero: the foundation loses stiffness as u increases. It is relatively easy to verify that with a nonlinearity F such that F'' increases as $|u|$ moves away from zero there are no homoclinic energy minimizers¹.

¹This can be verified by taking a sequence of functions of the form $u_n(t) = A_n \eta(t/n) \sin t$, where η is a fixed cut-off function (e.g. $\eta(x) = \exp(-x^2)$) and A_n is chosen to satisfy the constraint (2.1). For such a sequence the strain energy converges to 2λ , while by Fourier analysis it can be shown that

Recently interest has turned to foundations that have a destiffening character for small u , but restiffen as u becomes large. The function F with $b > 0$ (cases (f), (c), and (b)) is an example of such a foundation. The character of the homoclinic solutions is fundamentally different in the cases $0 \leq b < 2/9$ (case (f)) and $b > 2/9$ (case (b)). To illustrate this, consider the branch of primary solutions that bifurcates from $p = 2$ and continues for decreasing values of p (Figure 2). For $0 \leq b < 2/9$ this branch continues past $p = -2$ to $p = -\infty$, being the unique branch of solutions for $p \leq -2$ by Theorem 2, case (f). (The behaviour of the branch for $0 \leq p < 2$ is not proven rigorously, but there is ample numerical and asymptotic evidence to support the conjectures made above). As b passes $2/9$, and the right-hand well crosses the zero axis, the branch of solutions for $p \leq -2$ disappears through a widening of the profile of the solutions; numerical evidence suggests that the profile converges to two heteroclinic orbits, connected in a multi-bump fashion, and the techniques of this paper could be used to prove this conjecture.




Figure 2. As b passes $2/9$, the curve bifurcating from $p = 2$ changes character (figure taken from [16]).

At the same time, the branch with $-2 < p < 2$ disconnects at $p = -2$ and performs an oscillatory movement, as shown in Figure 2. The results of Theorem 2 for this situation can best be viewed in Figure 3. For a detailed discussion of this transition, as well as other examples of systems in which such phenomena occur, see [15, 16].

for every non-zero $u \in H^2(\mathbb{R})$,

$$\frac{1}{2} \int u''^2 + \frac{1}{2} \int u^2 > \int u'^2.$$

Since $F'' \geq 1$ and therefore $F(u) \geq u^2/2$, this implies that the sequence u_n converges to a strain energy level lower than that of every admissible fixed profile u .




Figure 3. Graphical interpretation of (part of) Theorem 2.

3. PRELIMINARY LEMMAS AND A PRIORI BOUNDS

The following simple lemma has been proved many times; see for instance [18] or [13].

Lemma 3. *Let u be a solution of (1.1). If u is monotone and bounded for $t \geq 0$, then the constant $u_\infty = \lim_{t \rightarrow \infty} u(t)$ is a solution of (1.1).*

An important tool in this paper is the following a priori bound, which appears in [13] and generalizes a result from [11]. Define I as the smallest closed interval such that the equation

$$F(u) = s$$

has exactly two solutions for all $s \in F(\mathbb{R} \setminus I)$ (see Figure 4).




Figure 4. The function F and the interval I .

Lemma 4. *Let $p \leq 0$ and let u be a non-constant solution of (1.1) satisfying $H(u) = 0$. If u is bounded on \mathbb{R} , then $u(\mathbb{R}) \subset I$.*

The proof of Theorem 1 is a direct consequence of Lemma 4, since for a one-well potential F we have $I = \{0\}$.

In the double-well case, there is a unique \bar{u} at which F has a local maximum. A direct consequence of Theorem 1 is

Corollary 5. *Let u be a non-constant solution and set $R = u(\mathbb{R})$. Then R contains a neighbourhood of \bar{u} .*

If this were not the case, one could change the function F outside of R and create a single-well potential; thus the existence of the solution u would contradict Theorem 1.

By this corollary, if u is a non-constant solution, then either

1. u is monotone; this implies the existence of two equilibria, and that u is a heteroclinic orbit connecting the two; or
2. u has local maxima above \bar{u} and/or local minima below \bar{u} . These we call ‘high’ maxima and ‘low’ minima, and will serve in the sections below.

Case (e) in Theorem 2 follows readily from this remark. There are no equilibria; therefore if u is a solution, then it has a high local maximum. Let this maximum be attained at $t = 0$; from (1.2) we have $F(u(0)) = u''^2/2 \geq 0$. By the form of F this implies that $u(0) \notin I$, a contradiction.

4. CASE (C)

Besides proving Theorem 2 for case (c), we give in this section a brief description of the monotonicity property enjoyed by solutions of (1.1). For the uniqueness and non-existence results that we prove, only a small subset of the available techniques is necessary; the reader can find examples of complete treatments within this framework in [2, 3, 4].

We draw the u, u'' -plane for case (c) (Figure 5). Although this is a two-dimensional projection of the four-dimensional state space, we will call it the ‘phase plane’ because of the similarity of the methods employed here to phase plane analysis for two-dimensional autonomous ODE’s. Note that in case (c), where $b = 2/9$, the function F and equation (1.1) are invariant under the reflection $u \mapsto 3 - u$.

The choice of the variables u and u'' stems from the Hamiltonian condition $H = 0$, which can be written as

$$u' \left(u''' + \frac{p}{2} u' \right) = V(u, u'') := \frac{1}{2} u''^2 - F(u). \quad (4.1)$$

If $V = 0$, then either $u' = 0$ or $u''' = -pu'/2$. Therefore, if an orbit intersects the set $\{V = 0\}$ at $t = t_0$, there are two possibilities: either

1. the velocity (u', u''') is non-zero, and the direction of the orbit (given by the ratio u'''/u') is either ∞ (vertical) or $-p/2$; or,
2. $u' = u''' = 0$, and u is symmetric about t_0 . This point is then called a turning point. In the phase plane the orbit returns along the same way as it arrived.

If an orbit is at a turning point on $\{V = 0\}$, then the direction of the orbit (both in positive and in negative time, since the two orbits coincide) is fixed by the equation. Figure 5(b) shows these directions as arrows. Note that at the points where the cone is tangent to the set $\{V = 0\}$ (called ‘magic points’, marked with γ and δ), the direction of the arrows changes side.





Figure 5. 5(a) The phase plane u, u'' , with the set $\{V = 0\}$; 5(b) An orbit with zero velocity leaves the set $\{V = 0\}$ in the direction indicated by the arrows (drawn here for $p < -2$).

The sign of V also implies a form of monotonicity. If $V < 0$, then by equation (4.1) the velocity (u', u''') lies in a double cone as indicated by the shaded area below.



As long as V remains strictly positive, the velocity can not change from one cone to the other. Consequently, if D is a bounded component of the set $\{V < 0\}$, then an orbit with $(u, u'') \in D$ at $t = t_0$ must converge to ∂D in both positive and negative time. Similar statements hold for the set $\{V > 0\}$.

To prove Theorem 2 for case (c), consider a non-constant solution u , and let u attain a high maximum at $t = 0$, so that $u(0) \geq \bar{u}$, $u'(0) = 0$, and $u''(0) \leq 0$. This implies that the orbit is at a point on the curve Γ in Figure 6. The shaded vertical line on the left of this figure is the left boundary of I ; a bounded solution can not pass this line. If the orbit has zero velocity on Γ , then it enters the set A (see the arrows in Figure 5(b)); if it has non-zero velocity, then it enters A either in positive or in negative time. In both cases the monotonicity cone property implies that the orbit remains in A and, since it is bounded, eventually leaves I . This is a contradiction.

Note that the hypothesis $p \leq -2$ is essential to this argument: as p crosses the value -2 , the ‘magic point’ marked δ in Figure 5(b) crosses the intersection at $u = 3$. Therefore orbits with zero velocity on the curve between δ and the intersection point enter the inner region $\{V < 0\}$ instead of A . Here the argument given above breaks down (and in fact Theorem 2 does not hold for this case [11, 12]).




Figure 6. The left boundary of A is given by the direction of the cone ($u'''/u' = -p/2$). The direction of this cone implies that all bounded orbits passing through Γ leave I .

The argument above proves that an orbit has no high maximum, and similarly by the symmetry $F(u) = F(3 - u)$ we conclude that it has no low minimum either. This implies that it is monotone, and therefore a heteroclinic orbit connecting the two equilibria, 0 and 3. For the uniqueness of this heteroclinic orbit we refer to [13].

This concludes the proof of case (c) of Theorem 2.

5. CASES (B) AND (F)

For cases (b) and (f) we shall prove Theorem 2 in several steps. The first consists of the following Lemma; other parts of the proof for these cases follow in Sections 6 and 7.

Lemma 6. *In cases (b) and (f) for $p \leq -2$, a bounded non-constant solution has no local minima.*

Proof. Lemma 6 uses the same idea as the previous section; both cases (b) and (f) are similar to case (c) in the left half of the phase plane (Figure 7). We will discuss the proof only for (b), since the argument is nearly identical for (f).




Figure 7. The phase planes in cases (b) and (f).

Suppose that a non-constant solution u attains a low local minimum at $t = 0$. Following the argument used in the previous section we conclude that the orbit intersects the line marked Γ in Figure 8, and enters the set A (if necessary after inverting time).

The boundedness of the orbit implies that it leaves the set I , which is a contradiction. \square




Figure 8.

6. CASES (A), (B), AND (D)

The common aspect of these three cases lies in the right half of the phase plane (see Figure 9).




Figure 9. The phase planes in cases (a), (b), and (d).

Lemma 7. *In cases (a), (b) and (d) there exists $q_1 \leq 0$, such that for $p \leq q_1$ a bounded non-constant solution has no high local maxima. In case (a) a bounded non-constant solution has no low local minima either.*

Proof. We discuss case (b) first, and subsequently return to the other two cases.

Consider the curve Γ in Figure 10. If this curve is considered as the graph of a function, $u'' = g(u)$, then the derivative g' is bounded from above by a constant (say) M . Setting $q_1 = -2M$, it follows that for $p < q_1$ the tangent to Γ nowhere has the direction $-p/2$, and therefore there are no magic points (see Section 4) on Γ . This implies that orbits starting with zero velocity on Γ enter the set A .

If $p = q_1$, then there is exactly one point on Γ where the cone and Γ are tangent. By a continuity argument it follows that orbits starting from that point also enter A .




Figure 10.

Repeating the argument from previous lemmas, we find that the existence of a high local maximum implies that the orbit enters A from Γ , in positive or negative time, which in turn leads to a contradiction. This proves the Lemma for case (b); the argument for case (d) is similar.

In case (a) this argument can be applied twice, both to a high maximum and to a low minimum. In general the ensuing values of q_1 will be different. \square

In these three cases we can now conclude the proof of Theorem 2. For case (a), we note that there are no equilibria, and that therefore any bounded solution has high maxima and local minima. This contradicts Lemma 7. For case (d), since there are no equilibria, there must be a high local maximum; by Lemma 7 this contradicts the boundedness of the solution.

For case (b), Lemmas 6 and 7 give the nonexistence of non-constant solutions, for values of p in the combined range $(-\infty, -2] \cup (-\infty, q_1)$. To complete the statement of Theorem 2 we need to show that $q_1 > -2$:

Lemma 8. *For all $1/4 > b > 2/9$ we have $q_1 > -2$.*

Proof. As in the proof of Lemma 7 we let M denote the maximum of g' on the curve Γ ; we need to prove that $M < 1$ for all $b > 2/9$. Since g is decreasing for $u > u_1$, we only need to concentrate on $\bar{u} < u < u_1$.

Let $u_1 = (1 + \sqrt{1 - 4b})/2b$ be the location of the second well. Note that $F''(u_1) = u_1 - 2 < 1$. We write F in the form

$$F(u) - F(u_1) = \frac{F''(u_1)}{2}(u - u_1)^2 + h(u),$$

where

$$h(u) = \frac{F'''(u_1)}{6}(u - u_1)^3 + \frac{F''''(u_1)}{24}(u - u_1)^4.$$

Note that $h(u) < 0$ and $h''(u) < 0$ for $\bar{u} < u < u_1$. We also define

$$\hat{F}(u) = F(u_1) + \frac{F''(u_1)}{2}(u - u_1)^2 = F(u) - h(u).$$

Let $g'(u)$ be maximal in $u = \tilde{u}$. There is a uniquely defined \hat{u} such that $\hat{F}'(\hat{u}) = F'(\tilde{u})$; since $h'' < 0$ it follows that $u_1 > \hat{u} > \tilde{u}$, and it is straightforward to show that $\hat{F}(\hat{u}) < F(\tilde{u})$. We then have

$$g'(\tilde{u}) = \frac{F'(\tilde{u})}{\sqrt{2F(\tilde{u})}} < \frac{\hat{F}'(\hat{u})}{\sqrt{2\hat{F}(\hat{u})}} = \frac{F''(u_1)(\hat{u} - u_1)}{\sqrt{2F(u_1) + F''(u_1)(\hat{u} - u_1)^2}} < \sqrt{F''(u_1)}.$$

Since $F''(u_1) < 1$ for $2/9 < b < 1$, the result is proved. \square

7. CASES (F) AND (H)

The uniting characteristic of these two cases is again a half of the phase plane (Figure 11). We define the closed convex set C as in the figures. The work of Hofer and




Figure 11. The phase planes in cases (f) and (h).

Toland [2], as well as many others, shows the existence of an even homoclinic orbit for $p \leq -2$ in case (f), and of a periodic orbit for $p \leq 0$ in case (h). Both do not leave the set C . Note that the set C is strictly convex for this function F .

In [5] it was proved for $b = 0$ that any homoclinic orbit with $p \leq -2$ is even. Here we generalize this result to $b > 0$, and to more general bounded orbits. The result in [5] is based on a monotonicity argument that requires the function $f(u) - u$ to be non-increasing. For $b = 0$ this is the case for all $u \geq 0$; for positive b , u is restricted to the set $W = [0, 2/3b]$.

Lemma 9. *Let $p \leq -2$ and let u be a bounded solution of (1.1) such that the range of u is a subset of W . Then u is even about any maximum or minimum.*

Proof. Let $\lambda \in (0, 1]$ satisfy $\lambda + \lambda^{-1} = -p$. Let u have a maximum at $t = 0$, and suppose, to force a contradiction, that $u'''(0) > 0$. Set $v(x) = u(-x)$; then we have

$$\begin{aligned} u(0) &= v(0), & u'(0) &= v'(0), \\ (u'' - \lambda u)(0) &= (v'' - \lambda v)(0), & (u''' - \lambda u')(0) &> (v''' - \lambda v')(0). \end{aligned}$$

There exists a $\sigma > 0$ such that $z = u - v > 0$ on $(0, \sigma)$. Set $w = u'' - v'' - \lambda(u - v)$. We then have

$$w(0) = 0, \quad w'(0) > 0,$$

and

$$w'' - \lambda^{-1}w = -(f(u) - u) + f(v) - v > 0 \quad \text{on } (0, \sigma),$$

so that $w > 0$ on $(0, \sigma)$. This implies that

$$z'' - \lambda z > 0 \quad \text{on } (0, \sigma), \tag{7.1}$$

so that $z(\sigma) > 0$; consequently $z > 0$ on $(0, \sigma]$, and by reasoning by contradiction we find that we can take $\sigma = \infty$. Going back to (7.1) we find that $z(t) \rightarrow \infty$ as $t \rightarrow \infty$, which contradicts the boundedness of u and v . \square

We shall show, starting with the lemma below, that in fact no bounded solution can leave the set C :

Lemma 10. *In cases (f) and (h) there exists $q_2 < 0$ such that for $p \leq q_2$ a bounded solution can not have a high local maximum outside C . In case (h) the same holds for a low local minimum.*

Proof. Consider the phase plane of case (f), in Figure 12. Let u have a high local maximum outside C , that therefore lies on the curve Γ_1 . There exists a $q_2 < 0$ such that any half-line starting on Γ_1 and extending in the direction $(-1, q_2/2)^T$ leaves I before intersecting Γ_2 . For $p \leq q_2$ the same is true for any orbit that enters A from Γ_1 . Using the same arguments as before, and noting that orbits with zero velocity that start on Γ_1 enter A , this implies a contradiction. \square




Figure 12.

Lemma 11. *In cases (f) and (h) for $p \leq -2$, if an orbit has a local maximum on ∂C , then the orbit is even about the maximum and lies within C for all time. In case (h) the same is true for a local minimum.*

Proof. In case (f) the orbit has no low local minima by Lemma 6, so that $u(t) \rightarrow 0$ as $t \rightarrow \pm\infty$ and u is monotone whenever $u \leq \bar{u}$. Since $C \subset [0, 3] \subset [0, 2/3b]$ we therefore have $u(\mathbb{R}) \subset W$ so that the orbit is even by Lemma 9.




Figure 13. The arrows in (a) indicate the direction of orbits that start with zero velocity on ∂C . The direction of orbits starting in α and β follows from a continuity argument.

We identify on ∂C three (open) sections Γ_1 , Γ_2 , and Γ_3 , and the two ‘magic points’ α and β (Figure 13(a)). We concentrate on the maximum of the orbit, which lies on $\Gamma_2 \cup \Gamma_3 \cup \{\beta\}$, and leaves ∂C with zero velocity since the orbit is even. Note that an orbit starting in α has a local minimum and therefore is not a candidate.

If the orbit starts on Γ_2 , then it remains inside the cone shown in Figure 13(b) and can only exit C through Γ_1 or Γ_3 . Figure 14 shows how both of these alternatives lead to a non-monotone orbit; exit via Γ_1 yields a local minimum on Γ_1 , and exit through Γ_3 implies that the orbit can not tend to zero without u taking negative values. Therefore this case leads to a contradiction.




Figure 14.

If the orbit starts on $\Gamma_3 \cup \{\beta\}$, then the non-monotonicity mentioned above again applies, leading to a contradiction.

A similar argument applies to case (h). □

We now conclude the proof of Theorem 2 for the remaining cases. For case (f), by Lemmas 10 and 11, for $p \leq \min\{q_2, -2\}$ a bounded orbit is even, remains inside C , and is homoclinic to zero. The uniqueness for orbits of this type is proven in [3].

In case (h) a bounded orbit also is even and remains inside C , and by slightly adapting the uniqueness proof of [3] these orbits can also be shown to be unique.

Case (g) is a special example of case (h), one in which the set C is reduced to one point. This point is the unique stationary point, and by the uniqueness for case (h) the unique bounded solution. This concludes the proof of Theorem 2. \square

8. GENERALIZATIONS

As an example of a possible generalization of Theorem 2 we consider here equation (1.1) with nonlinearity $F(u) = a + u^2/2 - u^4/4 + bu^6/6$ (Figure 15). This nonlinearity is interesting in the context of structural mechanics and structural geology (see Section 2 and [16]) because of the symmetry around zero. In this application the emphasis is on solutions that are homoclinic or heteroclinic, so we only consider $a = 0$. As before, the Hamiltonian is always assumed zero.




Figure 15. The nonlinearity $F(u) = a + u^2/2 - u^4/4 + bu^6/6$.

Fixing $a = 0$, the phase plane contains two convex regions C_{\pm} for $0 < b < 3/16$ that open up as b passes $3/16$ (Figure 16). This is analogous to the procession of cases (f)-(c)-(b) of Figure 1.




Figure 16. As b passes the value $3/16$ the convex regions C_{\pm} open up.

Applying the same ideas as above we find

Theorem 12. 1. When $0 < b < 3/16$, there are two solutions homoclinic to zero for any $p \leq -2$; when $p \leq q_1 \leq -2$, these are unique in the class of bounded solutions.

2. When $b = 3/16$, there are two heteroclinic solutions for $p \leq -4$, one connecting -2 and 0 and one connecting 0 and 2 (and their mirror images under $t \mapsto -t$). The only other bounded solutions are the three equilibrium solutions.
3. When $b > 3/16$ there is a $q_2 \leq -2$ such that there are no bounded solutions for $p \leq p_2$.

Proof. The simplest case is $b > 3/16$ (Figure 17). Here the argument is very similar to that in Section 6 for case (b): there is a number $q_2 \leq 0$, given by the geometry of the curve Γ , such that if $p \leq q_2$ then the monotonicity cone points inwards from any point on Γ . The rest of the argument is identical. The final statement is that there are no bounded orbits for $p \leq q_2$. Note that in contrast to the case considered above we have $q_2 \leq -2$ for all $b > 3/16$, since the direction of Γ at the origin is $(\pm 1, \pm 1)^T$. When $b = 3/16$ the phase plane is a variation on case (c) discussed above (Figure 18).




Figure 17.

Any non-monotone bounded solution has either a low local minimum (i.e. below \bar{u}) or a high local maximum (i.e. above $-\bar{u}$). Concentrating on the high local maximum, and using the convexity of the regions C_{\pm} , the monotonicity cone points into A from any point on Γ , provided $p \leq -4$ (the value of -4 is related to the direction of Γ at $(\pm 2, 0)$ which is $(\pm 1, \pm 2)$). Consequently the only bounded orbits are those that are monotone; these connect two equilibria, and by [2, 3] these exist for the connections $-2 \leftrightarrow 0$ and $0 \leftrightarrow 2$. These are unique within the class of orbits that remain inside C_- or C_+ . With similar arguments as above it follows that any bounded orbit must remain inside $C_- \cup C_+$. There is a theoretical possibility of a third connection, between -2 and 2 , provided the orbit does not leave the set $C_- \cup C_+$; this is excluded for $p < -2$ by the imposed directions of the orbit at the origin (since the orbit has to pass the origin with non-zero velocity, the direction u'''/u' is either infinity or $-p/2$; in both cases this is not possible for orbits remaining inside $C_- \cup C_+$).

For $0 < b < 3/16$ the results of [2, 3] give existence of two homoclinics for $p \leq -2$, again one in each of the regions C_{\pm} , which are unique among solutions that remain inside C_{\pm} . Lemma 10 and its proof hold unchanged for this situation, so that there is a $q_1 < 0$ such that for $p \leq q_1$ there are no bounded orbits with a maximum on Γ_1




Figure 18.

(see Figure 19). For orbits having a high local maximum on Γ_2 or Γ_3 the argument is similar to that of Lemma 11: such orbits are monotone by Lemma 9, and if an orbit leaves $C_- \cup C_+$ then this monotonicity is violated. \square




Figure 19.

Remark 3. When $b = 3/16$, the case of all wells having equal depth, the value $p = -4$ marks the passing of magic points through the two outer equilibria. Therefore, using the techniques of [4] one expects the appearance for $p = -4^+$ of a large number of solutions that are homoclinic to ± 3 . Therefore the condition $p \leq -4$ is sharp for the uniqueness result.

Acknowledgements. The author wishes to thank Boris Buffoni, Alan Champneys, and John Toland, for the many and fruitful discussions that led to this paper, as well as the Centre for Nonlinear Mechanics, Bath, for providing such a stimulating environment. In addition, gratitude is due to Jan Bouwe van den Berg for the mention of Lemma 4. This work was partially supported by the EPSRC, through grant GR/L17177 of the Applied Nonlinear Mathematics Initiative.

References

1. J. F. Toland. Solitary wave solutions for a model of the two-way propagation of water waves in a channel. *Math. Proc. Cambridge Philos. Soc.*, 90:343–360, 1981.
2. H. Hofer and J. F. Toland. On the existence of homoclinic, heteroclinic, and periodic orbits for a class of indefinite Hamiltonian systems. *Math. Annalen*, 268:387–403, 1984.
3. C. J. Amick and J. F. Toland. Homoclinic orbits in the dynamic phase-space analogy of an elastic strut. *European J. Appl. Math.*, 3:97–114, 1992.
4. A. R. Champneys and J. F. Toland. Bifurcation of a plethora of multi-modal homoclinic orbits for autonomous Hamiltonian systems. *Nonlinearity*, 6:665–772, 1993.
5. B. Buffoni, A. R. Champneys, and J. F. Toland. Bifurcation and coalescence of a plethora of homoclinic orbits for a Hamiltonian system. *J. Dyn. Stab. Sys.*, 8:221–281, 1996.
6. B. Buffoni. Periodic and homoclinic orbits for Lorentz-Lagrangian systems via variational methods. *Nonlinear Anal.*, 26(3):443–462, 1996.
7. B. Buffoni and E. Séré. A global condition for quasi-random behaviour in a class of conservative systems. *Comm. Pure Appl. Math.*, 49:285–305, 1996.
8. W. D. Kalies and R. C. A. M. van der Vorst. Multitransition homoclinic and heteroclinic solutions of the Extended Fisher-Kolmogorov equation. *J. Differential Equations*, 131:209–228, 1996.
9. W. D. Kalies, J. Kwapisz, and R. C. A. M. van der Vorst. Homotopy classes for stable connections between Hamiltonian saddle-focus equilibria. *Comm. Math. Phys.*, 193(2):337–371, 1998.

10. L. A. Peletier and W. C. Troy. Spatial patterns described by the Extended Fisher-Kolmogorov (EFK) equation: Kinks. *Differential Integral Equations*, 8(7):1437–1460, 1995.
11. L. A. Peletier and W. C. Troy. Chaotic spatial patterns described by the Extended Fisher-Kolmogorov equation. *J. Diff. Eq.*, 129:458–508, 1996.
12. L. A. Peletier and W. C. Troy. Spatial patterns described by the Extended Fisher-Kolmogorov (EFK) equation: Periodic solutions. *SIAM J. Math. Anal.*, 28:1317–1353, 1997.
13. G. J. B. van den Berg. Uniqueness of solutions for the Extended Fisher-Kolmogorov equation. Technical Report W 97-24, University of Leiden, 1997.
14. L. A. Peletier, A. I. Rotatiu-Bruma, and W. C. Troy. Pulse-like spatial patterns described by higher-order model equations. *J. Diff. Eq.*, 150:124–187, 1998.
15. P. D. Woods and A. R. Champneys. Heteroclinic tangles and homoclinic snaking in the unfolding of a degenerate reversible Hamiltonian Hopf bifurcation. *Physica D*. To appear.
16. G. W. Hunt, A. R. Champneys, P. D. Woods, M. A. Peletier, M. Ahmer Wadee, C. J. Budd, and G. L. Lord. Cellular buckling in long structures. To appear.
17. M. A. Peletier. Sequential buckling: A variational analysis. To appear.
18. J. Hadamard. Sur certaines propriétés des trajectoires en dynamique. *J. Math. Ser. 5*, (3):331–387, 1896.