



MORPHOLOGY, ULTRASTRUCTURE, AND MOLECULAR PHYLOGENY OF *WANGODINIUM SINENSE* GEN. ET SP. NOV. (GYMNODINIALES, DINOPHYCEAE) AND REVISITING OF *GYMNODINIUM DORSALISULCUM* AND *GYMNODINIUM IMPUDICUM*<sup>1</sup>

Zhaohe Luo

Third Institute of Oceanography, SOA, Xiamen 361005, China

Zhangxi Hu , Yingzhong Tang 

CAS Key Laboratory of Marine Ecology and Environmental Sciences, Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

Kenneth Neil Mertens 

Ifremer, LER BO, Station de Biologie Marine, Place de la Croix, BP40537, F-29185, Concarneau Cedex, France

Chui Pin Leaw, Po Teen Lim

Bachok Marine Research Station, Institute of Ocean and Earth Sciences, University of Malaya, 16310, Bachok, Kelantan, Malaysia

Sing Tung Teng

Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300, Kota Samarahan, Sarawak, Malaysia

Lei Wang, and Haifeng Gu<sup>2</sup> 

Third Institute of Oceanography, SOA, Xiamen 361005, China

The genus *Gymnodinium* includes many morphologically similar species, but molecular phylogenies show that it is polyphyletic. Eight strains of *Gymnodinium impudicum*, *Gymnodinium dorsalisulcum* and a novel *Gymnodinium*-like species from Chinese and Malaysian waters and the Mediterranean Sea were established. All of these strains were examined with light microscopy, scanning electron microscopy and transmission electron microscopy. SSU, LSU and internal transcribed spacers rDNA sequences were obtained. A new genus, *Wangodinium*, was erected to incorporate strains with a loop-shaped apical structure complex (ASC) comprising two rows of amphiesmal vesicles, here referred to as a new type of ASC. The chloroplasts of *Wangodinium sinense* are enveloped by two membranes. Pigment analysis shows that peridinin is the main accessory pigment in *W. sinense*. *Wangodinium* differs from other genera mainly in its unique ASC, and additionally differs from *Gymnodinium* in the absence of nuclear chambers, and from *Lepidodinium* in the absence of Chl *b* and nuclear chambers. New morphological information was provided for *G. dorsalisulcum* and *G. impudicum*, e.g., a short sulcal intrusion in *G. dorsalisulcum*; nuclear chambers in *G. impudicum* and *G. dorsalisulcum*; and a

chloroplast enveloped by two membranes in *G. impudicum*. Molecular phylogeny was inferred using maximum likelihood and Bayesian inference with independent SSU and LSU rDNA sequences. Our results support the classification of *Wangodinium* within the *Gymnodiniales* sensu stricto clade and it is close to *Lepidodinium*. Our results also support the close relationship among *G. dorsalisulcum*, *G. impudicum*, and *Barrufeta*. Further research is needed to assign these *Gymnodinium* species to *Barrufeta* or to erect new genera.

**Key index words:** apical structure complex; cyst; dinoflagellate; *Lepidodinium*; nuclear chambers; pigment; pyrenoid; ultrastructure

**Abbreviations:** ASC, apical structure complex; AV, amphiesmal vesicles; BI, Bayesian inference; BPP, Bayesian posterior probabilities; BS, bootstrap support; DAPI, 4',6-diamidino-2-phenylindole dihydrochloride; DMF, N,N-dimethylformamide; GTR, general time-reversible; MCMC, Markov chain Monte Carlo; ML, maximum likelihood; Mv-chl *a*, monovinyl chlorophyll *a*; NFC, nuclear fibrous connector; RAXML, Randomized Axelerated Maximum Likelihood; Tchl *a*, total chlorophyll *a*

<sup>1</sup>Received 16 January 2018. Accepted 14 August 2018. First Published Online 25 August 2018. Published Online 19 September 2018, Wiley Online Library (wileyonlinelibrary.com).

<sup>2</sup>Author for correspondence: e-mail guhaifeng@tio.org.cn. Editorial Responsibility: S. Lin (Associate Editor)

Many athecate dinoflagellates have been classified in the genus *Gymnodinium*, which originally encompassed gymnodinioid species with a cingulum