

TROPICAL AGRICULTURAL SCIENCE

Journal homepage: http://www.pertanika.upm.edu.my/

Biorisk Assessment of Antibiotic-Resistant Pathogenic Bacteria Isolated from Swiftlet Houses in Sarawak

Sui Sien Leong^{1*}, Lihan Samuel², Hwa Chuan Chia², Jui Heng Roland Kueh³ and Yee Min Kwan⁴

¹Department of Animal Science and Fishery, Faculty of Agricultural and Food Sciences, Universiti Putra Malaysia, Nyabau Road, 97008 Bintulu, Sarawak, Malaysia

²Department of Molecular Biology, Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia

³Department of Forestry Science, Faculty of Agricultural and Food Sciences, Universiti Putra Malaysia, Nyabau Road, 97008 Bintulu, Sarawak, Malaysia

⁴Department of Crop Science, Faculty of Agricultural and Food Sciences, Universiti Putra Malaysia, Nyabau Road, 97008 Bintulu, Sarawak, Malaysia

ABSTRACT

The occurrence of antibiotic resistance in pathogens is a growing concern globally. Development of multiple antibiotic-resistant bacteria has overwhelmed new medical advancement and threatens patients with untreatable infections. The qualitative risk assessment study was carried out to investigate the relative effects of the main determinants of antibiotic-resistant pathogenic bacteria and to estimate the risk of the emergence and spread of antibiotic resistance among humans in the swiftlet's faeces and its indoor air to human health. The methodology applied focused mainly on three main sections namely the hazard identification, exposure assessment, and hazard assessment. Sources of data for bio risk assessment include published literature, data from on-going research projects and data collected from the industry. The results showed that the prevalence of isolating

ARTICLE INFO

Article history: Received: 22 May 2018 Accepted: 30 August 2018 Published: 25 February 2019

E-mail addresses: leongsuisien@upm.edu.my (Sui Sien Leong) lsamuel@unimas.my (Lihan Samuel) chiahwachuan@gmail.com (Hwa Chuan Chia) roland@upm.edu.my (Jui Heng Roland Kueh) yeemink@upm.edu.my (Yee Min Kwan) * Corresponding author Gram-positive bacteria were higher in swiftlet houses. Over half of the pathogenic bacterial isolates were multidrug-resistant to a wide range of commonly used antibiotics such as *Bacillus*, *Enterococcus*, *Escherichia coli*, *Staphylococcus*, *Lysinibacillus*, *Paenibacillus and Sporosarcina*. 80% of the bacteria isolates showed high MAR index of over 0.2. These emerging pathogenic

© Universiti Putra Malaysia Press