

Running head: ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS**1 A longitudinal study on the relationship between aerobic endurance and lower body
2 strength in Italian sedentary Older Adults**

3

4 *Magistro Daniele PhD*^{1,2}, *Candela Filippo PhD*², *Paolo Riccardo Brustio*², *Liubicich Monica*
5 *Emma PhD*³, *Rabaglietti Emanuela PhD*²

6 ¹Institute of Development, Aging and Cancer, IDAC; Department of Functional Brain Imaging,
7 Tohoku University, Japan

8 ²Department of Psychology, University of Torino, Italy

9 ³University of Torino, Motor Science Research Center, University School of Motor and Sport
10 Science

11

12

13 Magistro Daniele (corresponding author)

14 Department of Functional Brain Imaging, IDAC, Tohoku University, Japan

15 4-1, Seiryomachi, Aobaku, Sendai, Miyagi, 980-8575, JAPAN

16 danielemagistro@gmail.com

17 daniele.magistro@idac.tohoku.ac.jp

18 Department of Psychology, University of Torino, Italy

19 Via Verdi 10, 10124, Torino, Italy

20 Phone: 0039 011 6702793 Fax: 0039 011 6702790

21

22 Candela Filippo

23 Department of Psychology, University of Torino, Italy

24 Via Verdi 10, 10124, Torino, Italy

25 Phone: 0039 011 6702793 Fax: 0039 011 6702790

26 candela@agx.it

27

28 Paolo Riccardo Brustio

29 Department of Psychology, University of Torino, Italy

30 Via Verdi 10, 10124, Torino, Italy

31 Phone: 0039 011 6702793 Fax: 0039 011 6702790

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 *paoloriccardo.brustio@unito.it*

2

3 **Liubicich Monica Emma**

4 University School of Motor and Sport Science (SUISM), Motor Science Research Center, University of
5 Torino, Italy

6 Piazza Bernini 12, 10143 Torino, Italy

7 Phone: 0039 011 7411727

8 *monica.liubicich@unito.it*

9

10 **Rabaglietti Emanuela**

11 Department of Psychology, University of Torino, Italy

12 Via Verdi 10, 10124, Torino, Italy

13 Phone: 0039 011 6702793 Fax: 0039 011 6702790

14 *emanuela.rabaglietti@unito.it*

15

16

17

18

19

20

21

22

23

24

25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

A Longitudinal Study on the Relationship between Aerobic Endurance and Lower Body Strength in Italian Sedentary Older Adults.

Abstract

Functional ageing processes are characterised by a loss of performance capabilities for most physiological systems, such as aerobic endurance and lower body strength, which are important for independent living and active ageing. The present study examines the direction of influence between aerobic endurance and lower body strength over time in Italian sedentary older adults. A three-wave longitudinal model was tested using cross-lagged analysis for 202 individuals aged over 65 years (mean = 73.92, SD = 5.84; 140 females). Analysis revealed that aerobic endurance and lower body strength declines over time. In addition, greater aerobic endurance positively affected lower body strength over time; however, the converse was true only during the first period (first 6 months). These findings emphasise the importance of these relationships for the design and implementation of effective physical intervention for the older adults.

Keywords: older adults, longitudinal study, cross lag

1
2

Introduction

3 Epidemiological studies show that the population of older adults is increasing in Western
4 societies (Leon, 2011). Ageing is a primary risk factor for the development and progression
5 of most chronic degenerative diseases, and an ageing population presents major health, social
6 and economic burdens (WHO, 2002). As populations become both older and more sedentary,
7 it becomes critical to identify protective factors for sustaining high levels of physical and
8 psychological functioning in the older adults. Regular physical activity can play an important
9 role in preventive gerontology by maintaining general health and physical efficiency (Garber
10 et al., 2011). Decrease in physical activity is strongly associated with the frequency of
11 chronic diseases in the 20th century (Beswick et al., 2008). In addition, regular physical
12 activity is essential for healthy ageing (Chodzko-Zajko et al., 2009), and many studies report
13 the positive effect of physical activity on the physical (Blinded for review) and cognitive
14 performance of older adults (Colcombe & Kramer, 2003; Blinded for review). One of the
15 biggest challenges faced by healthcare professionals and policy makers is encouraging older
16 people to adopt physically and socially active lifestyles.

17 In general, ageing reduces performance in most physiological systems, even in the
18 absence of discernible disease (Chodzko-Zajko et al., 2009). This functional decline affects a
19 broad range of tissues and organ systems; this cumulatively impacts the activities of daily
20 living and physical independence (Chodzko-Zajko et al., 2009). The decline in aerobic
21 capacity and skeletal muscle performance are two important aspects of physiological ageing
22 (Menz, Lord, & Fitzpatrick, 2003; Wallerstein et al., 2012). In fact, variations in these aspects
23 are important determinants in the older adults for both functional abilities (Binder et al.,
24 1999; Guralnik et al., 2000) and exercise tolerance (Toth, Gardner, Ades, & Poehlman,
25 1994). The ability of older people to function independently is largely dependent on the

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 maintenance of sufficient aerobic capacity and strength to perform daily activities (Chodzko-
2 Zajko et al., 2009).

3 Reduction in aerobic capacity is the principal cause in older people for decreases in
4 maximum oxygen uptake and in maximum organic performance capacity (Garber et al.,
5 2011). A decline in maximal aerobic exercise capacity of 10% per decade occurs during the
6 adult years, (Hawkins & Wiswell, 2003) and accelerates in later years. This age-associated
7 decline in aerobic capacity is accentuated by co-morbidities that are common in the older
8 adults, such as cardiac, pulmonary and peripheral artery diseases (Olshansky, Hayflick, &
9 Carnes, 2002). Recent longitudinal studies (Fleg et al., 2005; Hollenberg, Yang, Haight, &
10 Tager, 2006; Jackson, Sui, Hebert, Church, & Blair, 2009) have shown declines in the aerobic
11 capacity of 20%–25% per decade in healthy individuals over 70 years in age. Although
12 aerobic capacity decreases with decreased physical activity, similar rates of decline in aerobic
13 capacity are observed in sedentary and highly active older adults (Katzel, Sorkin, & Fleg,
14 2001). However, the maximum aerobic capacity of extremely active people is substantially
15 higher than that of their age sedentary peers (Fleg et al., 2005). When the energy requirement
16 for an activity exceeds a person's aerobic capacity, the individual can no longer perform that
17 activity. Thus, the ability to maintain a high aerobic capacity is a major determinant of an
18 older adult's functional independence (Fleg & Strait, 2012).

19 Ageing is accompanied by a progressive loss of muscle mass, which has been associated
20 with both lower muscle-force-production capacity and inability to perform the activities of
21 daily living (Wallerstein et al., 2012; Menz et al., 2003). Cross-sectional data (Hunter,
22 McCarthy, & Bamman, 2004) indicate that strength begins to decline slowly into the fifth
23 decade of life, and subsequently declines more rapidly, at 12%–15% per decade. This decline
24 appears to be greatest in the lower extremities (Aoyagi & Shepard, 1992; Judge, 2003) and is
25 associated with a reduction in the size of muscle fibres and muscle fibre loss (Cadore et al.,

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 2011) as well as in maximum voluntary neural activation of muscle (Hakkinen, Alen,
2 Kallinen, Newton, & Kraemer, 2000). This additionally results in a substantial impairment of
3 sensorimotor information exchange, resulting in a decreased quality of intermuscular and
4 intramuscular coordination. As a result, functional losses in strength and increasing gait
5 uncertainties are experienced. Correspondingly, risk increases for acute problems arising
6 from falls and injuries and chronic recurrent and degenerative illnesses (Faulkner, Larkin,
7 Claflin, & Brooks, 2007).

8 Although aerobic endurance and lower body strength decrease with ageing, it is not clear
9 how these aspects are related in older adults populations, i.e. whether aerobic endurance
10 affects lower body strength or vice versa. Several studies report a significant correlation
11 between aerobic endurance, gait velocity and mobility (i.e. Fiser et al., 2010; Rikli & Jones,
12 2013). Other studies determined relationships between lower body strength, gait velocity and
13 mobility (Hicks et al., 2012; Protas & Tissier, 2009). However, to the best of our knowledge,
14 no studies have investigated the relationship between aerobic endurance and lower body
15 strength in the older adults in a longitudinal prospective. Knowledge about how these two
16 variables influence each other over time can lead to the development of more specific and
17 effective intervention strategies.

18 To understand the aging process, longitudinal studies provide many advantages relative to
19 cross-sectional studies, as between-person age comparisons cannot provide a basis for
20 disentangling changes due to ageing from stable individual characteristics, average between
21 person trends, or decline effects, for example. In fact, longitudinal research could provide a
22 framework for advancing aging theory and studies (Baltes 1979). Some issues motivate the
23 emphasis for a longitudinal study to understand the ageing process. For example, physical
24 and cognitive change reflects development and involves external influences that interact with
25 both of these processes within individuals (Nilsson et al., 2004). Second, evidence for the

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 importance of the life span developmental perspective is provided in findings showing that
2 early patterns of development are predictive of later changes (Hayward & Gorman, 2004). In
3 fact, we expect that complex interactive effects are associated with multiple causal influences,
4 with cumulative effects of risk factors, and perhaps with age-specific causal action (Rutter,
5 1988; Seeman, Singer, Rowe, & McEwen, 2001). Also, inferences regarding change and
6 causal processes may be obscured if aging processes (time) is not taken into account (Vaupel
7 & Yashin, 1985).

8 The goal of this study was to examine the relationship and determine any reciprocal
9 influences between aerobic endurance and lower body strength in Italian sedentary older
10 adults over time. Sedentary living is defined as a way of living or a lifestyle that requires
11 minimal physical activity and that encourages inactivity through limited choices,
12 disincentives, and/or structural or financial barriers (Chodzko-Zajko et al., 2009). However, it
13 is well known that promoting the participation of sedentary elderly people in physical
14 trainings implemented in places such as traditional gyms with specific facilities can be
15 problematic, particularly in Italy (National Institute of Statistics, 2010). For this reason, it is
16 important to adapt the physical training to the specific characteristic of the interest population.

17 The relationship was investigated using a longitudinal methodology and analysed by
18 longitudinal cross-lagged modelling. Longitudinal modelling has three primary advantages
19 that make its use particularly desirable (Menard, 2002; Burkholder & Harlow, 2003): First, it
20 allows researchers to discover patterns of covariation among variables by observe their
21 behaviour over time; Second, researchers can test models that include data collected at
22 multiple time points and thus allow testing both directions of potential causality that are more
23 difficult to test in a cross-sectional design. Finally, longitudinal data analysis provides
24 estimates on relative construct stability by analysing the relation between subsequent
25 measurements of the same variable. There are some studies of how cross-lagged analysis was

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 used to determine temporal relations among measures for example about: perceptual
2 processing speed and intelligence (Deary, 1995,); physical and mental health (Hays,
3 Marshall, Wang, & Sherbourne, 1994); reading and mathematics skills development (Onatsu-
4 Arvilommi & Nurmi, 2000); emotional distress and cigarette smoking in adolescents
5 (Orlando, Ellickson, & Jinnett, 2001). However, there are no studies that could be found that
6 use the cross-lagged analysis to examine the relationship and determine any reciprocal
7 influences between aerobic endurance and lower body strength in a longitudinal prospective.

8 In a cross-lagged design, information for each variable assessed at each time point is
9 analysed. The figure 1 lists the four models that we analysed. The first model tested the
10 stability of aerobic endurance and lower body strength at different times, without cross-
11 lagged paths. The second model tested the effect of aerobic endurance on lower body strength
12 over time. The third model tested the effects of lower body strength on aerobic endurance.
13 The fourth model tested the reciprocal effects of aerobic endurance and lower body strength
14 over time with a cross-lagged structural equation model.

15 These models were designed to answer the following questions:

16 1) How do aerobic endurance and lower body strength change with time and does a
17 relationship exists between the two aspects?

18 2) What is the direction of influence between the two factors in Italian sedentary older
19 adults over time?

20 As already suggested in the literature (Weineck, 2007), we expected that aerobic
21 endurance is associated with lower body strength. Thus, we expected to find that sedentary
22 older adults with good aerobic endurance are more likely to have better lower body strength
23 than older adults with lower level of both variables and vice versa.

24 We expected that this study would facilitate the design of specific training that could be
25 used to address one or more physical abilities to improve the physical status of the older

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 adults. To design an effective trial and training, it is necessary to understand how physical
2 ability, as determined by aerobic endurance and lower body strength, of the older adults
3 develop in a longitudinal perspective. This is particularly relevant for sedentary older adults
4 who are not used to even moderate physical activity. Also, despite the relevance of this topic,
5 we found no studies that investigated the relation between aerobic endurance and lower body
6 strength in sedentary older adults over time.

7

8

Methods

9 A longitudinal research design was used in this study with three waves. All participants
10 were tested three times in 1 year (6 months between tests) at Baseline, Time1 (T1) and Time2
11 (T2). This research is a part of the larger longitudinal 'Act on Ageing' project conducted in
12 the Piedmont region, Italy.

13

Participants

14 Participant were 202 sedentary individuals over 65 years of age (mean = 73.92, SD =
15 5.84; 140 females) who live independently in the Piedmont region, Italy. Descriptive
16 characteristics of participants are presented in Table 1. Participants were classified as having
17 a sedentary lifestyle if they did not report participating in regular moderate or vigorous
18 physical activity for the last 5 years. The eligibility criteria for participation were age over 65
19 years, ability to walk without assistance and a Mini Mental State Examination (MMSE) score
20 higher than 26 (this score indicates the absence of cognitive impairment). Exclusion criteria
21 were myocardial infarction and/or coronary artery bypass grafting valvular surgery within 1
22 year, uncontrolled diabetes or hypertension, orthopaedic impairment or upper or lower
23 extremity fracture within past 6 months or participation in another study, even if the study
24 was of a different kind. These criteria were selected to exclude impairments that could
25

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 negatively affect the test and to guarantee the health and safety of participants. Lists of older
2 adults obtained from the Health Office of the Piedmont region and general practitioners were
3 used to contact potential participants by telephone. After recruitment, individuals were
4 required to obtain written approval from their primary-care physician to be eligible for the
5 study. All participants were retired at the time of the study and were advised to maintain their
6 normal daily routines. The Ethical Committee of the University of Torino approved the study,
7 and all participants were informed that participation in the study was voluntary and
8 confidential. All participants gave their written informed consent in accordance with Italian
9 law.

10

11 **Measures**

12 Demographic data (age, gender, marital status and level of education) were self-reported.
13 Each participant was assessed at the baseline time point for height, weight, and cognitive
14 function (Folstein, Folstein, & McHugh, 1975).

15 We extracted the aerobic endurance and lower body strength tests from the Senior Fitness
16 Test battery (SFT) (Rikli & Jones, 2013a). The SFT measures the underlying physical
17 parameters associated with functional ability, and reflect usual age-related changes in
18 physical performance. Also, the SFT is able to detect physical changes due to ageing process
19 or training. The SFT meets scientific standards for validity and reliability (Rikli and Jones,
20 1999; Rikli & Jones, 2013a, 2013b). Moreover, these tests are easy, safe and quick to
21 administer. As well, SFT test are feasible for use in common clinical, community and home
22 settings meeting the concept of ecological perspective of research and health promotion
23 (McLaren & Hawe, 2005).

24 Aerobic endurance and lower body strength tests were performed in the morning of the
25 same day, with an hour of rest between the tests.

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 Aerobic endurance, defined as the capacity to perform large-muscle activity over a
2 defined period of time (Weineck, 2007), was assessed with the 6-min walking test (Rikli &
3 Jones, 2013a), i.e. the maximum distance that a person can walk in 6 min. Test- retest
4 reliability of the measuring was .94 (Rikli & Jones, 2013b). A walkway was marked with
5 cones to indicate the area of the test. The participants were instructed to walk as far as
6 possible in 6 minutes, without running along a 60 metres walkway. The distance covered by
7 each participant was measured in metres. Scores can be obtained for individuals of all ability
8 levels—from borderline frail individuals, who can walk only a few feet in 6 min, to the
9 highly fit, who can cover several hundred metres in the same time.

10 Lower body strength was assessed using the 30-s chair–stand test (Jones, Rikli, & Beam,
11 1999; Rikli & Jones, 2013a). Test- retest reliability was .89 (Rikli & Jones, 2013b). In this
12 context, strength is the ability to generate force to overcome inertia or a load (Zatsiorsky,
13 2008). Lower body strength is an important aspect of fitness in older adults because of its role
14 in daily activity. This test is effective for detecting normal age-related decline in strength
15 (Bohannon, Shove, Barreca, Masters, & Sigouin, 2007) and the effect of physical training in
16 older adults (Carvalho, Marques, & Mota, 2004). The test involves counting the number of
17 times a person with their arms folded over their chest can come to a full stand from a seated
18 position within 30 s. The armless chair was placed against a wall for support and safety and
19 was adapted to improve or reduce the angle of 90° at the hips and knees (Rikli & Jones,
20 2013b).

21

22 Table 1

23

24 **Statistical Analyses**

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 We initially determined whether there was a statistical difference in the results obtained
2 from the three waves by repeated measures analyses of variance (ANOVA). Also, in order to
3 highlight the change over time, we computed the percentage of change by dividing the
4 average gain by the baseline mean of our measures (Thomas, Nelson, Silverman, &
5 Silverman, 2011).

6 To test the relations between aerobic endurance and lower body strength, we used a
7 cross-lagged analysis using MPlus version 6 with a maximum likelihood method, which is
8 recommended for small sample sizes (Olsson, Foss, Troye, & Howell, 2000). For missing
9 data, we used full information maximum likelihood approach, which performs better than
10 list-wise and pair-wise deletion methods (Schafer & Graham, 2002).

11 The models presented in Figure 1 were tested. Concerning the fit indices, we examined
12 the chi-square, comparative fit index (CFI) and standardised root-mean-square residual
13 (SRMR). A model is considered to fit well when the model chi-square is not significant. In
14 addition, a CFI value of 0.95 or higher suggests good model fit (Hu & Bentler, 1998). The
15 Root Mean Square Error of Approximation (RMSEA), and Standardized Root Mean Square
16 Residual (SRMR) values of 0.08 or lower are expected for an adequate model fit (Brown &
17 Cudeck, 1993; Hu & Bentler, 1998). Nested models were further compared using chi-square
18 difference test.

19

20

Results

Aerobic endurance

22 As seen in Table 2, the means of aerobic endurance for the three waves are as follows:
23 Baseline, mean = 416.7 meters, SD = 79.9; T1, mean = 392.7 meters, SD = 110.4; T2, mean
24 = 377 meters, SD = 112.24. Repeated measure ANOVA suggested that aerobic endurance
25 decrease significantly from Baseline to T1 and from T1 to T2 [$F(1, 161) = 22.88, p <$

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 0.0001]. Moreover, the results showed a relative decrease of 9.5% over time of aerobic
2 endurance.

3 **Lower body strength**

4 Lower body strength also changed over time (Table 2). The means of lower body strength
5 for the three waves are as follows: Baseline, mean = 10 times, SD = 3; T1, mean = 9 times,
6 SD = 2; T2, mean = 8 times, SD = 2. Lower body strength decreased significantly from
7 Baseline to T1 and from T1 to T2 [$F(1, 161) = 60.36, p < 0.0001$]. Also for lower body
8 strength the results showed a relative decrease over time of 20%.

9

10 **Cross-lagged analysis between aerobic endurance and lower body strength**

11 First, we tested the four models presented in Figure 1. As seen from Table 3, only the
12 reciprocal-effects model achieved a decent fit. The lower-body-strength-driven model had a
13 significantly better fit compared to the stability model, and the aerobic-endurance-driven
14 model revealed a marginally significantly better fit than the stability model. In addition, the
15 reciprocal effects model fitted the data significantly better than the lower-body-strength-
16 driven and aerobic-endurance-driven models. In conclusion, the findings suggest that the
17 reciprocal-effects model represent the relationship between lower body strength and aerobic
18 endurance over time relatively better than the alternative models. Figure 2 shows the
19 structural part of the model (for the measurement part see Table 3). Overall, the relationship
20 of both aerobic endurance and lower body strength were relatively stable from Baseline to T1
21 ($\beta = 0.54$, and $\beta = 0.70, p < 0.0001$, respectively) and T1 to T2 ($\beta = 0.94$, and $\beta = 0.62, p <$
22 0.001 , respectively). For cross-lagged paths, there were three statistically significant effects.
23 Aerobic endurance at Baseline positively predicted lower body strength at T1 ($\beta = 0.21, p <$
24 0.0001), suggesting that higher aerobic endurance at Baseline was associated with higher
25 levels of lower body strength 6 months later. Furthermore, aerobic endurance at T1

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 significantly predicted lower body strength at T2 ($\beta = 0.37, p < 0.0001$), suggesting that
2 higher aerobic endurance at T1 was associated with higher levels of lower body strength 6
3 months later. Regarding the effect of lower body strength, lower limb strength at Baseline
4 significantly predicted aerobic endurance at T1 ($\beta = 0.17, p < 0.0001$), suggesting that higher
5 lower body strength at Baseline was associated with higher aerobic endurance at T1, 6
6 months later. However, lower body strength at T1 did not significantly predict aerobic
7 endurance at T2 ($\beta = 0.04$). In sum, the effect of aerobic endurance on lower limb strength
8 was constant over time. In contrast, the effect of lower body strength was significant only for
9 the first period.

10

11

Discussion

12

13

14

15

16

17

18

The present study aimed to examine the relationship between aerobic endurance and lower body strength in sedentary older adults. We tested four alternative models aimed at assessing the association between aerobic endurance and lower body strength. We found that the best way to represent this relationship was to have bidirectional links between aerobic endurance and lower body strength. The reciprocal-effects model best represented the complex relationship between aerobic endurance and lower body strength. Aerobic endurance and lower body strength affect each other in sedentary older adults.

19

20

21

Understanding the reciprocal associations between aerobic endurance and lower body strength is important for designing physical training specifically for the older adults, as well as for designing studies investigating physical intervention for the older adults.

22

23

24

25

Our longitudinal data show that, aerobic endurance and lower body strength are affected by time. Our findings are supported by studies from Chodzko-Zajko and colleagues (2009) and Hollman (2007), which indicated these aspects of physical functioning decrease over time. Furthermore, our results in accordance to those reported in previous longitudinal studies

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 (Hollenberg, Yang, Haight, & Tager, 2006; Hughes et al., 2001), confirm that there is a
2 decline of aerobic endurance and lower body strength across the age span in healthy persons,
3 and that this decline is related to life style habits. It is clear that a sedentary lifestyle,
4 especially for the older adults, can be highly debilitating, resulting in the loss of self-
5 sufficiency and health. In contrast, physical activity can be a protective factor for good
6 physical and psychological conditions, even for individuals with frail physical or
7 psychosocial states (Daley & Spinks, 2000). In general, physical functioning decreases with
8 age (Shea, Park, & Braden, 2006; Voelck-er-Rehage & Alberts, 2005), and specific
9 parameters, such as strength, endurance and balance, are predictors of dependence in the
10 activities of daily living (Guralnik et al., 2000). The capacity of older adults to maintain
11 muscle strength and power is an essential factor in maintaining independence (Rantanen,
12 2003). A threshold level of muscle strength is required to perform the basic activities of daily
13 living and to participate in activities designed to maintain aerobic endurance.

14 Second, our model showed the relationship between aerobic endurance and lower body
15 strength over time in sedentary older adults (Table 2) is very important because it shows how
16 these fundamental elements of physical functioning influenced each other. As can be seen
17 from the model, the level of aerobic endurance significantly affected lower body strength
18 over time. In addition, lower body strength initially affected aerobic endurance significantly.
19 In fact, we found that older adults who had high level of aerobic endurance also had a high
20 level of lower limb strength, and this relationship was steady over time. Furthermore, the
21 older adults who had a good level of aerobic endurance at the Baseline were more likely to
22 maintain a good level of lower limb strength after 6 and 12 months. Aerobic–endurance
23 activity is an effective form of exercise for increasing muscle strength and power in older
24 individuals (Caserotti, Aagaard, & Puggaard, 2008; Latham, Bennett, Stretton, & Anderson,
25 2004; Macaluso & De Vito, 2004; Sayers, 2007). Specificity of training suggests that

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 endurance training has little effect on the development of muscle strength or power in healthy
2 individuals (Nader, 2006). Moreover, the decreased physical functioning of the older adults
3 allows greater initial gains in muscle function as a result of aerobic–endurance training
4 because of a large adaptation potential (Fleck & Kraemer, 2004). Aerobic endurance training,
5 e.g. cycle-ergometry or walking, is sufficient to increase lower body strength and power in
6 older individuals as reported by a study of Lovell and colleagues (2010).

7 The relation between aerobic endurance and lower body strength has important practical
8 consequences, because during the design of physical activity training for older adults is
9 crucial to know how the physical variables influence each other. To design an effective
10 training for the older adults, it is first necessary to know how aerobic endurance and lower
11 body strength of the older adults develop over time. Our study suggests that combined
12 training focusing on both aspects might be most effective than either form of training alone in
13 counteracting the detrimental effects of a sedentary lifestyle on the health and physical
14 functioning. However, training that focuses only on one of these aspects, aerobic endurance
15 or lower limb strength, could indirectly result in improvement of the other

16 The small sample size of our study makes it difficult to generalise our results to larger
17 contexts. For example, our sample size did not allow us to determine the relationship for
18 subgroups based on age or gender. Furthermore, the present study examines only two time
19 periods. Thus, a longer trial with more time points and a larger sample size might be more
20 informative. Finally, it would be interesting to compare this relationship in groups of
21 sedentary as well as physically active older adults.

22 Despite these limitations, our study highlights that the relation between aerobic endurance
23 and lower body strength might not be represented properly without considering the reciprocal
24 effects. Moreover, it extends the investigation of this relationship for sedentary older adults.
25 These findings might be able to clarify the integrative mutual action over time of aerobic

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 endurance and lower limb strength in older adults. Indeed, setting realistic goals and
- 2 designing activity protocols that take scientific evidence into consideration might lead to
- 3 constructing and implementing interventions that are useful for improving and promoting
- 4 active ageing for the Italian sedentary older adults.
- 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

References

- Aoyagi, Y., & Shephard, R. J. (1992). Aging and muscle function. *Sports medicine*, *14*(6), 376–396.
- Baltes, P. B., & Nesselroade, J. R. (1979). History and rationale of longitudinal research. *Longitudinal research in the study of behavior and development*, *1*, 59.
- Beswick, A. D., Rees, K., Dieppe, P., Ayis, S., Gooberman-Hill, R., Horwood, J., et al. (2008). Complex interventions to improve physical function and maintain independent living in elderly people: A systematic review and meta-analysis. *Lancet*, *371*, 725–735. doi:10.1016/S0140-6736(08)60342-6
- Binder, E.F., Birge, S.J., Spina, R., Ehsani, A. A., Brown, M., Sinacore, D. R., & Kohrt, W. M. (1999). Peak aerobic power is an important component of physical performance in older women. *Journal of Gerontology Series A: Biological Sciences and Medical Sciences*, *54*(7), M353–M356.
- Bohannon, R.W., Shove, M. E., Barreca, S. R., Masters, L. M., & Sigouin, C. S. (2007). Five-repetition sit-to-stand test performance by community-dwelling adults: A preliminary investigation of times, determinants, and relationship with self-reported physical performance. *Isokinetics and Exercise Science*, *15*, 77–81.
- Brown, M. W., & Cudeck, R. (1993). Alternative ways of assessing model fit. In K. Bollen & S. Long (Eds.). *Testing structural equation models*. Newbury Park, NJ: Sage.
- Burkholder, G. J., & Harlow, L. L. (2003). An illustration of a longitudinal cross-lagged design for larger structural equation models. *Structural Equation Modeling*, *10*(3), 465-486.
- Cadore, E. L., Pinto, R. S., Alberton, C. L., Pinto, S. S., Lhullier, F. L. R., Tartaruga, M. P., ... Krueger, L. F. M. (2011). Neuromuscular economy, strength, and endurance in

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 healthy elderly men. *Journal of Strength & Conditioning Research*, 25(4), 997–1003.
2 doi:10.1519/JSC.0b013e3181d650ba
- 3 Blinded for review
- 4 Carvalho, M. J., Marques, E., & Mota, J. (2004). Effects of multi component training on
5 functional fitness in older adults. *Journal of Aging and Physical Activity*, 55, 41–48.
- 6 Caserotti, P., Aagaard, P., & Puggaard, L. (2008). Changes in power and force generation
7 during coupled eccentric-concentric versus concentric muscle contraction with
8 training and aging. *European Journal of Applied Physiology*, 103, 151–161.
- 9 Chodzko-Zajko, W. J., Proctor, D. N., Fiatarone Singh, M. A., Minson, C. T., Nigg, C. R.,
10 Salem, G. J., & Skinner, J. S. (2009). American College of Sports Medicine position
11 stand. Exercise and physical activity for older adults. *Medicine & Science in Sports &
12 Exercise*, 41(7), 1510–1530.
- 13 Colcombe, S., & Kramer A. F. (2003). Fitness effects on the cognitive function of older
14 adults: a meta-analytic study. *Psychological Science*, 14, 125–130. doi:10.1111/1467-
15 9280.t01-1-01430.
- 16 Daley, M. J., & Spinks, W. L. (2000). Exercise, mobility and aging. *Sports Medicine*, 29, 1-
17 12. doi:10.2165/00007256-200029010-00001
- 18 Deary, I. J. (1995). Auditory inspection time and intelligence: What is the direction of
19 causation? *Developmental Psychology*, 31(2), 237.
- 20 Fleck, S. J., & Kraemer, W. J. (2004). *Designing resistance training programs* (3rd ed.).
21 Champaign, IL: Human Kinetics.
- 22 Fleg, J. L., Morrell, C. H., Bos, A.G., Brant, L. J., Talbot, L. A., Wright, J. G., & Lakatta, E.
23 G. (2005). Accelerated longitudinal decline of aerobic capacity in healthy older
24 adults. *Circulation* 112, 674–682.

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 Fleg, J. L., & Strait, J. (2012). Age-associated changes in cardiovascular structure and
2 function: a fertile milieu for future disease. *Heart failure reviews*, *17*(4-5), 545–554.
- 3 Fiser, W. M., Hays, N. P., Rogers, S. C., Kajkenova, O., Williams, A. E., Evans, C. M., &
4 Evans, W. J. (2010). Energetics of walking in elderly people: factors related to gait
5 speed. *The Journals of Gerontology Series A: Biological Sciences and Medical*
6 *Sciences*, *65*(12), 1332–1337.
- 7 Folstein, M., Folstein, S., & McHugh, P. R. (1975). Mini-Mental State: A practical method
8 for grading the cognitive state of patients for the clinician. *Journal of Psychiatric*
9 *Research*, *12*, 89–198. doi:10.1016/0022-3956(75)90026-6.
- 10 Faulkner, J. A., Larkin, L. M., Claflin, D. R., & Brooks, S. V. (2007). Age-related changes in the
11 structure and function of skeletal muscles. *Clinical and Experimental Pharmacology and*
12 *Physiology*, *34*(11), 1091–1096.
- 13 Garber, C. E., Blissmer, B., Deschenes, M. R., Franklin, B. A., Lamonte, M. J., Lee, I. M., ...
14 Swain, D. P. (2011). American College of Sports Medicine position stand. Quantity
15 and quality of exercise for developing and maintaining cardiorespiratory,
16 musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for
17 prescribing exercise. *Medicine & Science in Sports & Exercise*, *43*(7), 1334–1359.
- 18 Guralnik, J.M., Ferrucci, L., Pieper, C. F., Leveille, S. G., Markides, K. S. & Ostir, G. V.
19 (2000). Lower extremity function and subsequent disability: Consistency across
20 studies, predictive models, and value of gait speed alone compared with the Short
21 Physical Performance Battery. *Journal of Gerontology Series A: Biological Sciences*
22 *Medical Sciences*, *55*(4), M221–M231. doi:10.1093/gerona/55.4.M221.
- 23 Häkkinen, K., Alen, M., Kallinen, M., Newton, R. U., & Kraemer, W. J. (2000).
24 Neuromuscular adaptation during prolonged strength training, detraining and re-
25 strength-training in middle-aged and elderly people. *European Journal of Applied*

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 *Physiology*, 83(1), 51–62. Hawkins, S. A., & Wiswell, R. A. (2003). Rate and
2 mechanism of maximal oxygen consumption decline with aging. *Sports Medicine*,
3 33(12), 877–888.
- 4 Hays, R. D., Marshall, G. N., Wang, E. Y. I., & Sherbourne, C. D. (1994). Four-year cross-
5 lagged associations between physical and mental health in the Medical Outcomes
6 Study. *Journal of consulting and clinical psychology*, 62(3), 441.
- 7 Hayward, M. D., & Gorman, B. K. (2004). The long arm of childhood: The influence of
8 early-life social conditions on men’s mortality. *Demography*, 41(1), 87-107.
- 9 Hicks, G. E., Shardell, M., Alley, D. E., Miller, R. R., Bandinelli, S., Guralnik, J., Lauretani,
10 F., Simonsick, E. M., & Ferrucci, L. (2012). Absolute strength and loss of strength as
11 predictors of mobility decline in older adults: the InCHIANTI study. *The Journals of*
12 *Gerontology Series A: Biological Sciences and Medical Sciences*, 67(1), 66–73.
- 13 Hollenberg, M., Yang, J., Haight, T. J., & Tager, I. B. (2006). Longitudinal changes in
14 aerobic capacity: implications for concepts of aging. *Journal of Gerontology Series A:*
15 *Biological Sciences and Medical Sciences*, 61, 851–858.
- 16 Hu, L., & Bentler, P. M. (1998). Fit indices in covariance structure modeling: Sensitivity to
17 underparameterized model misspecification. *Psychological Methods*, 3(4), 424–453.
- 18 Hughes, V. A., Frontera, W. R., Wood, M., Evans, W. J., Dallal, G. E., Roubenoff, R., &
19 Singh, M. A. F. (2001). Longitudinal muscle strength changes in older adults
20 influence of muscle mass, physical activity, and health. *The Journals of Gerontology*
21 *Series A: Biological Sciences and Medical Sciences*, 56(5), B209-B217.
- 22 Hunter, G. R., McCarthy, J. P., & Bamman, M. M. (2004). Effects of resistance training on
23 older adults. *Sports Medicine*, 34(5), 329–348.

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 Jackson, A. S., Sui, X., Hebert, J. R., Church, T. S., & Blair S. N. (2009). Role of lifestyle
2 and ageing on the longitudinal change in cardiorespiratory fitness. *Archives of*
3 *Internal Medicine, 169*, 1781–1787.
- 4 Jones, C. J., Rikli, R. E. & Beam, W. C. (1999). A 30-s chair-stand test as a measure of lower
5 body strength in community-residing older adults. *Research Quarterly for Exercise*
6 *and Sport, 70*(2), 113–119.
- 7 Judge, J. O. (2003). Balance training to maintain mobility and prevent disability. *American*
8 *Journal of Preventive Medicine, 25*(3), 150–156.
- 9 Katzel, L. I., Sorkin, J. D., & Fleg, J. L. (2001). A comparison of longitudinal changes in
10 aerobic fitness in older endurance athletes and sedentary men. *Journal of the*
11 *American Geriatrics Society, 49*(12), 1657–1664.
- 12 Latham, N. K., Bennett, D. A., Stretton, C. M., & Anderson, C. S. (2004). Systematic review
13 of progressive resistance strength training in older adults. *Journal of Gerontology*
14 *Series A: Biological Sciences and Medical Sciences, 59*, 48–61.
- 15 Leon, D. A. (2011). Trends in European life expectancy: a salutary view. *International*
16 *Journal of Epidemiology, 40*(2), 271–277.
- 17 Lovell, D. I., Cuneo, R., & Gass, G. C. (2010). Can aerobic training improve muscle strength
18 and power in older men? *Journal of Aging and Physical Activity, 18*, 14–26.
- 19 Macaluso, A., & De Vito, G. (2004). Muscle strength, power and adaptations to resistance
20 training in older people. *European Journal of Applied Physiology, 91*(4), 450–472.
- 21 Blinded for review
- 22 McLaren, L., & Hawe, P. (2005). Ecological perspectives in health research. *Journal of*
23 *Epidemiology and Community Health, 59*(1), 6-14.
- 24 Menard, S. (2002). *Longitudinal research* (Vol. 76): Sage.

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 Menz, H.B., Lord, S.R., & Fitzpatrick, R.C. (2003). Age - related differences in walking
2 stability. *Age Ageing*, 32(2), 137–142 doi:10.1093/ageing/32.2.137
- 3 Nader, G.A. (2006). Concurrent strength and endurance training: from molecules to man.
4 *Medicine & Science in Sports & Exercise*, 38, 1965–1970.
- 5 National Institute of Statistics. (2010). Annuario statistico italiano 2010 [Statistical Italian
6 yearbook - 2010] [Data file]. Roma: ISTAT. Retrieved from <http://www.istat.it>
- 7 Nilsson, L.-G., Adolfsson, R., Bäckman, L., de Frias, C. M., Molander, B., & Nyberg, L.
8 (2004). Betula: A prospective cohort study on memory, health and aging. *Aging*
9 *Neuropsychology and Cognition*, 11(2-3), 134-148.
- 10 Olshansky, S. J., Hayflick, L., & Carnes, B. A. (2002). Position statement on human aging.
11 *Journal of Gerontology Series A Biological Sciences Medical Sciences*, 57(8), B292–
12 B297. doi:10.1093/gerona/57.8.B292.
- 13 Olsson, U. H., Foss, T., Troye, S. V., & Howell, R. D. (2000). The performance of ML, GLS,
14 and WLS estimation in structural equation modeling under conditions of
15 misspecification and nonnormality. *Structural Equation Modeling*, 7, 557–595.
- 16 Onatsu-Arviolommi, T., & Nurmi, J.-E. (2000). The role of task-avoidant and task-focused
17 behaviors in the development of reading and mathematical skills during the first
18 school year: A cross-lagged longitudinal study. *Journal of educational psychology*,
19 92(3), 478.
- 20 Orlando, M., Ellickson, P. L., & Jinnett, K. (2001). The temporal relationship between
21 emotional distress and cigarette smoking during adolescence and young
22 adulthood. *Journal of consulting and clinical psychology*, 69(6), 959.
- 23 Protas, E. J., & Tissier, S. (2009). Strength and speed training for elders with mobility
24 disability. *Journal of aging and physical activity*, 17(3), 257.

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 Rantanen, T. (2003). Muscle strength, disability and mortality. *Scandinavian Journal of*
2 *Medicine & Science in Sports, 13*, 3–8.
- 3 Rikli, R., & Jones, C. J. (2013a). *Senior Fitness Test Manual*. Champaign, IL: Human
4 Kinetics.
- 5 Rikli, R. E., & Jones, C. J. (2013b). Development and validation of criterion-referenced
6 clinically relevant fitness standards for maintaining physical independence in later
7 years. *The Gerontologist, 53*(2), 255–267.
- 8 Rutter, M. E. (1988). *Studies of psychosocial risk: The power of longitudinal data:*
9 Cambridge University Press.
- 10 Sayers, S.P. (2007). High-speed power training: a novel approach to resistance training in
11 older men and women. A brief review and pilot study. *Journal of Strength and*
12 *Conditioning Research, 21*, 518–526.
- 13 Schafer, J. L., & Graham, J. W. (2002). Missing data: Our view of the state of the art.
14 *Psychological methods, 7*(2), 147.
- 15 Seeman, T. E., McEwen, B. S., Rowe, J. W., & Singer, B. H. (2001). Allostatic load as a
16 marker of cumulative biological risk: MacArthur studies of successful
17 aging. *Proceedings of the National Academy of Sciences, 98*(8), 4770-4775.
- 18 Shea, C. H., Park, J. H., & Braden, H. W. (2006). Age-related effects in sequential motor
19 learning. *Physical Therapy Journal, 86*, 478–488.
- 20 Thomas, J. R., Nelson, J. K., & Silverman, S. J. (2011). *Research methods in physical*
21 *activity*: Human Kinetics.
- 22 Toth, M. J., Gardner, A. W., Ades, P. A., & Poehlman, E. T. (1994). Contribution of body
23 composition and physical activity to age-related decline in peak VO₂ in men and
24 women. *Journal of Applied Physiology, 77*, 647–652.

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

- 1 Wallerstein, L. F., Tricoli, V., Barroso, R., Rodacki, A. L. F., Russo, L., Aihara, A. Y., ...
2 Ugrinowitsch, C. (2012). Effects of strength and power training on neuromuscular
3 variables in older adults. *Journal of Aging and Physical Activity*, 20(2), 171–185.
- 4 Weineck, J. (2007). *Optimal Training*. Balingen: SpittaVerlag GmbH & Co. KG.
- 5 World Health Organization (WHO). (2002). *Active ageing: A policy framework*. Geneva,
6 Switzerland: Author.
- 7 Vaupel, J. W., & Yashin, A. I. (1985). Heterogeneity's ruses: some surprising effects of
8 selection on population dynamics. *The American Statistician*, 39(3), 176-185.
- 9 Voelcker-Rehage, C., & Alberts, J.L. (2005). Age-related changes in grasping force
10 modulation. *Experimental Brain Research*, 166(1), 61–70. doi: 10.1007/s00221-005-
11 2342-6
- 12 Zatsiorsky, V.M. (2008). Biomechanics of Strength and Strength Training. In P. V. Komi
13 (Ed.), *Strength and Power in Sport*, Second Edition Oxford, UK: Blackwell Science
14 Ltd.
- 15

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1
2

3 *Table 1: Socio-demographic characteristics of study participants*

		<i>N</i>	<i>%</i>
Gender (<i>N</i>)	Women	140	69
	Men	62	31
Level of education (<i>N</i>)	Low	134	66
	High	68	34
Family condition	Never married	19	9
	Married	102	51
	Widow/Widower	56	28
	Divorced	25	12
Past job	Manual labour	156	77
	Non-manual labour	46	23
Age		M = 73.92	SD = 5.84
BMI score		M = 29.6	SD = 5.3
MMSE (score 0–30)		M = 28.8	SD = 1.5

4 M = Mean; SD = standard deviation

5

6

7

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1 *Table 2: Intercorrelations among, and means (M) and standard deviations (SD) of aerobic*
 2 *endurance and lower limb strength variables*

	1	2	3	4	5	6
1. Aerobic Endurance T1	--					
2. Lower body strength T1	.45***	--				
3. Aerobic Endurance T2	.65***	.48***	--			
4. Lower body strength T2	.50***	.76***	.60***	--		
5. Aerobic Endurance T3	.65***	.50***	.97***	.63***	--	
6. Lower body strength T3	.53***	.69***	.76***	.84***	.79***	--
M	416.7	10	392.7	9	377	8
SD	79.9	3	110.4	2	112.2	2

* $p < .05$; ** $p < .01$; *** $p < .001$

6 *Table 3: Model fit for competitive models*

Model	χ^2	df	p	CFI	RMSEA	SRMR	Model Comparison	$\Delta\chi^2$	df	p
A) Stability model	98.343	8	.0001	.908	.236	.151	-			
B) Aerobic endurance driven	18.343	6	.0054	.987	.101	.067	A	80	2	.0001
C) Lower body strength driven	74.398	6	.0001	.931	.238	.099	A	23.945	2	.0001
D) Reciprocal effects model	5.788	4	.2156	.998	.047	.009	B-C	12.555/68.61	2	0.002/0001


ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1


2

3 *Figure 1: Alternative models for the relationship between aerobic endurance and lower body*
4 *strength*

5 a) *Stability model*



8 b) *Aerobic endurance driven model*




ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS

1

2

c) *Lower body strength driven model*

3




4

5

6

d) *Reciprocal effects model*

7



8

9


10

11 *Figure 2: Structural model of the reciprocal effects between aerobic endurance and lower*

12 *body strength*

13

ENDURANCE AND LOWER BODY STRENGTH IN OLDER ADULTS



1

2

3

* $p < .05$; ** $p < .01$; *** $p < .001$

4

5