

IRIS A<sub>per</sub>TO



UNIVERSITÀ  
DEGLI STUDI  
DI TORINO

**This is the author's final version of the contribution published as:**

D. Nicodemo, A. Pauciullo, A. Castello, E. Roldan, M. Gomendio, G. Cosenza, V. Peretti, A. Perucatti, G.P. Di Meo, L. Ramunno, L. Iannuzzi, J. Rubes, D. Di Berardino. X-Y Sperm Aneuploidy in 2 Cattle (*Bos taurus*) Breeds as Determined by Dual Color Fluorescent in situ Hybridization (FISH). *Cytogenet Genome Res* 2009;126:217–225.

DOI: 10.1159/000245923

**The publisher's version is available at:**

<https://www.karger.com/Article/Pdf/245923>

**When citing, please refer to the published version.**

**Link to this full text:**

<https://www.karger.com/Article/Pdf/245923>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

---

iris-AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

1 **X-Y Sperm Aneuploidy in 2 Cattle (*Bos taurus*) Breeds as Determined by Dual Color**  
2 **Fluorescent in situ Hybridization (FISH)**

3

4 D. Nicodemo <sup>a</sup>, A. Pauciullo <sup>a</sup>, A. Castello <sup>b</sup>, E. Roldan <sup>b</sup>, M. Gomendio <sup>b</sup>, G. Cosenza <sup>a</sup>, V. Peretti <sup>c</sup>,  
5 A. Perucatti <sup>d</sup>, G.P. Di Meo <sup>d</sup>, L. Ramunno <sup>a</sup>, L. Iannuzzi <sup>d</sup>, J. Rubes <sup>e</sup>, D. Di Berardino <sup>a</sup>

6

7

8 <sup>a</sup> Department of Soil, Plant, Environment and Animal Production Sciences, University of Naples  
9 ‘Federico II’, Portici , Italy;

10 <sup>b</sup> Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Cientificas, Madrid ,  
11 Spain;

12 <sup>c</sup> Department of Animal Science and Food Inspection, University of Naples ‘Federico II’

13 <sup>d</sup> Research National Council (CNR), ISPAAM, Laboratory of Animal Cytogenetics and Gene  
14 Mapping, Naples , Italy;

15 <sup>e</sup> Veterinary Research Institute, Brno , Czech Republic

16

17

## 1 **Abstract**

2 The present study was undertaken to investigate aneuploidy rates in the sperm populations of 2 cattle  
3 (*Bos taurus*) breeds by using dual color fluorescent in situ hybridization (FISH) with Xcen and Y  
4 chromosome-specific painting probes, obtained by chromosome microdissection and DOPPCR.  
5 Frozen semen from 10 Italian Friesian and 10 Italian Brown testing bulls was used for the  
6 investigation. For each bull, more than 5,000 sperm were analyzed, for a total of 52,586 and 51,342  
7 sperm cells for the 2 breeds, respectively. The present study revealed – in both breeds – a  
8 preponderance of the Y-bearing sperm compared to the X-bearing sperm. Within each breed, a  
9 statistically significant variation in the various classes of aneuploidy (XX, YY and XY) was found:  
10 differences were found in the Friesian breed among the 3 diploidy classes, and in the Brown breed,  
11 among the 3 disomy classes ( $p \leq 0.05$ ) as well as among the 3 diploidy classes ( $p \leq 0.01$ ). However,  
12 the 2 breeds did not differ significantly in the overall mean rates of X-Y aneuploidy (disomy +  
13 diploidy) which amounts to 0.162% in the Italian Friesian and 0.142% in the Italian Brown. When  
14 meiosis I (MI) and II (MII) errors were compared, statistically significant differences ( $p \leq 0.01$ ) were  
15 found in the disomy classes and in both breeds, whereas the differences between diploidy classes  
16 were not significant. Compared to humans, a lower level of aneuploidy has been found in the domestic  
17 species analysed so far. The present study contributes to the establishment of a baseline level of  
18 aneuploidy in the sperm populations of 2 cattle breeds which could be used for monitoring future  
19 trends of reproductive health, especially in relation to environmental changes and mutagens.

20

## 21 **Introduction**

22 Sperm chromosomes of domestic animals have been investigated by conventional cytogenetic  
23 methods since 1987 [Creighton and Houghton, 1987; Tateno and Mikamo, 1987] by using a quite  
24 expensive and laborious ‘pig or cattle/hamster’ heterospecific in vitro fertilization system.

25 The first report on the application of FISH directly on pig sperm was published by Kawarasaki et al.  
26 [1995], by using a male specific DNA probe; other contributions were made on pig, by using

1 molecular probes for chromosomes Y and 1 [Kawarasaki et al., 1996; Parrilla et al., 2003], or flow-  
2 sorted chromosomes X-Y [Kawarasaki et al., 1998], and on cattle, by using molecular probes for  
3 chromosome Y [Kobayashi et al., 1999], cosmid PL44 for the X and painting probe for the Y  
4 [Hassanane et al., 1999] or BACs for chromosome X and the repetitive sequence BRY4a for the Y  
5 [Piumi et al., 2001]. Chromosome-specific painting probes produced by chromosome microdissection  
6 or chromosome sorting followed by DOP-PCR were also used in the pig [Rubes et al., 1999] as well  
7 as in cattle [Rens et al., 2001; Révay et al., 2002] and in cattle, river buffalo, sheep and goat [Di  
8 Berardino et al., 2004]. Recently, Bonnet-Garnier et al. [2006] used bovine chromosome-specific  
9 painting probes for chromosomes and 29 to study meiotic segregation of the translocated  
10 chromosomes in sperm.

11 While the majority of these papers demonstrated the usefulness of the FISH technique as a method  
12 for validating sperm sexing technologies, only a few of them investigated aneuploidy rates. Since  
13 chromosomal abnormalities in the germ cells are one of the most important causes of embryonic  
14 mortality [King, 1990], and since we do not have sufficient information about the real impact of  
15 aneuploidy in the various domestic species and breeds, we aimed to further, at least partially, this  
16 knowledge.

17

## 18 **Material and Methods**

### 19 *Chromosome Microdissection and Probe Preparation*

20 Metaphase cells for the production of probes ‘via’ microdissection were prepared according to  
21 standard cytogenetic techniques [Iannuzzi and Di Berardino, 2008]. For microdissection, the fixed  
22 lymphocyte suspension was spread onto a precleaned 24 ! 60-mm coverslip which was then air dried  
23 and treated for GTG-banding.

24 The Xcen probe was produced by isolating the pericentromeric region, corresponding to the  
25 centromere and to the region Xp14–p11 of the standardized GTG-banded karyotype; the Y probe was  
26 produced by scraping the whole chromosome. Microdissected chromosomes were amplified

1 following the protocol of Engelen et al. [1998]. Probes were labeled with dUTP-11-digoxigenin  
2 (Xcen) and dUTP-16-biotin (Y) (both from Roche), respectively, in a second DOP-PCR reaction,  
3 using 2  $\mu$ l of product from the first reactions as template.

4

#### 5 *Semen Samples*

6 Cryopreserved semen samples belonging to the Italian Friesian and Italian Brown breeds of cattle  
7 were obtained from artificial insemination centers and private farms. For each breed, frozen semen  
8 from 10 animals was used for this study. All animals showed normal semen parameters and were all  
9 karyologically normal ( $2n = 60$ )

10

#### 11 *In situ Hybridization*

12 The Xcen and Y probes were hybridized simultaneously on metaphase plates for validation and  
13 subsequently on decondensed sperm. Decondensation of sperm nuclei was performed following the  
14 method described by Han et al. [1992]. Probes were precipitate in the presence of 10  $\mu$ g of salmon  
15 sperm DNA and 10  $\mu$ g of calf thymus DNA (both from Sigma), dissolved in 15  $\mu$ l of hybridization  
16 solution (50% formamide in 2  $\times$  SSC + 10% dextran sulphate; both from Sigma), denatured at 72  $^{\circ}$  C  
17 for 10 min and incubated at 37  $^{\circ}$  C for 90 min. Metaphase and sperm preparations were denatured for  
18 2 and 5 min respectively in a solution of 70% formamide in 2  $\times$  SSC (pH 7.0) at 72  $^{\circ}$  C. The slides  
19 were hybridized in a moist chamber at 37  $^{\circ}$  C overnight. After hybridization and slide washing, the  
20 biotin-labeled probe was revealed using a green Alexa 488 fluorochrome conjugated to streptavidin  
21 (Invitrogen), and the digoxigenin-labeled probe using a red Rhodamine fluorochrome conjugated to  
22 an anti-digoxigenin antibody from sheep (Roche). Slides were counterstained with DAPI (0.24  $\mu$ g/ml;  
23 Sigma) in Antifade (Vector Lab).

24

#### 25 *Fluorescence Analysis and Scoring*

1 The slides were observed at 100 ! magnification with a Leica DMRA fluorescence microscope  
2 equipped with DAPI, FITC, Cy3 specific filters, the DAPI/FITC/TXRD triple filter, and phase-  
3 contrast optics. Digital images were captured using the Leica QFISH software. At least 5,000 sperm  
4 nuclei were examined for each animal. The scoring was carried out using strict scoring criteria  
5 [Robbins et al., 1995]. Overlapped cells and those with ambiguous FISH signals were not scored. For  
6 each nucleus, type (X or Y) and number of sex chromosomes were analyzed. Sperm with one signal  
7 (green or red) were scored as normal haploid; spermatozoa with 2 signals were classified as disomic  
8 (XX, YY and XY depending on the 2 signal colors). Diploid sperm were distinguished from disomic  
9 sperm on the basis of their size [Joseph et al., 1984]. Since sperm decondensation might not be  
10 uniform along the slide, size comparison was made only within the same microscopic field where the  
11 diploid sperm were found. In any case, phase-contrast optics was used to check for the presence of  
12 the tail. Furthermore, sperm nuclei without signal were scored to calculate hybridization efficiency.

13

#### 14 *Validation of the Data*

15 In the present study, disomies and diploidies were differentiated according to the size of the nuclei;  
16 to verify if this could lead to errors in the estimation of aneuploidy, we performed an additional  
17 hybridization experiment on a limited sample of 6 animals (3 for each breed, previously analyzed  
18 with Xcen and Y probes) by using a probe for chromosome 6 [Habermann et al., 2005], labeled in  
19 red, and the 2 probes for X-Y chromosomes, both labeled in green. Hybridization conditions were the  
20 same as above. Ten thousand sperm were scored for each subject, in order to calculate the frequencies  
21 of diploidy (without distinguishing between XX, XY and YY diploidy), disomy for sex chromosome  
22 (without distinguishing between XX, XY and YY disomy) and disomy for chromosome 6. The results  
23 were then compared with those obtained on the same subjects using only Xcen and Y probes.

24

#### 25 *Statistical Analysis*

1 The following statistics were used: the  $\chi^2$  test with Yates' corrections for interindividual differences;  
2 the Kruskal-Wallis and the Mann-Whitney tests were used for multiple comparisons and for class  
3 differences.

4

## 5 **Results**

6 Figure 1 shows the Xcen and Y painting probes, DAPI staining and their diagrammatic representation.

7 Figure 2 shows disomic and diploid XX, XY and YY sperm after FISH staining.

8

### 9 *Interindividual Variations within the Breed ( table 1 )*

10 Totally, more than 100,000 sperm were examined with Xcen and Y probes, and more than 50,000  
11 sperm analysed for each breed (10 bulls). The hybridization efficiency was high, around 98–99% in  
12 all cases, except for subject 5 of the Friesian breed, where the hybridization efficiency was 92.34%.

13 A deviation from the expected 1: 1 ratio between the X- and Y-bearing sperm was found in both  
14 breeds in favour of the Y chromosome ( $p \neq 0.05$ ); such a deviation was particularly evident on subject  
15 8 of the Friesian breed. In the Friesian breed the differences among individuals for each class of  
16 aneuploidy (XX, XY, YY) were not statistically significant. In the Brown breed, differences among  
17 the animals were again not significant, except for the XY disomy ( $p < 0.05$ ).

18

### 19 *XX, YY, XY Class Comparison within the Breed ( table 2 )*

20 In the Friesian breed, statistically significant differences ( $p < 0.05$ ) were found only among the  
21 diploidy classes, whose frequencies were 0.015%–0.006%–0.029%, whereas in the Brown breed,  
22 differences were found not only among the 3 diploidy classes (0.018%–0.004%–0.041%) ( $p < 0.01$ )  
23 but also among the 3 disomy ones (0.022%–0.045%–0.012%) ( $p < 0.05$ ).

24 To analyze possible differences in the occurrence of errors during meiosis I (XY disomic/diploid  
25 sperm) or meiosis II (XX-YY disomic/diploid sperm) we applied the Mann-Whitney test. Meiotic  
26 errors giving rise to disomies were significantly ( $p \neq 0.01$ ) more frequent in MII than in MI (0.081%

1 vs. 0.031% in the Friesian, 0.067% vs. 0.012% in the Brown). Concerning the diploidy, the differences  
2 between MI and MII meiotic errors were not statistically significant. The overall frequency (disomies  
3 + diploidies) of errors in MII was higher than the frequency of errors in MI in both breeds ( $p < 0.05$ ).

4

#### 5 *Interbreed Comparison ( table 3 )*

6 The 2 breeds analyzed were basically similar in the overall frequency of X-Y chromosome aneuploidy  
7 (0.162% vs. 0.142%, respectively for Friesian and Brown). However, while in the Friesian breed the  
8 disomies (0.112%) were more frequent than diploidies (0.050%) ( $p \neq 0.01$ ), in the Brown the 2 types  
9 of abnormalities were basically similar (0.079% and 0.063%, respectively). The different classes of  
10 diploidy were equally represented between the 2 breeds; differences ( $p \neq 0.05$ ) were found at the level  
11 of XX disomy (0.044% vs. 0.022%) and XY disomy (0.031% vs. 0.012%).

12

#### 13 *Validation of the Data and Disomy of Chromosome 6 ( table 4 , fig. 3 )*

14 The analysis of data from the second set of slides on 6 subjects revealed a substantial similarity in the  
15 frequencies of diploidy (0.029% vs. 0.025%) and sex chromosomes disomy (0.139% vs. 0.142%)  
16 detected using the 2 different sets of probes. Disomy for chromosome 6 had an average frequency of  
17 0.080% (range 0.038–0.116%).

18

## 19 **Discussion**

20 The results of the present study indicated that in both breeds there was a significant preponderance  
21 of the Y-bearing sperm compared to the X. This finding is similar to that previously reported by  
22 Hassanane et al. [1999] on the Swedish Friesian breed. Unfortunately, due to the paucity of studies  
23 on this topic, we do not know whether this finding is a sporadic occurrence or a consistent  
24 phenomenon which is ongoing in the Friesian breed, but certainly cannot be explained as due to the  
25 low intensity of any of the 2 signals, as could happen when using small-size cosmid or BACs. In the  
26 present case, in fact, the probes we used provide quite strong signals being of the 'painting' type. It



1 is therefore necessary to expand such investigations in order to provide more information on this  
2 aspect. The present study also indicated a quite low interindividual variability in the frequencies of  
3 the different aneuploidy classes. Since the 2 breeds analysed in the present study are highly selected,  
4 the consequent great genetic uniformity could explain the low variability observed. The analysis of  
5 unselected breeds (or genetic types) would clarify if this low variability is characteristic of the species  
6 *Bos taurus* or is a consequence of the genetic uniformity due to selection. The analysis 'within' each  
7 breed showed several differences. In the Italian Friesian, disomies were more frequent than diploidies,  
8 but while the former were equally represented in the 3 classes, for the latter XY diploidies showed a  
9 higher frequency compared to YY. On the contrary, in the Brown breed, disomies and diploidies had  
10 the same frequencies in both cases, while the frequencies of the different classes were different. In  
11 this breed, disomic YY sperm had a higher frequency compared to XY disomic sperm. Also in this  
12 breed, like in the Italian Friesian, XY diploidies were the most represented. In order to explain this  
13 difference, abnormalities were classified according to their origin as arising from errors in meiosis I  
14 (XY disomic/diploid sperm) or meiosis II (XX-YY disomic/diploid sperm). Disomies were more  
15 frequent due to errors occurring during MII in both breeds, while diploidies originated from errors in  
16 MI or in MII with the same frequencies. This means that diploidy can originate with the same  
17 frequencies as a consequence of chromatid or chromosomal nondisjunction events, while, at least for  
18 sex chromosomes, disomies originate mainly from chromatid nondisjunction.

19 Usually, the reduced pairing of sex chromosomes during MI is considered responsible for their higher  
20 aneuploidy incidence compared to autosomes [Shi and Martin, 2000]. However, in humans, while  
21 some authors confirmed this finding [Robbins et al., 1995; Downie et al., 1997], others [Spriggs et  
22 al., 1995; Rubes et al., 2005] did not find differences in the frequencies of MI and MII errors, thus  
23 indicating that other factors could be responsible for the higher incidence of sex chromosome  
24 aneuploidy in sperm. Our results seem to confirm the latter hypothesis, showing that X and Y  
25 disomies originate mainly from chromatid nondisjunction, indicating that, in this species, the reduced  
26 pairing of X and Y is responsible for only a small part of the total disomies. The interbreed

1 comparison ( table 3 ) showed that the 2 breeds did not differ from each other in the overall rate of X  
2 and Y aberrant spermatozoa (disomic + diploid) which were 0.162% in the Friesian and 0.142% in  
3 the Brown. The diploidy rate was very similar (0.050% vs. 0.062%), while for the disomy rate Italian  
4 Friesian showed a higher frequency than Italian Brown (0.112% vs. 0.079%), but this difference was  
5 not significant for  $p = 0.05$ . Due to the paucity of scientific reports on aneuploidy rates in sperm of  
6 cattle, we could compare the present results only to those previously reported by Hassanane et al.  
7 [1999] and by Di Berardino et al. [2004] ( table 3 ). Concerning the Italian Friesian breed, close  
8 similarities can be noticed between the results of the present study and those previously reported by  
9 Hassanane et al. [1999] in the Swedish Friesian breed: in fact, the X-Y disomy rates are 0.112% and  
10 0.125%, respectively, whereas the X-Y diploidy rates are 0.050% and 0.045%, respectively. Such  
11 agreement in the results can be further explained by the fact that the Friesian breed of cattle is highly  
12 selected everywhere, around the world, and the connections between countries could have led to a  
13 homogenization of the population from a genetic point of view. From table 3 it can also be seen that  
14 the aneuploidy rates detected in cattle breeds, so far, are also comparable to those reported in several  
15 other breeds of pig [Rubes et al., 1999]. The YY disomy rates in cattle, in fact, vary between 0.029%  
16 and 0.045%, quite close to the value observed in the Bohemian Fleishy breed (0.058%), being lower  
17 than those found in the Large White (0.112%), White Improved (0.125%) and Pietrain (0.170%), but  
18 higher than those of the Landrace (0.006%) and Duroc (0.009%). Surprisingly, the YY diploidy rate  
19 found in cattle is between 0.004% and 0.024%, much less compared to the corresponding values  
20 observed in the pig breeds, where the YY diploidy rate varies from 0.133% to 0.350%.

21 In order to compare cattle with other domestic species, we averaged the mean aneuploidy values  
22 previously reported in the Swedish Friesian [Hassanane et al., 1999] with those reported in the present  
23 study on the Italian Friesian and Italian Brown breeds ( table 5 ). The mean rates for cattle (*Bos*  
24 *taurus*) resulted in 0.105% for disomy, 0.052% for diploidy and 0.157% for the total aneuploidy rate.  
25 These values were then compared to those of the river buffalo ( *Bubalus bubalis* , river type), sheep  
26 (*Ovis aries*) and goat (*Capra hircus*) reported by Di Berardino et al. [2004], pig (*Sus scrofa*

1 *domestica*) reported by Rubes et al. [1999], mouse (*Mus musculus*) reported by Adler et al. [1996]  
2 and humans (*Homo sapiens*) calculated as average values [Griffin et al., 1995; Martin et al., 1995;  
3 Robbins et al., 1995; Spriggs et al., 1995; Downie et al., 1997; Rubes et al., 2005]. By taking human  
4 as reference: (a) the total disomy rates found in the domestic species investigated, so far, were all  
5 lower, being comprised of 0.080% (mouse) to 0.164% (goat); (b) the total diploidy rates were higher  
6 only in the goat (0.229%) and lower in the others, being comprised of 0.033% (sheep) and 0.177%  
7 (pig); (c) the total aneuploidy (disomy + diploidy) rates were all lower, being comprised of 0.033%  
8 (sheep) to 0.393% (goat). The fact that in the domestic species the aneuploidy rates are lower  
9 compared to human could, at least in part, be explained by the fact that domestic animals have been  
10 strongly selected for fertility, and this could have led to a reduction in the frequencies of aberrant  
11 spermatozoa. To verify this hypothesis, it could be useful to investigate wild species belonging to the  
12 same family as the domestic ones, in order to clarify the effects of the artificial selection on the sperm  
13 aneuploidy rate. In addition, domestic animals have been targeted by FISH only recently and,  
14 therefore, only few data are available, whereas in humans such studies are more abundant and  
15 consistent. The present work focused on X and Y sperm aneuploidy, using sperm size as a criteria for  
16 distinguishing between disomic and diploid sperm. Size comparison was made only within the same  
17 microscopic field where the diploid sperm were found and phase-contrast optics was used to check  
18 for the presence of the tail, to be certain that it was not a round cell. To verify if this kind of approach  
19 could determine an 'under' or an 'over' estimation in the frequencies of the different classes of  
20 abnormalities, we performed an additional experiment by using an autosomal probe (chromosome 6)  
21 on a limited number of subjects.

22 As shown in table 4 , the data obtained from the 2 analyses were very similar, thus indicating that, at  
23 least under the experimental conditions used in the present study, the scoring criteria we used were  
24 sufficient to distinguish between disomic and diploid cells. In addition, disomy of chromosome 6 had  
25 an incidence of 0.080%, lower than disomy of the X-Y chromosomes in the animals tested, except  
26 for one. As in humans, it seems that the sex chromosomes are more prone to undergo non-disjunction

1 events. However, it will be necessary to extend the analysis to a higher number of animals and to use  
2 more probes, in order to test possible interchromosomal differences.

3 Since mammalian fertility is strongly affected by chromosomal abnormalities, which are responsible  
4 for nearly 70% of the embryonic mortality in humans [Hassold, 1998] as well as in domestic animals  
5 [King, 1990; Vanroose et al., 2000], further studies of germ cells (sperm and oocytes) should be  
6 implemented in order to better understand the genetic causes of aneuploidies and their impact on the  
7 reproductive and productive efficiency of domestic animals. Several studies [for a review, see  
8 Pacchierotti et al., 2007] showed that chemical substances, commonly used in medicine and  
9 agriculture (drug and pesticide) or present in the environment as pollutant, can increase the  
10 frequencies of germ cell aneuploidy in human and mouse. Domestic animals are often exposed to  
11 such substances through the farming environment or feedstuff; and the sperm-FISH assay can  
12 represent a useful tool to allow the identification of chemicals that can negatively affect the animal's  
13 health, thus reducing its reproductive efficiency.

14

## 15 **Acknowledgment**

16 This work was supported by the financial contribution of the Ministry of Agricultural and Forestry  
17 Politics (MiPAAF) of Rome (SpermovoFISH project n. 291/7303/06) which is gratefully  
18 acknowledged.

19

## 20 **References**

21 Adler ID, Bishop J, Lowe X, Schmid TE, Schriever-Schwemmer G: Spontaneous rate of sex  
22 chromosomal aneuploidies in sperm and offspring of mice: a validation of the detection of aneuploid  
23 sperm by fluorescence in situ hybridization. *Mutat Res* 372: 259–268 (1996).

24

1 Bonnet-Garnier A, Pinton A, Berland H, Khireddine B, Eggen A, et al: Sperm nuclei analysis of 1/29  
2 Robertsonian translocation carrier bulls using fluorescence in situ hybridization. *Cytogenet Genome*  
3 *Res* 112: 241–247 (2006).  
4  
5 Creighton P, Houghton JA: Visualization of pig sperm chromosomes by in vitro penetration of zona-  
6 free hamster ova. *J Reprod Fertil* 80: 619–622 (1987).  
7  
8 Di Berardino D, Vozdova M, Kubickova S, Cernohorska H, Coppola G, et al: Sexing river buffalo  
9 (*Bubalus bubalis* L.), sheep (*Ovis aries* L.), goat (*Capra hircus* L.), and cattle spermatozoa by double  
10 color fish using bovine (*Bos taurus* L.) X- and Y-painting probes. *Mol Reprod Dev* 67: 108–115  
11 (2004).  
12  
13 Downie SE, Flaherty SP, Swann NJ, Matthews CD: Estimation of aneuploidy for chromosomes 3, 7,  
14 16, X and Y in spermatozoa from 10 normospermic men using fluorescence in-situ hybridization.  
15 *Mol Hum Reprod* 3: 815–819 (1997).  
16  
17 Engelen JJ, Albrechts JC, Hamers GJ, Geraedts JP: A simple and efficient method for microdissection  
18 and microFISH. *J Med Genet* 35: 265–268 (1998).  
19  
20 Griffin DK, Abruzzo MA, Millie EA, Sheean LA, Feingold E, et al: Non-disjunction in human sperm:  
21 evidence for an effect of increasing paternal age. *Hum Mol Genet* 4: 2227–2232 (1995).  
22  
23 Habermann FA, Winter A, Olsaker I, Reichert P, Fries R: Validation of sperm sexing in the cattle  
24 (*Bos taurus*) by dual color fluorescence in situ hybridization. *J Anim Breed Genet* 122 Suppl 1: 22–  
25 27 (2005).  
26

1 Han TL, Webb GC, Flaherty SP, Correll A, Matthews CD, Ford JH: Detection of chromosome 17-  
2 and X-bearing human spermatozoa using fluorescence in situ hybridization. *Mol Reprod Dev* 33:  
3 189–194 (1992).

4

5 Hassanane M, Kovacs A, Laurent P, Lindblad K, Gustavsson I: Simultaneous detection of X and Y-  
6 bearing bull spermatozoa by double color fluorescence in situ hybridization. *Mol Reprod Dev* 53:  
7 407–412 (1999).

8

9 Hassold TJ: Nondisjunction in the human male; in Handel MA (ed): *Meiosis and gametogenesis*, pp  
10 383–406 (Academic Press, New York 1998).

11

12 Iannuzzi L, Di Berardino D: Tools of the trade: diagnostics and research in domestic animal  
13 cytogenetics. *J Appl Genet* 49: 357–366 (2008).

14

15 Joseph AM, Gosden JR, Chandley AC: Estimation of aneuploidy levels in human spermatozoa using  
16 chromosome specific probes and in situ hybridisation. *Hum Genet* 66: 234–238 (1984).

17

18 Kawarasaki T, Kohsaka T, Sone M, Yoshida M, Bamba K: Detection of Y-bearing porcine  
19 spermatozoa by in situ hybridization using digoxigenin-labeled, porcine male-specific DNA probe  
20 produced by polymerase chain reaction. *Mol Reprod Dev* 40: 455–459 (1995).


21

22 Kawarasaki SJ, Basrur PK, Stubbings RB, Hansen PJ, King WA: Chromosomal abnormalities in  
23 bovine embryos and their influence on development. *Biol Reprod* 54: 53–59 (1996).

24

- 1 Kawarasaki SJ, Welch GR, Long CR, Yoshida M, Johnson LA: Verification of flow cytometrically  
2 sorted X- and Y-bearing porcine spermatozoa and reanalysis of spermatozoa for DNA content using  
3 the fluorescent in situ hybridization (FISH) technique. *Theriogenology* 50: 625–635 (1998).
- 4
- 5 King WA: Chromosome abnormalities and pregnancy failure in domestic animals; in McFeely RA  
6 (ed): *Domestic Animal Cytogenetics*, pp 229–250 (Academic Press, New York 1990).
- 7
- 8 Kobayashi J, Kohsaka T, Sasada H, Umezu M, Sato E: Fluorescence in situ hybridization with Y  
9 chromosome-specific probe in decondensed bovine spermatozoa. *Theriogenology* 52: 1043–1054  
10 (1999).
- 11
- 12 Martin RH, Spriggs E, Ko E, Rademaker AW: The relationship between paternal age, sex ratios, and  
13 aneuploidy frequencies in human sperm, as assessed by multicolor FISH. *Am J Hum Genet* 57: 1395–  
14 1399 (1995).
- 15
- 16 Pacchierotti F, Adler I, Eichenlaub-Ritter U, Mailhes JB: Gender effects on the incidence of  
17 aneuploidy in mammalian germ cells. *Environ Res* 104: 46–69 (2007).
- 18
- 19


1 **Fig. 1.** Xcen and Y painting probes, DAPI staining and diagrammatic representation



2  
3




1 **Fig. 2.** Fluorescent in situ hybridization (FISH) on cattle sperm showing XX, XY and YY disomic  
2 and diploid sperm.



3  
4

1 **Fig. 3.** Fluorescent in situ hybridization (FISH) on cattle sperm using X-Y probe (green) and  
2 chromosome 6 probe (red) showing disomic (a) and diploid (b) sperm.  
3



4  
5

1 **Table 1.** Number and frequency (%) of X- and Y-bearing, disomic and diploid sperm in bulls of the  
 2 Italian Friesian (I.F.) and Italian Brown (I.B.) breeds of cattle

	X	Y	Disomic			Diploid			With signal	Without signal	Total sperm
			XX	YY	XY	XX	YY	XY			
I.F.											
1	2,819 (46.968)	3,154 (52.549)	1 (0.017)	5 (0.083)	1 (0.017)	1 (0.017)	0 (0)	0 (0)	5,981 (99.650)	21 (0.350)	6,002
2	2,411 (47.081)	2,623 (51.220)	4 (0.078)	5 (0.098)	2 (0.039)	1 (0.020)	0 (0)	0 (0)	5,046 (98.535)	75 (1.465)	5,121
3	2,643 (51.300)	2,487 (48.273)	5 (0.097)	2 (0.039)	2 (0.039)	1 (0.019)	0 (0)	1 (0.019)	5,141 (99.786)	11 (0.214)	5,152
4	2,509 (49.380)	2,564 (50.463)	2 (0.039)	0 (0)	1 (0.020)	0 (0)	0 (0)	0 (0)	5,076 (99.902)	5 (0.098)	5,081
5	2,420 (44.658)	2,574 (47.500)	2 (0.037)	1 (0.018)	2 (0.037)	1 (0.018)	0 (0)	4 (0.074)	5,004 (92.342)	415 (7.658)	5,419
6	2,257 (47.080)	2,468 (51.481)	0 (0)	2 (0.042)	1 (0.021)	1 (0.021)	1 (0.021)	1 (0.021)	4,731 (98.686)	63 (1.314)	4,794
7	2,510 (47.225)	2,761 (51.947)	2 (0.038)	0 (0)	1 (0.019)	0 (0)	1 (0.019)	2 (0.038)	5,277 (99.285)	38 (0.715)	5,315
8	2,380 (44.846)	2,878 (54.230)	4 (0.075)	2 (0.038)	2 (0.038)	0 (0)	1 (0.019)	2 (0.038)	5,269 (99.284)	38 (0.716)	5,307
9	2,472 (46.094)	2,854 (53.216)	3 (0.056)	1 (0.019)	2 (0.037)	1 (0.019)	0 (0)	2 (0.037)	5,335 (99.478)	28 (0.522)	5,363
10	2,372 (47.138)	2,645 (52.564)	0 (0)	1 (0.020)	2 (0.040)	2 (0.040)	0 (0)	3 (0.060)	5,025 (99.861)	7 (0.139)	5,032
All	24,793 (47.148)	27,008 (51.360)	23 (0.044)	19 (0.037)	16 (0.031)	8 (0.015)	3 (0.006)	15 (0.029)	51,885 (98.667)	701 (1.333)	52,586
I.B.											
1	2,548 (49.824)	2,505 (48.983)	2 (0.039)	4 (0.078)	0 (0)	1 (0.020)	0 (0)	1 (0.020)	5,061 (98.964)	53 (1.036)	5,114
2	2,474 (47.559)	2,669 (51.307)	0 (0)	3 (0.058)	0 (0)	1 (0.019)	0 (0)	2 (0.038)	5,149 (98.981)	53 (1.019)	5,202
3	2,444 (47.383)	2,632 (51.028)	0 (0)	6 (0.116)	2 (0.039)	0 (0)	0 (0)	0 (0)	5,084 (98.565)	74 (1.435)	5,158
4	2,635 (49.905)	2,589 (49.034)	2 (0.038)	1 (0.019)	0 (0)	2 (0.038)	0 (0)	3 (0.057)	5,232 (99.091)	48 (0.909)	5,280
5	2,505 (48.802)	2,576 (50.185)	2 (0.039)	1 (0.019)	0 (0)	2 (0.039)	0 (0)	2 (0.039)	5,088 (99.123)	45 (0.877)	5,133
6	2,374 (45.866)	2,714 (52.434)	1 (0.019)	2 (0.039)	0 (0)	0 (0)	0 (0)	2 (0.039)	5,093 (98.396)	83 (1.604)	5,176
7	2,246 (44.643)	2,715 (53.965)	3 (0.060)	1 (0.020)	0 (0)	0 (0)	0 (0)	6 (0.119)	4,971 (98.807)	60 (1.193)	5,031
8	2,327 (46.318)	2,664 (53.025)	1 (0.020)	3 (0.060)	1 (0.020)	0 (0)	1 (0.020)	1 (0.020)	4,998 (99.482)	26 (0.518)	5,024
9	2,345 (45.209)	2,796 (53.904)	0 (0)	1 (0.019)	0 (0)	1 (0.019)	1 (0.019)	1 (0.019)	5,145 (99.190)	42 (0.810)	5,187
10	2,415 (47.945)	2,590 (51.419)	0 (0)	1 (0.020)	3 (0.060)	2 (0.040)	0 (0)	3 (0.060)	5,014 (99.543)	23 (0.457)	5,037
All	24,313 (47.355)	26,450 (51.517)	11 (0.022)	23 (0.045)	6 (0.012)	9 (0.018)	2 (0.004)	21 (0.041)	50,835 (99.013)	507 (0.987)	51,342

1 **Table 2.** Statistical significance of the comparisons ‘within’ each breed in the frequency of the  
 2 different aneuploidy classes

	Italian Friesian			Italian Brown		
	%	com- parison	p	%	com- parison	p
<i>Disomy</i>						
XX (1)	0.044	1-2	NS	0.022	1-2	NS
YY (2)	0.037	2-3	NS	0.045	2-3	<0.05
XY (3)	0.031	1-3	NS	0.012	1-3	NS
Tot (4)	0.112	4-10	<0.01	0.079	4-10	NS
MI (5)	0.031	5-6	<0.01	0.012	5-6	<0.01
MII (6)	0.081	-		0.067	-	
<i>Diploidy</i>						
XX (7)	0.015	7-8	NS	0.018	7-8	NS
YY (8)	0.006	8-9	<0.05	0.004	8-9	<0.01
XY (9)	0.029	7-9	NS	0.041	7-9	NS
Tot (10)	0.050	-		0.063	-	
MI (11)	0.029	11-12	NS	0.041	11-12	NS
MII (12)	0.021			0.022		

1 **Table 3.** Interbreed variability in the frequency of X-Y sperm aneuploidy (disomy + diploidy)

Trait	Cattle			Pig					
	S.F.	I.F.	I.B.	L.W.	B.F.	L.	W.I.	D.	P.
No. animals	5	10	10	6	4	3	2	1	1
No. sperm/animal	10,000	5,000	5,000	10,000	10,000	10,000	10,000	10,000	10,000
Disomy XX	0.067	0.044	0.022						
Disomy YY	0.029	0.037	0.045	0.112	0.058	0.006	0.125	0.009	0.170
Disomy XY	0.029	0.031	0.012						
<b>Total dis. (a)</b>	<b>0.125</b>	<b>0.112</b>	<b>0.079</b>	<b>0.112<sup>a</sup></b>	<b>0.058<sup>a</sup></b>	<b>0.006<sup>a</sup></b>	<b>0.125<sup>a</sup></b>	<b>0.009<sup>a</sup></b>	<b>0.170<sup>a</sup></b>
Diploid XX	0.021	0.015	0.018						
Diploid YY	0.024	0.006	0.004						
Diploid XY	n.d.	0.029	0.041						
<b>Total dip. (b)</b>	<b>0.045</b>	<b>0.050</b>	<b>0.063</b>	<b>0.192</b>	<b>0.133</b>	<b>0.140</b>	<b>0.170</b>	<b>0.350</b>	<b>0.210</b>
(a) + (b)	0.170	0.162	0.142						
Reference <sup>b</sup>	(1)	(2)	(2)	(3)	(3)	(3)	(3)	(3)	(3)

2  
3 <sup>a</sup>Data available only for YY disomy. <sup>b</sup>(1) Hassanane et al. [1999]; (2) present study; (3) Rubes et al. [1999].

4 S.F. = Swedish Friesian; I.F. = Italian Friesian; I.B. = Italian Brown; L.W. = Large White; B.F. = Bohemian  
5 Fleshy; L = Landrace; W.I. = White Improved; D. = Duroc; P. = Pietrain.  
6

1 **Table 4.** Comparison of the disomy and diploidy frequencies using the two different sets of probes  
 2 and disomy frequency for chromosome 6.

Breed	Sub.	Disomic XY		Diploid		Disomic 6
		probes X+Y	probes XY+6	probes X+Y	probes XY+6	
I.F.	1	0.117	0.134	0.017	0.022	0.038
	2	0.215	0.233	0.020	0.019	0.077
	3	0.175	0.157	0.038	0.029	0.098
I.B.	4	0.117	0.136	0.040	0.038	0.097
	5	0.058	0.058	0.057	0.030	0.116
	6	0.155	0.136	0.000	0.009	0.058
Total		0.139	0.142	0.029	0.025	0.080

X+Y = Xcen labeled in red; Y labeled in green.

XY+6 = Xcen and Y labeled in green; 6 labeled in red.