

IRIS A<sub>per</sub>TO



UNIVERSITÀ  
DEGLI STUDI  
DI TORINO

This is the author's final version of the contribution published as:

E. Lumini; A. Orgiazzi; R. Borriello; P. Bonfante; V. Bianciotto. Disclosing arbuscular mycorrhizal fungal biodiversity in soil through a land-use gradient using a pyrosequencing approach. ENVIRONMENTAL MICROBIOLOGY. 12(8) pp: 2165-2179.

DOI: 10.1111/j.1462-2920.2009.02099.x

The publisher's version is available at:

<http://doi.wiley.com/10.1111/j.1462-2920.2009.02099.x>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/62186>

This full text was downloaded from iris - AperTO: <https://iris.unito.it/>

---

iris - AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

# Disclosing arbuscular mycorrhizal fungal biodiversity in soil through a land-use gradient using a pyrosequencing approach

## Authors

Erica Lumini, Alberto Orgiazzi, Roberto Borriello, Paola Bonfante, Valeria Bianciotto

First published: 17 November 2009

DOI: 10.1111/j.1462-2920.2009.02099.x

E-mail v.bianciotto@ipp.cnr.it; Tel. (+39) 11 6502927/51; Fax (+39) 11 6705962.

## Summary

The biodiversity of arbuscular mycorrhizal fungi (AMF) communities present in five Sardinian soils (Italy) subjected to different land-use (tilled vineyard, covered vineyard, pasture, managed meadow and cork-oak formation) was analysed using a pyrosequencing-based approach for the first time. Two regions of the 18S ribosomal RNA gene were considered as molecular target. The pyrosequencing produced a total of 10924 sequences: 6799 from the first and 4125 from the second target region. Among these sequences, 3189 and 1003 were selected to generate operational taxonomic units (OTUs) and to evaluate the AMF community richness and similarity: 117 (37 of which were singletons) and 28 (nine of which were singletons) unique AMF OTUs were detected respectively. Within the *Glomeromycota* OTUs, those belonging to the *Glomerales* order were dominant in all the soils. *Diversisporales* OTUs were always detected, even though less frequently, while *Archaeosporales* and *Paraglomerales* OTUs were exclusive of the pasture soil. Eleven OTUs were shared by all the soils, but each of the five AMF communities showed particular features, suggesting a meaningful dissimilarity among the *Glomeromycota* populations. The environments with low inputs (pasture and covered vineyard) showed a higher AMF biodiversity than those subjected to human input (managed meadow and tilled vineyard). A reduction in AMF was found in the cork-oak formation because other mycorrhizal fungal species, more likely associated to trees and shrubs, were detected. These findings reinforce the view that AMF biodiversity is influenced by both human input and ecological traits, illustrating a gradient of AMF communities which mirror the land-use gradient. The high number of sequences obtained by the pyrosequencing strategy has provided detailed information on the soil AMF assemblages, thus offering a source of light to shine on this crucial soil microbial group.

# Introduction

Metagenomic studies have recently provided new approaches that shed light on microbial communities in a variety of environments – e.g. sewage digestors, dental plaque, termite gut, deep mine drainages – ([Schloss and Handelsman, 2005](#); [Hugenholtz and Tyson, 2008](#)). Metagenomics provides a relatively unbiased view not only of the community structures (species richness and distribution), but also of the functional potential of a community ([van Elsas \*et al.\*, 2008](#)).

In this context, DNA pyrosequencing with the 454 GS-FLX platform, a rapid and relatively inexpensive sequencing technology that produces hundreds of thousands of short sequences, is being used more and more for prokaryotic metagenomic studies ([Edwards \*et al.\*, 2006](#); [Warnecke \*et al.\*, 2007](#)). This technique has recently been used to enumerate and compare soil bacterial diversity ([Roesch \*et al.\*, 2007](#); [Fulthorpe \*et al.\*, 2008](#)). However, so far, no pyrosequencing-based studies have been focused on soil fungi. Fungi are a crucial component of soil microbial communities, in which they function as decomposers, pathogens and mycorrhizal mutualists. Among the members of this last group, arbuscular mycorrhizal fungi (AMF) are the most important symbionts in many ecosystems ([Oehl \*et al.\*, 2003](#); [Öpik \*et al.\*, 2008](#); [Toljander \*et al.\*, 2008](#)). Arbuscular mycorrhizal fungi have been separated from all other major fungal groups and grouped in a monophyletic clade, named *Glomeromycota* ([Schüßler \*et al.\*, 2001](#)), within which four orders (*Glomerales*, *Diversisporales*, *Archaeosporales* and *Paraglomerales*) have been described ([Krüger \*et al.\*, 2009](#)).

Arbuscular mycorrhizal fungi play a key role in supplying phosphorus to plants, which in return receive plant carbon assimilates ([Girlanda \*et al.\*, 2007](#); [Smith and Read, 2008](#)). In addition to an improvement in plant nutrition, AM fungi protect their hosts from pathogens ([Poza and Azcon-Aguilar, 2007](#)) and affect plant growth traits ([Artursson \*et al.\*, 2006](#)). Furthermore, it has been suggested that mycorrhizal fungal diversity is a determinant of plant diversity ([van der Heijden \*et al.\*, 1998](#); [van der Heijden and Scheublin, 2007](#)). This concept has been a driving force in the investigation of AMF community diversity in different environments ([Öpik \*et al.\*, 2006](#); [Vallino \*et al.\*, 2006](#)) and up to 34 different AMF taxa have been found in a single habitat ([Öpik \*et al.\*, 2008](#)). Arbuscular mycorrhizal fungi taxa have a specific multidimensional niche that is determined by the plant species that are present at a site and by edaphic factors such as pH, moisture content, phosphorus (P) and nitrogen (N) availability. As a result, large between- and within-site variations in the composition of AMF communities have been described ([Burrows and Pflieger, 2002](#); [Klironomos and Hart, 2002](#)). However, a basic weakness of AMF community analyses is that many studies have been exclusively focused on roots, without considering that the extra-radical phase is at least as important as the intraradical mycelia for the nutrient supply of the host plant ([Horton and Bruns, 2001](#)). The current interest in extra-radical mycelium is increasing ([Croll \*et al.\*, 2009](#)) because the occurrence of anastomoses between genetically distinct mycelia suggests the existence of mycelial networks which connect different plant–host roots in soil ([Young, 2009](#)).

Here we present, to our knowledge, the first and the largest AMF soil-based sequence data set, obtained from high-throughput DNA pyrosequencing. Our investigation had two objectives: first to test the potentiality of pyrosequencing to investigate the diversity of AMF in a Mediterranean ecosystem, and second to characterize and compare AMF communities along a land-use gradient. The study area, located in Berchidda (Sardinia, Italy), in the past was covered by cork-oak forests (*Quercus suber* L). Over the years this vegetation has been subjected to intense usage for the extraction of cork and pasture. Therefore, today, it is possible to find different land-use units close together: tilled vineyard (coded TV), non-tilled cover cropped vineyard (CV), managed meadow (MM), pasture or grassland (PA) (dominated by pasture or grass species with a low tree density) and cork-oak formation (CO) (dominated by shrub cover and distributed cork-oak trees).

Using two couples of primers we obtained well over 10 000 eukaryotic rRNA 18S gene fragments from five Sardinian soils. The sequences were classified into operational taxonomic units (OTUs). The OTUs were created using the DOTUR software, which grouped the 18S rDNA sequences according to different similarity levels. This study has been focused on OTUs defined at the  $\geq 97\%$  similarity level (OTU<sub>0.03</sub>) to characterize the AMF communities. Although this distance cut-off is arbitrary and could be considered controversial, it has been used in many studies ([Schloss and Handelsman, 2005](#); [Huber et al., 2007](#)), and it facilitates comparisons with similar studies based on cloning and sequencing ([O'Brien et al., 2005](#)). The sequence data sets allowed us to make comparative analyses of AM fungal communities, taxa richness and coverage estimates as well as to obtain an overall description of the order-level diversity.

## Results

### Overall taxonomic richness

A total of 10 924 sequences were obtained: 6799 and 4125 with the AMV4.5NF/AMDGR and NS31/AMmix primer pair respectively. Only fragments of  $\geq 230$  bp in length were analysed for AMV4.5NF/AMDGR and of  $\geq 250$  bp for NS31/AMmix, leading to a total of 4192 sequences: 3189 (average length: 258 bp) and 1003 (average length: 274 bp) for AMV4.5NF/AMDGR and NS31-AMmix respectively ([Fig. 1A](#)).




Figure 1. Table (A) reports the number of sequences of each fungal phylum detected with the AMV4.5NF/AMDGR and NS31/AMmix primer pairs in all the Sardinian soils sampled. Picture (B) shows the overall and proportional distribution of the fungal phyla detected. Glomeromycota (blue) and Ascomycota (red) were the most abundant taxa found with the two couples of primers respectively.

In spite of the supposed AM primer specificity, some ‘contaminant’ sequences were detected, belonging to taxa different from *Glomeromycota*. The taxonomic distribution of the 18S sequences obtained with each primer pair in the Sardinian soils is shown in [Fig. 1A and B](#). The majority of eukaryotic sequences (2438 corresponding to 76.45% of the total) amplified with AMV4.5NF/AMDGR belong to *Glomeromycota*. With the same primer pair, the *Basidiomycota* relative abundance was 11.13%, while only 0.13% of the sequences was assigned to *Ascomycota*. Among the 1003 NS31/AMmix sequences, 448 sequences, corresponding to 44.67% of the total sequences, instead referred to *Ascomycota* and only 37.59% of the sequences (377) to *Glomeromycota*. The *Basidiomycota* relative frequency was 3.49%. The other two fungal phyla (*Chytridiomycota* and *Zygomycota*) were recovered less by both primer pairs: 0.10% with NS31/AMmix and 3.23% with AMV4.5NF/AMDGR, and 0.22% (AMV4.5NF/AMDGR) and 1.30% (NS31/AMmix) respectively ([Fig. 1](#)).

### Arbuscular mycorrhizal fungi community richness

Overall, 2815 *Glomeromycota* sequences were obtained: 2438 and 377 with AMV4.5NF/AMDGR and NS31/AMmix respectively. Considering each soil type, AMF sequences ranged from 188 (CO) to 1275 (PA) for the AMV4.5NF/AMDGR primers and from 12 (CO) to 200 (PA) for the NS31/AMmix primers ([Table 1](#)). Despite the different sequence numbers obtained, there was good agreement between the amplification results obtained from the two couples of primers, possibly reflecting an actual higher presence of *Glomeromycota* in the PA soil and a substantially lower occurrence in the CO soil.

Table 1. Number of *Glomeromycota* sequences detected with the AMV4.5NF/AMDGR and NS31/AMmix primer pairs.

	Number of <i>Glomeromycota</i> sequences	
	AMV4.5NF/AMDGR	NS31/AMmix
1. Numbers refer to the $\geq 230$ bp sequences for the first couple of primers and to $\geq 250$ bp for the second one.		
Tilled vineyard	429	17
Covered vineyard	292	66
Managed meadow	254	82
Pasture	1275	200
Cork-oak formation	188	12
Total	2438	377

The obtained AMF sequences were grouped into OTUs, representing groups of sequences at different similarity levels. Rarefaction curves were used to depict the effect of percentage similarity on the number of identified OTUs (Fig. 2). The patterns of the rarefaction curves relative to the two targeted 18S rDNA regions, amplified by the two primer sets, were similar: at high similarity levels (97%, OTU<sub>S0.03</sub> and 95%, OTU<sub>S0.05</sub>), neither curve reached the plateau, while at low similarity levels (85%, OTU<sub>S0.15</sub> and 80%, OTU<sub>S0.20</sub>) both the curves reached the plateau. This result is confirmed by a comparison of the observed (rarefaction index) and estimated (non-parametric ACE-Abundance base Coverage Estimator and Chao1 indices) OTUs at different levels of similarity: at the 80% and 85% sequence similarity levels, the number of observed OTUs was found to be equivalent, or close to those of the number of estimated OTUs (ACE and Chao1 indices), while at higher similarity levels, the ACE and Chao1 richness values were relatively far from the observed ones (Table 2).




Figure 2. Rarefaction curves for the AMV4.5NF/AMDGR and NS31/AMmix sequences at different similarity levels ranging from 80% to 97%. The 97% similarity level corresponds to OTU<sub>0.03</sub>, the 95% to OTU<sub>0.05</sub>, the 85% to OTU<sub>0.15</sub> and the 80% to OTU<sub>0.20</sub>.

Table 2. The ability of the two non-parametric richness estimators (ACE and Chao1) to predict the number of *Glomeromycota* OTUs at different levels of similarity in Sardinian soils is compared with the numbers of observed OTUs (rarefaction index).

	Observed OTUs					Estimated OTUs							
	Rarefaction					ACE				Chao1			
Similarity (%)	97	95	85	80	97	95	85	80	97	95	85	80	
AMV4.5NF/AMDGR	<b>117</b>	<b>69</b>	<b>24</b>	<b>14</b>	<b>161</b>	<b>85</b>	<b>33</b>	<b>14</b>	<b>156</b>	<b>79</b>	<b>32</b>	<b>14</b>	
NS31/AMmix	<b>28</b>	<b>20</b>	<b>8</b>	<b>4</b>	<b>35</b>	<b>23</b>	<b>9</b>	n.c.	<b>37</b>	<b>25</b>	<b>8</b>	n.c.	

1. The values are calculated using the DOTUR software for both couples of primers.
2. n.c., not calculated.

The 97% sequence similarity level, corresponding to OTU<sub>0.03</sub>, was used in the subsequent analyses. The 2438 AMV4.5NF/AMDGR *Glomeromycota* sequences comprised 117 unique OTU<sub>0.03</sub>, 37 of which were singletons that occurred only once in the entire data set. The remaining 80 OTUs ranged in abundance from 2 to 680 sequences. A total of 377 AMF sequences were obtained with the NS31/AMmix primers and 28 AMF OTU<sub>0.03</sub> were generated: nine of these OTUs were singletons, while the other 19 OTUs comprised 2–163 sequences. However, the rank-abundance diagrams (Fig. 3) of both the AMV4.5NF/AMDGR and NS31/AMmix sequences indicated that approximately 50% of the total sequences belonged to three OTU<sub>0.03</sub> (OTU numbers 1, 2 and 3 in Table S1) and two OTU<sub>0.03</sub> (OTU numbers 1 and 2 in Table S2) respectively.




Figure 3. Rank-abundance diagrams (plotting the number of intra-OTU<sub>0.03</sub> sequences as a function of the OTU rank) for *Glomeromycota* OTUs<sub>0.03</sub> detected with AMV4.5NF/AMDGR (A) and NS31/AMmix (B) at 3% level of dissimilarity. The diagram trend is analogous for both primer sets: on one hand, the first OTUs<sub>0.03</sub> cover about 50% of all sequences, on the other, a high number of singletons represent the ‘long tail’ of the diagram.

The total number of AMF OTU<sub>0.03</sub> identified with the two primer pair and their taxonomical distribution over the currently recognized *Glomeromycota* orders are shown in Fig. 4. The ratio of the four AMF orders is almost the same, even though the total number of OTU<sub>0.03</sub> generated using the two primer pairs is different: 117 with AMV4.5NF/AMDGR and 28 with NS31-AMmix. As the AMF sequences generated with the NS31/AMmix primers, particularly in the CO and in the TV (12 and 17 sequences respectively), were quite low, the AMF distribution over the five different environments was taken into account considering the sequences generated with the AMV4.5NF/AMDGR primers. Analysing these latter sequences, the number of OTU<sub>0.03</sub> generated for each environment was: 27 in the CO (188 total sequences), 30 in the MM (254 sequences), 43 in the CV (292 sequences), 55 in the TV (429 sequences) and 74 in the PA (1275 sequences) (Fig. 5A and B). Figure 6 showed the proportion of the different *Glomeromycota* orders, which was comparable over the Sardinian soils: the *Glomerales* frequencies were always the highest in terms of both the OTU<sub>0.03</sub> (from 71.6% to 81.5% of the total OTU<sub>0.03</sub>) and sequence numbers (from 81.1% to 95.7% of the total sequences) respectively. In terms of OTU<sub>0.03</sub>, *Diversisporales* instead represented 18.5% of the CO soil community, 21.6% of the PA, 25.5% of the TV, 25.6% of the CV and 26.7% of the MM of the total sequence numbers (Fig. 6A). In terms of sequences, *Diversisporales* ranged from 4.3% (CO) to 18.9% (MM) (Fig. 6B). *Archaeosporales* and *Paraglomerales* were only found in the PA soil and accounted for 5.4% and 0.5%, and 1.4% and 0.2% of the total OTU<sub>0.03</sub> and sequences respectively (Fig. 6A and B).




Figure 4. The AMV4.5NF/AMDGR sequences (A) overall generate 117 OTU<sub>0.03</sub> and the NS31/AMmix sequences (B) generate 28 OTU<sub>0.03</sub>. OTU<sub>0.03</sub> are separated according to their taxonomy: OTU<sub>0.03</sub> belonging to Glomerales represent the majority for both data sets. An OTU<sub>0.03</sub> amenable to Paraglomerales (purple) was found with the AMV4.5NF/AMDGR primer set.






Figure 5. Distribution of different Glomeromycota orders detected with the AMV4.5NF/AMDGR primer set in each of the five Sardinian environments. A. The y-axis indicates the number of OTUs<sub>0.03</sub> assigned to each Glomeromycota order. B. The y-axis indicates the number of sequences assigned to each Glomeromycota order. The total numbers of OTUs<sub>0.03</sub> and sequences found in each environment are shown in brackets. TV, tilled vineyard; CV, covered vineyard; MM, managed meadow; PA, pasture; CO, cork-oak formation.




Figure 6. Proportional distribution of the different Glomeromycota orders detected with the AMV4.5NF/AMDGR primer set in each of the five Sardinian environments. A. The y-axis indicates the proportion of OTUs<sub>0.03</sub> assigned to each Glomeromycota order. B. The y axis indicates the proportion of sequences assigned to each Glomeromycota order. TV, tilled vineyard; CV, covered vineyard; MM, managed meadow; PA, pasture; CO, cork-oak formation.

A comparison of the observed and estimated OTU<sub>S0.03</sub> relative to the five soil types, with the non-parametric ACE and Chao1 indices, is shown in Fig. 7. With both indices, the number of observed OTU<sub>S0.03</sub> is close to the one estimated for the CV. For the other four environments, instead, the number of observed OTU<sub>S0.03</sub> is always lower than the richness estimated by the ACE and Chao1 indices.




Figure 7. Comparison between the AMV4.5NF/AMDGR OTU<sub>S0.03</sub> observed and estimated with the two non-parametric estimator indices ACE (A) and Chao1 (B). The bars represent the 95% confidence intervals for the estimated OTU<sub>S0.03</sub> number. TV, tilled vineyard; CV, covered vineyard; MM, managed meadow; PA, pasture; CO, cork-oak formation.

Shannon–Weaver biodiversity indices ( $H'$ ) were also calculated, taking into account both the number and the relative proportions of taxa in a community, to compare AMF diversity in the different soil types. The  $H'$  indices for the OTUs detected in each environment with the AMV4.5NF/AMDGR primers at 97% sequence similarity, and the corresponding 95% intervals of confidence are as follows:  $1.37 \pm 0.23$  (CO),  $1.55 \pm 0.18$  (MM),  $2.43 \pm 0.14$  (TV),  $2.50 \pm 0.13$  (CV) and  $2.75 \pm 0.08$  (PA).

## Comparison of AMF community memberships and structures

A comparison of the community structures detected using the two couples of primers is shown in Fig. 8. Eleven and one OTU<sub>S0.03</sub>, detected with the AMV4.5NF/AMDGR and NS31/AMmix primers respectively, were shared by all the soils. Among the 11 common OTU<sub>S0.03</sub> (OTU numbers 1, 2, 4, 13, 15, 16, 19, 23, 24, 28 and 38 in Table S1), nine belonged to *Glomerales* and two to *Diversisporales* and they accounted for 1379 out of 2438 sequences representing 56.6% of the total sequences. The only common OTU<sub>S0.03</sub> detected by the NS31/AMmix primers (OTU number 1 in Table S2) belonged to *Glomerales* and accounted for 163 out of 337 sequences (44.1% of the total sequences). This OTU<sub>S0.03</sub> was also the most abundant and commonly identified OTU<sub>S0.03</sub> when the other primers were used (OTU number 1 in Table S1), on the basis of a pairwise comparison between some sequences of each of these two OTU<sub>S0.03</sub> (data not shown). These two OTU<sub>S0.03</sub> (OTU number 1 in Table S1 and in Table S2) most likely correspond to a *Glomus intraradices*-related species.




Figure 8. Venn diagrams comparing the OTUs<sub>0.03</sub> memberships detected with the AMV4.5NF/AMDGR (A) and NS31/AMmix (B) primers in the five land-use soil types. The relative total percentage of OTUs<sub>0.03</sub> is reported in brackets. The size of each sample component is based on the total number of OTUs<sub>0.03</sub> for that sample relative to the other samples. TV, tilled vineyard; CV, covered vineyard; MM, managed meadow; PA, pasture; CO, cork-oak formation.

Pairwise Jaccard similarity indices were calculated to evaluate the overlap among the AMF assemblages detected in the five environments with the AMV4.5NF/AMDGR primer pair (Table 3). The TV/PA and TV/CV populations exhibited the highest Jaccard similarity indices: 0.39 and 0.38 respectively. The lowest similarities were found between the cork-oak AMF community and the other four soil populations, with indices ranging from 0.21 (CO/TV) to 0.27 (CO/MM).

Table 3. Jaccard similarity coefficient of AMF communities detected with the AMV4.5NF/AMDGR couple of primers in the five Sardinian soils.

	JACCARD similarity index				
	Tilled vineyard	Covered vineyard	Managed meadow	Pasture	Cork-oak formation
1. This index ranges from 0, complete dissimilarity, to 1, complete similarity between two communities. The values reported in the table refer to AMF communities detected at 97% of similarity (OTU <sub>S0.03</sub> ).					
Cork-oak formation	0.24240	0.20690	0.26670	0.21690	1
Pasture	0.38710	0.27170	0.30000	1	
Managed meadow	0.34920	0.37740	1		
Covered vineyard	0.38030	1			
Tilled vineyard	1				

## Discussion

This pyrosequencing study has allowed us to shed light on the diversity of the AMF that thrive in a Mediterranean ecosystem, where different land-use and plant cover types lead to the creation of a soil and vegetation gradient: from an old-growth cork-oak formation to two vineyards (one tilled and one covered with natural plant species), passing through a pasture and a managed meadow. This landscape can be considered of ecological interest as it offers a dynamic mosaic of diverse habitats.

The 454 GS-FLX platform has demonstrated the potentiality of a high-throughput technology to investigate AMF communities and has revealed that 117 *Glomeromycota* OTUs were present in the soils, that *Glomerales* were the dominant order, and – at a lower taxon level – that the *Glomus intraradices* related species were the overwhelming majority, irrespective of the environment.

## Methodological considerations

The first critical point of a metagenomic study, in particular in a complex environment such as soil, concerns the sampling, which should ideally cover the entire biodiversity. In the present work, in order to be sure that representative AMF communities were sampled, we decided to pool the polymerase chain reaction (PCR) products that were independently amplified from the five soil samples from each location (TV, CV, MM, PA and CO). Although this approach does not consider replicates, it offers a reliable overview of the AMF community present in each location ([Renker et al., 2006](#)).

Another important aspect that influences the reliability of the results is the technique that is used to analyse the biodiversity. From this point of view, high-throughput technologies, such as pyrosequencing, represents a powerful instrument which is now available in metagenomics. The GS-FLX System potentially produces a huge number of sequences, but due to the design of the experiment and the numerous short reads, only 4192, out of the about 10 000 sequences that were

obtained, were maintained and analysed. Although the number of obtained sequences is lower than that declared for 454 pyrosequencing, it far exceeds the value of fungal sequences obtained until now in fungal biodiversity studies in soil ([O'Brien et al., 2005](#); [Hempel et al., 2007](#)). The relatively low number of produced sequences is due to the availability of only five lanes out of 16 in the pyrosequencing plate and to the contemporary sequencing of two amplicon mixtures of different length. In particular, the presence of long (> 300 bp) and short fragments led to a general decrease in the obtainable sequence number, because the long fragments (NS31/AMmix) inhibit the emulsion-PCR step ([Margulies et al., 2005](#)) at the expense of short amplicons (AMV4.5NF/AMDGR). The primer pair choice affects the sequence number to a great extent: the AMV4.5NF/AMDGR couple resulted to be the best, in terms of total sequence number, percentage of *Glomeromycota* sequences and spectrum of *Glomeromycota* detected, for AMF soil community analysis. The remarkable difference between sequences obtained with each of the two primer sets is due to the length of the amplified fragments. However, the NS31/AMmix primer set was chosen because a high number of its targeted sequences are deposited in GenBank, even though this primer set did not exactly fit the characteristics required for the GS-FLX pyrosequencing platform. Moreover, a mix of primers AM (AM1, AM2 and AM3) in conjunction with NS31 ([Santos-Gonzalez et al., 2007](#); [Toljander et al., 2008](#)) was used in order to target a wider range of AMF taxonomical groups. The obtained results confirmed that this primer set can also amplify non-target organisms ([Douhan et al., 2005](#); [Alguacil et al., 2008](#)), especially in a complex environment such as soil, where they have been used for the first time.

A primer pair designed on the SSU region, AML1–AML2 ([Lee et al., 2008](#)), which guarantees a good coverage of *Glomeromycota* taxa, is actually available, but it amplifies a fragment of 795 bp, therefore exceeding the suitable length of both the GS-FLX Standard and the New Titanium Series Reagents. Other regions currently used to infer AMF phylogeny were discarded for this metagenomic study because they show some negative aspects. For example, internal transcribed spacer (ITS) and large subunit (LSU) regions, can offer a resolution at a lower taxonomic level (species) than the SSU region. However, it was decided not to use ITS and LSU because of some critical aspects. The ITS region is too variable to be confidentially aligned using 250-bp-long sequences, which is the average length obtainable with the GS-FLX standard kit. For the LSU region the major concern was about the lower number of sequences deposited in GenBank in comparison with the SSU sequences. In conclusion, it seems that a completely satisfactory AMF primer couple is not yet available and that the use of multiple sets is – at the moment – a good strategy to overcome problems.

The limited length (200–300 bp) of fragments that can be obtained with the 454 GS-FLX platform allowed us to identify the AMF at the highest taxonomic levels: we therefore revealed differences among the AMF communities of the five soil types rather than identifying the AMF species. Another difficulty in AMF biodiversity studies at a species level concerns the fact that the majority of AMF sequences deposited in databases are labelled as ‘unknown’ or ‘uncultured’ *Glomeromycota*, because they are obtained from molecular analysis of plant roots, without a corresponding morphospecies description. Such a high proportion of ‘known as-sequence-only’ taxa reflects the accumulation of molecular diversity data of AM fungi as a result of the increasing number of studies of *Glomeromycota* in natural ecosystems ([Öpik et al., 2008](#)). However, more matches with known AM fungal species could be expected when more effort is directed towards sequencing DNA from morphologically characterized AMF isolates. In the last years, several groups have dedicated a great deal of effort to describing new taxa from spore morphology coupled with molecular analyses ([Blaszkowski et al., 2008](#); [Stockinger et al., 2009](#)). In addition, sequencing using new strategies, which lead to longer reads (GS-FLX System with New Titanium Series Reagents), together with new primer availability ([Krüger et al., 2009](#)) will allow studies to be conducted at a species level.

As in many other fungal biodiversity studies ([Hunt et al., 2004](#); [Santos-Gonzalez et al., 2007](#)), we had to make a critical decision on an appropriate sequences divergence cut-off to define the OTUs. The sequence-based definition of species as clusters of sequences differing by at most 3% of sites was adopted for both the fungi and bacteria. While such a cut-off is generally regarded as canonical for bacterial 16S sequences ([Hanage et al., 2006](#); [Konstantinidis and Tiedje, 2007](#); [Elshahed et al., 2008](#)), a similar agreement does not exist for fungal rDNA. In particular, the *Glomeromycota* rRNA gene can show different variants within the same species and even within the same spore ([Lanfranco et al., 1999](#); [Sanders, 2004](#); [Börstler et al., 2008](#)). In our study, blast results were found to be consistent for several 97% sequence identity AMF OTUs and we therefore decided to use 97% sequence similarity as the cut-off level. In previous studies, the same sequence identity level was considered a reliable threshold to discriminate possible AMF species ([Santos-Gonzalez et al., 2007](#)).

## **Mediterranean soils harbour diverse AMF assemblages dominated by *Glomerales***

The ecological theory predicts that heterogeneous landscapes should involve higher species diversity levels than homogenous areas since structurally complex habitats provide more niches and ways of exploiting resources ([Wardle et al., 2004](#); [Lekberg et al., 2007](#)). Habitat heterogeneity is especially high in typical Mediterranean landscapes ([da Silva et al., 2009](#)). Starting from natural ecosystems dominated by cork-oak woodlands, a mix of silviculture and traditional grazing practices has led to a dynamic agro-forest mosaic of different habitats, which might harbour a high below-ground diversity of AMF and consequently shape plant biodiversity and ecosystem functioning ([van der Heijden et al., 1998](#); [Maherali and Klironomos, 2007](#); [van der Heijden et al., 2008](#)). Given the increasing interest in soil metagenomes and because, so far, most studies, with a single exception ([Hempel et al., 2007](#)), have been based solely on AM spores separated from soil samples and/or mycorrhizal roots, we have explored the potentiality of a 454 GS-FLX platform for a high-throughput study on AMF biodiversity starting from DNA directly extracted from bulk soil, which contains a large proportion of DNA from AM extra-radical mycelium and spores ([Johnson et al., 2003](#); [Gryndler et al., 2006](#)). The possibility of studying AMF biodiversity with this new technique is particularly interesting, from an ecological point of view, if we consider that the Sardinian environment is characterized by a land-use gradient, which ranges from old-growth cork-oak woodland to intermediate or high land-use intensification levels, including pasture, vineyards and managed meadows.

The AMF communities of the investigated soils were found to be characterized by a dominance of *Glomerales*. Many studies have reported *Glomus* spp. as the most widespread species in many ecologically different environments: from natural woodlands to high input managed agro-ecosystems ([Helgason et al., 2002](#); [Oehl et al., 2005](#); [Hijri et al., 2006](#)). *Glomerales* was the most abundant taxon in terms of OTU<sub>0.03</sub> and sequence number: rank-abundance diagrams have indicated that only a few taxa accounted for 50% of the total sequences recovered, while many other OTU<sub>0.03</sub> were represented by only a few, or even a single sequence. This suggested a striking dominance of some taxa over many rare taxa ('tails' in rank-abundance diagrams). Although some of these rare taxa do possibly really occur, others are likely to be artefacts due to the intrinsic error rate of pyrosequencing, which could lead to an overestimation of the biodiversity, as stated by some recent articles related to 454 pyrosequencing biases. Indeed it has been shown that the presence of long homopolymers in the sequenced fragments may result in frequent miscalls: either insertion or deletions ([Quinlan et al., 2008](#); [Kunin et al., 2009](#)). However, interest in developing new algorithms able to manage large data set and eliminating sequencing mistakes is growing in order to overcome this problem ([Quince et al., 2009](#)).

The most commonly encountered fungus (OTU number 1 for both primer pairs) possibly corresponds to a *Glomus* species, which is presumably related to the *Glomus intraradices* group (*Glomus* group Ab). *Glomus intraradices* contains several cryptic taxa with differences in various ecological

properties ([Croll et al., 2009](#)): this is in agreement with the observation that the AMV4.5NF/AMDGR primer pair amplified sequences that could be amenable to *G. intraradices*, but were grouped in different OTUs. Many isolates of this species have been detected in different locations throughout the world, of both stable and disturbed ecosystems ([Öpik et al., 2006](#); [Appoloni et al., 2008](#)) and in many host species ([Helgason et al., 2007](#)), suggesting that this AM fungal species has a generalist and ruderal lifestyle (disturbance tolerance) as it produces large numbers of spores and extra-radical mycelium ([Jansa et al., 2003](#); [Öpik et al., 2006](#)). Our analysis on soil DNA is in agreement with such behaviour and points out the dominance of this species, not only in plant roots, but also in the soil environment.

The presence, in all five examined soils, of *Diversisporales* OTU<sub>S0.03</sub>, which is thought to produce substantial amounts of external mycelium and dense hyphal clusters ([Hart and Reader, 2002](#)) and to prefer sandy soils ([Duponnois et al., 2001](#); [Lekberg et al., 2007](#)), is also consistent with both the ecology of this fungal group and the features of the Berchidda area. *Paraglomerales* and *Archaeosporales* were instead only found in the PA soil. Some molecular studies on AM communities ([Hijri et al., 2006](#); [Vallino et al., 2006](#); [Alguacil et al., 2008](#)) have reported the difficulty of detecting *Paraglomerales* and *Archaeosporales* in root samples, while [Hempel and colleagues \(2007\)](#) observed that they could be dominant in certain soils. In the current study, only a few OTU<sub>S0.03</sub> and sequences amenable to these taxa were found and these were only found in the PA soil. This finding would seem to suggest either a scanty occurrence of these orders in the soils or a likely influence of some natural traits, such as those generally associated with grasslands, on the spread of these fungi in ecosystems ([Hempel et al., 2007](#)). However, another possible reason for the exclusive detection of *Paraglomerales* and *Archaeosporales* in the PA soil is the high number of sequences obtained from this environment. The high value of the Shannon–Weaver biodiversity index (*H'*), together with the assemblages found in the PA soil, along with the high number of sequences obtained would suggest a higher biodiversity of *Glomeromycota* in the PA, compared with the other four systems. The high AMF diversity in this environment is in agreement with a trend that has already been demonstrated in many studies ([Öpik et al., 2006](#); [Öpik et al., 2008](#)): soils subjected to human input (TV and MM) have a lower AMF richness than low-input soils (CV and PA).

As already observed in many bacterial metagenomic studies on soil ([Roesch et al., 2007](#); [Fulthorpe et al., 2008](#)), and despite the high number of sequences obtained, in particular with the AMV4.5NF/AMDGR primers, the actual extent of total AMF diversity possibly remains largely undiscovered. This is true for four out of the five environments investigated, while for the CV, the number of observed OTU<sub>S0.03</sub> is close to the estimated number, indicating good sampling efficiency.

## **Do AMF communities mirror the vegetation community?**

The description of the AMF communities in an area characterized by different land-uses can also allow some considerations to be made on the overlap of the *Glomeromycota* assemblages. Eleven out of the 117 and one out of the 28 OTU<sub>S0.03</sub> obtained with the two primer pairs were found to be common to all the soils. On the basis of a pairwise comparison between some sequences of each of these two OTU<sub>S0.03</sub> (data not shown), the only common OTU<sub>0.03</sub> revealed with the NS31/AMmix probably corresponds to OTU<sub>0.03</sub> number 1 obtained with AMV4.5NF/AMDGR. These OTU<sub>S0.03</sub> are likely related to the *G. intraradices* group (*Glomus* group Ab) and they accounted for the highest number of sequences (163 and 680 sequences respectively). This finding confirms the generalistic and likely dominant nature of this group not only in plant roots but also in soil. Despite this overlap, each of the five soil AMF communities exhibited particular features, as indicated by both the occurrence of OTU<sub>S0.03</sub> unique to each soil type, and the low Jaccard similarity indices obtained in the pairwise comparison.

Of the five AMF communities studied, the CO deserves particular mention. Although the CO is a natural ecosystem, its Shannon–Weaver index was the lowest ( $H' = 1.37$ ) and the number of AMF OTUs<sub>0.03</sub> was relatively low, probably because the fungal communities in this environment had shifted from AMF to other mycorrhizal fungal species associated with shrub (*Erica arborea* L and *Arbutus unedo* L) and tree species (*Quercus suber* L). This hypothesis is supported by the high number of sequences belonging to *Sebacinales* (data not shown), a *Basidiomycota* order establishing ericoid mycorrhiza (Selosse *et al.*, 2007), which were obtained with the AMV4.5NF/AMDGR primer pair. The particular composition of the CO soil AMF community was also indicated by the low Jaccard indices obtained when comparing this community with the other four soil AMF communities. This suggests that the plant coverage of this ecosystem, which is characterized by trees and shrubs, has determined not only a reduction in AMF presence but also a typical, unique community structure.

## Conclusion

A metagenomic approach applied to AMF fungi and based on a gene of taxonomic interest, like the ribosomal gene, has generated a high number of AMF sequences that far exceeded the number of AMF fungal sequences so far obtained in soil. The obtained data have allowed us to precisely describe the AMF communities that proliferate in a Mediterranean environment, where different land-use and plant cover types lead to the creation of a soil and vegetation gradient. With the exception of a generalist fungus, *Glomus intraradices*, only a few other AMF taxa were found in common in the five environments, suggesting fungal assemblage specificity. The environments with a low input (PA and CV) showed a higher AMF biodiversity than those subjected to human input (MM and TV). A reduction in AMF, replaced by other mycorrhizal fungal species more likely associated to trees and shrubs, was found in the CO. These findings suggest that AMF biodiversity is influenced by both human input and ecological traits. This study poses many questions on the functionality of the AMF communities described, first of all whether the most frequently represented taxa are also the most functionally active. Metagenomic approaches based on soil RNA could help explain not only what species are present, but also what benefits are given to their host plants.

## Experimental procedures

### Study site

The study area is located in the northern hills of Sardinia, Italy. The Berchidda site (Olbia-Tempio) (40°30'13.37"N 9°47'00.56"E) is made up of hydromorphic and granitic soil with a loamy sand texture. The altitude ranges from 275 m to 300 m. This area is referred to as a mesomediterranean, subhumid phytoclimatic belt with annual rainfall averages of 862 mm (5% summer rainfall percentage), while the mean temperature is 13.8°C (Bacchetta *et al.*, 2004). The soil has pH values that range from 5.0 to 6.5.

In the past, the Berchidda area was covered by cork-oak forests which were subjected to intense usage for the extraction of cork and pasture. Today, there are five dominant soil use types: TV, CV, MM, PA (dominated by grass species with a low tree density) and CO (dominated by shrub cover and distributed cork-oak trees) (Fig. S1).

### Soil sampling, DNA extraction, PCR and preparation of the amplicon libraries

In May 2007, five soil core samples (5 cm Ø and 20 cm depth) were taken from each of the five locations. The 25 soil samples were independently packed in ice upon collection and transported to the labs for DNA extraction. The soil samples were sieved (2 mm) to remove fine roots and large



organic debris and stored at  $-80^{\circ}\text{C}$ . Twenty-five soil DNA extractions were performed from at least 0.5 g of mixed soil, using the FastDNA Kit (MP Biomedicals, LLC, Fountain Pkwy, Solon, OH, USA).

Two couples of primers: AMV4.5NF/AMDGR ([Sato et al., 2005](#)) and NS31 with a mixture of equal amounts of the reverse AM1, AM2 and AM3 (AMmix) ([Santos-Gonzalez et al., 2007](#)), were used to amplify an 18S rRNA gene fragment for the 454 GS-FLX pyrosequencing platform. A total of 50 independent PCR amplifications were performed: 25 with AMV4.5NF/AMDGR and 25 with the NS31/AMmix primer set. The DNA amplifications were performed from an equivalent amount of DNA to that found in 2.5 g of each soil type. In order to make sure that the representative AMF communities were sampled, we thought that pooling the PCR products independently amplified from the five soil samples, obtained from the same location (TV, CV, MM, PA and CO), could be the best strategy to overcome the lack of replicates due to the reduced number of available lanes on the pyrosequencing plate (5 out of 16).

The AMV4.5NF/AMDGR primer pair was chosen because it showed suitable characteristics for the GS-FLX System: it amplified a target sequence of approximately 300 bp including the SSU rDNA variable domain V4 from a broad spectrum of *Glomeromycota* (*Glomerales*, *Diversisporales* and *Archaeosporales*) ([Sato et al., 2005](#)). Although the NS31/AMmix primers had not been previously used on soil DNA, they were selected because they are frequently used in AMF biodiversity studies: many targeted sequences are therefore easily found in the databases. The general fungal primer AM1 was designed to amplify fungal DNA ([Helgason et al., 1998](#)). This primer, in conjunction with the universal eukaryotic primer NS31 ([Simon et al., 1992](#)), has been found to amplify AM fungal DNA sequences from field-collected roots ([Helgason et al., 1998](#); [Vallino et al., 2006](#); [Alguacil et al., 2008](#)), although it yielded some mismatches at the priming site with taxa belonging to *Glomus* group B and *Glomus* group C (*Diversisporaceae*). For this reason, the exclusive use of AM1 is limiting in a biodiversity study. Therefore, AM2 and AM3 primers were added: these are modifications of the AM1 primer and are designed to amplify DNA from taxa not detected by AM1 (*Glomus* group B and *Glomus* group C) ([Santos-Gonzalez et al., 2007](#); [Toljander et al., 2008](#)).

In order to perform 454 pyrosequencing with the GS-FLX System, the sequences of these oligonucleotides included the 454 Life Science A or B sequencing adapters (19 bp) fused to the 5' primer ends (in brackets):

- A-NS31 5'-(GCCTCCCTCGCGCCATCAG)TTGGAGGGCAAGTCTGGTGCC-3'
- B-AM1 5'-(GCCTTGCCAGCCCGCTCAG)GTTTCCCGTAAGGCGCCGAA-3'
- B-AM2 5'-(GCCTTGCCAGCCCGCTCAG)GTTTCCCGTAAGGTGCCAAA-3'
- B-AM3 5'-(GCCTTGCCAGCCCGCTCAG)GTTTCCCGTAAGGTGCCGAA-3'
- A-AMV4.5NF 5'-(GCCTCCCTCGCGCCATCAG)AAGCTCGTAGTTGAATTTTCG-3'
- B-AMDGR 5'-(GCCTTGCCAGCCCGCTCAG)CCCAACTATCCCTATTAATCAT-3'

The PCRs contained 17.1  $\mu\text{l}$  of sterile water, 2.5  $\mu\text{l}$  10 $\times$  of reaction buffer (Sigma), 2.5  $\mu\text{l}$  of each deoxyribonucleotide triphosphate (dNTP 2.0  $\mu\text{M}$ ), 0.5  $\mu\text{l}$  of each primer (10  $\mu\text{M}$ ), 0.4  $\mu\text{l}$  of DNA polymerase (High Fidelity Taq, Roche) and 2  $\mu\text{l}$  of DNA template in a final volume of 25  $\mu\text{l}$ .

The DNA was amplified using a T3000 thermal cycler (Biometra, Göttingen, DE). The following programme was used for DNA amplification: initial denaturation at  $94^{\circ}\text{C}$  for 3 min, followed by 35 cycles of denaturation at  $94^{\circ}\text{C}$  for 45 s, annealing at  $60^{\circ}\text{C}$  for 45 s, extension at  $72^{\circ}\text{C}$  for 1 min and a final extension at  $72^{\circ}\text{C}$  for 7 min with a ramp of  $3^{\circ}\text{C s}^{-1}$ .

Twenty-five independent amplifications (five for each soil type) were conducted for the AMV4.5NF/AMDGR and NS31/AMmix couples respectively. The PCR products obtained with the two primer pairs were purified with the Agencourt® AMPure® Kit (Beckman Coulter, CA, USA) and pooled to generate 10 samples (five for AMV4.5NF/AMDGR and five for NS31/AMmix). The quality of these samples was assessed through: (i) gel electrophoresis of 5 µl subsamples on 1.5% agarose gel; (ii) evaluation of the AD260/280 ratio calculated using the ND-1000 Spectrophotometer NanoDrop® (Thermo Scientific, Wilmington, DE); and (iii) analysis with the Experion™ System (Bio-Rad, Hercules, CA, USA), using a DNA1K Chip.

In order to create equimolar mixtures of multiple amplicons (amplicon libraries) for 454 pyrosequencing, the 10 pooled samples were quantified by the ND-1000 Spectrophotometer NanoDrop® and five final amplicon libraries (TV, CV, MM, PA and CO), containing 10<sup>10</sup> molecules/µl of each primer set amplification, were generated. The samples were stored at -20°C and sent to BMR Genomics s.r.l. (Padua, IT) for pyrosequencing by means of a Genome Sequencer FLX System platform (454 Life Science Branford, CT, USA). The samples were processed together with other soils (C. Murat, V. Bianciotto, S. Daghino, M. Girlanda, A. Lazzari, E. Lumini *et al.*, unpublished) and they occupied five lines out of the 16 available in the GS-FLX System.

### **Alignment, clustering and statistical analyses of the OTU richness**

All the sequences were analysed after trimming off the adapter sequences. According to [Wommack and colleagues \(2008\)](#) and after a preliminary blast analysis where small sequences (50–199 bp) did not show any significant similarity, the AMV4.5NF/AMDGR sequences shorter than 230 bp and those shorter than 250 bp were eliminated for the NS31/AMmix primers.

Sequences from the five land use-units were aligned using MUSCLE 3.6 ([Edgar, 2004](#)) with default parameters. The alignments were manually edited and distance matrices were constructed using DNAdist from the phylip suite of programmes, version 3.6, with default parameters ([Felsenstein, 2005](#)). These pairwise distances were used as input for DOTUR ([Schloss and Handelsman, 2005](#)) in order to cluster the sequences into OTUs of a defined sequence identity. The OTUs were defined according to their different sequence similarity values, which spanned 80%–97% sequence identity. Although these distance cut-offs were arbitrary and can be considered controversial, a 97% sequence similarity level, corresponding to 0.03 (OTU<sub>0.03</sub>), was chosen in this study, according to the conventional definition of a microbial ‘species’ ([Rosselló-Mora and Amann, 2001](#); [Konstantinidis and Tiedje, 2007](#)). Blast searches were carried out, to test for within-OTU consistency, with sequences within the first 10 most abundant AMV4.5NF/AMDGR and NS31/AMmix OTUs at 97% sequence similarity: these searches yielded the same best blast hit for all the sequences within each OTU. Consensus sequences, obtained for each OTU<sub>0.03</sub> using CAP3 ([Huang and Madan, 1999](#)), were used as queries for the blast searches in the GenBank database. A conservative approach was followed for the fungal species identification, considering only identifications with a ≥ 200 blast score value reliable and labelling all the others as ‘unknown organisms’. We used OTU<sub>0.03</sub> from DOTUR rather than taxonomic assignments based on blast analyses because not all the sequences matched a known sequence in the database and the use of sequence similarities prevented uncertainties associated with fungal taxonomy and classification.

In order to analyse the richness and diversity of the AM fungal communities in the Sardinian soils, only sequences belonging to *Glomeromycota* were selected for the subsequent analyses and the others were discarded. Arbuscular mycorrhizal fungi sequences were used to generate new distance matrices through DNAdist and these were then used as DOTUR input to generate AMF OTUs containing sequences of different genetic distance values based on a furthest-neighbour algorithm. The use of DOTUR also allowed us to obtain a variety of diversity richness estimators (rarefaction curve, bias

corrected Chao1 richness and abundance-based coverage estimator ACE). Consensus sequences of each AMF OTU<sub>0.03</sub> were queried against the GenBank using blast. The blast results were used to calculate the relative abundance of each *Glomeromycota* order within each land-use unit. Similarity among the AMF communities of the five soil-use types was determined using SONS, a software which uses the OTU data obtained from DOTUR to estimate the overlap between pairs of communities and which calculates the Jaccard similarity index (determined as the ratio of the number of OTUs shared and the total number of OTUs in both samples). This index, which ranges from 0 (complete dissimilarity) to 1 (complete similarity), is the simplest parameter to measure the ratio of shared species without considering abundance data.

The Shannon–Weaver diversity index, the rarefaction index and the non-parametric ACE and Chao1 indices were also calculated for each land-use type in order to compare the observed and estimated AMF richness and to assess sampling efficiency in the five soils.

### **Nucleotide sequence accession numbers**

The sequences analysed in this study have been deposited in the EMBL database under accession numbers [FN386789](#)–390831.

### **Acknowledgements**

Special thanks are due to M. Girlanda for her careful manuscript reading and her always valuable suggestions. We also thank R. Pastorelli for her help with the soil collection, P. Schloss for his encouraging and enthusiastic emails, S. Ghignone and C. Murat for their bioinformatic consulting and M. Novero for her help in the image editing. The authors would like to thank R. Francaviglia, P.P. Roggero and his team for their contribution to collect the Sardinian soils in the frame of the Italian National Project SOILSINK (FISR). The P.B. research was funded by the SOILSINK Project (FISR) and Compagnia di San Paolo. V.B. research was funded by IPP-CNR (AG-P02-006, Biodiversity Project–Torino) and ENDURE project (European Network of Excellence). E.L and R.B. were supported by SOILSINK Project grants. We also thank the two anonymous reviewers for their helpful comments.

## References

- Alguacil, M.M., Lumini, E., Roldán, A., Salinas-Garcia, J.R., Bonfante, P., and Bianciotto, V. (2008) The impact of tillage practices on arbuscular mycorrhizal fungal diversity in subtropical crops. *Ecol Appl* 18: 527–536.
- Appoloni, S., Lekberg, Y., Tercek, M.T., Zabinski, C.A., and Redecker, D. (2008) Molecular community analysis of arbuscular mycorrhizal fungi in roots of geothermal soils in Yellowstone National Park (USA). *Microb Ecol* 56: 649–559.
- Artursson, V., Finlay, R.D., and Jansson, J.K. (2006) Interactions between arbuscular mycorrhizal fungi and bacteria and their potential for stimulating plant growth. *Environ Microbiol* 8: 1–10.
- Bacchetta, G., Bagella, S., Biondi, E., Farris, E., Filigheddu, R., and Mossa, L. (2004) A contribution to the knowledge of the order *Quercetalia ilicis* Br.-Bl. ex Molinier 1934 of Sardinia. *Fitosociologia* 41: 29–51.
- Blaszkowski, J., Czerniawska, B., Wubet, T., Schäfer, T., Buscot, F., and Renker, C. (2008) *Glomus irregulare*, a new arbuscular mycorrhizal fungus in the Glomeromycota. *Mycotaxon* 106: 247–267.
- Börstler, B., Raab, P.A., Thiéry, O., Morton, J.B., and Redecker, D. (2008) Genetic diversity of the arbuscular mycorrhizal fungus *Glomus intraradices* as determined by mitochondrial large subunit rRNA gene sequences is considerably higher than previously expected. *New Phytol* 180: 452–465.
- Burrows, R.L., and Pflieger, F.L. (2002) Arbuscular mycorrhizal fungi respond to increasing plant diversity. *Can J Bot* 80: 120–130.
- Croll, D., Giovannetti, M., Koch, A.M., Sbrana, C., Ehinger, M., Lammers, P.J., and Sanders, I.R. (2009) Nonspecific vegetative fusion and genetic exchange in the arbuscular mycorrhizal fungus *Glomus intraradices*. *New Phytol* 181: 924–937.
- Douhan, G.W., Petersen, C., Bledsoe, C.S., and Rizzo, D.M. (2005) Contrasting root associated fungi of three common oak-woodland plant species based on molecular identification: host specificity or non-specific amplification? *Mycorrhiza* 15: 365–372.
- Duponnois, R., Plenchette, C., Thioulouse, J., and Cadet, P. (2001) The mycorrhizal soil infectivity and arbuscular mycorrhizal fungal spore communities in soils of different aged fallows in Senegal. *Appl Soil Ecol* 17: 239–251.
- Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32: 1792–1797.
- Edwards, R.A., Rodriguez-Brito, B., Wegley, L., Haynes, M., Breitbart, M., Peterson, D.M., et al. (2006) Using pyrosequencing to shed light on deep mine microbial ecology. *BMC Genomics* 7: 57.
- Van Elsas, J.D., Costa, R., Jansson, J., Sjöling, S., Bailey, M., Nalin, R., et al. (2008) The metagenomics of disease-suppressive soils – experiences from the METACONTROL project. *Trends Biotechnol* 26: 591–601.
- Elshahed, M.S., Youssef, N.H., Spain, A.M., Sheik, C., Najar, F.Z., Sukharnikov, L.O., et al. (2008) Novelty, uniqueness patterns of rare members of the soil biosphere. *Appl Environ Microbiol* 74: 5422–5428.
- Felsenstein, J. (2005) *phylip* (Phylogeny Inference Package), Version 3.6. Distributed by the Author. Seattle, WA, USA: Department of Genome Sciences, University of Washington.
- Fulthorpe, R.R., Roesch, L.F.W., Riva, A., and Triplett, E.W. (2008) Distantly sampled soils carry few species in common. *ISME J* 2: 901–910.
- Girlanda, M., Perotto, S., and Bonfante, P. (2007) Mycorrhizal Fungi: their habitats and nutritional strategies. In *The Mycota IV Environmental and Microbial Relationships*, 2nd edn. Berlin-Heidelberg, Germany: Springer-Verlag, pp. 229–256.

- Gryndler, M., Larsen, J., Hrselova, H., Rezacova, V., Gryndlerova, H., and Kubat, J. (2006) Organic and mineral fertilization, respectively, increase and decrease the development of external mycelium of arbuscular mycorrhizal fungi in a long-term field experiment. *Mycorrhiza* 16: 159–166.
- Hanage, W.P., Fraser, C., and Spratt, B.G. (2006) Sequences, sequence clusters and bacterial species. *Philos T R Soc B* 361: 1917–1927.
- Hart, M.M., and Reader, R.J. (2002) Host plant benefit from association with arbuscular mycorrhizal fungi: variation due to differences in size of mycelium. *Biol Fert Soils* 36: 357–366.
- Van Der Heijden, M.G.A., and Scheublin, T.R. (2007) Functional traits in mycorrhizal ecology: their use for predicting the impact of arbuscular mycorrhizal fungal communities on plant growth and ecosystem functioning. *New Phytol* 174: 244–250.
- Van Der Heijden, M.G.A., Klironomos, J.N., Ursic, M., Moutoglis, P., Streitwolf-Engel, R., Boller, T., et al. (1998) Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* 396: 69–72.
- Van Der Heijden, M.G.A., Bardgett, R.D., and Van Straalen, N.M. (2008) The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol Lett* 11: 296–310.
- Helgason, T., Daniell, T.J., Husband, R., Fitter, A.H., and Young, J.P.W. (1998) Ploughing up the wood-wide web? *Nature* 394: 431.
- Helgason, T., Merryweather, J.W., Denison, J., Wilson, P., Young, J.P.W., and Fitter, A.H. (2002) Selectivity and functional diversity in arbuscular mycorrhizas of co-occurring fungi and plants from a temperate deciduous woodland. *J Ecol* 90: 371–384.
- Helgason, T., Merryweather, J.W., Young, J.P.W., and Fitter, A.H. (2007) Specificity and resilience in the arbuscular mycorrhizal fungi of a natural woodland community. *J Ecol* 94: 623–630.
- Hempel, S., Renker, C., and Buscot, F. (2007) Differences in the species composition of arbuscular mycorrhizal fungi in spore, root and soil communities in a grassland ecosystem. *Environ Microbiol* 9: 1930–1938.
- Hijri, I., Sykorova, Z., Oehl, F., Ineichen, K., Mäder, P., Wiemken, A., and Redecker, D. (2006) Communities of arbuscular mycorrhizal fungi in arable soils are not necessarily low in diversity. *Mol Ecol* 15: 2277–2289.
- Horton, T.R., and Bruns, T.D. (2001) The molecular revolution in ectomycorrhizal ecology: peeking into the black-box. *Mol Ecol* 10: 1855–1871.
- Huang, X., and Madan, A. (1999) CAP3: a DNA sequence assembly program. *Genome Res* 9: 868–877.
- Huber, J.A., Welch, D.B., Morrison, H.G., Huse, S.M., Neal, P.R., Butterfield, D.A., and Sogin, M.L. (2007) Microbial population structures in the deep marine biosphere. *Science* 318: 97–100.
- Hugenholtz, P., and Tyson, G.W. (2008) Microbiology: metagenomics. *Nature* 455: 481–483.
- Hunt, J., Boddy, L., Randerson, P.F., and Rogers, H.J. (2004) An evaluation of 18S rDNA approaches for the study of fungal diversity in grassland soils. *Microb Ecol* 47: 385–395.
- Jansa, J., Mozafar, A., Kuhn, G., Anken, T., Ruh, R., Sanders, I.R., and Frossard, E. (2003) Soil tillage affects the community structure of mycorrhizal fungi in maize roots. *Ecol Appl* 13: 1164–1176.
- Johnson, D., Vandenkoornhuyse, P., Leake, J.R., Gilbert, L., Booth, R.E., Grime, J.P., et al. (2003) Plant communities affect arbuscular mycorrhizal fungal diversity and community composition in grassland microcosm. *New Phytol* 161: 503–515.
- Klironomos, J.N., and Hart, M.M. (2002) Colonization of roots by arbuscular mycorrhizal fungi using different sources of inoculum. *Mycorrhiza* 12: 181–184.
- Konstantinidis, K.T., and Tiedje, J.M. (2007) Prokaryotic taxonomy and phylogeny in the genomic era: advancements and challenges ahead. *Curr Opin Microbiol* 10: 504–509.

- Krüger, M., Stockinger, H., Krüger, C., and Schüßler, A. (2009) DNA-based species-level detection of Glomeromycota: one PCR primer set for all arbuscular mycorrhizal fungi. *New Phytol* 183: 212–223.
- Kunin, V., Engelbrektsen, A., Ochman, H., and Hugenholtz, P. (2009) Wrinkles in the rare biosphere: pyrosequencing errors lead to artificial inflation of diversity estimates. *Environ Microbiol* (in press): doi: 10.1111/j.1462-2920.2009.02051.x.
- Lanfranco, L., Delpero, M., and Bonfante, P. (1999) Intrasporal variability of ribosomal sequences in the endomycorrhizal fungus *Gigaspora margarita*. *Mol Ecol* 8: 37–45.
- Lee, J., Lee, S., and Young, J.P.W. (2008) Improved PCR primers for the detection and identification of arbuscular mycorrhizal fungi. *FEMS Microbiol Ecol* 65: 339–349.
- Lekberg, Y., Koide, R.T., Rohr, J.R., Aldrich-Wolfe, L., and Morton, J.B. (2007) Role of niche restrictions and dispersal in the composition of arbuscular mycorrhizal fungal communities. *J Ecol* 95: 95–105.
- Maherali, H., and Klironomos, J.N. (2007) Influence of phylogeny on fungal community assembly and ecosystem functioning. *Science* 316: 1746–1748.
- Margulies, M., Egholm, M., Altman, W.E., Attiya, S., Bader, J.S., Bemben, L.A., et al. (2005) Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437: 376–380.
- O'Brien, H., Parrent, J., Jackson, J., Moncalvo, J., and Vilgalys, R. (2005) Fungal community analysis by large-scale sequencing of environmental samples. *Appl Environ Microbiol* 71: 5544–5550.
- Oehl, F., Sieverding, E., Ineichen, K., Mäder, P.M., Boller, T., and Wiemken, A. (2003) Impact of land use intensity on the species diversity of arbuscular mycorrhizal fungi in agroecosystems of central Europe. *Appl Environ Microbiol* 69: 2816–2824.
- Oehl, F., Sieverding, E., Ineichen, K., Ris, E.A., Boller, T., and Wiemken, A. (2005) Community structure of arbuscular mycorrhizal fungi at different soil depths in extensively and intensively managed agroecosystems. *New Phytol* 165: 273–283.
- Öpik, M., Moora, M., Liira, J., and Zobel, M. (2006) Composition of root-colonizing arbuscular mycorrhizal fungal communities in different ecosystems around the globe. *J Ecol* 94: 778–790.
- Öpik, M., Moora, M., Zobel, M., Saks, U., Wheatley, R., Wright, F., and Daniell, T. (2008) High diversity of arbuscular mycorrhizal fungi in a boreal herb-rich coniferous forest. *New Phytol* 179: 867–876.
- Pozo, M.J., and Azcon-Aguilar, C. (2007) Unraveling mycorrhiza-induced resistance. *Curr Opin Plant Biol* 10: 393–398.
- Quince, C., Lanzén, A., Curtis, T.P., Davenport, R.J., Hall, N., Head, I.M., et al. (2009) Accurate determination of microbial diversity from 454 pyrosequencing data. *Nat Methods* 6: 639–641.
- Quinlan, A.R., Stewart, D.A., Strömberg, M.P., and Marth, G.T. (2008) Pyrobayes: an improved base caller for SNP discovery in pyrosequences. *Nat Methods* 5: 179–181.
- Renker, C., Weißhuhn, K., Kellner, H., and Buscot, F. (2006) Rationalizing molecular analysis of field-collected roots for assessing diversity of arbuscular mycorrhizal fungi: to pool, or not to pool, that is the question. *Mycorrhiza* 16: 525–531.
- Roesch, L.F.W., Fulthorpe, R.R., Riva, A., Casella, G., Hadwin, A.K.M., Kent, A.D., et al. (2007) Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME J* 1: 283–290.
- Rosselló-Mora, R., and Amann, R. (2001) The species concept for prokaryotes. *FEMS Microbiol Rev* 25: 39–67.
- Sanders, I.A. (2004) Plant and arbuscular mycorrhizal fungal diversity – are we looking at the relevant level of diversity and are we using the right techniques? *New Phytol* 164: 415–418.
- Santos-Gonzalez, J.C., Finlay, R.D., and Tehler, A. (2007) Seasonal dynamics of arbuscular mycorrhizal fungal communities in roots in a seminatural grassland. *Appl Environ Microbiol* 73: 5613–5623.

- Sato, K., Suyama, Y., Saito, M., and Sugawara, K. (2005) A new primer for discrimination of arbuscular mycorrhizal fungi with polymerase chain reaction-denature gradient gel electrophoresis. *Grassl Sci* 51: 179–181.
- Schloss, P.D., and Handelsman, J. (2005) Introducing DOTUR, a computer program for defining operational taxonomic units and estimating species richness. *Appl Environ Microbiol* 71: 1501–1506.
- Schüßler, A., Schwarzott, D., and Walker, C. (2001) A new fungal phylum, the Glomeromycota: phylogeny and evolution. *Mycol Res* 105: 1413–1421.
- Selosse, M.A., Setaro, S., Glatard, F., Richard, F., Urcelay, C., and Weiß, M. (2007) Sebaciniales are common mycorrhizal associates of Ericaceae. *New Phytol* 174: 864–878.
- Da Silva, P.M., Carlos, A.S.A., Niemela, J., Sousa, J.P., and Serrano, A.R.M. (2009) Cork-oak woodlands as key-habitats for biodiversity conservation in Mediterranean landscapes: a case study using rove and ground beetles (Coleoptera: Staphylinidae, Carabidae). *Biodivers Conserv* 18: 605–619.
- Simon, L.M., Lalonde, T.D., and Bruns, T.D. (1992) Specific amplification of 18S fungal ribosomal genes from vesicular arbuscular endomycorrhizal fungi colonising roots. *Appl Environ Microbiol* 58: 291–295.
- Smith, S.E., and Read, D.J. (2008) *Mycorrhizal Symbiosis*, 3rd edn. New York, USA: Academic Press.
- Stockinger, H., Walker, C., and Schüßler, A. (2009) ‘*Glomus intraradices* DAOM197198’, a model fungus in arbuscular mycorrhiza research, is not *Glomus intraradices*. *New Phytol* 183: 1176–1187.
- Toljander, J.F., Santos-González, J.C., Tehler, A., and Finlay, R.D. (2008) Community analysis of arbuscular mycorrhizal fungi and bacteria in the maize mycorrhizosphere in a long-term fertilization trial. *FEMS Microbiol Ecol* 65: 323–338.
- Vallino, M., Massa, N., Lumini, E., Bianciotto, V., Berta, G., and Bonfante, P. (2006) Assessment of arbuscular mycorrhizal fungal diversity in roots of *Solidago gigantea* growing on a polluted soil in northern Italy. *Environ Microbiol* 8: 971–983.
- Wardle, D., Bardgett, R., Klironomos, J., Setälä, H., Van Der Putten, W., and Wall, D. (2004) Ecological linkages between aboveground and belowground biota. *Science* 304: 1629–1633.
- Warnecke, F., Luginbühl, P., Ivanova, N., Ghassemian, M., Richardson, T.H., Stege, J.T., et al. (2007) Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite. *Nature* 450: 560–565.
- Wommack, K.E., Bhavsar, J., and Ravel, J. (2008) Metagenomics: read length matters. *Appl Environ Microbiol* 74: 1453–1463.
- Young, J.P.W. (2009) Kissing cousins: mycorrhizal fungi get together. *New Phytol* 181: 751–753.