

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51590020>

Primary duodenal follicular lymphoma: 6-years complete remission after combined radio-immunotherapy

Article in *Acta gastro-enterologica Belgica* · June 2011

Source: PubMed

CITATION

1

READS

94

8 authors, including:



Pierfrancesco Franco

Università degli Studi di Torino

126 PUBLICATIONS 706 CITATIONS

SEE PROFILE



Andrea Riccardo Filippi

Università degli Studi di Torino

91 PUBLICATIONS 786 CITATIONS

SEE PROFILE



Patrizia Ciammella

Azienda Ospedaliera Santa Maria Nuova di R...

58 PUBLICATIONS 427 CITATIONS

SEE PROFILE



Corrado Tarella

Università degli Studi di Torino

63 PUBLICATIONS 1,944 CITATIONS

SEE PROFILE

All content following this page was uploaded by [Pierfrancesco Franco](#) on 29 November 2016.

The user has requested enhancements of the downloaded file.

Primary duodenal follicular lymphoma : 6-years complete remission after combined radio-immunotherapy

Pierfrancesco Franco¹, Andrea Riccardo Filippi¹, Patrizia Ciammella¹, Angela Botticella¹, Agnieszka Namysl-Kaletka^{2,1}, Alberto De Crescenzo³, Corrado Tarella³, Umberto Ricardi¹

(1) Department of Medical and Surgical Sciences, Radiation Oncology Unit, University of Torino, Ospedale S. Giovanni Battista, Turin, Italy; (2) Department of Radiotherapy, Center of Oncology, Maria-Sklodowska-Curie Memorial Institute, Gliwice, Poland; (3) Hematology and Cell Therapies Unit, University of Torino, Ospedale Umberto I, Turin, Italy.

Abstract

Primary gastrointestinal lymphoma (PGL) is known to account for 40% of all extranodal non-Hodgkin's lymphomas (NHLs) and between 4% to 12% of all NHLs. The small intestine is the site of presentation in 20-30% of cases, with the terminal ileum usually involved. Duodenal localizations have always been thought to be rare, but are presently growing in incidence. We herein report on a case of Stage IV primary duodenal FCL, located to the second portion of the duodenum with concomitant minimal bone marrow involvement. The patient was frontline approached with a conservative combined modality treatment consisting of 4 weekly infusions of the chimeric human-murine IgG1 mono-clonal antibody against the B-cell surface antigen CD-20, Rituximab (375 mg/m²) and consolidation 3D conformal external beam radiotherapy up to a total dose of 36 Gy given into 20 fractions to the involved duodenal portion. Six years after treatment has been completed, the patient is free from disease with no treatment-related toxicity. (*Acta gastroenterol. belg.*, 2011, 74, 337-342).

Key words : radiotherapy, follicular, duodenal, lymphoma, rituximab.

Introduction

Primary gastrointestinal lymphoma (PGL) is known to account for 1 to 4% of all gastro-intestinal malignancies (1). Moreover, the gastro-intestinal tract as a site of origin for non-Hodgkin's lymphomas (NHLs) includes from 4 to 12% of them and 40% of all extranodal NHLs (2). While the stomach is the most common site of involvement, accounting for approximately 50-60% of all cases, the small intestine is the primary location in 20-30%. It often occurs within the terminal ileum, whereas duodenal localizations have always been thought to be rare, but are presently growing in incidence (2-4). Concerning histology, high-grade diffuse large B-cell lymphoma and mucosa-associated lymphoid tissue lymphoma (MALT lymphoma) represent the vast majority of PGLs ; follicular cell lymphoma (FCL) accounts for only 1-7% of them (4-7). Primary FCL of the duodenum is a rare disease that has been recently recognized as a distinct clinico-pathologic entity. A few studies of duodenal FCL have shown that it has features similar to those of conventional nodal follicular lymphoma with an indolent clinical course and immuno-histological features undistinguishable from nodal follicular lymphoma (8). We herein report on a case of Stage IV primary duodenal FCL, located to the second portion of the duodenum with

concomitant minimal bone marrow involvement. The treatment approach consisted of combined immuno-radiation which achieved a long lasting complete remission.

Case report

In 2003, a 42 aged healthy women was referred to our institutional hospital, due to the recent appearance of epigastric discomfort usually worsening shortly after food intake, accompanied by nausea sensation and sporadic vomiting. Patient's previous medical history was silent, apart from a minimal change in bowel habits and a slight weight loss during the last four months. Physical examination could not reveal any remarkable finding, with no superficial lymphadenopathy detectable. Moreover no hepato-splenomegaly or cutaneous lesions could be highlighted. Complete blood count and serum biochemistry were performed, but all findings were within normal limits. However, serum anti-Helicobacter Pylori antibodies were demonstrated as positive (65 U/ml). Thereby, to confirm clinical suspicious, an upper gastrointestinal endoscopy was performed, with the evidence of a duodenal granular flat elevated mucosal lesion, irregularly shaped and composed of whitish, coarsely lumped, polypoid raised granules (Fig. 1a). The patient subsequently underwent a punch biopsy of the lesion showing the presence of neoplastic follicles composed predominantly of small-to-medium-sized cleaved cells (centrocytes) with scattered small round lymphocytes. Immunostaining showed that atypical cells were positive for antibodies directed towards the B-lineage antigens such as L-26, CD 10, CD 20, Cd79a, CD 75 and BCL-2, (intensely positive in the nodule center) and negative for CD5 and CD3, CD23 and cyclin D1. A diagnosis of lymphoma was made and it was classified according to the Revised European-American Lymphoma Classification as a follicle center lymphoma,

Correspondence to : Pierfrancesco Franco, Radiation Oncology, University of Torino, Via Genova, 3, 10126 Torino, Italy.
E-mail : pierfrancesco.franco@gmail.com

Submission date : 09/06/2009

Acceptance date : 07/06/2010

follicular, grade 1. Moreover, no evidence of *Helicobacter Pylori* could be detected after gastrointestinal endoscopy.

A complete staging work-up was performed. Endoscopic ultrasonography (EUS) demonstrated a substantial thickening of the mucosal and submucosal layers of the 2nd portion of duodenal wall with the synchronous evidence of multiple hypoechoic granules. A complete enteroscopy procedure was performed with the ingestion of a wireless capsule in the shape of videocapsule endoscopy (VCE). Lesions were uniquely located within the duodenum. A computed tomography scans of the chest, abdomen and pelvis failed to show any lymph node enlargement. Bone marrow aspiration showed a minimal invasion by small sized cleaved cells. The tumour was staged as IV due to bone marrow involvement (according to Musshoff's modification of the Ann Arbor classification) (9). The patient was first approached with immunotherapy including a weekly infusion of the chimeric human-murine IgG1 monoclonal antibody against the B-cell surface antigen CD-20, Rituximab (375 mg/m²). After completion of immunotherapy, an upper gastrointestinal endoscopy

was repeated in order to evaluate FCL response to rituximab. The examination demonstrated a partial response of the duodenal lesion, diminishing in size and thickening (Fig. 1b). In order to consolidate rituximab results, the patient underwent a full course of external beam radiation therapy delivered with a three-dimensional conformal field arrangement (Fig. 2). The conventionally fractionated total dose was up to 36 Gy given in 20 fractions. Localization of the lesion was facilitated by the placement of a metallic clip in the peri-lesional region during the endoscopic restaging procedure (Fig. 1c). No acute toxicity could be recorded. One month after RT had been completed, a re-evaluation endoscopy showed a complete disease remission with the total disappearance of the duodenal lesion (Fig. 1d). The first bone marrow aspiration, performed at six months, showed the persistence of minimal residual disease (MRD) with nested Polymerase Chain Reaction (PCR) for Bcl-2/Ig-H translocation. At one year, a second biopsy demonstrated the achievement of a molecular remission (MR) status. Six years after treatment the patient is still in complete maintained MR. No treatment-related toxicity has yet been recorded up to last follow-up examination.




Fig. 1. — Granular flat elevated mucosal vegetation of the duodenum at diagnostic upper gastrointestinal endoscopy (a) ; partial remission of the macroscopic lesion after 4 cycles of Rituximab (b) ; metallic clip placed during restaging endoscopy (c) ; complete remission after 36 Gy external beam radiation therapy (d).




Fig. 2. — Antero-posterior view on digitally reconstructed radiographies (a,b) ; evidence of metallic clip (arrow) during simulation fluoroscopy (c) ; anterior electronic portal image with radio-opaque clip (arrow) acquired during treatment (d).

Discussion

Early reports have stated that duodenal lymphomas are a rare clinicopathological entity, since gastro-intestinal tract (GI) lymphomas mainly involve the stomach and the colorectum. However follicular cell histology is a frequent finding among them (3,10). LeBrun *et al.* described 31 cases of follicular cell lymphomas (FCL) of the GI tract in which lesions were predominantly located within the small intestine (55%), showing a peculiar predilection for the terminal ileum. No mention was made about the exact number of duodenal cases (8). More recently, Yoshino *et al.* reported a high incidence of follicular duodenal lymphoma (4). In their series, 8 out of 222 cases (3.6%) arose from the GI mucosa. Among them, 5 out of 8 cases belonged to the duodenum. When these data have been updated, 52 extranodal follicular lymphomas out of 550 (10%) were described ; among them 40 cases were located within the duodenum. The authors observed that this finding might be due to the usual exploration of duodenal tract during routine endoscopies which is of common practice among the Japanese population (11). Duodenal FCL seems to share

some common characteristics with nodal ones such as the indolent nature, the female predominance, the occurrence in the fourth or fifth decade of life and a supposed excellent outcome (12). Moreover, a predilection for the second portion of the duodenum in the region surrounding the ampulla of Vater has been noted by some authors. Several hypotheses have been raised in order to explain this characteristic. Firstly, a regional quantitative difference in the amount of intestinal lymphoid tissue and secondly, a possible role of bile salts and pancreatic juice in acting as a carcinogen stimulation (13). Adjunctively, regional lymphnode involvement has frequently been found in duodenal FCLs, even if the primary lesion only involves the submucosal layer. This may be due to the regional anatomical peculiarities with intricate muscular layers allowing lymphoma cells to easily infiltrate regardless of biological intrinsic aggressiveness or systemic diffusion tendency (14). Treatment options have been various such as surgical resection, radiation therapy (RT), chemotherapeutic agents (CT), immunomodulator, eradication therapy and several therapeutic combinations (Table I) but a combined CT-immunotherapy approach could be considered as a standard option (15). Generally

Table 1. — Overview of all reported cases

Reference	Pts	Sex	Duodenal	Grade	Stage	Therapy	Outcome Portion
CT-Rituximab							
Ahmed <i>et al.</i> (19)	1	F	2 nd	1	IIA	CVP ; R	CR at 2 years
Esaki <i>et al.</i> (20)	1	M	1 st - 2 nd	NR	IIA	R-CHOP	PR at 6 months
Aguiar <i>et al.</i> (21)	1	NA	NA	NA	I	R	CR
Tsujioka <i>et al.</i> (22)	8	4F/4M	Ampulla of Vater	1	IAE	R-CHOP	7/8 CR at 39 months
Isogai <i>et al.</i> (23)	1	NA	NA	NA	NA	R-CHOP	NA
Al-Salman <i>et al.</i> (24)	1	M	Whole jejunum/3 rd	NR	IV	R-CHOP	CR after therapy
CT-Rituximab-RT							
Zenda <i>et al.</i> (25)	1	F	Ampulla of Vater	1	I	R-CHOP ; 4Gy RT	CR after therapy
Exclusive surgery							
Tanaka <i>et al.</i> (14)	1	F	3 rd	1	IA	PD	NR
Matsuzawa (26)	1	F	2 nd	3	III	PD Adjuvant CT	CR at 2 years
Yoshino <i>et al.</i> (15)	5	F	2 nd	1	I-II	S	Alive at 2-50 months
Surgery + RT							
Misdraji <i>et al.</i> (13)	1	F	Ampulla of Vater	1	II	10 Gy Pre-op RT PD 36 Gy Post-op RT	CR after 5 months
Eradication therapy							
Higuchi <i>et al.</i> (27)	1	M	2 nd /jejunum	1	I	LAC	SD at 5 months
Toyoda <i>et al.</i> (28)	1	M	Ampulla of Vater	2	IE	LAC	CR at 2 years
Exclusive radiation							
Takamura <i>et al.</i> (29)	1	M	2 nd – 3 rd	1	IA	30 Gy RT	CR at 3 years
Miscellaneous							
Bende <i>et al.</i> (12)	3	3 F	Duodenum	NR	IE	2 RT only 1 CT only	NED at 24-60 months
Born <i>et al.</i> (30)	1	F	Whole/jejunum	1	IA	None	NR
Nakase <i>et al.</i> (31)	1	F	Papilla of Vater	NR	IA	NR	NR
Mori <i>et al.</i> (32)	1	NA	NA	NA	NA	NA	NA
Tang <i>et al.</i> (33)	1	M	2 nd	NR	I	None	SD
Ridondo-Cerezo (34)	1	F	2 nd	NR	NR	Laparoscopy/CT	NR
Katsuki <i>et al.</i> (35)	1	NA	NA	NA	Early	S	NA
Takikawa <i>et al.</i> (36)	1	NA	NA	NA	NA	S	NA
Sasaki <i>et al.</i> (37)	1	NA	NA	NA	NA	R	NA
Sato <i>et al.</i> (11)	40	17 F 23 M	38 2 nd 2 2 nd + 3 rd	4 G2 36 G1 3 III E	30 IE 7 IIE	13 O 5 PD 3 ppD 2 CT 2 R-CT 5 E 1 E + R-CT 1 CT-RT 8 NR	13 AWD 2-61 mo 5 NED 17-50 mo 3 NED 13-32 mo 2 AWD at 4-5 mo 1 NED at 24 mo 1 AWD at 12 mo 5 AWD at 6-60 mo 1 NED at 12 mo 1 NED at 18 mo NR
Shia <i>et al.</i> (38)	10	7 M 3 F	7 duodenum 3 duodenum/jejunum	6 G1 4 G2	6 IE 4 IIE	3 O 2 S 1 S + CT 3 CT 1 CT/RT	AWD at 2-69 mo 1 NED at 33 mo/1 DOD NED at 73 mo 2 NED at 10-55mo 1 AWD at 6 mo 1 AWD at 44 mo

F : female ; M : male ; NA : not available ; NR : not reported ; R : rituximab ; CVP : cyclophosphamide, vincristine, prednisone ; CHOP : cyclophosphamide, doxorubicin, vincristine, prednisone ; CR : complete remission ; PR : partial remission ; SD : stable disease ; NED : no evidence of disease ; AWD : alive with disease ; LAC : lansoprazole, amoxicillin, clarithromycin ; O : observation ; CT : chemotherapy ; RT : radiotherapy ; S : surgery ; PD : pancreatoduodenectomy ; ppPD : pylorus preserving pancreatoduodenectomy ; E : eradication ; NR : no response.

speaking, FCL is usually characterised by an indolent course and a median survival of 10 years or even more. The vast majority of strategies employed in FL patients with curative intent achieves complete remission in most patients. Unfortunately, the clinical course is usually characterised by multiple relapses. Regardless of the initial treatment chosen, most patients eventually die of lymphoma (16). The use of combination CT guaranteed a major impact in terms of high complete remission rates but did not change the overall survival pattern. High dose therapy with autologous bone marrow transplantation, given either to consolidate remission or as an upfront strategy, may improve remission duration, but it has not shown to improve survival. On the contrary, incumbent toxicity in terms of myelodysplastic syndromes has been described. Most studies suggest that early treatment dose not appear to prolong the survival of asymptomatic patients. In this context, the observation from Stanford of an equal survival between asymptomatic patients managed with expectancy until symptoms appear and those deemed to need therapy immediately properly led to the widespread acceptance of the so called 'benign neglect'(15). More recently, the anti CD-20 monoclonal antibody Rituximab proved to be effective against CD-20 positive low-grade follicular lymphoma, with high response rates in combination with conventional CT and even in CT-refractory patients. Adjunctively, treatment-related toxicity has been reported to be mild and primarily limited to infusion-related events (17). Our group previously reported on the use of rituximab and radiation in localized nodal and extranodal FCL (18), with promising results either in terms of disease control (both local and systemic) and treatment-related toxicity. In this context, in order to combine an effective local treatment with a systemic approach able to clear bone marrow from neoplastic clones, with a concomitant attention towards a low toxicity profile we chose to treat the patient with combined rituximab and radiation. In this way we have been able to perform a conservative approach, avoiding both surgical excision (which might have been associated with a high morbidity considering the anatomical site involved) and CT-related side effects and sequelae with an excellent clinical outcome. The sequential approach (rituximab prior to consolidation radiotherapy) lead to the possibility of a shrinkage of the duodenal lesion allowing the reduction of both radiation treatment volumes and doses, diminishing the probability of RT-related acute and chronic effects.

References

1. KAWAI T. Lymphoma arising in mucosa associated lymphoid tissue of the duodenal bulb. *J. Gastroenterol.*, 1998, **33** : 97-101.
2. DOMIZIO P., OWEN R.A., SHEPERD N.A., TALBOT I.C., NORTON A.J. Primary lymphoma of the small intestine. *Am. J. Surg. Pathol.*, 1993, **17** : 429-42.
3. CIRILLO M., FEDERICO M., CURCI G., TAMBORRINO E., PICCININI L., SILINGARDI V. Primary gastrointestinal lymphoma: a clinicopathological study of 58 cases. *Haematologica*, 1992, **77** : 156-61.

4. YOSHINO T., MIYAKE K., ICHIMURA K., MANNAMI T., OHARA N., HAMAZAKI S., AKAGI T. Increased incidence of follicular lymphoma of the duodenum. *Am. J. Surg. Pathol.*, 2000, **24** : 688-93.
5. SHIA J., TERUYA-FELDSTEIN J., PAN D., HEGDE A., KLIMSTRA D.S., CHAGANTI R.S., QIN J., PORTLOCK C.S., FILIPPA D.A. Primary follicular lymphoma of the gastro-intestinal tract : a clinical and pathologic study of 26 cases. *Am. J. Surg. Pathol.*, 2002, **26** : 216-24.
6. BENDE R.J., SMIT L.A., BOSSENBROEK J.G., AARTS W.M., SPAARGAREN M., DE LEVAL L., BOECKXSTAENS G.E., PALS S.T., VAN NOESEL C.J. Primary follicular lymphoma of the small intestine : alpha4beta7 expression and immunoglobulin configuration suggest an origin from local antigen-experienced B cells. *Am. J. Pathol.*, 2003, **162** : 105-13.
7. GOODLAD J.R., MACPHERSON S., JACKSON R., BATSTONE P., WHITE J. on behalf of the Scotland & Newcastle Lymphoma Group. Extranodal follicular lymphoma : a clinicopathological and genetic analysis of 15 cases arising at non-cutaneous extranodal sites. *Histopathology*, 2004, **44** : 268-73.
8. LEBRUN D.P., KAMEL O.W., CLEARLY M.L., DORFMAN R.F., WARNKE R.A. Follicular lymphomas of the gastrointestinal tract : pathologic features in 31 cases and bcl-2 oncogenic protein expression. *Am. J. Pathol.*, 1992, **140** : 1327-35.
9. MUSSHOF K. Clinical staging classification of non-Hodgkin's lymphomas. *Strahlentherapie*, 1977, **153** : 218-21.
10. NNAJEM A.Z., PORCARO J.L., RUSH B.F. Primary non-Hodgkin's lymphoma of the duodenum : case report and literature review. *Cancer*, 1984, **54** : 895-8.
11. SATO Y., ICHIMURA K., TANAKA T., TAKATA K., MORITO T., SATO H., KONDO E., YANAI H., OHARA N., OKA T., YOSHINO T. Duodenal follicular lymphomas share common characteristics with mucosa-associated lymphoid tissue lymphomas. *J. Clin. Pathol.*, 2008, **61** : 377-81.
12. BENDE R.J., SMIT L.A., BOSSENBROEK J.G., AARTS W.M., SPAARGAREN M., DE LEVAL L., BOECKXSTAENS G.E., PALS S.T., VAN NOESEL C.J.M. Primary follicular lymphoma of the small intestine. 4B7 expression and immunoglobulin configuration suggest an origin from local antigen-experienced B cells. *Am. J. Pathol.*, 2003, **162** : 105-13.
13. MISDRAJI J., DEL CASTILLO C.F., FERRY J. Follicle center lymphoma of the ampulla of Vater presenting with jaundice. *Am. J. Surg. Pathol.*, 1997, **140** : 1327-35.
14. TANAKA S., ONOUE G., FUJIMOTO T., KOSAKA T., YAMASAKI H., YASUI Y., KONAGA E., TERAMOTO N., OKADA H., MIZUNO M., SHIRATORI Y. A case of primary follicular lymphoma in the duodenum confined to the mucosa layer. *J. Clin. Gastroenterol.*, 2002, **35** : 285-6.
15. ADVANI R., ROSENBERG S.A., HORNING S.J. Stage I and II follicular non-Hodgkin's lymphoma : long-term follow-up of no initial therapy. *J. Clin. Oncol.*, 2004, **22** : 1454-9.
16. YOSHINO T., MIYAKE K., ICHIMURA K., MANNAMI T., OHARA N., HAMAZAKI S., AKAGI T. Increased incidence of follicular lymphoma of the duodenum. *Am. J. Surg. Pathol.*, 2000, **24** : 688-93.
17. HAINSWORTH J.D. First-line and maintenance treatment with rituximab for patients with indolent non-Hodgkin's lymphoma. *Semin. Oncol.*, 2003, **30** : 9-15.
18. RICARDI U., FILIPPI A.R., CUTTICA A., CARACCILO D., ZANNI M., BAZZAN M., FREILONE R., FRANCO P., BOCCADORO M., TARELLA C. Rituximab followed by localized radiation therapy in limited stage indolent non-Hodgkin's lymphoma : preliminary results of a pilot study. *J. Clin. Oncol.*, 2005, **23** (16S) : 6696.
19. AHMED S., SINGH A., KRAUSS J., WENTZ T., GUNARATNAM N.T. Successful treatment of refractory low -grade duodenal lymphoma with Rituximab, an Anti-CD20 monoclonal antibody. *Am. J. Clin. Oncol.*, 2003, **26** : 408-10.
20. ESAKI M., MATSUMOTO T., NAKAMURA S., SUEKANE H., OHJI Y., YAO T., IIDA M. Capsule endoscopy findings in intestinal follicular lymphoma. *Endoscopy*, 2007, **39** : E86-E87.
21. AGUIAR-BUJANDA D., QUINONES-MORALES I., CAMACHO-GALAN R., LLORCA-MARTINEZ I., RIVERO-VERA J.C., BOHN-SARMIENTO U., AGUIAR-MORALES J. Primary duodenal follicular lymphoma successfully treated with rituximab. *Clin. Transl. Oncol.*, 2007, **9** : 471-2.
22. TSUJIOKA T., WADA H., YATA K., KONDO T., SUEMORI S., TOKUNAGA H., OHMORI K., KUBO Y., NAKANISHI H., MIKAMI M., HARUMA K., SADAHIRA Y., SUGIHARA T. Clinical analysis of eight patients with primary follicular lymphoma in the duodenum. *Rinsho Ketsueki*, 2007, **48** : 134-9.
23. ISOGAI K., YAMAMOTO E., YAMAOKA S., TARASAWA I., MURAKAMI R., MORIMOTO I., OBATA T., MORI M., ENDO T., IMAI K. A case of follicular lymphoma of the duodenum treated with the combined therapy of anti-CD 20 antibody and chemotherapy. *Nippon Shokakibyō Gakkai Zasshi*, 2006, **103** : 301-9.

24. AL-SALMAN J., SALIB H., BOONSWANG P. Successful treatment of gastrointestinal follicular lymphoma with Rituxan and combination chemotherapy. *Med. Oncol.*, 2001, **18** : 277-83.
25. ZENDA T., MASUNAGA T., FUWA B., OKADA T., ONTACHI Y., KONDO Y., NAKAO S., MINATO H. Small follicular lymphoma arising near the ampulla of Vater : a distinct subtype of duodenal lymphoma ? *Int. J. Gastrointest. Cancer*, 2005, **36** : 113-9.
26. MATSUZAWA M., SHIMODAIRA K., NAKAMURA N., OCHI Y., HOSAKA S., KIYOSAWA K., AKAMATSU T., SUGIYAMA A., OTA H., KATSUYAMA T. A case of primary follicular lymphoma of the duodenum with BCL-2 gene rearrangement. *J. Gastroenterol.*, 2003, **38** : 512-3.
27. HIGUCHI K., KOMATSU K., WAKAMATSU H., KAWASAKI H., MURATA M., MIYAZAKI K., OIKAWA K., OHWADA M., NANJO H., OTAKA M., WATANABE S., KOMATSU K. Small Intestine follicular lymphoma with multiple tumor formations diagnosed by double-balloon enteroscopy. *Intern. Med.*, 2007, **46** : 705-9.
28. TOYODA H., YAMAGUCHI M., NAKAMURA S., NAKAMURA T., KIMURA M., SUZUKI H., MUKAI K., SAWA H., KAWAMURA K., SHIKU H. Regression of primary lymphoma of the ampulla od Vater after eradication of *Helicobacter pylori*. *Gastrointest. Endosc.*, 2001, **54** : 92-6.
29. TAKAMURA M., NARISAWA R., MARUYAMA Y., YOKOYAMA J., FUKUHARA Y., KAWAI H., YAMAGIWA S., SATO Y., YASUNOBU M., SUGIMURA K., ICHIDA T., AOYAGI Y. A primary follicular lymphoma of the duodenum treated successfully with radiation therapy. *Intern. Med.*, 2006, **45** : 309-11.
30. BORN P., VIETH M., STOLTE M. Follicular lymphoma of the duodenum. *Endoscopy*, 2007, **39** : E39.
31. NAKASE H., MATSUURA M., MIKAMI S., CHIBA T. Magnified endoscopic view of primary follicular lymphoma at the duodenal papilla. *Intern. Med.*, 2007, **46** : 141-2.
32. MORI Y., KATSURADA T., ITOH T., KATO M., ASAKA M. A case of small intestinal follicular lymphoma. *Rinsho Shokaki Naika*, 2005, **20** : 1614-16.
33. TANG Z., JING W., LINDEMAN N., HARRIS N.L., FERRY J.A. One patient, two lymphomas. Simultaneous primary gastric marginal zone lymphoma and primary duodenal follicular lymphoma. *Arch. Pathol. Lab. Med.*, 2004, **128** : 1035-8.
34. REDONDO-CEREZO E., DE BENITO L., SERRENO-EGEA A., GOMEZ C.J., PEREZ G., PEREZ J.I., GARCIA-CANO J., MORILLAS J., GONZALEZ J.A., PEREZ-SOLA A. Unusual presentation and diagnosis of extraintestinal follicular lymphoma. *Dig. Dis. Sci.*, 2004, **49** : 1513-7.
35. KATSUKI S., NOJIRI S., OHMI R. Primary early follicular lymphoma of the duodenum, report of a case. *I to Cho (Stomach and Intestine)*, 1988, **33** : 1665-70.
36. TAKIKAWA H., HIKI N., TATSUTOMI Y. A case of follicular lymphoma in the duodenum successfully treated with surgical resection after eradication of *Helicobacter pylori*. *Prog. Dig. Endosc.*, 2002, **61** : 98-9.
37. SASAKI M., YAKUSHIJIN Y., HATO T., KUSHIDA Y., SHIRAHISHI T., SHIRAHISHI S. A case of follicular lymphoma of the duodenum treated by anti-CD-20 antibody (rituximab). *Gastroenterol. Endosc.*, 2004, **46** : 151-7.
38. SHIA J., TERUYA-FELDSTEIN J., PAN D., HEDGE A., KLIMSTRA D.S., CHAGANTI R.S., QIN J., PORTLOCK C.S., FILIPPA D.A. Primary follicular lymphoma of the gastrointestinal tract : a clinical and pathologic study of 26 cases. *Am. J. Surg. Pathol.*, 2002, **26** : 216-24.