



UNIVERSITÀ DEGLI STUDI DI TORINO

The final publication is available at Springer via
<http://link.springer.com/article/10.1007%2Fs00394-011-0256-6>

Assessment of iron absorption in mice by ICP-MS measurements of ⁵⁷Fe levels

Veronica Fiorito^{1,2,*}, Simonetta Geninatti Crich^{1,3,*}, Lorenzo Silengo^{1,2}, Fiorella Altruda^{1,2}, Silvio Aime^{1,3,°} and Emanuela Tolosano^{1,2,°}

¹Molecular Biotechnology Center, University of Torino, Torino, Italy

²Department of Genetics, Biology and Biochemistry, University of Torino, Torino, Italy

³Department of Chemistry IFM, University of Torino, Torino, Italy

* and ° Equally contributed

Corresponding author:

Emanuela Tolosano, PhD

Molecular Biotechnology Center

Dept. Genetics, Biology and Biochemistry

Via Nizza 52

10126 Torino, Italy

Phone: +39-011-6706423

Fax: +39-011-6706432

email: emanuela.tolosano@unito.it

Keywords: iron absorption, mice, ICP-MS

Funding: this work was supported by the Italian Ministry of University and Research to E.T. and F.A., and by the Regione Piemonte to F.A.

Abstract

Background. The study of iron metabolism is essential in nutritional sciences as iron deficiency is one of the most common nutritional deficiencies in humans, and represents a serious health problem worldwide. The mouse is utilized as a unique and powerful model for the identification and characterization of genes involved in iron metabolism, and for studying the pathogenesis of iron disorders. Thus, sophisticated and sensitive techniques have been developed to study iron metabolism in this animal model. In particular, iron absorption has been studied in mice by using the radioisotopes ^{55}Fe and ^{59}Fe in tied-off or dissected and everted duodenal segments. Nevertheless, several drawbacks discourage the extended use of these approaches.

Methods and Results. Here we report the use of the stable isotope ^{57}Fe to measure iron absorption in mice. We show that after oral administration of ^{57}Fe -containing solutions it is possible to measure both duodenal iron retention and duodenal iron transfer to specific organs, using inductively coupled plasma mass spectrometry (ICP-MS). As ^{57}Fe is administered orally, no surgical operation is needed before the end of the experiment, thus allowing the measurement of iron absorption under physiologic conditions. Moreover, the use of ICP-MS for ^{57}Fe detection ensures high sensitivity and provides quantitative data. Finally, the use of a stable isotope enables the measurement of both iron absorption and histologic and/or biochemical analyses in the same animal.

Conclusions. The use of ^{57}Fe to measure iron absorption in mice therefore represents an alternative to radioisotope-based methods, providing a new tool to extend our knowledge on the mechanism of iron absorption.

Introduction

Iron is an essential nutrient for all forms of life as it participates in many biological processes, including oxygen transport, as well as behaving as a cofactor to several enzymes. In humans, iron deficiency represents a common health problem both in Western and developing countries. The main causes of iron deficiency are poor absorption of iron by the body, inadequate daily intake of iron, pregnancy, growth spurts or blood loss due to trauma, infection, malaria or hematological disorders. Conversely, iron is toxic when in excess, and therefore body iron levels are tightly regulated. As the body cannot actively excrete iron, the body iron content is regulated at the absorption step when it is taken up by duodenal villus enterocytes. Main stimuli for iron absorption are represented by changes in body iron stores, and according to the requirements of the developing erythroid mass [1].

Most of our knowledge on iron homeostasis relies on studies carried out on mice. Several genetic models are available, including both natural and gene targeting-generated mutations of the main genes involved in iron absorption, recycling, storage and utilization [2,3,4,5,6]. The mouse also represents a unique model to identify novel players in iron homeostasis [7,8,9]. Finally, it has been shown that the occurrence, in mice, of genetic mutations responsible for human iron disorders, for example hemochromatosis and anemia, fully recapitulate the human diseases, thus stressing the importance of mouse models for studying the pathogenetic mechanisms of these diseases and for testing new therapies [10,11].

Up until now, iron absorption has been measured in mice by using the tied-off duodenal technique along with the radioisotopes ^{55}Fe and ^{59}Fe [12,13]. ^{55}Fe is a beta emitter which allows the determination of mucosal uptake after administration of an oral dose of iron or incubation of a duodenal segment in the presence of the radioisotope, whereas ^{59}Fe is a gamma and beta emitter that,

besides measurement of mucosal uptake, enables the determination of iron transferred from the duodenum to the rest of the body. The main disadvantage of using ^{59}Fe is that gamma emissions have a significant penetrating and shallow external exposure hazard, thus requiring laboratories specifically equipped for handling animals treated with this isotope. Here we report an alternative approach to assess iron uptake and distribution in mice, based on the use of the stable iron isotope ^{57}Fe , detected by inductively coupled plasma mass spectrometry (ICP-MS). ^{57}Fe is one of the four naturally-occurring isotopes of iron, and accounts for 2.119% of total iron, the other isotopes being ^{54}Fe (5.845%), ^{56}Fe (91.754%) and ^{58}Fe (0.282%) [14]. Natural iron isotope variations in the blood may serve to identify differences in intestinal iron absorption between individuals and genotypes [15]. Moreover, ^{57}Fe and ^{58}Fe have already been used as tracers in nutritional studies on humans due to their low natural abundance [16,17,18].

Herein a procedure is reported to assess both duodenal mucosal retention and mucosal transfer of ^{57}Fe to specific organs, following administration of an oral dose of ^{57}Fe in mice.

Materials and methods

Animals

Wild-type and Hfe-null mice [19] in the 129Sv genetic background were fed on a standard diet (4RF25 GLP, Mucedola, Settimo Milanese, Milano, Italy) containing 8 mg/kg and 292 mg/kg heme-iron and inorganic iron respectively, and received water ad lib. In the low-iron experiments, mice were given a AIN93G iron-deficient diet (Mucedola, Settimo Milanese, Milano, Italy) containing 54 mg/kg total iron for 2 weeks.

All experiments were approved by the animal ethical committee of the University of Torino (Italy).

Treatment

The stable iron isotope ^{57}Fe (^{57}Fe at 94% enrichment; Frontier Scientific Inc., Logan, Utah USA) was used as tracer. A 0.4 mol/L solution of $^{57}\text{FeSO}_4$ was prepared by overnight dissolution of 22.85 g $^{57}\text{Fe}/\text{L}$ in 0.4 mol/L H_2SO_4 (Sigma Aldrich, Milano, Italy). The obtained $^{57}\text{FeSO}_4$ solution was stored at 4°C. Before its use, 87.7 mg sucrose and 0.83 mg ascorbic acid per mg iron were added to the $^{57}\text{FeSO}_4$ solution to yield to a final concentration of 20 mmol/L ^{57}Fe , 5.38 mmol/L ascorbic acid and 10% sucrose .

As a negative control, an analogous solution with no tracer was prepared.

Both the ^{57}Fe -labelled and the control solution were adjusted to pH7 by adding the required volume of 1 mol/L NaOH. To assess the in vivo tissue absorption of ^{57}Fe after oral administration of ^{57}Fe -containing solutions, overnight-fasted mice were orally treated with 20 μl of ^{57}Fe -labelled solution. Control mice received vehicle solutions. During treatment, mice received water ad lib. Tissues were collected at varying times following treatment. Control mice represented the “0” time point of the treatment.

Tissue Collection

Mice were anaesthetized with Avertin (2,2,2-tribromoethanol; Sigma-Aldrich, Milano, Italy) at a dose of 2 mg/kg body weight .

Blood was collected from mice by cardiac puncture. Briefly, following a wash in heparin-containing solution, the needle of a syringe was inserted at the base of the sternum into the thoracic cavity and ventricular blood was slowly withdrawn. Samples were then centrifuged at 4000 rpm for 6 minutes and supernatants were recovered, weighed and stored at -20°C prior to proceeding with the analyses.

Duodenums, livers and kidneys were excised after transcardial perfusion of mice with 0.1 mol/L phosphate-buffered saline (PBS), washed in PBS, blotted dry, weighed and stored at -20°C prior to analyses. Tissue perfusion with PBS was necessary to reduce background signal in ICP-MS measurements of ⁵⁷Fe absorption, considering the high amount of physiological iron present in blood.

Bone marrow was obtained from the hind legs of the mice. Briefly, immediately before surgery, mice were sacrificed. Tibias and femurs were dissected from surrounding muscles and tendons, and flushed with PBS by inserting the needle of a syringe into the ends of the bones. The flow-through was then centrifuged at 1200 rpm for 5 minutes and the pellet was weighed and stored at -20°C prior to proceeding with sample analyses.

Tissue analyses by ICP-MS

⁵⁷Fe isotope and total Fe were determined using inductively coupled plasma mass spectrometry (ICP-MS) (Element-2; Thermo-Finnigan, Rodano (MI), Italy) at medium mass resolution ($M/\Delta M \sim 4000$). This resolution is unable to eliminate isobaric interferences of ⁵⁴Cr and ⁵⁸Ni on ⁵⁸Fe and ⁵⁴Fe isotopes. For this reason the isotope ⁵⁷Fe was selected as tracer. Sample digestion was performed with 2 ml of concentrated HNO₃ (70%) by means of microwave heating at 160°C for 20 minutes (Milestone MicroSYNTH Microwave labstation equipped with an optical fiber temperature control and HPR-1000/6M six position high-pressure reactor, Bergamo, Italy). After digestion, the volume of each sample was brought to 3 ml with ultrapure water and the sample was analyzed by ICP-MS. A natural abundance iron standard solution was analyzed during sample runs in order to check changing in the systematic bias. The calibration curve was obtained using four iron absorption standard solutions (Sigma-Aldrich) in the range 0.2–0.05 µg/ml.

Calculations

After treatment with ^{57}Fe -labelled or control solution, the amount (expressed as μg of iron per g of wet tissue) of total iron ($_{\text{tot}}\text{Fe}$) and of ^{57}Fe isotope was determined for each tissue of interest by ICP-MS. For each tissue, the calculated amount of naturally-occurring ^{57}Fe was subtracted from the measured value in order to get a measure of the amount of iron retained by the tissue upon the treatment with ^{57}Fe -labelled solution. The amount of naturally-occurring ^{57}Fe was calculated considering the average percentage of ^{57}Fe under basal conditions ($\%^{57}\text{Fe}$) in wild-type mice with respect to total iron observed ($_{\text{tot}}\text{Fe}$), as shown in the example reported in Table 1.

Tissue iron measurement by colorimetric method

For tissue iron determination by colorimetric measurement, the BPS-based method was performed, using 4,7-diphenyl-1,10-phenantroline disulphonic acid (BPS) as chromogen [20]. Briefly, 0,1 g of dry tissue was incubated overnight in a mixture of trichloroacetic (10%) and hydrochloric (3 N) acids, and 20 μl of supernatant reduced with thioglycolic acid (Sigma-Aldrich, Milano, Italy) and acetic acid-acetate buffer (pH 4.5). Ferrous iron content was determined spectrophotometrically at 535 nm following the addition of BPS and incubation for 1 hour at 37°C.

Statistical Analysis

Results were expressed as mean \pm SEM. Statistical analyses were performed using one-way or two-way analysis of variance followed by the Bonferroni correction for multiple group comparisons. An unpaired Student's *t*-test was used when only two groups were compared. A *P* value of less than 0.05 was regarded as significant.

Results

Assessment of natural abundance of ^{57}Fe in mouse tissues

The preliminary experimental work for the ^{57}Fe method in mice is summarized as follows.

Firstly, after defining the experimental set-up for tissue collection (for details see “Materials and methods”), the reliability of total iron and ^{57}Fe detection by mass spectrometry in samples recovered from wild-type mice at basal conditions was assessed. Duodenum, liver and kidney tissues were dissected and subjected to ICP-MS and spectrophotometric analyses.

A good correlation between iron levels measured by mass spectrometry, and the standard colorimetric method using BPS as a chromogen (Figure 1A) was found. As expected total liver iron in Hfe-null mice suffering from hemochromatosis resulted as being significantly higher than in wild-type animals (Figure 1B).

In wild-type mice, the amount of ^{57}Fe with respect to total iron ($\%^{57}\text{Fe} /_{\text{tot}}\text{Fe}$) was constant in the different tissues analyzed, i.e serum, duodenum, liver, kidney and bone marrow, and reached a value of about 2.3%, very close to the reported natural percentage of ^{57}Fe in the earth [14] (Figure 1C).

Measurement of ^{57}Fe retention by duodenal mucosa after oral iron administration

The relatively low natural abundance of ^{57}Fe offers the possibility of exploiting this isotope as an easy detectable tracer upon an oral administration of solutions enriched in ^{57}Fe .

To assess the in vivo uptake of ^{57}Fe by the duodenal mucosa after oral administration of ^{57}Fe -containing solutions, 2-month old wild-type mice were treated with 20 μl of a solution containing ^{57}Fe , and duodenum tissue was collected at different times following the treatment. A comparable volume of vehicle solution was given to control mice.

Thirty minutes after treatment, a considerably high amount of ^{57}Fe was detected in the duodenal mucosa of treated mice compared to controls, and the quantity of ^{57}Fe retained by duodenum further increased ninety minutes after the oral administration (Figure 2A).

We used Hfe-null mice as positive controls, known to have enhanced duodenal iron uptake [19], and wild-type mice fed with an iron deficient diet for 2 weeks, that display increased expression of duodenal iron transporters [21]. When ^{57}Fe was administered to Hfe-null mice, at sixty minutes following treatment, the amount of ^{57}Fe detected in the duodenal mucosa was already higher than in wild-type animals (Figure 2B). The same was observed in wild-type mice fed on an iron-deficient diet (Figure 2C).

Determination of the amount of ^{57}Fe delivered to different organs

After uptake from the duodenal lumen, iron captured by enterocytes is usually partly stored in ferritins, and partly exported into the bloodstream through a process involving the iron exporter ferroportin1, located at the basal membrane of enterocytes. Once in the circulation, iron is captured by the iron carrier transferrin and transported to peripheral organs, primarily to the bone marrow [1,22].

To test the possibility that orally administered ^{57}Fe absorbed by the duodenum and subsequently delivered to peripheral organs could be detected by mass spectrometry analyses, we treated wild-type mice as described above and collected the livers and bone marrow at different times following treatment. A slight increase in ^{57}Fe levels in the liver was already evident 30 minutes after administration of iron, and further increased up to 90 minutes following treatment (Figure 3A). Furthermore, two days after treatment, a significant amount of tracer was detected in the bone marrow (Figure 3B). On the other hand, the transfer of ^{57}Fe to the kidney, an organ only marginally involved in iron handling, was negligible compared to the liver and bone marrow (Figure 3C).

As expected, the transfer of ^{57}Fe to the liver of Hfe-null mice and of wild-type mice fed on an iron-deficient diet was strongly enhanced compared to wild-type animals and well detectable 60 minutes after oral administration of iron-containing solutions (Figure 3D and 3E).

Discussion

In this work a novel procedure to measure iron absorption in mice using the stable ^{57}Fe isotope has been described.

Upon administration of an oral dose of a ^{57}Fe -containing solution, a considerably high amount of ^{57}Fe was detected in the duodenal mucosa of mice. Moreover, the transfer of a significant amount of ^{57}Fe to peripheral organs, such as the liver and bone marrow was assessed. As stable iron isotopes exist in nature in fixed ratios, corrections have to be made for background levels when they are used as tracers in absorption studies. For this purpose, corrections for the natural abundance of the stable isotope ^{57}Fe have been considered during the analysis of experiments. Moreover, a series of controls, such as the assessment of iron absorption in Hfe-null mice, and in animals fed on an iron-deficient diet, confirmed the reliability of the proposed method. The sensitivity of the ICP-MS technique allows detection of relatively low amounts of ^{57}Fe corresponding to an experimental detection limit of about 40 ng $^{57}\text{Fe}/\text{g}$ tissue.

The procedure described in the present work therefore represents a good technique to assess iron absorption in mice and, along with the classical methods, allows determination of both mucosal retention and mucosal transfer of iron, distinguishing between the two steps of iron absorption, thus offering a valuable alternative to the approaches that have been used in laboratory practices so far.

The use of orally administered ^{57}Fe to measure iron absorption in mice shows a series of advantages respect to other techniques reported in the literature [23].

Firstly, the use of a stable iron isotope instead of a radioactive one shows obvious benefits in terms of safety and ease of handling, as the use of ^{57}Fe does not require special laboratory equipment or particular skills for the operator.

Secondly, the cost of ^{57}Fe -enriched substances is considerably lower than that of radioactive iron-containing compounds, thus representing an affordable tracer for many research centres.

Moreover, the use of ICP-MS for ^{57}Fe detection ensures high sensitivity and the use of a calibration curve during measurements provides quantitative data.

Another advantage associated with the herein reported method deals with the possibility of carrying out several kinds of analyses on the tissue specimen recovered from animals treated with ^{57}Fe . Indeed, the limited handling of the mouse during the experiment, and the safety of the employed tracer allow the use of excised tissues partly for ICP-MS analyses and partly for any other kind of tests, i.e. biochemical and histological analyses.

Finally, while radioactive tracers are not suitable for multi-element studies because of difficulties due to overlapping energy spectra, the absorption of different stable isotopes can be measured by ICP-MS simultaneously.

A weak point of this methodology is the impossibility to detect the amount of absorbed iron in the entire carcass, which the use of radioactive compound allows [24,25,26]. Of course, separated analyses on different organs of interest can be performed in order to reach a global assessment of iron distribution in the entire organism.

In conclusion, the present work proposes a sensitive, safe, and reliable method to measure iron absorption in mice, thus providing an alternative technique to that reported so far and offering a new methodology to improve the study of iron absorption mechanisms.

Acknowledgments

We thank Antonello Pietrangelo for Hfe-null mice and Radhika Srinivasan for editing of the manuscript.

Conflict of Interest Disclosures

The authors declare no conflicts of interest

References

1. Hentze MW, Muckenthaler MU, Galy B, Camaschella C (2010) Two to tango: regulation of Mammalian iron metabolism. *Cell* 142: 24-38.
2. Andrews NC (2008) Forging a field: the golden age of iron biology. *Blood* 112: 219-230.
3. Bernstein SE (1987) Hereditary hypotransferrinemia with hemosiderosis, a murine disorder resembling human atransferrinemia. *J Lab Clin Med* 110: 690-705.
4. Edwards JA, Hoke JE (1972) Defect of intestinal mucosal iron uptake in mice with hereditary microcytic anemia. *Proc Soc Exp Biol Med* 141: 81-84.
5. Fleming MD, Trenor CC, 3rd, Su MA, Foernzler D, Beier DR, et al. (1997) Microcytic anaemia mice have a mutation in Nramp2, a candidate iron transporter gene. *Nat Genet* 16: 383-386.
6. Trenor CC, 3rd, Campagna DR, Sellers VM, Andrews NC, Fleming MD (2000) The molecular defect in hypotransferrinemic mice. *Blood* 96: 1113-1118.
7. Nicolas G, Bennoun M, Devaux I, Beaumont C, Grandchamp B, et al. (2001) Lack of hepcidin gene expression and severe tissue iron overload in upstream stimulatory factor 2 (USF2) knockout mice. *Proc Natl Acad Sci U S A* 98: 8780-8785.
8. Wang RH, Li C, Xu X, Zheng Y, Xiao C, et al. (2005) A role of SMAD4 in iron metabolism through the positive regulation of hepcidin expression. *Cell Metab* 2: 399-409.
9. Meynard D, Kautz L, Darnaud V, Canonne-Hergaux F, Coppin H, et al. (2009) Lack of the bone morphogenetic protein BMP6 induces massive iron overload. *Nat Genet* 41: 478-481.
10. Camaschella C (2005) Understanding iron homeostasis through genetic analysis of hemochromatosis and related disorders. *Blood* 106: 3710-3717.
11. Iolascon A, De Falco L, Beaumont C (2009) Molecular basis of inherited microcytic anemia due to defects in iron acquisition or heme synthesis. *Haematologica* 94: 395-408.
12. Laftah AH, Raja KB, Beaumont N, Simpson RJ, Deacon A, et al. (2004) The effects of inhibition of haem biosynthesis by griseofulvin on intestinal iron absorption. *Basic Clin Pharmacol Toxicol* 94: 161-168.
13. Masaratana P, Laftah AH, Latunde-Dada GO, Vaulont S, Simpson RJ, et al. (2011) Iron absorption in hepcidin1 knockout mice. *Br J Nutr* 105: 1583-1591.
14. Walczyk T, von Blanckenburg F (2002) Natural iron isotope variations in human blood. *Science* 295: 2065-2066.
15. Krayenbuehl PA, Walczyk T, Schoenberg R, von Blanckenburg F, Schulthess G (2005) Hereditary hemochromatosis is reflected in the iron isotope composition of blood. *Blood* 105: 3812-3816.
16. Chen Z, Griffin IJ, Plumlee LM, Abrams SA (2005) High resolution inductively coupled plasma mass spectrometry allows rapid assessment of iron absorption in infants and children. *J Nutr* 135: 1790-1795.
17. Ellen G.H.M. van den Heuvel WvD GS (1997) Methods to measure iron absorption in humans: a review. *Food Rev Int* 13 (1): 91-102.
18. Young MF, Glahn RP, Ariza-Nieto M, Inglis J, Olbina G, et al. (2009) Serum hepcidin is significantly associated with iron absorption from food and supplemental sources in healthy young women. *Am J Clin Nutr* 89: 533-538.
19. Zhou XY, Tomatsu S, Fleming RE, Parkkila S, Waheed A, et al. (1998) HFE gene knockout produces mouse model of hereditary hemochromatosis. *Proc Natl Acad Sci U S A* 95: 2492-2497.

20. Pieroni L, Khalil L, Charlotte F, Poynard T, Piton A, et al. (2001) Comparison of bathophenanthroline sulfonate and ferene as chromogens in colorimetric measurement of low hepatic iron concentration. *Clin Chem* 47: 2059-2061.
21. Dupic F, Fruchon S, Bensaid M, Loreal O, Brissot P, et al. (2002) Duodenal mRNA expression of iron related genes in response to iron loading and iron deficiency in four strains of mice. *Gut* 51: 648-653.
22. Hentze MW, Muckenthaler MU, Andrews NC (2004) Balancing acts: molecular control of mammalian iron metabolism. *Cell* 117: 285-297.
23. Wienk KJ, Marx JJ, Beynen AC (1999) The concept of iron bioavailability and its assessment. *Eur J Nutr* 38: 51-75.
25. Frazer DM, Wilkins SJ, Becker EM, Murphy TL, Vulpe CD, et al. (2003) A rapid decrease in the expression of DMT1 and Dcytb but not Ireg1 or hephaestin explains the mucosal block phenomenon of iron absorption. *Gut* 52: 340-346.
26. Laftah AH, Latunde-Dada GO, Fakih S, Hider RC, Simpson RJ, et al. (2009) Haem and folate transport by proton-coupled folate transporter/haem carrier protein 1 (SLC46A1). *Br J Nutr* 101: 1150-1156.
27. Owen GM, Kirkman S, Williams M, Jacobs A (1967) A simple method for measuring iron absorption using a scintiscanner for whole-body counting. *Br J Radiol* 40: 866-868.

Table 1. Calculation of ^{57}Fe retention in tissues. An example is given considering a measurement of 20 $\mu\text{g/g}$ for total iron and of 5 $\mu\text{g/g}$ for ^{57}Fe . The calculated amount of naturally occurring ^{57}Fe was subtracted from the measured value in order to get a measure of the amount of iron retained by the tissue upon treatment with the ^{57}Fe -labelled solution.

medium % ^{57}Fe under basal conditions	$_{\text{tot}}\text{Fe}$ ($\mu\text{g/g}$) observed after treatment	^{57}Fe ($\mu\text{g/g}$) observed after treatment	expected naturally occurring ^{57}Fe ($\mu\text{g/g}$)	^{57}Fe retained (^{57}Fe observed - ^{57}Fe expected)
2.4%	20	5	0.48 (2.4% of 20)	$5 - 0.48 = 4.52$

Figure Legends

Figure 1. Assessment of naturally-occurring total iron and ^{57}Fe in mouse tissues. (A) Total tissue iron content in the duodenum, liver and kidney of 2 month-old wild-type mice, measured by the BPS-based colorimetric method or by ICP-MS. Data represent mean \pm SEM; n=10. (B) Total iron content in the liver of 2 month-old wild-type and Hfe-null mice measured by ICP-MS. Values are expressed as μg iron/g tissue. Data represent mean \pm SEM; n=5; ***=P<0.001. (C) Percentage of naturally-occurring ^{57}Fe in different wild-type tissues. Values are expressed as a percentage of ^{57}Fe with respect to total iron. Data represent mean \pm SEM; n=9.

Figure 2. Measurement of ^{57}Fe retention in the duodenal mucosa after oral treatment. (A) ^{57}Fe retention in the duodenal mucosa of wild-type mice measured by ICP-MS 30, 60 and 90 minutes after oral treatment with a solution containing 20 mmol/L ^{57}Fe . Control mice were treated with a vehicle solution and represented the “0” time point of the treatment in the experiment. Values are expressed as μg ^{57}Fe / g tissue. Data represent mean \pm SEM; n=9 for each experimental point; *=P<0.05, ***=P<0.001. (B) ^{57}Fe retention in the duodenal mucosa of wild-type and Hfe-null mice measured by ICP-MS 60 minutes after oral treatment with a solution containing 20 mmol/L ^{57}Fe . Control wild-type and Hfe-null mice were treated with a vehicle solution and represented the “0” time point of the treatment in the experiment. Values are expressed as μg ^{57}Fe / g tissue. Data represent mean \pm SEM; n=4 for each experimental point; *=P<0.05, **=P<0.01 (comparing control mice with the corresponding group of ^{57}Fe -treated mice), #=P<0.05 (comparing the two genotypes). (C) ^{57}Fe retention in the duodenal mucosa of wild-type mice fed a standard or an iron-free (IF) diet, measured by ICP-MS 60 minutes after oral treatment with a solution containing 20 mmol/L ^{57}Fe . Control mice

were treated with a vehicle solution and represented the “0” time point of the treatment in the experiment. Values are expressed as $\mu\text{g } ^{57}\text{Fe}/ \text{g tissue}$. Data represent mean \pm SEM; n=4 for each experimental point; ***=P<0.001 (comparing control mice with the corresponding group of ^{57}Fe -treated mice), ###=P<0.001 (comparing mice fed on the two different diets).

Figure 3. Determination of the amount of ^{57}Fe delivered to different organs. (A,C) ^{57}Fe retention in the liver (A) or in the kidney (C) of wild-type mice measured by ICP-MS 30, 60 and 90 minutes after oral treatment with a solution containing 20 mmol/L ^{57}Fe . Control mice were treated with a vehicle solution and represented the “0” time point of the treatment in the experiment. Values are expressed as $\mu\text{g } ^{57}\text{Fe}/ \text{g tissue}$. Data represent mean \pm SEM; n=9 for each experimental point; **=P<0.01. (B) ^{57}Fe retention in the bone marrow of wild-type mice measured by ICP-MS 48 hours after oral treatment with a solution containing 20 mmol/L ^{57}Fe . Control mice were treated with a vehicle solution and represented the “0” time point of the treatment in the experiment. Values are expressed as $\mu\text{g } ^{57}\text{Fe}/ \text{g tissue}$. Data represent mean \pm SEM; n=9 for each experimental point; *=P<0.05. (D) ^{57}Fe retention in the liver of wild-type and Hfe-null mice measured by ICP-MS 60 minutes after oral treatment with a solution containing 20 mmol/L ^{57}Fe . Control wild-type and Hfe-null mice were treated with vehicle solution and represented the “0” time point of the treatment in the experiment. Values are expressed as $\mu\text{g } ^{57}\text{Fe}/ \text{g tissue}$. Data represent mean \pm SEM; n=4 for each experimental point; *=P<0.05 (comparing control mice with the corresponding group of ^{57}Fe -treated mice), ###=P<0.001 (comparing the two genotypes). (E) ^{57}Fe retention in the liver of wild-type mice fed a standard or an iron-free (IF) diet measured by ICP-MS 60 minutes after oral treatment with a solution containing 20 mmol/L ^{57}Fe . Control mice were treated with a vehicle solution and represented the “0” time point of the treatment in the experiment. Values are expressed as $\mu\text{g } ^{57}\text{Fe}/ \text{g tissue}$. Data represent mean \pm SEM; n=4 for each

experimental point; **= $P < 0.01$ (comparing control mice with the corresponding group of ^{57}Fe -treated mice), ##= $P < 0.01$ (comparing mice fed on the two different diets).

Figure 1




Figure 2




Figure 3

