



Possibility of a Dark Matter Interpretation for the Excess in Isotropic Radio Emission Reported by ARCADE

N. Fornengo,^{1,2} R. Lineros,³ M. Regis,^{1,2} and M. Taoso³

¹*Dipartimento di Fisica Teorica, Università di Torino, I-10125 Torino, Italy*

²*Istituto Nazionale di Fisica Nucleare, Sezione di Torino, I-10125 Torino, Italy*

³*IFIC, CSIC–Universidad de Valencia, Edificios Institutos, Apartado Correos 22085, E46071 Valencia, Spain*

(Received 4 August 2011; revised manuscript received 20 October 2011; published 29 December 2011)

The ARCADE 2 Collaboration has recently measured an isotropic radio emission which is significantly brighter than the expected contributions from known extra-galactic sources. The simplest explanation of such excess involves a “new” population of unresolved sources which become the most numerous at very low (observationally unreached) brightness. We investigate this scenario in terms of synchrotron radiation induced by weakly interacting massive particle (WIMP) annihilations or decays in extra-galactic halos. Intriguingly, for light-mass WIMPs with a thermal annihilation cross section, the level of expected radio emission matches the ARCADE observations.

DOI: [10.1103/PhysRevLett.107.271302](https://doi.org/10.1103/PhysRevLett.107.271302)

PACS numbers: 95.35.+d, 95.85.Bh

The detection of a nongravitational signal of dark matter (DM) would be one of the greatest pillars of modern physics, simultaneously confirming our views of cosmology, astrophysics, and particle physics. This possibility might not be far from realization if DM is in the form of weakly interacting massive particles (WIMPs), which currently are the most investigated class of DM candidates [1]. Signatures of this scenario include a multiwavelength spectrum associated with radiative emissions involving electrons and positrons generated in WIMP annihilations or decays (for a recent review on this topic see, e.g., Ref. [2]).

Recently, the balloon-borne experiment ARCADE 2 (Absolute Radiometer for Cosmology, Astrophysics and Diffuse Emission) [3] reported radio measurements of the sky temperature at frequencies ranging from 3 to 90 GHz. Observations have been performed on a region which is roughly an annulus centered at $(l, b) = (70, 0)$ with a radius and width of 30 and 20 deg, respectively [4].

An isotropic component can be isolated from the ARCADE data by subtracting foreground Galactic emission [5]. Surprisingly, the level of the remaining flux (which has been interpreted in terms of extra-galactic sky temperature) is about 5–6 times larger than the total contribution from the extra-galactic radio sources detected in current surveys [6,7]. Even extrapolating the source number counts to lower (unreached) brightness, such excess still remains.

Most sources of systematic effects which could explain the ARCADE excess have been ruled out [5]. An astrophysical galactic origin appears to be rather unlikely (see discussions in Refs. [4,8]). Indeed, free-free emission has been excluded based on the spectral shape, and diffuse Galactic synchrotron foreground is estimated using two different methods (namely, a cosecant dependence on Galactic latitude and the correlation between radio

and atomic line emissions), which agree well with each other.

The observed isotropic temperature can be fitted by the cosmic microwave background (CMB) blackbody contribution plus a power law:

$$T(\nu) = T_0 \frac{h\nu/(kT_0)}{\exp[h\nu/(kT_0)] - 1} + T_s \left(\frac{\nu}{\text{GHz}} \right)^\alpha, \quad (1)$$

where $T_0 = 2.729 \pm 0.004$ K [5] is the CMB thermodynamic temperature. Performing analogous analyses on past surveys at 22, 45, 408, and 1420 MHz similar results are obtained, and fitting all data simultaneously the ARCADE Collaboration derived $\alpha = -2.62 \pm 0.04$ and $T_s = 1.19 \pm 0.14$ K [5].

Such a level of cosmic radio background does not have an immediate explanation in standard astrophysical scenarios. In Ref. [8], radio supernovae, radio quiet quasars, and diffuse emission from intergalactic medium and clusters (as well as a missed flux from well-known sources) have been considered, concluding that none of them can significantly contribute. A new population of numerous and faint radio sources (able to dominate source counts around μJy flux) has to be introduced [8,9]. Ordinary star-forming galaxies with a radio to far-infrared (FIR) flux ratio which increases significantly with redshift can in principle offer a solution. On the other hand, this possibility is strongly constrained by multiwavelength observations. Indeed, the radio to far-infrared emission has to be increased by a factor of 5 above what is observed in local galaxies [10], while current measurements show very mild evolution, at least up to $z \sim 2-3$ [12]. An explanation of the ARCADE excess through radiative emission of secondary electrons in star-forming galaxies would overproduce the gamma-ray background from pion decays [13]. The same is true also for primary electrons unless such putative galaxies have extremely low gas density (and, in turn,

low ratio of primary electrons to pions) or extremely efficient proton escape. The picture that seems to emerge from ARCADE measurements [5,8] and subsequent interpretations [8,9,12,13] suggests the need for a population of numerous and faint synchrotron sources generated by primary electrons with a hard spectrum and with no or very faint correlated mechanisms at infrared and gamma-ray frequencies.

In our current understanding of structure clustering, any luminous source is embedded in a DM halo, and therefore extra-galactic DM halos can be seen as the most numerous source population. The flux induced by WIMP annihilations or decays is predicted to be very faint. It is associated with primary electrons and positrons generated as the final state of annihilation or decay, and WIMP models with large annihilation or decay branching ratios into leptons induce hard spectra of e^+/e^- with very faint gamma-ray counterpart (and, of course, no straightforward thermal emission). Therefore, WIMP sources represent an ideal candidate to fit the ARCADE excess, and in this Letter we quantitatively investigate such a possibility.

Assuming a one-to-one relation between the mass M of extra-galactic DM halos and the intrinsic luminosity \mathcal{L} of the source, the total isotropic intensity per solid angle at a given frequency ν is given by (for a more detailed derivation of equations considered in the following, see, e.g., Refs. [14,15])

$$\nu I_\nu = \frac{c\nu}{4\pi} \int \frac{dz}{(1+z)H(z)} \int_{M_c} dM \frac{dn}{dM}(M, z) \mathcal{L}(E, z, M), \quad (2)$$

where z is the redshift, H is the Hubble rate, M_c is the minimum mass of an emitting halo, and the luminosity \mathcal{L} is a function of the redshifted energy $E = E_\nu(1+z)$ with $E_\nu = h\nu$. The luminosity function, including also the contribution of substructures within the DM halo, can be written as

$$\mathcal{L} = (1-f)^a \int_0^{R_v} d^3r \frac{d\hat{N}_i}{dE} + \int dM_s \frac{dn_s}{dM_s} \int_0^{R_v} d^3r \frac{d\hat{N}_i}{dE}, \quad (3)$$

where R_v is the virial radius of the DM profile ρ and f is the fraction of halo mass in substructures (with $a = 1$ and $a = 2$ for decaying and annihilating DM, respectively). In Eqs. (2) and (3), $dn/dM(M)$ and $dn_s/dM_s(f, M_s, M)$ denote the mass function of the DM halo and of substructures, respectively. For synchrotron emission,

$$\frac{d\hat{N}_i}{dE_\nu} = 2 \int_{m_e}^{M_\chi} dE' P_{\text{syn}}(\nu, B, E') n_e, \quad (4)$$

where m_e is the electron mass, P_{syn} is the synchrotron power [16], B denotes the magnetic field, and ν is the frequency of emission [as opposed to frequency of observation in Eq. (2)]. The electron or positron equilibrium

number density n_e is obtained solving a transport equation for e^-/e^+ injected by DM with an energy spectrum set by dN_e/dE_e [15]. The source terms of this equation for annihilating and decaying DM are

$$Q_a = \frac{(\sigma_a v)}{2M_\chi^2} \rho^2 \frac{dN_e}{dE_e}, \quad Q_d = \frac{\rho}{\tau_d M_\chi} \frac{dN_e}{dE_e}, \quad (5)$$

where M_χ is the mass of the DM particle, $(\sigma_a v)$ is the nonrelativistic annihilation cross section, and τ_d the decay rate.

For what concerns the ‘‘astrophysical’’ parameters, we focus on two benchmark cases, which are fairly realistic (for a detailed discussion of impact of astrophysical uncertainties, see Ref. [15]). We adopt the halo mass function dn/dM from Ref. [17], recent N -body simulation results for concentration of halos [18], a DM distribution inside halos following a Navarro-Frenk-White (NFW) profile [19], and the minimum halo mass is set to $M_c = 10^6 M_\odot$ (model A, in order to consider only objects for which we can guess a reasonably large magnetic field), and to $M_c = 10^{-6} M_\odot$ (model B). The contribution from substructures is modeled such that $f = 10\%$ of the total mass is in substructures and $dn_s/d\ln(M_s) \propto 1/M_s$, which leads to a boost in the signal of $bf_{\text{sub}} \simeq 7$ (model A) and no boost in model B. The magnetic field is assumed to be constant in space and time with magnitude $B = 10 \mu\text{G}$ (model A) and $B = 2 \mu\text{G}$ (model B), and e^+/e^- are assumed to radiate at the same place where they are injected. [Notice that for the DM candidate of specific interest here (i.e., light or inducing a soft spectrum of e^+/e^-), electrons are mostly emitted at GeV energies, namely, are injected at energies relevant for radiation at GHz frequencies. Therefore they do not travel significant distances before radiating, while electrons emitted at larger energies would take some time to cool down, and diffusion and escape time would become much more relevant.] The normalization of the emission roughly decreases by an order of magnitude going from $B = 10 \mu\text{G}$ to $B = 1 \mu\text{G}$, increases by 2 orders of magnitude going from $M_c = 10^6 M_\odot$ to $M_c = 10^{-6} M_\odot$, scales linearly with bf_{sub} , and is mildly dependent on the halo mass function, concentration, and profile of DM. (For example, the case of an isothermal cored profile leads to a reduction in the intensity by, roughly, a factor of 1.5.)

The excess spectrum reported by ARCADE and described in Eq. (1) is rather hard, and requires a hard electron or positron spectrum dN_e/dE_e . This can be produced by DM scenarios with a large branching ratio of annihilation or decay into leptons. For illustrative purposes, we chose the $\mu^+ - \mu^-$ channel. To reproduce the absolute normalization of the excess with a ‘‘thermal’’ annihilation rate $(\sigma_a v) = 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ in our benchmark model A, we need a WIMP with $M_\chi = 10 \text{ GeV}$ ($M_\chi = 25 \text{ GeV}$ in model B), which, even though it induces a slightly softer spectrum than the best-fit power law, provides a reasonable agreement with the

data, as shown in Fig. 1. For this benchmark case we have $\chi^2/\text{d.o.f.} = 26.9/13$. The actual best fit for model A is obtained with a mass $M_\chi \sim 30$ GeV, ($\chi^2/\text{d.o.f.} = 14.3/13$), but at the price of increasing the cross section by 1 order of magnitude. The fact that light DM, in the 10 GeV mass range, can fairly well reproduce the ARCADE excess, without the need of unrealistically large DM overdensities, is particularly interesting, especially in light of recent claims of signals compatible with a DM interpretation from direct detection experiments (DAMA [20], CoGeNT [21], and CRESST [22]), that can in fact be accommodated with a ~ 10 GeV WIMP [23]. In the case of the ARCADE excess, the best option to explain the effect in terms of DM annihilation requires a light DM particle which annihilates mainly into leptons, and therefore that does not couple dominantly to quarks (coupling relevant to the direct detection scattering cross section). Nevertheless, it is not very difficult, from the model-building point of view, to foresee a model where a DM candidate, which annihilates mainly into leptons, still has a relatively large scattering cross section off the nuclei. For a concrete example, see, e.g., Ref. [24]. Note also that the radio emission in the Milky Way halo induced by WIMPs' fitting ARCADE data can either easily satisfy constraints (for cored galactic DM profiles) or be close to a possible detection in the central region of our Galaxy (for cuspy profiles) [25].

Similar conclusions on the viability of a light “leptophilic” DM particle in explaining the ARCADE data can




FIG. 1 (color online). Extra-galactic radio background as derived by ARCADE [5], together with three possible interpretations of the low-frequency (< 10 GHz) excess in terms of WIMP annihilations (blue, green, and orange curves, see text for details). The astrophysical source contribution estimated from number counts (red line), the CMB contribution (black dotted line), and a best-fit power law of the excess (black dashed line) are also reported [6].

also be drawn in the decaying case: for a DM mass of $M_\chi = 10$ GeV, the excess is reproduced if the lifetime is $\tau_d = 3 \times 10^{27}$ s (with a curve similar to the one shown for the annihilating case).

Since the ARCADE excess can be explained by DM annihilation or decay in terms of sizable production of electrons and positrons, emissions of x rays and gamma rays by means of inverse-Compton (IC) processes on interstellar radiation fields (here we include CMB only) and direct production of gamma rays from the DM particle annihilation [either by production of neutral pions or by final state radiation (FSR)] are present and have to be checked against available bounds. For the benchmark cases considered, these multiwavelength constraints are easily satisfied, as shown for x rays and γ rays in Fig. 2.

In Fig. 1, we also show the case of a more “classic” WIMP candidate with 100 GeV mass and hadronic annihilation channel ($b\bar{b}$ pair in the shown benchmark). This scenario is less appealing than previous cases. The spectrum is relatively too soft in order to reproduce well the ARCADE data and the fit is worse ($\chi^2/\text{d.o.f.} = 49.5/13$) than for DM annihilating into leptons; note that the excess is sizable up to at least 3 GHz (although not clearly visible in Fig. 1 due the smallness of the error bars and the scale of the plot), so a viable explanation has to roughly overlap the dashed best-fit curve up to those frequencies. Moreover, since now the DM mass is larger, the required boost factor is accordingly larger (by a factor of 20 in this specific case), which can stem from a larger annihilation cross section related to a nonstandard formation of DM relic density, from a Sommerfeld enhancement, or from a larger




FIG. 2 (color online). X-ray and gamma-ray fluxes for the three benchmark WIMP cases shown in Fig. 1. The CHANDRA [32] bound in the x-ray band and the COMPTTEL [33] and FERMI [34] extra-galactic gamma-ray fluxes are shown.




FIG. 3 (color online). Differential number counts for AGNs (dashed line), star-forming galaxies (dotted line), and the same 10 GeV benchmark DM model (solid lines) shown in Figs. 1 and 2. For DM, we consider a case such that all substructures are resolved, and an opposite case where all substructures are unresolved. For data and astrophysical models, see [11] and references therein.

contribution of substructures with respect to what is considered here (see, e.g., [26] for further details on possible boost factors). Heavier DM with hadronic annihilation or decay final states is also more strongly constrained by the γ -ray channel, as can be seen in Fig. 2.

As a further analysis on the radio emission arising from light DM annihilation or decay, able to adapt to the ARCADE excess, we show in Fig. 3 the differential number counts of sources at 1.4 GHz. If we assume all substructures to be unresolved, they mainly boost the signal of large and bright halos (since the latter host more subhalos). On the contrary, if all substructures are assumed to be resolved, counts drop much more slowly at low brightness. To highlight uncertainties related to the possibility of resolving substructures in the future, these two extreme cases are shown in Fig. 3 as solid lines, for the same 10 GeV DM benchmark of Fig. 1. As discussed above, the key point for our analysis is that in both scenarios the number of DM sources definitely becomes dominant over astrophysical contributions (AGN, star-forming galaxies) at the sub- μ Jy level. The contribution of star-forming galaxies, which is dominant over AGN emission at low fluxes, decreases more rapidly (assuming FIR-radio correlation holds at all redshift) than the expected contribution from DM, in both resolved and unresolved substructures. From Fig. 3 we notice that the flattening at low brightness in current data, although it can be easily accounted for by standard astrophysical populations (for a review see, e.g., Ref. [11]), nevertheless could be well fitted by a DM model between the two extreme cases presented in Fig. 3.

In summary, we discussed the possibility that synchrotron emission induced by WIMP annihilations can account for the isotropic radio component measured by the ARCADE 2 Collaboration. Although galactic or extra-galactic astrophysical interpretations of the excess cannot be excluded, they currently present some puzzling issues [4,6,8]. Under reasonable assumptions for clustering, we found that light-mass WIMPs producing hard-spectrum electrons and positrons (as in the case of leptonic annihilation channels) in extra-galactic halos with a “thermal” annihilation rate can fit the excess and satisfy constraints at other wavelengths. A population of sources which can generally explain ARCADE measurements has to become the most numerous at brightness around μ Jy, so it will certainly be studied in detail by SKA [27], and possibly also by its precursors, ASKAP [28] and MeerKAT [29]. If the excess is due to extra-galactic DM, a clear discovery of a nongravitational signal of DM might not be far away. A dedicated study of closest and brightest (in terms of DM-induced signal) objects with current radio telescopes (e.g., ATCA [30] and EVLA [31]) can start to probe this scenario in the near future.

We acknowledge research grants funded jointly by Ministero dell’Istruzione, dell’Università e della Ricerca (MIUR), by Università di Torino, and by Istituto Nazionale di Fisica Nucleare within the *Astroparticle Physics Project* (MIUR Contract No. PRIN 2008NR3EBK; INFN Grant No. FA51). R.L. and M.T. were supported by the EC contract UNILHC PITN-GA-2009-237920, by the Spanish grants FPA2008-00319 and MultiDark CSD2009-00064 (MICINN), and PROMETEO/2009/091 (Generalitat Valenciana). N.F. acknowledges support of the Spanish MICINN Consolider Ingenio 2010 Programme under Grant No. MULTIDARK CSD2009-00064. This work was partly completed at the Theory Division of CERN in the context of the TH-Institute “Dark Matter Underground and in the Heavens” (DMUH11). R.L. and M.T. are supported by a MultiDark Fellowship.

-
- [1] For reviews see, e.g., G. Jungman, M. Kamionkowski, and K. Griest, *Phys. Rep.* **267**, 195 (1996).
 - [2] S. Profumo and P. Ullio, [arXiv:1001.4086](https://arxiv.org/abs/1001.4086).
 - [3] J. Singal *et al.*, *Astrophys. J.* **730**, 138 (2011).
 - [4] A. Kogut *et al.*, *Astrophys. J.* **734**, 4 (2011).
 - [5] D.J. Fixsen *et al.*, [arXiv:0901.0555](https://arxiv.org/abs/0901.0555).
 - [6] M. Seiffert *et al.*, [arXiv:0901.0559](https://arxiv.org/abs/0901.0559).
 - [7] M. Gervasi *et al.*, *Astrophys. J.* **682**, 223 (2008).
 - [8] J. Singal *et al.*, *Mon. Not. R. Astron. Soc.* **409**, 1172 (2010).
 - [9] T. Vernstrom *et al.*, [arXiv:1102.0814](https://arxiv.org/abs/1102.0814).
 - [10] Radio synchrotron sources are associated with young stars and therefore strongly correlated with the star formation rate of galaxies. Empirically, a nearly linear far IR-radio correlation is reasonably well established (at least for high

- brightness and low redshift); see, e.g., the review in Ref. [11].
- [11] G. De Zotti *et al.*, *Astron. Astrophys. Rev.* **18** (2010).
- [12] P.P. Ponente *et al.*, *Mon. Not. R. Astron. Soc.* **418**, 691 (2011).
- [13] B. C. Lacki, *Astrophys. J.* **729**, L1 (2011).
- [14] L. Zhang and G. Sigl, *J. Cosmol. Astropart. Phys.* **09** (2008) 027.
- [15] N. Fornengo, R. Lineros, M. Regis, and M. Taoso (to be published).
- [16] G. Rybicki and A.P. Lightman, *Radiative Processes in Astrophysics* (John Wiley & Sons, New York, 1979).
- [17] R.K. Sheth and G. Tormen, *Mon. Not. R. Astron. Soc.* **308**, 119 (1999).
- [18] J.C. Munoz-Cuartas *et al.*, arXiv:1007.0438.
- [19] J.F. Navarro, C.S. Frenk, and S.D.M. White, *Astrophys. J.* **490**, 493 (1997).
- [20] R. Bernabei *et al.*, *Eur. Phys. J. C* **56**, 333 (2008); *Eur. Phys. J. C* **67**, 39 (2010).
- [21] C.E. Aalseth *et al.*, *Phys. Rev. Lett.* **107**, 141301 (2011).
- [22] G. Angloher *et al.*, arXiv:1109.0702.
- [23] T. Schwetz and J. Zupan, *J. Cosmol. Astropart. Phys.* **08** (2011) 008; R. Foot, *Phys. Lett. B* **703**, 7 (2011); P.J. Fox *et al.*, arXiv:1107.0717; D. Hooper and C. Kelso, *Phys. Rev. D* **84**, 083001 (2011); M. Farina *et al.*, arXiv:1107.0715; E. Del Nobile, C. Kouvaris, and F. Sannino, *Phys. Rev. D* **84**, 027301 (2011); M.T. Frandsen *et al.*, *Phys. Rev. D* **84**, 041301 (2011); G.B. Gelmini, arXiv:1106.6278; P. Belli *et al.*, *Phys. Rev. D* **84**, 055014 (2011).
- [24] M. S. Boucenna and S. Profumo, *Phys. Rev. D* **84**, 055011 (2011).
- [25] N. Fornengo, R. Lineros, M. Regis, and M. Taoso, arXiv:1110.4337.
- [26] J. Lavalley *et al.*, *Astron. Astrophys.* **479**, 427 (2008); M. Lattanzi and J. Silk, *Phys. Rev. D* **79**, 083523 (2009); M. Schelke *et al.*, *Phys. Rev. D* **74**, 083505(2006).
- [27] S. Rawlings, R. Schilizzi, arXiv:1105.5953; see also <http://www.skatelescope.org/>.
- [28] S. Johnston *et al.* (ASKAP Collaboration), *Pub. Astron. Soc. Aust.* **24**, 174 (2007); see also <http://www.atnf.csiro.au/projects/askap/>.
- [29] R.S. Booth *et al.*, arXiv:0910.2935; see also <http://www.ska.ac.za/meerkat/>.
- [30] <http://www.narrabri.atnf.csiro.au/>
- [31] <https://science.nrao.edu/facilities/evla>
- [32] R.C. Hickox and M. Markevitch, *Astrophys. J.* **661**, L117 (2007).
- [33] G. Weidenspointner *et al.*, *AIP Conf. Proc.* **510**, 467 (2000).
- [34] A.A. Abdo *et al.* (Fermi-LAT Collaboration), *Phys. Rev. Lett.* **104**, 101101 (2010).