

HEAVY QUARKS IN NUCLEUS–NUCLEUS
COLLISIONS: FROM RHIC TO LHC*A. BERAUDO^{a,b}, W.M. ALBERICO^{c,d}, A. DE PACE^d, A. MOLINARI^{c,d}
M. MONTENO^d, M. NARDI^d, F. PRINO^d^aCentro Studi e Ricerche “Enrico Fermi”, Piazza del Viminale 1, Roma, Italy^bPhysics Department, Theory Unit, CERN, 1211 Genève 23, Switzerland^cDipartimento di Fisica Teorica dell’Università di Torino
via P. Giuria 1, 10125 Torino, Italy^dIstituto Nazionale di Fisica Nucleare, Sezione di Torino
via P. Giuria 1, 10125 Torino, Italy*(Received July 26, 2011)*

We present a study of the heavy-flavor dynamics in nucleus–nucleus collisions. The initial (hard) production of c and b quarks is taken from NLO pQCD predictions. The presence of a hot medium (a Quark-Gluon Plasma described by hydrodynamics) affects the final spectra of open-charm (beauty) hadrons and their decay electrons with respect to what found in pp collisions. The propagation of c and b quarks in the plasma is based on a picture of multiple uncorrelated random collisions, described by a relativistic Langevin equation. A microscopic evaluation of the transport coefficients is provided within a pQCD approach (with proper resummation of medium effects). Results for the final spectra of heavy-flavor hadrons and decay-electrons are given, with particular emphasis on R_{AA} and v_2 .

DOI:10.5506/APhysPolBSupp.4.653

PACS numbers: 12.38.Mh, 25.75.Cj, 11.10.Wx

1. Introduction

Heavy flavor spectra are an interesting probe of the medium formed in heavy-ion collisions: c and b quarks are produced in hard processes during the crossing of the two nuclei and the QCD flavor conservation allows to “tag” final-state charm and beauty hadrons, so that the color charge and mass of the parent parton propagating in the plasma is known. The interaction between the heavy quarks and the medium leads to a modification

* Presented at the Workshop “Excited QCD 2011”, Les Houches, France, February 20–25, 2011.

of the final spectra with respect to the pp case. Different mechanisms and approaches were proposed in the literature to account for such an effect: *e.g.* medium-induced gluon-radiation [1, 2], scattering mediated by resonant states [3], calculations based on the Boltzmann equation [4]. As explained at length in [5, 6] we describe the heavy-quark dynamics through the relativistic Langevin equation. This amounts to assume that the heavy-quark momentum, crossing the plasma, gets changed due to multiple uncorrelated random collisions. For independent studies based on the Langevin equation see for instance Refs. [7, 8, 9].

2. The relativistic Langevin equation

The relativistic Langevin equation [10] is used to study the propagation of c and b quarks in the QGP. The variation of the heavy-quark momentum in the time-interval Δt

$$\frac{\Delta p^i}{\Delta t} = -\eta_D(p)p^i + \xi^i(t), \quad (2.1)$$

is given by the sum of a *deterministic* friction term and a *stochastic* noise term $\xi^i(t)$, which is completely determined by its two-point temporal correlator

$$\langle \xi^i(t)\xi^j(t') \rangle = b^{ij}(\mathbf{p})\delta(t-t'), \quad \text{with } b^{ij}(\mathbf{p}) \equiv \kappa_L(p)\hat{p}^i\hat{p}^j + \kappa_T(p)(\delta^{ij} - \hat{p}^i\hat{p}^j). \quad (2.2)$$

The latter involves the transport coefficients $\kappa_T(p) \equiv \frac{1}{2} \frac{\langle \Delta p_T^2 \rangle}{\Delta t}$ and $\kappa_L(p) \equiv \frac{\langle \Delta p_L^2 \rangle}{\Delta t}$, representing the average transverse and longitudinal squared-momentum acquired per unit time by the quark due to the collisions in the medium.

The transport coefficients $\kappa_{T/L}$ are evaluated according to the procedure presented in [5]. We introduce an intermediate cutoff $|t|^* \sim m_D^2$ ($t \equiv (P'-P)^2$) separating hard and soft scatterings. The contribution of hard collisions ($|t| > |t|^*$) is evaluated through a kinetic pQCD calculation of the processes $Q(P)q_{i/\bar{i}} \rightarrow Q(P')q_{i/\bar{i}}$ and $Q(P)g \rightarrow Q(P')g$. On the other hand, in soft collisions ($|t| < |t|^*$) the exchanged gluon feels the presence of the plasma. A resummation of medium effects is thus required and this is provided by the Hard Thermal Loop approximation. The final result is given by the sum of the two contributions $\kappa_{T/L}(p) = \kappa_{T/L}^{\text{hard}}(p) + \kappa_{T/L}^{\text{soft}}(p)$ and its explicit expression can be found in Ref. [5]. In Fig. 1 we display the behavior of $\kappa_{T/L}$ for c and b quarks. The sensitivity to the intermediate cutoff $|t|^*$ is quite small, supporting the validity of the approach.




Fig. 1. The c (left panel) and b (right panel) momentum diffusion coefficients $\kappa_{T/L}(p)$ resulting from the sum of the soft and hard contributions. The sensitivity to the intermediate cutoff $|t|^* \sim m_D^2$ is very mild. The curves refer to the temperature $T = 400$ MeV, with the coupling g evaluated at the scale $\mu = (3/2)\pi T$.

3. Heavy flavor in pp and heavy-ion collisions

The procedure to generate final-state heavy-flavor spectra can be factorized into some independent steps. Details can be found in [5].

- *Initial production of $Q\bar{Q}$ pairs in hard pQCD processes.* An initial sample of c and b quarks is generated using the POWHEG code [11], which implements pQCD at NLO. In the AA case nuclear PDFs are employed [12]. Heavy quarks are then distributed in the transverse plane according to the nuclear overlap function $dN/d\mathbf{x}_\perp \sim T_{AB}(x, y) \equiv T_A(x+b/2, y)T_B(x-b/2, y)$ and given a k_T -broadening depending on the crossed thickness of nuclear matter.
- *Langevin evolution in the fireball.* In the AA case at the proper-time $\tau \equiv \sqrt{t^2 - z^2} = \tau_0$ one starts following the stochastic Langevin dynamics of the quarks in the plasma. The expansion of the background medium is described by ideal/viscous hydrodynamics [13, 14, 15].
- *Hadronization and decays.* Heavy quarks are made hadronize (around the critical energy density ϵ_c in the AA case) using Peterson fragmentation functions [16] with branching fractions into hadrons taken from Refs. [17, 18]. Finally, each hadron is forced to decay into electrons with PYTHIA [19], using updated tables of branching ratios [20].

4. Numerical results

Our predictions for the cross-sections of heavy-flavor electrons in pp collisions at RHIC ($\sqrt{s} = 200$ GeV) and LHC ($\sqrt{s} = 5.5$ TeV) are given in Fig. 2. PHENIX results are well reproduced. Going from RHIC to LHC one observes a sizable increase of the cross-section, a hardening of the spectrum and the growth of the relative contribution of electrons from b decays.




Fig. 2. Invariant differential cross-section of electrons $[(e^+ + e^-)/2]$, from heavy-flavor decay in pp collisions at $\sqrt{s} = 200$ GeV (left panel) and $\sqrt{s} = 5.5$ TeV (right panel). The curves refer to electrons from c or b quarks generated by POWHEG. PHENIX data are extremely well reproduced.




Fig. 3. The R_{AA} of heavy-flavor hadrons and decay-electrons in minimum-bias AA collisions at RHIC (0–92% of the total hadronic cross-section, left panel) and LHC at 5.5 TeV (0–90%, right panel) for viscous hydrodynamics.

For AA collisions effects of the Langevin evolution were studied through the nuclear modification factor $R_{AA}(p_T) \equiv (dN/dp_T)^{AA} / \langle N \rangle_{\text{coll}} (dN/dp_T)^{pp}$ and the elliptic flow coefficient $v_2(p_T) \equiv \langle \cos(2\phi) \rangle_{p_T}$. Here, we display results obtained using viscous hydrodynamics, for few representative cases among the ones explored in [5]. In Fig. 3 we show our findings for the $R_{AA}(p_T)$ of heavy-flavor hadrons and their decay-electrons at RHIC and LHC at 5.5 TeV. A general agreement with PHENIX data [21] can be attained. Notice that the higher temperatures reached at LHC lead to a larger quenching of the spectra; however in the inclusive ($e_c + e_b$) result the effect is partially compensated by the stronger weight acquired by the b contribution. In Fig. 4 the centrality dependence of R_{AA} is displayed. The overall agreement with PHENIX data looks satisfactory.




Fig. 4. The nuclear modification factor of heavy-flavor electrons at RHIC (a) and LHC at 5.5 TeV (b) for viscous hydrodynamics ($\tau_0 = 1$ fm), obtained by integrating the electron yields over the indicated momentum ranges, as a function of N_{part} . The centrality dependence found by the PHENIX experiment [21] is nicely reproduced.




Fig. 5. (a) The minimum-bias anisotropy parameter v_2 for heavy-flavor electrons at RHIC for viscous hydrodynamics. PHENIX data [21,22] are compared with the outcomes of our calculations for electrons from c and b quarks and their combination. (b) As in panel (a), but for LHC and two choices of τ_0 .

In Fig. 5 we display our results for v_2 . Even assuming a very rapid thermalization, our findings tend to underestimate the experimental data: perturbative cross-sections favor small-angle scattering. However, in the p_T regime covered by the experimental measurements hadronization could play an important role; coalescence with quarks of the medium, displaying a sizable elliptic flow, could contribute to enhance the heavy-flavor v_2 [3].

5. Conclusions

We studied the dynamics of c and b quarks in the medium produced in heavy-ion collisions. In our approach, based on the relativistic Langevin equation, the quarks interact with the medium through multiple uncorrelated random momentum-exchanges, which would asymptotically drive them to thermal equilibrium. The heavy-quark transport coefficients were evaluated within a weak-coupling framework. Our procedure leads to results in reasonable agreement with the available experimental data, which extend up to quite high p_T . This could even suggest to reconsider the role of collisional energy-loss in the suppression of light-hadron spectra.

REFERENCES

- [1] M. Djordjevic, M. Gyulassy, *Nucl. Phys.* **A733**, 265 (2004).
- [2] N. Armesto, C.A. Salgado, U.A. Wiedemann, *Phys. Rev.* **D69**, 114003 (2004).
- [3] H. van Hees, V. Greco, R. Rapp, *Phys. Rev.* **C73**, 034913 (2006).
- [4] J. Uphoff *et al.*, arXiv:1104.2437 [hep-ph].
- [5] W.M. Alberico *et al.*, *Eur. Phys. J.* **C71**, 1666 (2011) [arXiv:1101.6008 [hep-ph]].
- [6] W.M. Alberico *et al.*, *J. Phys. Conf. Ser.* **270**, 012029 (2011) [arXiv:1007.4170 [hep-ph]]; arXiv:1009.2434 [hep-ph]. *Nucl. Phys.* **A855**, 404 (2011) [arXiv:1011.0400 [hep-ph]].
- [7] G.D. Moore, D. Teaney, *Phys. Rev.* **C71**, 064904 (2005).
- [8] P.B. Gossiaux, R. Bierkanndt, J. Aichelin, *Phys. Rev.* **C79**, 044906 (2009).
- [9] Y. Akamatsu, T. Hatsuda, T. Hirano, *Phys. Rev.* **C79**, 054907 (2009).
- [10] A. Beraudo *et al.*, *Nucl. Phys.* **A831**, 59 (2009).
- [11] S. Frixione, G. Ridolfi, P. Nason, *J. High Energy Phys.* **0709**, 126 (2007).
- [12] K.J. Eskola, H. Paukkunen, C.A. Salgado, *J. High Energy Phys.* **0904**, 065 (2009).
- [13] P.F. Kolb, J. Sollfrank, U. Heinz, *Phys. Rev.* **C62**, 054909 (2000).
- [14] P. Romatschke, U. Romatschke, *Phys. Rev. Lett.* **99**, 172301 (2007).

- [15] M. Luzum, P. Romatschke, *Phys. Rev.* **C78**, 034915 (2008).
- [16] C. Peterson *et al.*, *Phys. Rev.* **D27**, 105 (1983).
- [17] http://www-zeus.desy.de/zeus_papers/ZEUS_PAPERS/DESY-05-147.ps
- [18] http://www.slac.stanford.edu/xorg/hfag/osc/PDG_2009/#FRAC
- [19] T. Sjostrand, S. Mrenna, P.Z. Skands, *J. High Energy Phys.* **0605**, 026 (2006).
- [20] K. Nakamura *et al.* [Particle Data Group], *J. Phys. G* **37**, 075021 (2010).
- [21] A. Adare *et al.* [PHENIX Collaboration], [arXiv:1005.1627](https://arxiv.org/abs/1005.1627) [nucl-ex].
- [22] A. Adare *et al.* [PHENIX Collaboration], *Phys. Rev. Lett.* **98**, 172301 (2007).