

The logo for IRIS AperTO, featuring the text "IRIS" in a large, white, serif font, "Aper" in a smaller, white, sans-serif font, and "TO" in a large, white, serif font, all set against a red rectangular background.

UNIVERSITÀ  
DEGLI STUDI  
DI TORINO

This is the author's final version of the contribution published as:

Garibaldi A.; Bertetti D.; Poli A.; Gullino M.L.. Powdery mildew  
(*Golovinomyces orontii*) on Creeping Bellflower (*Campanula rapunculoides*)  
in Italy.. PLANT DISEASE. 96 pp: 291-291.

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/128399>

This full text was downloaded from iris - AperTO: <https://iris.unito.it/>

---

iris - AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

**Powdery mildew (*Golovinomyces orontii*) on Creeping Bellflower (*Campanula rapunculoides*) in Italy.** A. Garibaldi, D. Bertetti, A. Poli and M.L. Gullino, Centre of Competence AGROINNOVA, University of Torino, Via Leonardo da Vinci 44, 10095 Grugliasco, Italy.

Creeping (June) bellflower (*Campanula rapunculoides*) is an herbaceous plant belonging to the Campanulaceae family. It has showy flowers, very much appreciated for gardens and landscaping. During the summer 2011, 6-9-month-old plants grown in a garden near Biella (northern Italy) showed signs and symptoms of an unknown powdery mildew. The adaxial leaf surfaces were covered with white mycelia and conidia, while the abaxial surfaces were less infected. As the disease progressed, infected leaves turned yellow and wilted. Mycelia were also observed on stems, petioles and flower calyxes of inflorescences. Seventy percent of plants were diseased. Conidia were hyaline, elliptical to ovoid (sometimes doliform), borne in short chains (up to 3 conidia per chain), and measured 27 to 42 (34) × 16 to 24 (19) µm. Conidiophores were erect, with a cylindrical foot cell measuring 64 to 105 (80) × 11 to 12 (11) µm, followed by two shorter cells, measuring 17 to 24 (20) × 11 to 15 (13) µm. Fibrosin bodies were absent. Chasmothecia were not observed in the collected samples. The Internal Transcribed Spacer (ITS) region of rDNA was amplified using the primers ITS1F/ITS4 and sequenced (3) (GenBank Accession No. JN639855). The 405-bp amplicon had 98% homology with the sequence of *Golovinomyces orontii* GQ183948. Pathogenicity was confirmed through inoculation by gently pressing diseased leaves onto leaves of healthy *C. rapunculoides* plants. Three plants were inoculated, while the same number of non-inoculated plants served as a control. Plants were maintained outside at temperatures from 10 to 26°C. Fifteen days after inoculation, symptoms and signs of powdery mildew developed on inoculated plants. The conidial morphology of the powdery mildew fungus developed on inoculated plants was identical to the conidial morphology observed in the original fungus. Non-inoculated plants remained healthy. The pathogenicity test was carried out twice. *Golovinomyces orontii*, has been reported on *C. rapunculoides* in several eastern European countries as well as in Switzerland and Germany (1,2). This is the first report of the disease in Italy. The economic importance of this disease is at present limited in Italy, due to limited planting of this host.

*References:* (1) A. Bolay. *Crytog. Helv.*, 20:1, 2005. (2) U. Braun. *The Powdery Mildews (Erysiphales) of Europe*. Gustav Fischer Verlag, 337 pp, 1995. (3) T. J. White *et al.* *PCR Protocols: A Guide to Methods and Applications*. M. A. Innis *et al.*, eds. Academic Press, San Diego, 1990.