

IRIS A_{per}TO



UNIVERSITÀ
DEGLI STUDI
DI TORINO

This is the author's final version of the contribution published as:

Garibaldi A.; Bertetti D.; Poli A.; Gullino M.L.. First report of Fruit rot in pear caused by *Botryosphaeria dothidea* (Moug. : Fr.) Ces. & De Not. in Italy.. PLANT DISEASE. 96 pp: 910-910.

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/130119>

This full text was downloaded from iris - AperTO: <https://iris.unito.it/>

iris - AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

1 **First Report of Postharvest Fruit Rot in Pear Caused by *Botryosphaeria dothidea* (moug. Ex**
2 **Fr.) Ces. & De Not in Italy.** A. Garibaldi, D. Bertetti, A. Poli, and M. L. Gullino, Centre of
3 Competence for the Innovation in the Agro-Environmental Sector (AGROINNOVA) Via
4 Leonardo da Vinci 44, 10095 Grugliasco, Italy.

5
6 Pear (*Pyrus communis* L.) is widely grown in Italy, the leading producer in Europe. In summer
7 2011, a previously unknown rot was observed on fruit of an old variety “Spadoncina” in a garden
8 in Torino Province (northern Italy). The decayed area of fruit was soft, brown, slightly sunken,
9 surrounded by a margin irregular and circular. The internal decayed area appeared rotten and
10 brown. Rotted fruit eventually fell down. Fragments (approximately 2 mm) were taken from the
11 margin of the internal diseased tissues, cultured on potato dextrose agar (PDA) and incubated at
12 temperatures between 20-28°C, under alternating light and darkness. Colonies of the fungus
13 initially appeared whitish, then turning to dark gray and produced a dark pigment into the
14 medium. After 25 days of growth, unicellular fusiform to elliptical hyaline conidia were
15 produced. Conidia had a slightly obtuse apex and a truncated base and measured 16-24 x 5-7
16 (average 20.1 x 5.7) µm (length to width ratios were 2.8 to 4.6 with average of 3.5). The
17 morphological characteristics are similar to that of the fungus *Botryosphaeria dothidea* (4). The
18 Internal Transcribed Spacer (ITS) region of rDNA was amplified using the primers ITS1/ITS4,
19 and sequenced. BLAST analysis (1) of the 473 bp segment showed a 100% similarity with the
20 sequence of *B. dothidea* (GeneBank accession FM955378). The nucleotide sequence has been
21 assigned the GenBank Accession JQ418493. Pathogenicity tests were performed by inoculating
22 six pear fruits of the same cultivar after surface-disinfecting in 1% sodium hypochlorite and
23 wounding. Mycelial disks (8 mm diameter), obtained from PDA cultures of one strain, were

1 placed on wounds. Six control fruits were inoculated with plain PDA. Fruits were incubated at
2 $25\pm 1^{\circ}\text{C}$. The first symptoms developed 2 days after the artificial inoculation. After 5 days, the
3 rot was very evident and *B. dothidea* was consistently reisolated. Non-inoculated fruit remained
4 healthy. The pathogenicity test was performed twice. *B. dothidea* was identified on *P. communis*
5 in the US (2), South Africa, New Zealand and Japan (3). To our knowledge, this is the first report
6 of the presence of *B. dothidea* on pear in Italy, as well as in Europe. In Italy, the economic
7 importance of the disease on pear fruit is at present limited, although the pathogen could
8 represent a risk for this crop.

9

10 References: (1) S.F. Altschul *et al.* Nucleic Acids Res., 25:3389, 1997. (2) L.F. Grand. Agr. Res.
11 Serv. Techn. Bull. 240:1, 1985. (3) T. Kobayashi. Index of fungi inhabiting woody plants in
12 Japan. Host, Distribution, and Literature. Zenkohu-Noson-Kyoiku Kyokai Publishing Co. Ltd,
13 1227 pages, 2007. (4) W.A. Sinclair, H.H. Lyon. Diseases of trees and shrubs. Second Edition,
14 Cornell University Press, Ithaca, N.Y., USA, 660 pages, 2004.

15