



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: March 1, 2013

ACCEPTED: March 18, 2013

PUBLISHED: April 5, 2013

Deformed $\mathcal{N} = 2$ theories, generalized recursion relations and S-duality

M. Billó,^a M. Frau,^a L. Gallot,^b A. Lerda^c and I. Pesando^a

^a *Università di Torino, Dipartimento di Fisica and
I.N.F.N. — sezione di Torino,
Via P. Giuria 1, I-10125 Torino, Italy*

^b *LAPTH, Université de Savoie, CNRS,
9, Chemin de Bellevue, 74941 Annecy le Vieux Cedex, France*

^c *Università del Piemonte Orientale, Dipartimento di Scienze e Innovazione Tecnologica and
I.N.F.N. — Gruppo Collegato di Alessandria — sezione di Torino,
Viale T. Michel 11, I-15121 Alessandria, Italy*

E-mail: billlo@to.infn.it, frau@to.infn.it,
laurent.gallot@lapp.in2p3.fr, lerda@to.infn.it, ipesando@to.infn.it

ABSTRACT: We study the non-perturbative properties of $\mathcal{N} = 2$ super conformal field theories in four dimensions using localization techniques. In particular we consider $SU(2)$ gauge theories, deformed by a generic ϵ -background, with four fundamental flavors or with one adjoint hypermultiplet. In both cases we explicitly compute the first few instanton corrections to the partition function and the prepotential using Nekrasov's approach. These results allow us to reconstruct exact expressions involving quasi-modular functions of the bare gauge coupling constant and to show that the prepotential terms satisfy a modular anomaly equation that takes the form of a recursion relation with an explicitly ϵ -dependent term. We then investigate the implications of this recursion relation on the modular properties of the effective theory and find that with a suitable redefinition of the prepotential and of the effective coupling it is possible, at least up to the third order in the deformation parameters, to cast the S-duality relations in the same form as they appear in the Seiberg-Witten solution of the undeformed theory.

KEYWORDS: Nonperturbative Effects, Extended Supersymmetry, Duality in Gauge Field Theories, Supersymmetric Effective Theories

ARXIV EPRINT: [1302.0686](https://arxiv.org/abs/1302.0686)

Contents

1	Introduction	1
2	The $\mathcal{N} = 2^*$ SU(2) theory	4
2.1	Instanton partition functions	5
2.2	Perturbative part	7
2.3	Generalized prepotential	8
2.4	Recursion relations	10
3	The $\mathcal{N} = 2$ SU(2) theory with $N_f = 4$	11
3.1	Instanton partition functions	11
3.2	Perturbative part	13
3.3	Generalized prepotential	13
3.4	Recursion relations	16
4	Modular anomaly equations and S-duality	17
4.1	S-duality at first order in $\epsilon_1\epsilon_2$	19
4.2	S-duality on the prepotential	22
5	Conclusions	23
A	Modular functions	24
B	The coefficient h_3 of the SU(2) $N_f = 4$ prepotential	26
C	Reformulating the modular anomaly equations	27
D	S-duality at order $(\epsilon_1\epsilon_2)^2$	29

1 Introduction

Four-dimensional field theories with rigid $\mathcal{N} = 2$ supersymmetry provide a remarkable arena where many exact results can be obtained; indeed $\mathcal{N} = 2$ supersymmetry, not being maximal, permits a great deal of flexibility but, at the same time, is large enough to guarantee full control. This fact was exploited in the seminal papers [1, 2] where it was shown that the effective dynamics of $\mathcal{N} = 2$ super Yang-Mills (SYM) theories in the limit of low energy and momenta can be exactly encoded in the so-called Seiberg-Witten (SW) curve describing the geometry of the moduli space of the SYM vacua. When the gauge group is SU(2), the SW curve defines a torus whose complex structure parameter is identified with the (complexified) gauge coupling constant τ of the SYM theory at low energy. This

coupling receives perturbative corrections at 1-loop and non-perturbative corrections due to instantons, and the corresponding effective action follows from a prepotential \mathcal{F} that is a holomorphic function of the vacuum expectation value a of the adjoint vector multiplet, of the flavor masses, if any, and of the dynamically generated scale in asymptotically free theories or of the bare gauge coupling constant τ_0 in conformal models (see for instance [3] for a review and extensions of this approach).

Recently, $\mathcal{N} = 2$ superconformal field theories (SCFT) have attracted a lot of attention. Two canonical examples of SCFT's are the $\mathcal{N} = 2$ SU(2) SYM theory with $N_f = 4$ fundamental hypermultiplets and the $\mathcal{N} = 2^*$ theory, namely a $\mathcal{N} = 2$ SYM theory with an adjoint hypermultiplet. In both cases, the β -function vanishes but, when the hypermultiplets are massive, the bare coupling τ_0 gets renormalized at 1-loop by terms proportional to the mass parameters. Besides these, there are also non-perturbative corrections due to instantons. As shown in [4] for the $\mathcal{N} = 2^*$ theory and more recently in [5] for the $N_f = 4$ theory, by organizing the effective prepotential \mathcal{F} as a series in inverse powers of a and by exploiting a recursion relation hidden in the SW curve, it is possible to write the various terms of \mathcal{F} as *exact* functions of the bare coupling. These functions are polynomials in Eisenstein series and Jacobi θ -functions of τ_0 and their modular properties allow one to show that the effective theory at low energy inherits the $Sl(2, \mathbb{Z})$ symmetry of the microscopic theory at high energy. In particular one can show [4, 5] that the S-duality map on the bare coupling, i.e. $\tau_0 \rightarrow -1/\tau_0$, implies the corresponding map on the effective coupling, i.e. $\tau \rightarrow -1/\tau$, and that the prepotential \mathcal{F} and its S-dual are related to each other by a Legendre transformation.

The non-perturbative corrections predicted by the SW solution can also be obtained directly via multi-instanton calculus and the use of localization techniques [6, 7].¹ This approach is based on the calculation of the instanton partition function after introducing two deformation parameters, ϵ_1 and ϵ_2 , of mass dimension 1 which break the four-dimensional Lorentz invariance, regularize the space-time volume and fully localize the integrals over the instanton moduli space on sets of isolated points, permitting their explicit evaluation. This method, which has been extensively applied to many models (see for instance [10–17]) can be interpreted as the effect of putting the gauge theory in a curved background, known as Ω -background [6, 7, 18], or in a supergravity background with a non-trivial graviphoton field strength, which are equivalent on the instanton moduli space [19, 20]. The resulting partition function $Z_{\text{inst}}(\epsilon_1, \epsilon_2)$, also known as Nekrasov partition function, allows one to obtain the non-perturbative part of the SYM prepotential according to

$$\mathcal{F}_{\text{inst}} = - \lim_{\epsilon_1, \epsilon_2 \rightarrow 0} \epsilon_1 \epsilon_2 \log Z_{\text{inst}}(\epsilon_1, \epsilon_2). \tag{1.1}$$

Actually, the Nekrasov partition function is useful not only when the ϵ parameters are sent to zero, but also when they are kept at finite values. In this case, in fact, the non-perturbative ϵ -deformed prepotential

$$F_{\text{inst}}(\epsilon_1, \epsilon_2) = -\epsilon_1 \epsilon_2 \log Z_{\text{inst}}(\epsilon_1, \epsilon_2) \tag{1.2}$$

¹See also [8, 9] for earlier applications of these techniques.

represents a very interesting generalization of the SYM one. By adding to it the corresponding (ϵ -deformed) perturbative part F_{pert} , one gets a generalized prepotential that can be conveniently expanded as follows

$$F_{\text{pert}} + F_{\text{inst}} = \sum_{n,g=0}^{\infty} F^{(n,g)} (\epsilon_1 + \epsilon_2)^{2n} (\epsilon_1 \epsilon_2)^g. \quad (1.3)$$

The amplitude $F^{(0,0)}$, which is the only one that remains when the ϵ -deformations are switched off, coincides with the SYM prepotential \mathcal{F} of the SW theory, up to the classical tree-level term. The amplitudes $F^{(0,g)}$ with $g \geq 1$ account instead for gravitational couplings and correspond to F-terms in the effective action of the form $F^{(0,g)} \mathcal{W}^{2g}$, where \mathcal{W} is the chiral Weyl superfield containing the graviphoton field strength as its lowest component. These terms were obtained long ago from the genus g partition function of the $\mathcal{N} = 2$ topological string on an appropriate Calabi-Yau background [21] and were shown to satisfy a holomorphic anomaly equation [22, 23] from which one can recursively reconstruct the higher genus contributions starting from the lower genus ones (see for instance [24, 25]). More recently, also the amplitudes $F^{(n,g)}$ with $n \neq 0$ have been related to the $\mathcal{N} = 2$ topological string and have been shown to correspond to higher dimensional F-terms of the type $F^{(n,g)} \Upsilon^{2n} \mathcal{W}^{2g}$ where Υ is a chiral projection of real functions of $\mathcal{N} = 2$ vector superfields [26], which also satisfy an extended holomorphic anomaly equation [27, 28]. By taking the limit $\epsilon_2 \rightarrow 0$ with ϵ_1 finite, one selects in (1.3) the amplitudes $F^{(n,0)}$. This limit, also known as Nekrasov-Shatashvili limit [29], is particularly interesting since it is believed that the $\mathcal{N} = 2$ effective theory can be described in this case by certain quantum integrable systems. Furthermore, in this Nekrasov-Shatashvili limit, using saddle point methods it is possible to derive a generalized SW curve [30, 31] and extend the above-mentioned results for the SYM theories to the ϵ -deformed ones.

By considering the Nekrasov partition function and the corresponding generalized prepotential for rank one SCFT's, in [32] a very remarkable relation has been uncovered with the correlation functions of a two-dimensional Liouville theory with an ϵ -dependent central charge. In particular, for the $SU(2)$ theory with $N_f = 4$ the generalized prepotential turns out to be related to the logarithm of the conformal blocks of four Liouville operators on a sphere and the bare gauge coupling constant to the cross-ratio of the punctures where the four operators are located; for the $\mathcal{N} = 2^* SU(2)$ theory, instead, the correspondence works with the one-point conformal blocks on a torus whose complex structure parameter plays the role of the complexified bare gauge coupling. Since the conformal blocks of the Liouville theory have well-defined properties under modular transformations, it is natural to explore modularity also on the four-dimensional gauge theory and try to connect it to the strong/weak-coupling S-duality, generalizing in this way the SW results when ϵ_1 and ϵ_2 are non-zero. On the other hand, one expects that the deformed gauge theory should somehow inherit the duality properties of the Type IIB string theory in which it can be embedded. Some important steps towards this goal have been made in [33] and also in [34] where it has been shown that the SW contour integral techniques remain valid also when both ϵ_1 and ϵ_2 are non-vanishing. To make further progress and gain a more quantitative understanding, it would be useful to know the various amplitudes $F^{(n,g)}$ in (1.3) as

exact functions of the gauge coupling constant and analyze their behavior under modular transformations, similarly to what has been done for the SW prepotential $F^{(0,0)}$ in [4, 5]. Recently, by exploiting the generalized holomorphic anomaly equation, an exact expression in terms of Eisenstein series has been given for the first few amplitudes $F^{(n,g)}$ of the $SU(2)$ theory with $N_f = 4$ and the $\mathcal{N} = 2^*$ $SU(2)$ theory in the limit of vanishing hypermultiplet masses [35]. This analysis has then been extended in [36] to the massive $\mathcal{N} = 2^*$ model in the Nekrasov-Shatashvili limit, using again the extended holomorphic anomaly equation, and in [37] using the properties of the Liouville toroidal conformal blocks in the semi-classical limit of infinite central charge. However, finding the modular properties of the deformed prepotential in full generality still remains an open issue.

In this paper we address this problem and extend the previous results by adopting a different strategy. In section 2, using localization techniques we explicitly compute the first few instanton corrections to the prepotential for the $\mathcal{N} = 2^*$ massive theory with gauge group $SU(2)$ in a generic ϵ -background. From these explicit results we then infer the exact expressions of the various prepotential coefficients and write them in terms of Eisenstein series of the bare coupling. Our results reduce to those of [4, 5] when the deformation parameters are switched off, and to those of [35–37] in the massless or in the Nekrasov-Shatashvili limits. The properties of the Eisenstein series allow us to analyze the behavior of the various prepotential terms under modular transformations and also to write a recursion relation that is equivalent to the holomorphic anomaly equation if one trades modularity for holomorphicity. The recursion relation we find contains a term proportional to $\epsilon_1\epsilon_2$, which is invisible in the SYM limit or in the Nekrasov-Shatashvili limit. In section 3 we repeat the same steps for the $\mathcal{N} = 2$ $SU(2)$ theory with $N_f = 4$ and arbitrary mass parameters and also in this case derive the modular anomaly equation in the form of a recursion relation. In section 4 we study in detail the properties of the generalized prepotential under S-duality, and show that when both ϵ_1 and ϵ_2 are different from zero, due to the new term in the recursion relation, the prepotential and its S-dual are not any more related by a Legendre transformation, an observation which has been recently put forward in [38] from a different perspective. We also propose how the relation between the prepotential and its S-dual has to be modified, by computing the first corrections in $\epsilon_1\epsilon_2$. Finally, in section 5 we conclude by showing that there exist suitable redefinitions of the prepotential and of the effective coupling from which we can recover the simple Legendre relation and write the S-duality relations in the same form as in the undeformed SW theory. The appendices contain some technical details and present several explicit formulas which are useful for the computations described in the main text.

2 The $\mathcal{N} = 2^*$ $SU(2)$ theory

The $\mathcal{N} = 2^*$ SYM theory describes the interactions of a $\mathcal{N} = 2$ gauge vector multiplet with a massive $\mathcal{N} = 2$ hypermultiplet in the adjoint representation of the gauge group. It can be regarded as a massive deformation of the $\mathcal{N} = 4$ SYM theory in which the β -function remains vanishing but the gauge coupling constant receives both perturbative and non-perturbative corrections proportional to the hypermultiplet mass. Using the lo-

calization techniques [6, 7] one can obtain a generalization of this theory by considering the ϵ -dependent terms in the Nekrasov partition function. In the following we only discuss the case in which the gauge group is $SU(2)$ (broken down to $U(1)$ by the vacuum expectation value a of the adjoint scalar of the gauge vector multiplet). We begin by considering the non-perturbative corrections.

2.1 Instanton partition functions

The partition function Z_k at instanton number k is defined by the following integral over the instanton moduli space \mathcal{M}_k :

$$Z_k = \int d\mathcal{M}_k e^{-S_{\text{inst}}} \tag{2.1}$$

where S_{inst} is the instanton moduli action of the $\mathcal{N} = 2^*$ theory. After introducing two deformation parameters ϵ_1 and ϵ_2 , the partition function Z_k can be explicitly computed using the localization techniques. In the case at hand, each Z_k can be expressed as a sum of terms in one-to-one correspondence with an ordered pair of Young tableaux of $U(k)$ such that the total number of boxes in the two tableaux is k . For example, at $k = 1$ we have the two possibilities: (\square, \bullet) and (\bullet, \square) ; at $k = 2$ we have instead the five cases: $(\square\square, \bullet)$, $(\bullet, \square\square)$, (\square, \bullet) , (\bullet, \square) , (\square, \square) ; and so on and so forth. Referring for example to the appendix A of [39] for details, at $k = 1$ one finds

$$\begin{aligned} Z_{(\square, \bullet)} &= \frac{(-\epsilon_1 + \tilde{m})(-\epsilon_2 + \tilde{m})(a_{12} + \tilde{m})(a_{21} + \tilde{m} - \epsilon_1 - \epsilon_2)}{(-\epsilon_1)(-\epsilon_2)a_{12}(a_{21} - \epsilon_1 - \epsilon_2)}, \\ Z_{(\bullet, \square)} &= \frac{(-\epsilon_1 + \tilde{m})(-\epsilon_2 + \tilde{m})(a_{21} + \tilde{m})(a_{12} + \tilde{m} - \epsilon_1 - \epsilon_2)}{(-\epsilon_1)(-\epsilon_2)a_{21}(a_{12} - \epsilon_1 - \epsilon_2)}, \end{aligned} \tag{2.2}$$

where $a_{uv} = a_u - a_v$ with $a_1 = -a_2 = a$, and

$$\tilde{m} = m + \frac{\epsilon_1 + \epsilon_2}{2} \tag{2.3}$$

is the equivariant mass parameter for the adjoint hypermultiplet in the ϵ -background [35, 40]. The 1-instanton partition function is therefore

$$Z_1 = Z_{(\square, \bullet)} + Z_{(\bullet, \square)} = \frac{(4m^2 - (\epsilon_1 - \epsilon_2)^2)(16a^2 - 4m^2 - 3(\epsilon_1 + \epsilon_2)^2)}{8\epsilon_1\epsilon_2(4a^2 - (\epsilon_1 + \epsilon_2)^2)}. \tag{2.4}$$

At $k = 2$ the relevant partition functions are

$$\begin{aligned} Z_{(\square\square, \bullet)} &= \frac{(-\epsilon_1 + \tilde{m})(-\epsilon_2 + \tilde{m})(-\epsilon_1 + \epsilon_2 + \tilde{m})(-2\epsilon_2 + \tilde{m})(a_{12} + \tilde{m})}{(-\epsilon_1)(-\epsilon_2)(-\epsilon_1 + \epsilon_2)(-2\epsilon_2)a_{12}} \\ &\quad \times \frac{(a_{12} + \tilde{m} + \epsilon_2)(a_{21} + \tilde{m} - \epsilon_1 - \epsilon_2)(a_{21} + \tilde{m} - \epsilon_1 - 2\epsilon_2)}{(a_{12} + \epsilon_2)(a_{21} - \epsilon_1 - \epsilon_2)(a_{21} - \epsilon_1 - 2\epsilon_2)}, \\ Z_{(\square, \square)} &= \frac{(-\epsilon_1 + \tilde{m})^2(-\epsilon_2 + \tilde{m})^2(a_{12} + \tilde{m} - \epsilon_1)(a_{12} + \tilde{m} - \epsilon_2)}{(-\epsilon_1)^2(-\epsilon_2)^2(a_{12} - \epsilon_1)(a_{12} - \epsilon_2)} \\ &\quad \times \frac{(a_{21} + \tilde{m} - \epsilon_1)(a_{21} + \tilde{m} - \epsilon_2)}{(a_{21} - \epsilon_1)(a_{21} - \epsilon_2)}. \end{aligned} \tag{2.5}$$

The contributions corresponding to the other Young tableaux at $k = 2$ can be obtained from the previous expressions with suitable redefinitions. In particular, $Z_{(\bullet, \square\square)}$ is obtained from $Z_{(\square\square, \bullet)}$ in (2.5) by exchanging $a_{12} \leftrightarrow a_{21}$, $Z_{(\square, \bullet)}$ is obtained by exchanging $\epsilon_1 \leftrightarrow \epsilon_2$, and finally $Z_{(\bullet, \square)}$ is obtained by simultaneously exchanging $a_{12} \leftrightarrow a_{21}$ and $\epsilon_1 \leftrightarrow \epsilon_2$. The complete 2-instanton partition function is

$$Z_2 = Z_{(\square\square, \bullet)} + Z_{(\bullet, \square\square)} + Z_{(\square, \bullet)} + Z_{(\bullet, \square)} + Z_{(\square, \square)} \tag{2.6}$$

but we refrain from writing its expression since it is not particularly inspiring. This procedure can be systematically extended to higher instanton numbers leading to explicit formulas for the instanton partition functions.

Following [7] we can cast these results in a nice and compact form. Indeed, defining

$$q = e^{\pi i \tau_0} \tag{2.7}$$

where $\tau_0 = \frac{\theta}{2\pi} + i \frac{4\pi}{g^2}$ is the complexified gauge coupling constant of the $\mathcal{N} = 2^*$ SYM theory, the grand-canonical instanton partition function

$$Z_{\text{inst}} = \sum_{k=0}^{\infty} q^{2k} Z_k \tag{2.8}$$

where $Z_0 = 1$, can be rewritten as

$$Z_{\text{inst}} = \sum_{(Y_1, Y_2)} q^{2|Y|} \prod_{i,j=1}^{\infty} \prod_{u,v=1}^2 \left[\frac{a_{uv} + \epsilon_1(i-1) - \epsilon_2 j}{a_{uv} + \epsilon_1(i-1 - \tilde{k}_{vj}) - \epsilon_2(j - k_{ui})} \times \frac{a_{uv} + \tilde{m} + \epsilon_1(i-1 - \tilde{k}_{vj}) - \epsilon_2(j - k_{ui})}{a_{uv} + \tilde{m} + \epsilon_1(i-1) - \epsilon_2 j} \right] \tag{2.9}$$

where the first line represents the contribution of the gauge vector multiplet and the second line that of the adjoint hypermultiplet. Here k_{ui} and \tilde{k}_{ui} denote, respectively, the number of boxes in the i -th row and in the i -th column of a Young tableaux Y_u and are related to the Dynkin indices of the corresponding representation. These quantities can be extended for any integer i with the convention that $k_{ui} = 0$ or $\tilde{k}_{ui} = 0$ if the i -th row or the i -th column of Y_u is empty. For example, for $Y_u = \square\square$ the k_{ui} 's are $(2, 0, 0, 0, \dots)$ and the \tilde{k}_{ui} 's are $(1, 1, 0, 0, \dots)$. Moreover we have

$$|Y| = \sum_{u,i} k_{ui} = \sum_{u,i} \tilde{k}_{ui} = k. \tag{2.10}$$

It is quite straightforward to check that the expressions (2.2) and (2.5) are reproduced by the compact formula (2.9) by selecting the appropriate Young tableaux.

Following Nekrasov's prescription, we can obtain the generalized non-perturbative pre-potential according to

$$F_{\text{inst}} = -\epsilon_1 \epsilon_2 \log Z_{\text{inst}} = \sum_{k=1}^{\infty} q^{2k} F_k. \tag{2.11}$$

In the limit $\epsilon_\ell \rightarrow 0$, the above expression computes the instanton contributions to the pre-potential of the $\mathcal{N} = 2^*$ SYM theory, while the finite ϵ -dependent terms represent further non-perturbative corrections. Notice that in the limit $\tilde{m} \rightarrow 0$, (2.9) correctly reduces to the partition function of the $\mathcal{N} = 4$ SYM theory, namely to the Euler characteristic of the instanton moduli space (see for instance [41]).

2.2 Perturbative part

The compact expression (2.9) allows us to “guess” the perturbative part of the partition function by applying the same formal reasoning of section 3.10 of [6]. Indeed, in (2.9) we recognize the following universal (i.e. k -independent but a -dependent) factor

$$\prod_{i,j=1}^{\infty} \prod_{\substack{u,v=1 \\ u \neq v}}^2 \frac{a_{uv} + \epsilon_1(i-1) - \epsilon_2 j}{a_{uv} + \tilde{m} + \epsilon_1(i-1) - \epsilon_2 j} \tag{2.12}$$

which, if suitably interpreted and regularized [6, 7], can be related to the perturbative part of the partition function of the $\mathcal{N} = 2^*$ theory in the ϵ -background. According to this idea, we are then led to write

$$F_{\text{pert}} = \epsilon_1 \epsilon_2 \sum_{\substack{u,v=1 \\ u \neq v}}^2 \sum_{i,j=1}^{\infty} \log \frac{a_{uv} + \epsilon_1(i-1) - \epsilon_2 j}{a_{uv} + \tilde{m} + \epsilon_1(i-1) - \epsilon_2 j}. \tag{2.13}$$

Using the following representation for the logarithm

$$\log \frac{x}{\Lambda} = - \frac{d}{ds} \left(\frac{\Lambda^s}{\Gamma(s)} \int_0^\infty \frac{dt}{t} t^s e^{-tx} \right) \Big|_{s=0} \tag{2.14}$$

where Λ is an arbitrary scale and summing over i and j , we can rewrite (2.13) as

$$F_{\text{pert}} = \epsilon_1 \epsilon_2 \sum_{\substack{u,v=1 \\ u \neq v}}^2 \left[\gamma_{\epsilon_1, \epsilon_2}(a_{uv}) - \gamma_{\epsilon_1, \epsilon_2}(a_{uv} + \tilde{m}) \right] \tag{2.15}$$

where (see also [18, 35])

$$\gamma_{\epsilon_1, \epsilon_2}(x) = \frac{d}{ds} \left(\frac{\Lambda^s}{\Gamma(s)} \int_0^\infty \frac{dt}{t} \frac{t^s e^{-tx}}{(e^{-\epsilon_1 t} - 1)(e^{-\epsilon_2 t} - 1)} \right) \Big|_{s=0}. \tag{2.16}$$

This function, which is related to the logarithm of the Barnes double Γ -function, can be easily computed by expanding for small values of ϵ_1 and ϵ_2 . As a result, F_{pert} becomes a series in inverse powers of a^2 whose first few terms are

$$F_{\text{pert}} = \frac{1}{4} (4m^2 - s^2) \log \frac{2a}{\Lambda} - \frac{(4m^2 - s^2)(4m^2 + 4p - s^2)}{768 a^2} - \frac{(4m^2 - s^2)(4m^2 + 4p - s^2)(4m^2 + 6p - 3s^2)}{61440 a^4}$$

$$\begin{aligned}
 & - \frac{(4m^2 - s^2)(4m^2 + 4p - s^2)(48m^4 + 176m^2p + 160p^2 - 88m^2s^2 - 204ps^2 + 51s^4)}{8257536 a^6} \\
 & + \mathcal{O}(a^{-8}).
 \end{aligned} \tag{2.17}$$

Here we have used (2.3) and introduced the convenient notation

$$s = \epsilon_1 + \epsilon_2 \quad , \quad p = \epsilon_1 \epsilon_2. \tag{2.18}$$

One can easily check that in the limit $\epsilon_\ell \rightarrow 0$ this expression correctly reproduces the 1-loop prepotential of the $\mathcal{N} = 2^*$ SU(2) gauge theory (see for instance [4]).

2.3 Generalized prepotential

The complete generalized prepotential is the sum of the classical, perturbative and non-perturbative parts. We now focus on the latter two terms which are directly related to the Nekrasov partition function. Just like the 1-loop piece (2.17), also the instanton terms (2.11) can be organized as a series expansion for small values of ϵ_1 and ϵ_2 , or equivalently for large values of a . Discarding a -independent terms, which are not relevant for the gauge theory dynamics, we write

$$F_{\text{pert}} + F_{\text{inst}} = h_0 \log \frac{2a}{\Lambda} - \sum_{\ell=1}^{\infty} \frac{h_\ell}{2^{\ell+1} \ell} \frac{1}{a^{2\ell}} \tag{2.19}$$

where the coefficients h_ℓ are polynomials in m^2 , s^2 and p which can be explicitly derived from (2.17) as far as the perturbative part is concerned, and from the instanton partition functions (2.9), after using (2.11), for the non-perturbative part. Here we list the first few of these coefficients up to three instantons:

$$h_0 = \frac{1}{4}(4m^2 - s^2), \tag{2.20}$$

$$h_1 = (4m^2 - s^2)(4m^2 + 4p - s^2) \left(\frac{1}{192} - \frac{1}{8}q^2 - \frac{3}{8}q^4 - \frac{1}{2}q^6 + \dots \right), \tag{2.21}$$

$$\begin{aligned}
 h_2 = & (4m^2 - s^2)(4m^2 + 4p - s^2) \left(\frac{4m^2 + 6p - 3s^2}{3840} - \frac{s^2}{8}q^2 \right. \\
 & \left. + \frac{12m^2 + 18p - 21s^2}{16}q^4 + \frac{8m^2 + 12p - 9s^2}{2}q^6 + \dots \right),
 \end{aligned} \tag{2.22}$$

$$\begin{aligned}
 h_3 = & (4m^2 - s^2)(4m^2 + 4p - s^2) \left(\frac{48m^4 + 176m^2p + 160p^2 - 88m^2s^2 - 204ps^2 + 51s^4}{172032} \right. \\
 & - \frac{3s^4}{32}q^2 - \frac{240m^4 + 1392m^2p + 1440p^2 - 3000m^2s^2 - 7548ps^2 + 3903s^4}{1024}q^4 \\
 & \left. - \frac{240m^4 + 1008m^2p + 960p^2 - 1080m^2s^2 - 2652ps^2 + 987s^4}{32}q^6 + \dots \right).
 \end{aligned} \tag{2.23}$$

It is interesting to notice that all h_ℓ 's are proportional to $(4m^2 - s^2)$ and, except for h_0 , also to $(4m^2 + 4p - s^2)$. Explicit expressions for h_ℓ with $\ell > 3$ can be systematically derived from the generalized prepotential but they are not needed for our considerations.

Building on previous results obtained in $\epsilon_\ell \rightarrow 0$ limit from the SW curve [4, 5] and on the analysis of [35] for the massless case, we expect that the expressions (2.20)–(2.23) are just the first terms in the instanton expansion of (quasi) modular functions of q . More precisely, we expect that h_ℓ are (quasi) modular functions of weight 2ℓ that can be written solely in terms of the Eisenstein series E_2 , E_4 and E_6 (see appendix A for our conventions and definitions). This is indeed what happens. In fact we have

$$h_0 = \frac{1}{4}(4m^2 - s^2), \tag{2.24}$$

$$h_1 = \frac{1}{2^6 \cdot 3}(4m^2 - s^2)(4m^2 + 4p - s^2) E_2, \tag{2.25}$$

$$h_2 = \frac{1}{2^9 \cdot 3^2 \cdot 5}(4m^2 - s^2)(4m^2 + 4p - s^2) \times \left[5(4m^2 + 6p - s^2)E_2^2 + (4m^2 + 6p - 13s^2)E_4 \right], \tag{2.26}$$

$$h_3 = \frac{1}{2^{14} \cdot 3^3 \cdot 5 \cdot 7}(4m^2 - s^2)(4m^2 + 4p - s^2) \times \left[35(80m^4 + 272m^2p + 240p^2 - 40m^2s^2 - 68ps^2 + 5s^4)E_2^3 + 84(16m^4 + 64m^2p + 60p^2 - 56m^2s^2 - 136ps^2 + 13s^4)E_4 E_2 + (176m^4 + 944m^2p + 960p^2 - 1816m^2s^2 - 4556ps^2 + 3323s^4)E_6 \right]. \tag{2.27}$$

By using the small q expansion of the Eisenstein series one can check that the explicit instanton contributions we have computed using localization techniques are correctly recovered from the previous formulas. We stress that the fact that the various instanton terms nicely combine into expressions involving only the Eisenstein series is not obvious a priori and is a very strong a posteriori test on the numerical coefficients appearing in (2.20)–(2.23). It is quite remarkable that the explicit instanton results at low k can be nicely extrapolated to reconstruct modular forms from which, by expanding in powers of q , one can obtain the contributions at any instanton number.

We can also organize the generalized prepotential according to (1.3) and obtain the amplitudes $F^{(n,g)}$ as a series in inverse powers of a^2 with coefficients that are polynomials in E_2 , E_4 and E_6 . The first few of such amplitudes are:

$$F^{(0,0)} = m^2 \log \frac{2a}{\Lambda} - \frac{m^4 E_2}{48 a^2} - \frac{m^6(5E_2^2 + E_4)}{5760 a^4} - \frac{m^8(175E_2^3 + 84E_2E_4 + 11E_6)}{2903040 a^6} + \dots, \tag{2.28}$$

$$F^{(1,0)} = -\frac{1}{4} \log \frac{2a}{\Lambda} + \frac{m^2 E_2}{96 a^2} + \frac{m^4(E_2^2 + E_4)}{1536 a^4} + \frac{m^6(175E_2^3 + 336E_2E_4 + 8E_6)}{2903040 a^6} + \dots \tag{2.29}$$

$$F^{(0,1)} = -\frac{m^2 E_2}{48 a^2} - \frac{m^4(5E_2^2 + E_4)}{2304 a^4} - \frac{m^6(11E_2^3 + 6E_2E_4 + E_6)}{41472 a^6} + \dots \tag{2.30}$$

$$F^{(2,0)} = -\frac{E_2}{768 a^2} - \frac{m^2(5E_2^2 + 9E_4)}{30720 a^4} + \frac{m^4(175E_2^3 + 588E_2E_4 + 559E_6)}{7741440 a^6} + \dots \tag{2.31}$$

$$F^{(1,1)} = \frac{E_2}{192 a^2} + \frac{m^2(25E_2^2 + 17E_4)}{23040 a^4} + \frac{m^4(385E_2^3 + 798E_2E_4 + 213E_6)}{1935360 a^6} + \dots \tag{2.32}$$

$$F^{(0,2)} = -\frac{m^2(5E_2^2 + E_4)}{3840 a^4} - \frac{m^4(160E_2^3 + 93E_2E_4 + 17E_6)}{414720 a^6} + \dots \tag{2.33}$$

Terms with higher values of n and g can be systematically generated without any difficulty from the h_ℓ 's given in (2.24)–(2.27). The first term $F^{(0,0)}$ represents the prepotential of the $\mathcal{N} = 2^*$ SU(2) gauge theory and its expression (2.28) agrees with that found in [4] from the SW curve (see also [5]). The other terms are generalizations of those considered in [35] and more recently in [37] in particular limits ($m \rightarrow 0$ or $\epsilon_2 \rightarrow 0$) where they drastically simplify.

2.4 Recursion relations

The generalized prepotential (2.19) is clearly holomorphic by construction but does not have nice transformation properties under the modular group since the coefficients h_ℓ explicitly depend on the second Eisenstein series E_2 which is not a good modular function. Indeed, under

$$\tau_0 \rightarrow \frac{a\tau_0 + b}{c\tau_0 + d} \quad \text{with } ad - bc = 1, \tag{2.34}$$

E_2 transforms inhomogeneously as follows

$$E_2(\tau_0) \rightarrow (c\tau_0 + d)^2 E_2(\tau_0) + (c\tau_0 + d) \frac{6c}{\pi i}. \tag{2.35}$$

Therefore, in order to have good modular properties we should replace everywhere E_2 with the shifted Eisenstein series $\widehat{E}_2 = E_2 + \frac{6}{\pi i(\tau_0 - \bar{\tau}_0)}$ at the price, however, of losing holomorphicity. This fact leads to the so-called holomorphic anomaly equation [22, 23] (see also [28]). On the other hand, in the limit $\bar{\tau}_0 \rightarrow \infty$, holding τ fixed so that $\widehat{E}_2 \rightarrow E_2$, we recover holomorphicity but lose good modular properties and obtain the so-called modular anomaly equation [42]. This equation can be rephrased in terms of a recursion relation for the h_ℓ coefficients which allows us to completely fix their dependence on E_2 .

To this aim let us consider the explicit expressions (2.24)–(2.27) and compute the derivatives of h_ℓ with respect to E_2 . With simple algebra we find

$$\frac{\partial h_\ell}{\partial E_2} = \frac{\ell}{12} \sum_{i=0}^{\ell-1} h_i h_{\ell-i-1} + \frac{\ell(2\ell-1)}{12} \epsilon_1 \epsilon_2 h_{\ell-1} \tag{2.36}$$

with the initial condition

$$\frac{\partial h_0}{\partial E_2} = 0. \tag{2.37}$$

We have explicitly checked this relation for several values of $\ell > 3$; we can thus regard it as a distinctive property of the ϵ -deformed $\mathcal{N} = 2^*$ low-energy theory. Notice that in the limit $\epsilon_\ell \rightarrow 0$, (2.36) reduces to the recursion relation satisfied by the coefficients of the prepotential of the $\mathcal{N} = 2^*$ SU(2) theory found in [4, 5] from the SW curve, and that the linear term in the right hand side disappears in the so-called Nekrasov-Shatashvili limit [29] where one of the two deformation parameters vanishes and a generalized SW curve can be introduced [30, 31].

Eq. (2.36) can be formulated also as a recursion relation for the amplitudes $F^{(n,g)}$ defined in (1.3). To see this, let us first extract the a -dependence and, in analogy with (2.19), write

$$F^{(n,g)} = - \sum_{\ell=1}^{\infty} \frac{f_{\ell}^{(n,g)}}{2^{\ell+1} \ell a^{2\ell}}, \quad (2.38)$$

so that

$$h_{\ell} = \sum_{n,g} f_{\ell}^{(n,g)} (\epsilon_1 + \epsilon_2)^{2n} (\epsilon_1 \epsilon_2)^g. \quad (2.39)$$

Notice that the coefficients $f_{\ell}^{(n,g)}$, which are polynomials in the hypermultiplet mass m , have mass dimensions $2(1 + \ell - n - g)$; therefore $f_{\ell}^{(n,g)} = 0$ if $n + g > \ell + 1$ (this condition can be easily checked on the explicit expressions (2.28)–(2.31)). These definitions must be supplemented by the “initial conditions”

$$f_0^{(0,0)} = m^2, \quad f_0^{(1,0)} = -\frac{1}{4}, \quad f_0^{(0,1)} = 0 \quad (2.40)$$

which are obtained from (2.24). Inserting (2.39) in the recursion relation (2.36), we obtain

$$\frac{\partial f_{\ell}^{(n,g)}}{\partial E_2} = \frac{\ell}{12} \sum'_{n_1, n_2; g_1, g_2} \left(\sum_{i=0}^{\ell-1} f_i^{(n_1, g_1)} f_{\ell-i-1}^{(n_2, g_2)} \right) + \frac{\ell(2\ell-1)}{12} f_{\ell-1}^{(n, g-1)} \quad (2.41)$$

where the $'$ means that the sum is performed over all n_1, n_2, g_1 and g_2 such that $n_1 + n_2 = n$ and $g_1 + g_2 = g$. Eq. (2.41) shows that the coefficients $f_{\ell}^{(n,g)}$ and hence the amplitudes $F^{(n,g)}$ are recursively related to those with lower values of n and g , similarly to what can be deduced from the holomorphic anomaly equation [35].

3 The $\mathcal{N} = 2$ SU(2) theory with $N_f = 4$

We now consider the $\mathcal{N} = 2$ SU(2) theory with N_f flavor hypermultiplets in the fundamental representation of the gauge group. When $N_f = 4$ the 1-loop β -function vanishes and the conformal invariance is broken only by the flavor masses m_f ($f = 1, \dots, 4$). Furthermore there are non-perturbative effects due to instantons which are nicely encoded in the exact SW solution [2]. We now discuss the generalizations of these effects induced by the ϵ_{ℓ} deformation parameters in the Nekrasov partition function, following the same path described in the previous section for the $\mathcal{N} = 2^*$ theory.

3.1 Instanton partition functions

Using localization techniques, we can express the instanton partition functions Z_k of the $N_f = 4$ theory as sums of terms related to pairs of Young tableaux. Referring again the appendix A of [39] for details, at $k = 1$ we have the following two contributions

$$\begin{aligned} Z_{(\square, \bullet)} &= \frac{1}{(-\epsilon_1)(-\epsilon_2)a_{12}(a_{21} - \epsilon_1 - \epsilon_2)} \prod_{f=1}^4 (a_1 + \tilde{m}_f), \\ Z_{(\bullet, \square)} &= \frac{1}{(-\epsilon_1)(-\epsilon_2)a_{21}(a_{12} - \epsilon_1 - \epsilon_2)} \prod_{f=1}^4 (a_2 + \tilde{m}_f) \end{aligned} \quad (3.1)$$

where

$$\tilde{m}_f = m_f + \frac{\epsilon_1 + \epsilon_2}{2} \tag{3.2}$$

is the equivariant hypermultiplet mass in the ϵ -background.² Summing the two terms (3.1) and setting $a_1 = -a_2 = a$, we get

$$\begin{aligned} Z_1 &= Z_{(\square, \bullet)} + Z_{(\bullet, \square)} \\ &= -\frac{1}{2\epsilon_1\epsilon_2} \left[a^2 + \sum_{f < f'} m_f m_{f'} + (\epsilon_1 + \epsilon_2) \sum_f m_f + \frac{3}{4}(\epsilon_1 + \epsilon_2)^2 + \frac{4 \text{Pfm}}{4a^2 - (\epsilon_1 + \epsilon_2)^2} \right] \end{aligned} \tag{3.3}$$

where $\text{Pfm} \equiv m_1 m_2 m_3 m_4$. The partition function Z_1 contains a part proportional to a^2 , a part which does not depend on a and a part containing a^2 in the denominator. In all a -dependent terms the flavor masses m_f always occur in $\text{SO}(8)$ -invariant combinations and thus in all such terms we can always express the mass dependence using the following quadratic, quartic and sextic $\text{SO}(8)$ invariants:

$$\begin{aligned} R &= \frac{1}{2} \sum_f m_f^2, \\ T_1 &= \frac{1}{12} \sum_{f < f'} m_f^2 m_{f'}^2 - \frac{1}{24} \sum_f m_f^4, \\ T_2 &= -\frac{1}{24} \sum_{f < f'} m_f^2 m_{f'}^2 + \frac{1}{48} \sum_f m_f^4 - \frac{1}{2} \text{Pfm}, \\ N &= \frac{3}{16} \sum_{f < f' < f''} m_f^2 m_{f'}^2 m_{f''}^2 - \frac{1}{96} \sum_{f \neq f'} m_f^2 m_{f'}^4 + \frac{1}{96} \sum_f m_f^6. \end{aligned} \tag{3.4}$$

The a -independent terms in (3.3), instead, are not invariant under the $\text{SO}(8)$ flavor group. Indeed, neither $\sum_f m_f$ nor $\sum_{f < f'} m_f m_{f'}$ respect this invariance. Notice, however, that a -independent terms in the partition functions, and hence in the prepotential, are irrelevant for the gauge theory dynamics, and thus can be neglected. We will always do so in presenting our results.

The explicit expressions of the partition functions Z_k for $k > 1$ are rather cumbersome; nevertheless it is possible to write the (grand-canonical) instanton partition function in a quite compact way using the connection with the Young tableaux. Indeed, denoting as x the instanton counting parameter and using the same notations introduced in section 2.1, we have

$$\begin{aligned} Z_{\text{inst}} &= \sum_{k=0}^{\infty} x^k Z_k \\ &= \sum_{(Y_1, Y_2)} x^{|Y|} \prod_{i,j=1}^{\infty} \left[\prod_{u,v=1}^2 \frac{a_{uv} + \epsilon_1(i-1) - \epsilon_2 j}{a_{uv} + \epsilon_1(i-1 - \tilde{k}_{vj}) - \epsilon_2(j - k_{ui})} \right. \\ &\quad \left. \times \prod_{u=1}^2 \prod_{f=1}^4 \frac{a_u + \tilde{m}_f + \epsilon_1(i-1) - \epsilon_2(j - k_{ui})}{a_u + \tilde{m}_f + \epsilon_1(i-1) - \epsilon_2 j} \right]. \end{aligned} \tag{3.5}$$

²Note that in [39] the hypermultiplet masses were denoted $-m_f$.

Here the second line represents the contribution of the gauge vector multiplet and the last line that of the fundamental hypermultiplets. From this expression, by selecting the appropriate Young tableaux, one can obtain the various terms of the instanton partition function and their dependence on the ϵ_ℓ parameters.

3.2 Perturbative part

Also in the $N_f = 4$ theory one can deduce the perturbative contribution to the partition function from the “universal” factor of the grand-canonical instanton partition function (3.5), namely

$$\prod_{i,j=1}^{\infty} \left[\prod_{\substack{u,v=1 \\ u \neq v}}^2 (a_{uv} + \epsilon_1(i-1) - \epsilon_2 j) \prod_{u=1}^2 \prod_{f=1}^4 \frac{1}{(a_u + \tilde{m}_f + \epsilon_1(i-1) - \epsilon_2 j)} \right] \quad (3.6)$$

which, of course, requires a suitable interpretation and regularization. Using this observation and following again [6, 7], we can write the perturbative part of the prepotential as

$$F_{\text{pert}} = \epsilon_1 \epsilon_2 \sum_{i,j=1}^{\infty} \left[\sum_{\substack{u,v=1 \\ u \neq v}}^2 \log \frac{a_{uv} + \epsilon_1(i-1) - \epsilon_2 j}{\Lambda} - \sum_{u=1}^2 \sum_{f=1}^4 \log \frac{a_u + \tilde{m}_f + \epsilon_1(i-1) - \epsilon_2 j}{\Lambda} \right] \quad (3.7)$$

which, after using (2.14) and summing over i and j , becomes

$$F_{\text{pert}} = \epsilon_1 \epsilon_2 \left[\sum_{\substack{u,v=1 \\ u \neq v}}^2 \gamma_{\epsilon_1 \epsilon_2}(a_{uv}) - \sum_{u=1}^2 \sum_{f=1}^4 \gamma_{\epsilon_1 \epsilon_2}(a_u + \tilde{m}_f) \right] \quad (3.8)$$

where the function $\gamma_{\epsilon_1 \epsilon_2}$ is defined in (2.16). Expanding the latter for small values of ϵ_1 and ϵ_2 and discarding as usual a -independent terms, we obtain

$$\begin{aligned} F_{\text{pert}} = & -\log 16 a^2 + \frac{1}{2}(4R - s^2 + p) \log \frac{a}{\Lambda} \\ & - \frac{1}{96 a^2} \left[16R^2 - 96T_1 - 8R(s^2 - 2p) + s^4 - 4s^2 p + 3p^2 \right] \\ & - \frac{1}{3840 a^4} \left[128R^3 - 1920RT_1 + 768N - 160(R^2 - 6T_1)(s^2 - 2p) \right. \\ & \left. + R(56s^4 - 224s^2 p + 192p^2) - 6s^6 + 36s^4 p - 63s^2 p^2 + 30p^3 \right] + \mathcal{O}(a^{-6}) \end{aligned} \quad (3.9)$$

where s and p are as in (2.18). One can easily check that in the limit $\epsilon_\ell \rightarrow 0$ this expression correctly reproduces the 1-loop prepotential of the $\mathcal{N} = 2$ SU(2) $N_f = 4$ theory (see for instance [3]). Notice the presence of a term proportional to a^2 which corresponds to a finite 1-loop renormalization of the gauge coupling constant.

3.3 Generalized prepotential

We now consider the instanton corrections to the prepotential of the $N_f = 4$ theory which is given by

$$F_{\text{inst}} = -\epsilon_1 \epsilon_2 \log Z_{\text{inst}} = \sum_{k=1}^{\infty} x^k F_k. \quad (3.10)$$

At $k = 1$, up to a -independent terms, from (3.1) we simply get

$$F_1 = -\epsilon_1 \epsilon_2 Z_1 = \frac{a^2}{2} - \frac{2(T_1 + 2T_2)}{4a^2 - s^2} \quad (3.11)$$

The presence of an a^2 -term in F_1 signals that there is a renormalization of the gauge coupling constant at the 1-instanton level. Actually, the same thing occurs also at higher instanton numbers; indeed we find

$$F_2 = \frac{13 a^2}{64} + \dots, \quad F_3 = \frac{23 a^2}{192} + \dots, \quad F_4 = \frac{2701 a^2}{32678} + \dots \quad (3.12)$$

Combining these contributions with the perturbative one given by the first term in (3.8) and adding also the classical tree-level prepotential $F_{\text{cl}} = \log x a^2$, we obtain

$$\left(\log x - \log 16 + \frac{1}{2} x + \frac{13}{64} x^2 + \frac{23}{192} x^3 + \frac{2701}{32678} x^4 + \dots \right) a^2 \equiv \log q a^2 \quad (3.13)$$

that is a redefinition of the instanton expansion parameter and hence of the gauge coupling constant. By inverting this relation we get [43]

$$x = 16 q (1 - 8q + 44q^2 - 192q^3 + \dots) = \frac{\theta_2(q)^4}{\theta_3(q)^4} \quad (3.14)$$

(see appendix A for some properties of the Jacobi θ -functions). As pointed out for instance in [44], by setting

$$q = e^{i\pi\tau_0} \quad \text{with} \quad \tau_0 = \frac{\theta}{\pi} + i \frac{8\pi}{g^2}, \quad (3.15)$$

one can show that τ_0 is precisely the modular parameter of the SW curve for the $SU(2)$ $N_f = 4$ theory appearing in the original paper [2]. This gauge coupling constant receives only corrections proportional to the hypermultiplet masses and as such is the strict analogue³ of the gauge coupling constant τ_0 of the $\mathcal{N} = 2^*$ theory considered in section 2. On the other hand, using (3.14) one can show that x is a cross-ratio of the four roots of the original SW curve [2]. From now on we always present the results for the $N_f = 4$ theory in terms of q .

Expanding the complete prepotential $F_{\text{pert}} + F_{\text{inst}}$ as in (2.19), the first coefficients h_ℓ up to three instantons turn out to be

$$h_0 = \frac{1}{2}(4R - s^2 + p), \quad (3.16)$$

$$\begin{aligned} h_1 = \frac{1}{24} & \left[16(R^2 - 6T_1) - 8R(s^2 - 2p) + s^4 - 4s^2p + 3p^2 \right] + 32(T_1 + 2T_2) q \\ & - \left[16(R^2 + 6T_1) - 8R(s^2 - 2p) + s^4 - 4s^2p + 3p^2 \right] q^2 \\ & + 128(T_1 + 2T_2) q^3 + \mathcal{O}(q^4), \end{aligned} \quad (3.17)$$

$$\begin{aligned} h_2 = \frac{1}{240} & \left[128R^3 - 1920RT_1 + 768N - 160(R^2 - 6T_1)(s^2 - 2p) \right. \\ & \left. + R(56s^4 - 224s^2p + 192p^2) - 6s^6 + 36s^4p - 63s^2p^2 + 30p^3 \right] \end{aligned}$$

³Except for a customary overall factor of 2.

$$\begin{aligned}
& + 32s^2(T_1 + 2T_2)q \\
& + \left[768N + 384RT_1 - 64(R^2 + 6T_1)(s^2 - p) + 8R(4s^4 - 12s^2p + 5p^2) \right. \\
& \quad \left. - 4s^6 + 20s^4p - 25s^2p^2 + 6p^3 \right] q^2 \\
& - 128(T_1 + 2T_2)(16R - 11s^2 + 16p)q^3 + \mathcal{O}(q^4). \tag{3.18}
\end{aligned}$$

We worked out also the expressions for a few other h_ℓ 's with $\ell > 3$ which however become rapidly very cumbersome and thus we do not write them here (see for example appendix B where we give the detailed form of h_3).

As in the $\mathcal{N} = 2^*$ model, also in the $N_f = 4$ theory we can view the previous results as the first instanton terms of the expansion of (quasi) modular forms in powers of q . In this case, however, besides the Eisenstein series E_2 , E_4 and E_6 , also the Jacobi θ -functions θ_2 and θ_4 are needed. Matching the q -expansion of these modular functions with the explicit results (3.16)–(3.18), guided by what we already obtained in the $\epsilon_\ell \rightarrow 0$ limit [5] which suggests the existence of a recursion relation that fixes the E_2 dependence, we find⁴

$$h_0 = \frac{1}{2}(4R - s^2 + p), \tag{3.19}$$

$$h_1 = \frac{1}{24}(4R - s^2 + p)(4R - s^2 + 3p)E_2 - 4(T_1\theta_4^4 - T_2\theta_2^4), \tag{3.20}$$

$$\begin{aligned}
h_2 = & \frac{1}{144}(4R - s^2 + p)(4R - s^2 + 3p)(4R - s^2 + 4p)E_2^2 \\
& - \frac{4}{3}(4R - s^2 + 4p)(T_1\theta_4^4 - T_2\theta_2^4)E_2 \\
& + \frac{1}{720} \left[64R^3 - 80R^2(3s^2 - 4p) + 4R(27s^4 - 88s^2p + 49p^2) \right. \\
& \quad \left. - 13s^6 + 68s^4p - 94s^2p^2 + 30p^3 + 2304N \right] E_4 \\
& - \frac{8}{3}(R - s^2 + p) \left[T_1\theta_4^4(2\theta_2^4 + \theta_4^4) + T_2\theta_2^4(\theta_2^4 + 2\theta_4^4) \right]. \tag{3.21}
\end{aligned}$$

We have checked that a similar pattern occurs also in other h_ℓ 's with $\ell > 3$ (see appendix B).

Organizing the complete prepotential as an expansion in powers of s^2 and p as in (1.3), we can obtain the amplitudes $F^{(n,g)}$, the first few of which are

$$\begin{aligned}
F^{(0,0)} = & 2R \log \frac{a}{\Lambda} - \frac{R^2 E_2}{6a^2} + \frac{T_1\theta_4^4 - T_2\theta_2^4}{a^2} - \frac{R^3(5E_2^2 + E_4)}{180a^4} - \frac{NE_4}{5a^4} \\
& + \frac{RT_1\theta_4^4(2E_2 + 2\theta_2^4 + \theta_4^4)}{6a^4} - \frac{RT_2\theta_2^4(2E_2 + 2\theta_4^4 + \theta_2^4)}{6a^4} + \dots, \tag{3.22}
\end{aligned}$$

$$\begin{aligned}
F^{(1,0)} = & -\frac{1}{2} \log \frac{a}{\Lambda} + \frac{RE_2}{12a^2} + \frac{R^2(E_2^2 + E_4)}{48a^4} \\
& - \frac{T_1\theta_4^4(E_2 + 4\theta_2^4 + 2\theta_4^4)}{12a^4} + \frac{T_2\theta_2^4(E_2 - 4\theta_4^4 - 2\theta_2^4)}{12a^4} + \dots, \tag{3.23}
\end{aligned}$$

⁴As compared to the results presented in [5, 44], here all masses have been rescaled by a factor of $\sqrt{2}$, that is $m_f^{\text{there}} = \sqrt{2}m_f^{\text{here}}$ for all f .

$$\begin{aligned}
 F^{(0,1)} = & \frac{1}{2} \log \frac{a}{\Lambda} - \frac{RE_2}{6a^2} - \frac{R^2(2E_2^2 + E_4)}{36a^4} \\
 & + \frac{T_1\theta_4^4(2E_2 + 2\theta_2^4 + \theta_4^4)}{6a^4} - \frac{T_2\theta_2^4(2E_2 - 2\theta_4^4 - \theta_2^4)}{6a^4} + \dots, \quad (3.24)
 \end{aligned}$$

$$F^{(2,0)} = -\frac{E_2}{96a^2} - \frac{R(5E_2^2 + 9E_4)}{960a^4} + \dots, \quad (3.25)$$

$$F^{(1,1)} = \frac{E_2}{24a^2} + \frac{R(10E_2^2 + 11E_4)}{360a^4} + \dots, \quad (3.26)$$

$$F^{(0,2)} = -\frac{E_2}{32a^2} - \frac{R(95E_2^2 + 49E_4)}{2880a^4} + \dots, \quad (3.27)$$

$$F^{(3,0)} = \frac{5E_2^2 + 13E_4}{11520a^4} \dots, \quad F^{(2,1)} = -\frac{10E_2^2 + 17E_4}{2880a^4} \dots, \quad (3.28)$$

$$F^{(1,2)} = \frac{95E_2^2 + 94E_4}{11520a^4} \dots, \quad F^{(0,3)} = -\frac{2E_2^2 + E_4}{384a^4} \dots. \quad (3.29)$$

One can easily check that $F^{(0,0)}$ in (3.22) completely agrees with the prepotential of the $\mathcal{N} = 2$ $SU(2)$ $N_f = 4$ gauge theory as derived for example in [5]; notice also that in the massless limit, i.e. $R, N, T_1, T_2 \rightarrow 0$, these amplitudes drastically simplify and precisely match the results presented in [35]. It is also interesting to observe that in the Nekrasov-Shatshvili limit where one of the ϵ_ℓ 's vanishes, the amplitudes $F^{(n,0)}$ of the $N_f = 4$ theory reduce to those of the $\mathcal{N} = 2^*$ theory upon setting $T_1 = T_2 = N = 0$ and $R = m^2$. Indeed, with these positions and rescaling $a \rightarrow 2a$, the amplitudes $F^{(n,0)}$ in (3.22)–(3.29) become twice the corresponding amplitudes of the $\mathcal{N} = 2^*$ theory given in (2.28)–(2.33). This simple relation does not hold away from the Nekrasov-Shatashvili limit: the amplitudes $F^{(n,g)}$ with $g \neq 0$ are in fact intrinsically different in the two theories, as a consequence of their different gravitational structure.

3.4 Recursion relations

Looking at the explicit expressions (3.19)–(3.21), it is not difficult to realize that the E_2 dependence of the h_ℓ 's is quite simple and exhibits a recursive pattern. This points again to the existence of a recursion relation among the h_ℓ 's involving their derivatives with respect to E_2 . Indeed, following the same steps described in section 2.4, one can check that

$$\frac{\partial h_\ell}{\partial E_2} = \frac{\ell}{6} \sum_{i=0}^{\ell-1} h_i h_{\ell-i-1} + \frac{\ell(2\ell-1)}{6} \epsilon_1 \epsilon_2 h_{\ell-1} \quad (3.30)$$

with the initial condition

$$\frac{\partial h_0}{\partial E_2} = 0. \quad (3.31)$$

This recursion relation has exactly the same structure of that of the $\mathcal{N} = 2^*$ theory given in (2.36), the only difference being in the numerical coefficients. Notice that the coefficients of the quadratic terms can be matched by a rescaling of the h_ℓ 's, but those of the linear terms proportional to $\epsilon_1 \epsilon_2$ remain different for the two theories; this is another signal of their intrinsically different behavior when a generic ϵ -background is considered.

Finally, if we expand the h_ℓ 's as in (2.39) we can reformulate the recursion relation (3.30) in terms of the partial amplitudes $f_\ell^{(n,g)}$ and get

$$\frac{\partial f_\ell^{(n,g)}}{\partial E_2} = \frac{\ell}{6} \sum'_{n_1, n_2; g_1, g_2} \left(\sum_{i=0}^{\ell-1} f_i^{(n_1, g_1)} f_{\ell-i-1}^{(n_2, g_2)} \right) + \frac{\ell(2\ell-1)}{6} f_{\ell-1}^{(n, g-1)} \quad (3.32)$$

where the $'$ means that the sum is performed over all n_1, n_2, g_1 and g_2 such that $n_1 + n_2 = n$ and $g_1 + g_2 = g$. This relation has to be supplemented by the initial conditions

$$f_0^{(0,0)} = 2R, \quad f_0^{(1,0)} = -\frac{1}{2}, \quad f_0^{(0,1)} = \frac{1}{2} \quad (3.33)$$

obtained from (3.19).

In the next section we will analyze the implications of the recursion relations and in particular their consequences on the modular transformation properties of the generalized prepotential.

4 Modular anomaly equations and S-duality

The fact that the generalized prepotential can be written in terms of (quasi) modular functions of the bare coupling constant allows us to explore the modularity properties of the deformed theory and study how the $Sl(2, \mathbb{Z})$ symmetry of the microscopic high-energy theory is realized in the effective low-energy theory. In the following we will concentrate on the $SU(2)$ $N_f = 4$ theory, even if our analysis and our methods can be equally well applied to the $\mathcal{N} = 2^* SU(2)$ model. Furthermore, we will focus on the generator \mathcal{S} of the modular group, corresponding to the following transformation of the bare coupling constant

$$\mathcal{S} : \tau_0 \rightarrow -\frac{1}{\tau_0}. \quad (4.1)$$

Notice that this action implies that the instanton counting parameter x in the Nekrasov's partition of the $N_f = 4$ theory, defined in (3.14), transforms as follows

$$\mathcal{S} : x \rightarrow 1 - x, \quad (4.2)$$

as one can readily check from the properties of the Jacobi θ -functions. This type of transformation is consistent with the interpretation of x as a cross-ratio, a fact that is also exploited in the AGT correspondence with the Liouville conformal blocks [32]. As discussed in [2], in the $N_f = 4$ theory the modular group acts with triality transformations on the mass invariants (3.4); in particular one has

$$\mathcal{S} : R \rightarrow R, \quad T_1 \rightarrow T_2, \quad T_2 \rightarrow T_1, \quad N \rightarrow N. \quad (4.3)$$

In the deformed theory, these rules have to be supplemented by those that specify how $Sl(2, \mathbb{Z})$ acts on the equivariant deformation parameters. Adapting to the present case the considerations made in [45] for the ϵ -deformed conformal Chern-Simons theory in three

dimensions, we assume that \mathcal{S} simply exchanges ϵ_1 and ϵ_2 with each other.⁵ In particular this means that $s = \epsilon_1 + \epsilon_2$ and $p = \epsilon_1\epsilon_2$ are left unchanged, i.e.

$$\mathcal{S} : s \rightarrow s, \quad p \rightarrow p. \quad (4.4)$$

Using the rules (4.1) - (4.4) and the modular properties of the Eisenstein series and Jacobi θ -functions, it is easy to show that the coefficients h_ℓ of the generalized prepotential (see (3.21) and appendix B) transform as quasi-modular forms of weight 2ℓ with anomalous terms due to the presence of the second Eisenstein series E_2 , namely

$$\mathcal{S} : h_\ell(E_2) \rightarrow \tau_0^{2\ell} h_\ell\left(E_2 + \frac{6}{\pi i \tau_0}\right) = \tau_0^{2\ell} \sum_{k=0}^{\ell} \frac{1}{k!} \left(\frac{1}{2\pi i \tau_0}\right)^k D^k h_\ell(E_2) \quad (4.5)$$

where we have introduced the convenient notation $D \equiv 12\partial_{E_2}$. These transformation rules are formally identical to those of the underformed $N_f = 4$ theory derived in [5] from the SW curve; however, since the h_ℓ 's satisfy a modified recursion relation with a new term proportional to $\epsilon_1\epsilon_2$, the practical effects of (4.5) in the deformed theory are different from those of the undeformed case, as we shall see momentarily.

To proceed let us consider the pair made by a and its S-dual image $\mathcal{S}(a) \equiv a_D$, on which \mathcal{S} acts as follows [1, 2]:

$$\mathcal{S} : \begin{pmatrix} a \\ a_D \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} a \\ a_D \end{pmatrix} = \begin{pmatrix} a_D \\ -a \end{pmatrix}. \quad (4.6)$$

We therefore have

$$\mathcal{S}^2(a) = -a. \quad (4.7)$$

In the SW theory this relation is enforced by taking

$$\mathcal{S}(a) = \frac{1}{2\pi i} \frac{\partial \mathcal{F}}{\partial a} \quad (4.8)$$

where \mathcal{F} is the undeformed effective prepotential which is related to its S-dual by a Legendre transform:

$$\mathcal{F} - \mathcal{S}(\mathcal{F}) = 2\pi i a \mathcal{S}(a). \quad (4.9)$$

It seems natural to try the same thing also in the deformed theory. Recalling that the effective generalized prepotential is

$$F = \pi i \tau_0 a^2 + h_0 \log \frac{a}{\Lambda} - \sum_{\ell=1}^{\infty} \frac{h_\ell}{2^{\ell+1} \ell} \frac{1}{a^{2\ell}} \quad (4.10)$$

where the first term is the classical part, we therefore posit

$$\mathcal{S}(a) = \frac{1}{2\pi i} \frac{\partial F}{\partial a} = \tau_0 a + \frac{1}{2\pi i} \sum_{\ell=0}^{\infty} \frac{h_\ell}{2^\ell} \frac{1}{a^{2\ell+1}}. \quad (4.11)$$

⁵Note that this rule is consistent with the interpretation of ϵ_1 and ϵ_2 as fluxes of (complex) combinations of NS-NS and R-R 3-form field strengths in a Type IIB string theory realization which rotate among themselves under S-duality.

Applying the S-duality rules (4.1) and (4.5), we obtain

$$\begin{aligned} \mathcal{S}^2(a) &= -\frac{\mathcal{S}(a)}{\tau_0} + \frac{1}{2\pi i} \sum_{\ell=0}^{\infty} \frac{\tau_0^{2\ell} \left(h_\ell + \frac{1}{2\pi i \tau_0} Dh_\ell + \mathcal{O}(\tau_0^{-2}) \right)}{2^\ell (\mathcal{S}(a))^{2\ell+1}} \\ &= -a + \frac{1}{(2\pi i \tau_0)^2} \left[\sum_{\ell=0}^{\infty} \frac{Dh_\ell}{2^\ell} \frac{1}{a^{2\ell+1}} - \sum_{\ell, n=0}^{\infty} \frac{(2\ell+1) h_\ell h_n}{2^{\ell+n}} \frac{1}{a^{2\ell+2n+3}} \right] + \mathcal{O}(\tau_0^{-3}). \end{aligned} \tag{4.12}$$

The expression in square brackets can be simplified using the recursion relation (3.30); like in the undeformed theory, the quadratic terms in the h 's exactly cancel but, due to the new ϵ -dependent term in the deformed recursion relation, an uncanceled part proportional to $\epsilon_1 \epsilon_2$ remains. This simple calculation shows that in order to enforce the relation (4.7) when $\epsilon_1 \epsilon_2 \neq 0$, the standard definition (4.11) has to be modified by adding terms proportional to $\epsilon_1 \epsilon_2$ in the right hand side. In the sequel we will work out explicitly the first corrections and show how the relation (4.7) constrains the form of $\mathcal{S}(a)$.

4.1 S-duality at first order in $\epsilon_1 \epsilon_2$

To organize the calculation, we introduce a set of generating functions φ_ℓ for the coefficients h_ℓ according to

$$\varphi_0 = -h_0 \log \frac{a}{\Lambda} + \sum_{\ell=1}^{\infty} \frac{h_\ell}{2^{\ell+1} \ell} \frac{1}{a^{2\ell}}, \quad \varphi_{\ell+1} = -\partial_a \varphi_\ell \quad \text{for } \ell \geq 0. \tag{4.13}$$

In particular we have the following relations with the generalized prepotential F :

$$\varphi_0 = \pi i \tau_0 a^2 - F, \tag{4.14a}$$

$$\varphi_1 = -2\pi i \tau_0 a + \partial_a F, \tag{4.14b}$$

$$\varphi_2 = 2\pi i \tau_0 - \partial_a^2 F \equiv 2\pi i \tau_0 - 2\pi i \tau, \tag{4.14c}$$

where in the last line we have introduced the effective coupling τ .

As shown in appendix C, the φ_ℓ 's form a ring under the action of D due to the generalized modular anomaly equations (3.30). This ring structure will enable us to formally express everything as functions of the φ_ℓ 's with coefficients that may depend on the product $\epsilon_1 \epsilon_2$. Such a dependence is a consequence of the $\epsilon_1 \epsilon_2$ -term in the recursion relation (3.30) and is the only explicit dependence on the deformation parameters that will be relevant for our purposes, all the rest being implicit inside the φ_ℓ 's and the h_ℓ 's therein.

As argued above, the definition (4.11) for $\mathcal{S}(a)$, namely $\mathcal{S}(a) = \tau_0 a + \varphi_1/(2\pi i)$, has to be replaced by a new one containing terms proportional to $\epsilon_1 \epsilon_2$, i.e.

$$\mathcal{S}(a) = \tau_0 a + \frac{\varphi_1}{2\pi i} + \frac{X}{2\pi i} \tag{4.15}$$

with

$$X = \epsilon_1 \epsilon_2 X_1 + (\epsilon_1 \epsilon_2)^2 X_2 + \dots \tag{4.16}$$

where the X_ℓ 's have to be determined by imposing the constraint (4.7). Notice that in the undeformed theory a and $\mathcal{S}(a)$ can be regarded as a pair of canonically conjugated

variables [1, 2, 46]; when the $\epsilon_1\epsilon_2$ -deformation is turned on, the relations (4.15) and (4.16) have thus to be understood as a (sort of) semi-classical expansion [47]. Applying S-duality to (4.15), it is straightforward to obtain

$$\mathcal{S}^2(a) = -a + \frac{1}{2\pi i\tau_0} \left[(\tau_0 \mathcal{S}(\varphi_1) - \varphi_1) + (\tau_0 \mathcal{S}(X) - X) \right]; \quad (4.17)$$

therefore, in order to satisfy the relation (4.7), the expression in the square brackets above must vanish. Expanding this condition in $\epsilon_1\epsilon_2$, we obtain the following constraints

$$(\tau_0 \mathcal{S}(\varphi_1) - \varphi_1) \Big|_0 = 0, \quad (4.18a)$$

$$(\tau_0 \mathcal{S}(\varphi_1) - \varphi_1) \Big|_n + \sum_{k=1}^n (\tau_0 \mathcal{S}(X_k) - X_k) \Big|_{n-k} = 0 \quad (4.18b)$$

where the symbol $\Big|_n$ means taking the coefficient of $(\epsilon_1\epsilon_2)^n$ (notice that the φ_ℓ 's defined in (4.13) do not have any explicit dependence on $\epsilon_1\epsilon_2$ and thus $\varphi_\ell \Big|_n = \delta_{n0} \varphi_\ell$; for the same reason we also have $X_\ell \Big|_n = \delta_{n0} X_\ell$).

Let us now compute $\mathcal{S}(\varphi_1)$. Using the S-duality rules (4.1) and (4.5), and exhibiting temporarily the dependencies on E_2 and a which are the only relevant ones for our purposes, we obtain

$$\begin{aligned} \tau_0 \mathcal{S}(\varphi_1(E_2; a)) &= \sum_{\ell=0}^{\infty} \frac{\tau_0^{2\ell+1} h_\ell \left(E_2 + \frac{6}{\pi i\tau_0} \right)}{2^\ell (\mathcal{S}(a))^{2\ell+1}} = \sum_{\ell=0}^{\infty} \frac{h_\ell \left(E_2 + \frac{6}{\pi i\tau_0} \right)}{2^\ell \left(a + \frac{\varphi_1}{2\pi i\tau_0} + \frac{X}{2\pi i\tau_0} \right)^{2\ell+1}} \\ &= \varphi_1 \left(E_2 + \frac{6}{\pi i\tau_0}; a + \frac{\varphi_1}{2\pi i\tau_0} + \frac{X}{2\pi i\tau_0} \right) \\ &= e^{\frac{1}{2\pi i\tau_0} (D + \zeta \partial_a)} \varphi_1(E_2; a) \Big|_{\zeta = \varphi_1 + X}. \end{aligned} \quad (4.19)$$

Actually, this is a particular case of the more general result

$$\tau_0^\ell \mathcal{S}(\varphi_\ell) = e^{\frac{1}{2\pi i\tau_0} (D + \zeta \partial_a)} \varphi_\ell \Big|_{\zeta = \varphi_1 + X}. \quad (4.20)$$

Expanding the exponential and using the relation

$$(D + \zeta \partial_a)^n \varphi_\ell \Big|_{\zeta = \varphi_1 + X} = \sum_{k=0}^n (-1)^k \binom{n}{k} X^k (D + \zeta \partial_a)^n \varphi_{\ell+k} \Big|_{\zeta = \varphi_1}, \quad (4.21)$$

after some straightforward algebra to rearrange the various terms, we obtain

$$\tau_0^\ell \mathcal{S}(\varphi_\ell) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \left(\frac{X}{2\pi i\tau_0} \right)^n \Sigma^{(\ell+n)} \quad (4.22)$$

where the functions $\Sigma^{(\ell)}$ are defined by

$$\Sigma^{(\ell)} = e^{\frac{1}{2\pi i\tau_0} (D + \zeta \partial_a)} \varphi_\ell \Big|_{\zeta = \varphi_1}. \quad (4.23)$$

As shown in appendix C, these functions satisfy the remarkably simple relation

$$\Sigma^{(\ell+1)} = -\frac{\tau_0}{\tau} \partial_a \Sigma^{(\ell)} \tag{4.24}$$

that is a consequence of the ring properties obeyed by the functions φ_ℓ , which in turn are due to the modular anomaly equations (3.30). In view of this we can therefore rewrite (4.22) as

$$\tau_0^\ell \mathcal{S}(\varphi_\ell) = e^{\frac{\zeta}{2\pi i \tau}} \partial_a \Sigma^{(\ell)} \Big|_{\zeta=X} . \tag{4.25}$$

Note that the bare coupling τ_0 initially appearing in the right hand side of (4.20) has been dressed into the effective coupling τ .

We now exploit this result and proceed perturbatively in $\epsilon_1 \epsilon_2$ to obtain explicit expressions. At the zeroth order in $\epsilon_1 \epsilon_2$, from (4.25) we have

$$\tau_0 \mathcal{S}(\varphi_1) \Big|_0 = \Sigma^{(1)} \Big|_0 = \varphi_1 \tag{4.26}$$

where the last equality follows from (C.13). The constraint (4.18a) is therefore identically satisfied. This is no surprise since we have already shown that the relation (4.7) must be true up to terms proportional to $\epsilon_1 \epsilon_2$. At the first order in the deformation parameters we get instead

$$\tau_0 \mathcal{S}(\varphi_1) \Big|_1 = \Sigma^{(1)} \Big|_1 + \frac{X_1}{2\pi i \tau} \partial_a \Sigma^{(1)} \Big|_0 = \frac{\tau_0 \varphi_3}{2(2\pi i) \tau^2} - \frac{X_1 \varphi_2}{2\pi i \tau} \tag{4.27}$$

where we have used (C.13). Inserting this result into (4.18b) for $n = 1$, we obtain the following equation for X_1

$$\mathcal{S}(X_1) \Big|_0 = \frac{X_1}{\tau} - \frac{\varphi_3}{2(2\pi i) \tau^2} . \tag{4.28}$$

To solve it we take advantage of the S-duality relations at the zeroth order in $\epsilon_1 \epsilon_2$, which are formally the same of the SW theory. Using (4.24) - (4.26) it is easy to show that

$$\mathcal{S}(\varphi_2) \Big|_0 = \frac{\varphi_2}{\tau_0 \tau}, \quad \mathcal{S}(\varphi_3) \Big|_0 = \frac{\varphi_3}{\tau^3}, \tag{4.29}$$

and also that

$$\mathcal{S}(\tau) \Big|_0 = -\frac{1}{\tau} . \tag{4.30}$$

Equipped with these results, one can check that a solution to (4.28) is given by

$$X_1 = \frac{\varphi_3}{8\pi i \tau} . \tag{4.31}$$

In conclusion⁶ we find that the S-dual image of a , which obeys the constraint (4.7) up to terms of order $(\epsilon_1 \epsilon_2)^2$, is

$$\mathcal{S}(a) = \tau_0 a + \frac{\varphi_1}{2\pi i} + \epsilon_1 \epsilon_2 \frac{\varphi_3}{4(2\pi i)^2 \tau} + \mathcal{O}((\epsilon_1 \epsilon_2)^2) . \tag{4.32}$$

⁶Note that actually the solution (4.31) is not unique, since in principle one could add to X_1 a term Y_1 such that $\mathcal{S}(Y_1) = Y_1/\tau$. A detailed analysis shows that a term of this type is of the form $\alpha \varphi_3/(\tau^2 + 1)$ with α constant, which has different pole structure in τ with respect to X_1 . We therefore do not consider this possibility and take $\alpha = 0$.

Observing that $\varphi_3 = 2\pi i \partial_a \tau$, we can rewrite the $\epsilon_1 \epsilon_2$ -term above also as

$$\frac{\epsilon_1 \epsilon_2}{4(2\pi i)} \partial_a \log \left(\frac{\tau}{\tau_0} \right), \tag{4.33}$$

so that (4.32) becomes

$$2i\pi \mathcal{S}(a) = \partial_a F + \frac{\epsilon_1 \epsilon_2}{4} \partial_a \log \left(\frac{\tau}{\tau_0} \right) + \mathcal{O}((\epsilon_1 \epsilon_2)^2). \tag{4.34}$$

In the following we will investigate the implications of this result, while we refer to appendix D for its extension to the second order in $\epsilon_1 \epsilon_2$.

4.2 S-duality on the prepotential

An immediate consequence of the ϵ -correction in (4.34) is that $\mathcal{S}(a)$ is not simply proportional to the derivative of the prepotential F ; thus it is natural to expect that in the deformed theory the S-dual of the prepotential is not simply given by a Legendre transformation as it happens instead in the undeformed SW case (see (4.9)). In [38] the relation between F and $\mathcal{S}(F)$ has been conjectured to be a deformed Fourier transformation. Here we reach the same conclusion, even though from a different perspective since for us all deviations from the undeformed theory are parametrized by the product $\epsilon_1 \epsilon_2$.

We can compute the first ϵ -corrections to the relation between F and $\mathcal{S}(F)$ using the same methods of the previous subsection and the results given in appendix C. The starting point is the relation between φ_0 and the generalized prepotential (see (4.14a)); from this we get the useful identity

$$F - \mathcal{S}(F) = 2\pi i a \mathcal{S}(a) + \frac{\pi i}{\tau_0} (\mathcal{S}(a) - a\tau_0)^2 + \mathcal{S}(\varphi_0) - \varphi_0. \tag{4.35}$$

Using (4.15) and (4.22) for $\ell = 0$, we can easily rewrite the right hand side and obtain

$$\begin{aligned} F - \mathcal{S}(F) &= 2\pi i a \mathcal{S}(a) + \frac{\varphi_1^2}{2(2\pi i)\tau_0} + \Sigma^{(0)} - \varphi_0 - \frac{X}{2\pi i\tau_0} (\Sigma^{(1)} - \varphi_1) \\ &\quad + \frac{1}{2} \left(\frac{X}{2\pi i\tau_0} \right)^2 (\Sigma^{(2)} + 2\pi i\tau_0) + \dots \end{aligned} \tag{4.36}$$

where the dots stand for terms of order X^3 which are at least $\mathcal{O}((\epsilon_1 \epsilon_2)^3)$. Notice that the difference $\Sigma^{(1)} - \varphi_1$ is of order $\epsilon_1 \epsilon_2$, as one can see from (4.26); thus the linear term in X gives contributions $\mathcal{O}((\epsilon_1 \epsilon_2)^2)$, like the X^2 term. This means that the knowledge of the first order correction X_1 obtained in the previous subsection is enough to compute the correction to $F - \mathcal{S}(F)$ at order $(\epsilon_1 \epsilon_2)^2$. This mechanism actually works at all orders, namely the k -th order coefficient of $F - \mathcal{S}(F)$ does not depend on X_k but only on X_j with $j < k$.

Using the explicit value of X_1 given in (4.31) and the expressions for the $\Sigma^{(\ell)}$'s given in appendix C (see in particular (C.12) - (C.14)), after straightforward algebra we obtain

$$F - \mathcal{S}(F) = 2\pi i a \mathcal{S}(a) - \frac{\epsilon_1 \epsilon_2}{2} \log \left(\frac{\tau}{\tau_0} \right) + (\epsilon_1 \epsilon_2)^2 \left(\frac{1}{8} \frac{\varphi_4}{(2i\pi\tau)^2} + \frac{11}{96} \frac{\varphi_3^2}{(2i\pi\tau)^3} \right) + \mathcal{O}((\epsilon_1 \epsilon_2)^3). \tag{4.37}$$

This shows that the simple Legendre transform relation (4.9), which holds in the undeformed theory, does not work any more when $\epsilon_1\epsilon_2 \neq 0$, as argued, from a different point of view, in [38]. In the next section, however, we will show that with a suitable redefinition of the prepotential F and of the coupling constant τ it is possible to recover the standard Legendre transform relation also in the $\epsilon_1\epsilon_2$ -deformed theory.

5 Conclusions

The explicit first-order calculation of section 4.1 shows that $\mathcal{S}(a)$ is not simply the derivative of the prepotential F . However, it is still a total derivative, as is clear from (4.34). This feature is maintained also at the second order. Indeed, as shown in appendix D, the $(\epsilon_1\epsilon_2)^2$ correction X_2 can be chosen as

$$\begin{aligned} X_2 &= \frac{1}{16} \frac{\varphi_5}{(2\pi i\tau)^2} + \frac{23}{96} \frac{\varphi_3\varphi_4}{(2\pi i\tau)^3} + \frac{11}{64} \frac{\varphi_3^3}{(2\pi i\tau)^4} \\ &= \partial_a \left(-\frac{1}{16} \frac{\varphi_4}{(2\pi i\tau)^2} - \frac{11}{192} \frac{\varphi_3^2}{(2\pi i\tau)^3} \right), \end{aligned} \tag{5.1}$$

so that we can rewrite (4.15) in the following way

$$\mathcal{S}(a) = \frac{1}{2\pi i} \partial_a \hat{F} \tag{5.2}$$

with

$$\hat{F} = F + \frac{\epsilon_1\epsilon_2}{4} \log \left(\frac{\tau}{\tau_0} \right) - \left(\frac{\epsilon_1\epsilon_2}{4} \right)^2 \left(\frac{\varphi_4}{(2\pi i\tau)^2} + \frac{11}{12} \frac{\varphi_3^2}{(2\pi i\tau)^3} \right) + \mathcal{O}((\epsilon_1\epsilon_2)^3). \tag{5.3}$$

In the deformed theory the S-duality transformation of the effective coupling τ does not have a simple form; in fact, using (4.14c) and the transformation properties of φ_2 , it is easy to show that (see (D.7))

$$\mathcal{S}(\tau) = -\frac{1}{\tau} - \epsilon_1\epsilon_2 \left(\frac{1}{2} \frac{\varphi_4}{(2\pi i)^2\tau^3} + \frac{3}{4} \frac{\varphi_3^2}{(2\pi i)^3\tau^4} \right) + \mathcal{O}((\epsilon_1\epsilon_2)^2). \tag{5.4}$$

However, there is a modified effective coupling on which S-duality acts in a simple way. This is

$$\hat{\tau} \equiv \partial_a \mathcal{S}(a) = \frac{1}{2\pi i} \partial_a^2 \hat{F} = \tau \left[1 - \frac{\epsilon_1\epsilon_2}{4} \left(\frac{\varphi_4}{(2\pi i\tau)^2} + \frac{\varphi_3^2}{(2\pi i\tau)^3} \right) + \mathcal{O}((\epsilon_1\epsilon_2)^2) \right]. \tag{5.5}$$

One can easily check that

$$\mathcal{S}(\hat{\tau}) = \partial_{\mathcal{S}(a)} \mathcal{S}^2(a) = -\partial_{\mathcal{S}(a)} a = -\frac{1}{\hat{\tau}}. \tag{5.6}$$

Thus, in the effective ϵ -deformed theory it is $\hat{\tau}$, and not τ , that exhibits the same behavior of the bare coupling τ_0 under S-duality. We also observe that if one uses $\hat{\tau}$, the expression of the extended prepotential \hat{F} given in (5.3) simplifies and becomes

$$\hat{F} = F + \frac{\epsilon_1\epsilon_2}{4} \log \left(\frac{\hat{\tau}}{\tau_0} \right) + \left(\frac{\epsilon_1\epsilon_2}{4} \right)^2 \frac{\varphi_3^2}{12(2\pi i\hat{\tau})^3} + \mathcal{O}((\epsilon_1\epsilon_2)^3). \tag{5.7}$$

This result seems to suggest that it is possible to write the higher order corrections only in terms of φ_3 which, being the triple derivative of the prepotential, is proportional to the Yukawa coupling C_{aaa} , the rank-three symmetric tensor playing a crucial rôle in special geometry. Moreover, one can verify the simple Legendre transform relation

$$\widehat{F} - \mathcal{S}(\widehat{F}) = 2\pi i a \mathcal{S}(a) \tag{5.8}$$

up to terms of order $(\epsilon_1 \epsilon_2)^3$. It is quite natural to expect that this pattern extends also to higher orders.

Our detailed analysis shows that when both deformation parameters ϵ_1 and ϵ_2 are non-vanishing, besides the ϵ -dependent structures generated by the Nekrasov partition functions, the effective theory seems to require a new series of explicit $\epsilon_1 \epsilon_2$ -corrections in order to have S-duality acting in the proper way. These new corrections, being proportional to inverse powers of the coupling constant, appear to correspond to perturbative terms at higher loops and are absent in the Nekrasov-Shatashvili limit. We find remarkable that by using the modified prepotential \widehat{F} and the modified coupling $\widehat{\tau}$, all S-duality relations acquire the standard simple form as in the undeformed theory (see (5.2), (5.6) and (5.8)). It would be very nice to see whether these results admit an interpretation in the context of special geometry or in a more general geometric set-up that would allow us to go beyond the perturbative approach in the deformation parameters we have used in this paper. It would be interesting also to study the recursion relations obeyed by the functions φ_ℓ 's which can be iteratively obtained from the viscous Burgers equation satisfied by φ_1 (see (C.2)).

We conclude by observing that all S-duality formulas we have derived for the SU(2) $N_f = 4$ theory can be obtained for the $\mathcal{N} = 2^*$ SU(2) theory as well. The only difference in this case is that every explicit occurrence of $\epsilon_1 \epsilon_2$ has to be replaced by $(\epsilon_1 \epsilon_2)/2$ to take into account the different normalization of the viscous term in the modular anomaly recursion relation (see (2.36) as compared to (3.30)).

Acknowledgments

We thank Francesco Fucito and Francisco Morales for very useful discussions. This work was supported in part by the MIUR-PRIN contract 2009-KHZKRX.

A Modular functions

We collect here some useful formulas involving the modular functions we used.

θ -functions. The Jacobi θ -functions are defined as

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} (v|\tau) = \sum_{n \in \mathbb{Z}} q^{(n - \frac{a}{2})^2} e^{2\pi i (n - \frac{a}{2})(v - \frac{b}{2})}, \tag{A.1}$$

for $a, b = 0, 1$ and $q = e^{\pi i \tau}$. We simplify the notation by writing, as usual, $\theta_1 \equiv \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $\theta_2 \equiv \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\theta_3 \equiv \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, $\theta_4 \equiv \theta \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. The functions θ_a , $a = 2, 3, 4$, satisfy the “aequatio identica satis abstrusa”

$$\theta_3^4 - \theta_2^4 - \theta_4^4 = 0, \tag{A.2}$$

and admit the following series expansions

$$\begin{aligned}\theta_2(0|\tau) &= 2q^{\frac{1}{4}}(1 + q^2 + q^6 + q^{12} + \dots), \\ \theta_3(0|\tau) &= 1 + 2q + 2q^4 + 2q^9 + \dots, \\ \theta_4(0|\tau) &= 1 - 2q + 2q^4 - 2q^9 + \dots.\end{aligned}\tag{A.3}$$

η -function. The Dedekind η -function is defined by

$$\eta(q) = q^{\frac{1}{12}} \prod_{n=1}^{\infty} (1 - q^{2n}).\tag{A.4}$$

Eisenstein series. The first Eisenstein series can be expressed as follows:

$$\begin{aligned}E_2 &= 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^{2n} = 1 - 24q^2 - 72q^4 - 96q^6 + \dots, \\ E_4 &= 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) q^{2n} = 1 + 240q^2 + 2160q^4 + 6720q^6 + \dots, \\ E_6 &= 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^{2n} = 1 - 504q^2 - 16632q^4 - 122976q^6 + \dots,\end{aligned}\tag{A.5}$$

where $\sigma_k(n)$ is the sum of the k -th power of the divisors of n , i.e., $\sigma_k(n) = \sum_{d|n} d^k$. The series E_4 and E_6 are related to the θ -functions in the following way

$$E_4 = \frac{1}{2}(\theta_2^8 + \theta_3^8 + \theta_4^8), \quad E_6 = \frac{1}{2}(\theta_3^4 + \theta_4^4)(\theta_2^4 + \theta_3^4)(\theta_4^4 - \theta_2^4).\tag{A.6}$$

The series E_2 , E_4 and E_6 are connected among themselves by logarithmic q -derivatives according to

$$q\partial_q E_2 = \frac{1}{6}(E_2^2 - E_4), \quad q\partial_q E_4 = \frac{2}{3}(E_4 E_2 - E_6), \quad q\partial_q E_6 = E_6 E_2 - E_4^2.\tag{A.7}$$

Also the derivatives of the functions θ_a^4 have simple expressions:

$$q\partial_q \theta_2^4 = \frac{\theta_2^4}{3}(E_2 + \theta_3^4 + \theta_4^4), \quad q\partial_q \theta_3^4 = \frac{\theta_3^4}{3}(E_2 + \theta_2^4 - \theta_4^4), \quad q\partial_q \theta_4^4 = \frac{\theta_4^4}{3}(E_2 - \theta_2^4 - \theta_3^4).\tag{A.8}$$

Modular transformations. Under the $\text{Sl}(2, \mathbb{Z})$ modular transformation

$$\tau \rightarrow \tau' = \frac{a\tau + b}{c\tau + d} \quad \text{with } a, b, c, d \in \mathbb{Z} \quad \text{and} \quad ad - bc = 1,\tag{A.9}$$

the Eisenstein series E_4 and E_6 behave as modular forms of weight 4 and 6, respectively:

$$E_4(\tau') = (c\tau + d)^4 E_4(\tau), \quad E_6(\tau') = (c\tau + d)^6 E_6(\tau).\tag{A.10}$$

The series E_2 , instead, is a quasi modular form of degree 2:

$$E_2(\tau') = (c\tau + d)^2 E_2(\tau) + \frac{6}{i\pi} c(c\tau + d).\tag{A.11}$$

Under the generators \mathcal{T} and \mathcal{S} of the modular group the θ -functions and the Dedekind η function transform as follows

$$\begin{aligned}\mathcal{T} : \quad & \theta_3^4 \leftrightarrow \theta_4^4, \quad \theta_2^4 \rightarrow -\theta_2^4, \quad \eta \rightarrow e^{\frac{i\pi}{12}} \eta, \\ \mathcal{S} : \quad & \theta_2^4 \rightarrow -\tau^2 \theta_4^4, \quad \theta_3^4 \rightarrow -\tau^2 \theta_3^4, \quad \theta_4^4 \rightarrow -\tau^2 \theta_2^4, \quad \eta \rightarrow \sqrt{-i\tau} \eta.\end{aligned}\tag{A.12}$$

B The coefficient h_3 of the $SU(2)$ $N_f = 4$ prepotential

Here we give the explicit expression of the coefficient h_3 up to three instantons obtained using localization. It is given by

$$\begin{aligned}
 h_3 = & \frac{1}{1344} \left[768R^4 - 21504R^2T_1 + 12288RN + 26112T_1^2 - 6144T_1T_2 - 6144T_2^2 \right. \\
 & - (1792R^3 - 26880RT_1 + 10752N)(s^2 - 2p) + R^2(1568s^4 - 6272s^2p + 5376p^2) \\
 & - T_1(9408s^4 - 37632s^2p + 32256p^2) - R(496s^6 - 2976s^4p + 5248s^2p^2 - 2560p^3) \\
 & \left. + 51s^8 - 408s^6p + 1102s^4p^2 - 1144s^2p^3 + 357p^4 \right] + 24s^4(T_1 + 2T_2)q \\
 & - \frac{1}{4} \left[3840(T_1 + 2T_2)^2 - 1152(RT_1 + 2N)(5s^2 - 2p) \right. \\
 & + 48(R^2 + 6T_1)(16s^4 - 32s^2p + 7p^2) - 192R(2s^6 - 8s^4p + 7s^2p^2 - p^3) \\
 & \left. + 3(16s^8 - 96s^6p + 168s^4p^2 - 88s^2p^3 + 9p^4) \right] q^2 \\
 & + 32(T_1 + 2T_2) \left[160(R^2 + 6T_1) - 2R(280s^2 - 272p) \right. \\
 & \left. + 313s^4 - 856s^2p + 366p^2 \right] q^3 + \mathcal{O}(q^4). \tag{B.1}
 \end{aligned}$$

Following the procedure described in the main text, we can rewrite h_3 in terms of (quasi) modular functions according to

$$\begin{aligned}
 h_3 = & \frac{1}{3456} (4R - s^2 + p)(4R - s^2 + 3p) \left[80R^2 - 8R(5s^2 - 22p) + 5s^4 - 44s^2p + 99p^2 \right] E_2^3 \\
 & - \frac{1}{12} \left[80R^2 - 8R(5s^2 - 22p) + 5s^4 - 44s^2p + 99p^2 \right] (T_1\theta_4^4 - T_2\theta_2^4) E_2^2 \\
 & + \frac{1}{1440} (4R - s^2 + 6p) \left[64R^3 - 80R^2(3s^2 - 4p) + 4R(27s^4 - 88s^2p + 49p^2) \right. \\
 & \quad \left. - 13s^6 + 68s^4p - 94s^2p^2 + 30p^3 + 2304N \right] E_2 E_4 \\
 & - \frac{4}{3} (4R - s^2 + 6p)(R - s^2 + p) \left[T_1\theta_4^4(2\theta_2^4 + \theta_4^4) + T_2\theta_2^4(\theta_2^4 + 2\theta_4^4) \right] E_2 \\
 & + 8(T_1\theta_4^4 - T_2\theta_2^4)^2 E_2 + 16 \left[T_1^2(\theta_4^4 + 2\theta_2^4)\theta_4^8 - T_2^2(\theta_2^4 + 2\theta_4^4)\theta_2^8 - T_1T_2\theta_2^4\theta_4^4(\theta_2^4 - \theta_4^4) \right] \\
 & - \frac{1}{4} \left[16R^2 - 8R(5s^2 - 6p) + 21s^4 - 60s^2p + 31p^2 \right] (T_1\theta_4^4 - T_2\theta_2^4) E_4 \\
 & + \frac{1}{120960} \left[2816R^4 - 30464R^3s^2 + 39424R^3p + 67872R^2s^4 - 190848R^2s^2p \right. \\
 & + 94304R^2p^2 - 28400Rs^6 + 134112Rs^4p - 165664Rs^2p^2 + 47616Rp^3 \\
 & - 774144N(s^2 - p) + 331776NR - 552960(T_1^2 + T_1T_2 + T_2^2) \\
 & \left. + 3323s^8 - 22216s^6p + 46862s^4p^2 - 34584s^2p^3 + 6615p^4 \right] E_6. \tag{B.2}
 \end{aligned}$$

By expanding the modular functions in powers of q as shown in appendix A, one can recover the instanton terms in (B.1).

C Reformulating the modular anomaly equations

The modular anomaly equation (3.30) implies the following relation:

$$D\varphi_0 = \frac{1}{2}\varphi_1^2 + \frac{\epsilon_1\epsilon_2}{2}\varphi_2 \quad (\text{C.1})$$

where the functions φ_ℓ 's have been defined in (4.13). Since the operators D and $(-\partial_a)$ commute, by applying the latter to (C.1) and remembering that $\varphi_{\ell+1} = -\partial_a\varphi_\ell$, it is straightforward to obtain the action of D on any φ_ℓ and verify that these functions form a ring under it. For example, at the next step we obtain

$$D\varphi_1 = \varphi_1\varphi_2 + \frac{\epsilon_1\epsilon_2}{2}\varphi_3 = \varphi_1(-\partial_a\varphi_1) + \frac{\epsilon_1\epsilon_2}{2}\partial_a^2\varphi_1. \quad (\text{C.2})$$

which is, up to rescalings, the viscous Burgers equation

$$\partial_t u = u\partial_x u + \nu\partial_x^2 u \quad (\text{C.3})$$

with the viscosity ν proportional to $\epsilon_1\epsilon_2$.

In section 4 the S-duality requirements were formulated in terms of the quantities $\Sigma^{(\ell)}$ defined in (4.23), which we rewrite here for convenience as follows:

$$\Sigma^{(\ell)} = \sum_{n=0}^{\infty} \frac{1}{n!(2\pi i\tau_0)^n} P_n^{(\ell)} \quad (\text{C.4})$$

with

$$P_n^{(\ell)} = (D + \zeta\partial_a)^n \varphi_\ell \Big|_{\zeta=\varphi_1}. \quad (\text{C.5})$$

From this definition it is easy to show that the following relation holds:

$$P_n^{(\ell+1)} = -\partial_a P_n^{(\ell)} + n\varphi_2 P_{n-1}^{(\ell)}. \quad (\text{C.6})$$

In turn, this implies that

$$\Sigma^{(\ell+1)} = -\partial_a \Sigma^{(\ell)} + \frac{\varphi_2}{2\pi i\tau_0} \Sigma^{(\ell+1)}. \quad (\text{C.7})$$

Recalling that $2\pi i\tau_0 - \varphi_2 = 2\pi i\tau$ (see (4.14c)), this is tantamount to the recursion relation (4.24), namely

$$\Sigma^{(\ell+1)} = -\frac{\tau_0}{\tau} \partial_a \Sigma^{(\ell)}, \quad (\text{C.8})$$

which allows us to easily obtain the expression of any $\Sigma^{(\ell)}$ once $\Sigma^{(0)}$ is known.

To determine $\Sigma^{(0)}$ we start considering the quantities $P_n^{(0)}$. Clearly, from (C.5) we have $P_0^{(0)} = \varphi_0$. The next case is

$$P_1^{(0)} = (D + \zeta\partial_a)\varphi_0 \Big|_{\zeta=\varphi_1} = D\varphi_0 - \varphi_1^2 = -\frac{1}{2}\varphi_1^2 + \frac{\epsilon_1\epsilon_2}{2}\varphi_2 \quad (\text{C.9})$$

where the last step follows from (C.1). By further applications of the operator $(D + \zeta\partial_a)$, we get

$$\begin{aligned} P_2^{(0)} &= \frac{\epsilon_1\epsilon_2}{2}\varphi_2^2 + \left(\frac{\epsilon_1\epsilon_2}{2}\right)^2\varphi_4, \\ P_3^{(0)} &= \epsilon_1\epsilon_2\varphi_2^3 + \left(\frac{\epsilon_1\epsilon_2}{2}\right)^2(6\varphi_2\varphi_4 + 5\varphi_3^2) + \left(\frac{\epsilon_1\epsilon_2}{2}\right)^3\varphi_6, \end{aligned} \quad (\text{C.10})$$

with similar expressions for higher values of n . In fact, it is possible to derive the general expression of the $P_n^{(0)}$'s at the first orders in their explicit dependence on $\epsilon_1 \epsilon_2$:

$$\begin{aligned}
 P_n^{(0)} \Big|_0 &= \delta_{n,0} \varphi_0 - \delta_{n,1} \frac{\varphi_1^2}{2}, \\
 P_n^{(0)} \Big|_1 &= \frac{(n-1)!}{2} \varphi_2^n, \\
 P_n^{(0)} \Big|_2 &= \frac{(n-1)!}{4} \left(\frac{1}{2} \varphi_4 \left(\frac{\partial}{\partial \varphi_2} \right)^2 + \frac{5}{12} \varphi_3^2 \left(\frac{\partial}{\partial \varphi_2} \right)^3 \right) \varphi_2^n.
 \end{aligned} \tag{C.11}$$

Inserting these results into (C.4), we obtain

$$\begin{aligned}
 \Sigma^{(0)} \Big|_0 &= \varphi_0 - \frac{\varphi_1^2}{2(2\pi i \tau_0)}, \\
 \Sigma^{(0)} \Big|_1 &= -\frac{1}{2} \log \left(1 - \frac{\varphi_2}{2\pi i \tau_0} \right) = -\frac{1}{2} \log \frac{\tau}{\tau_0}, \\
 \Sigma^{(0)} \Big|_2 &= \frac{1}{8} \frac{\varphi_4}{(2\pi i \tau)^2} + \frac{5}{24} \frac{\varphi_3^2}{(2\pi i \tau)^3},
 \end{aligned} \tag{C.12}$$

where from the second equality on we have used that $2\pi i \tau_0 - \varphi_2 = 2\pi i \tau$.

Applying the recursion formula (C.8) we can easily get the explicit expressions for the $\Sigma^{(\ell)}$'s that are needed in the calculations presented in section 4 or in appendix D. They are, for $\ell = 1$:

$$\begin{aligned}
 \Sigma^{(1)} \Big|_0 &= \varphi_1, \\
 \Sigma^{(1)} \Big|_1 &= 2\pi i \tau_0 \left(\frac{1}{2} \frac{\varphi_3}{(2\pi i \tau)^2} \right), \\
 \Sigma^{(1)} \Big|_2 &= 2\pi i \tau_0 \left(\frac{1}{8} \frac{\varphi_5}{(2\pi i \tau)^3} + \frac{2}{3} \frac{\varphi_3 \varphi_4}{(2\pi i \tau)^4} + \frac{5}{8} \frac{\varphi_3^3}{(2\pi i \tau)^5} \right);
 \end{aligned} \tag{C.13}$$

for $\ell = 2$:

$$\begin{aligned}
 \Sigma^{(2)} \Big|_0 &= 2\pi i \tau_0 \left(\frac{\varphi_2}{2\pi i \tau} \right), \\
 \Sigma^{(2)} \Big|_1 &= (2\pi i \tau_0)^2 \left(\frac{1}{2} \frac{\varphi_4}{(2\pi i \tau)^3} + \frac{\varphi_3^2}{(2\pi i \tau)^4} \right);
 \end{aligned} \tag{C.14}$$

for $\ell = 3$:

$$\begin{aligned}
 \Sigma^{(3)} \Big|_0 &= (2\pi i \tau_0)^3 \left(\frac{\varphi_3}{(2\pi i \tau)^3} \right), \\
 \Sigma^{(3)} \Big|_1 &= (2\pi i \tau_0)^3 \left(\frac{1}{2} \frac{\varphi_5}{(2\pi i \tau)^4} + \frac{7}{2} \frac{\varphi_3 \varphi_4}{(2\pi i \tau)^5} + 4 \frac{\varphi_3^3}{(2\pi i \tau)^6} \right);
 \end{aligned} \tag{C.15}$$

for $\ell = 4$:

$$\Sigma^{(4)} \Big|_0 = (2\pi i \tau_0)^4 \left(\frac{\varphi_4}{(2\pi i \tau)^4} + 3 \frac{\varphi_3^2}{(2\pi i \tau)^5} \right); \tag{C.16}$$

and finally for $\ell = 5$:

$$\Sigma^{(5)} \Big|_0 = (2\pi i \tau_0)^5 \left(\frac{\varphi_5}{(2\pi i \tau)^5} + 10 \frac{\varphi_3 \varphi_4}{(2\pi i \tau)^6} + 15 \frac{\varphi_3^3}{(2\pi i \tau)^7} \right). \tag{C.17}$$

D S-duality at order $(\epsilon_1\epsilon_2)^2$

We start from the relation (4.18b) for $n = 2$, namely

$$\tau_0 \mathcal{S}(X_2) \Big|_0 - X_2 = -\tau_0 \mathcal{S}(\varphi_1) \Big|_2 - \tau_0 \mathcal{S}(X_1) \Big|_1. \quad (\text{D.1})$$

From (4.22) we read that

$$\tau_0 \mathcal{S}(\varphi_1) \Big|_2 = \Sigma^{(1)} \Big|_2 - \frac{X_1}{2\pi i \tau_0} \Sigma^{(2)} \Big|_1 - \frac{X_2}{2\pi i \tau_0} \Sigma^{(2)} \Big|_0 + \frac{1}{2} \frac{X_1^2}{(2\pi i \tau_0)^2} \Sigma^{(3)} \Big|_0. \quad (\text{D.2})$$

Substituting the expressions of the various $\Sigma^{(\ell)} \Big|_k$'s given in appendix C, and that of X_1 given in (4.31), we get

$$\tau_0 \mathcal{S}(\varphi_1) \Big|_2 = 2\pi i \tau_0 \left(\frac{1}{8} \frac{\varphi_5}{(2\pi i \tau)^3} + \frac{13}{24} \frac{\varphi_4 \varphi_3}{(2\pi i \tau)^4} + \frac{13}{32} \frac{\varphi_3^3}{(2\pi i \tau)^5} \right) - \frac{\varphi_2}{2\pi i \tau} X_2. \quad (\text{D.3})$$

On the other hand, from (4.31) it follows that

$$\tau_0 \mathcal{S}(X_1) = \frac{1}{8\pi i \tau_0^2} \frac{\tau_0^3 \mathcal{S}(\varphi_3)}{\mathcal{S}(\tau)}. \quad (\text{D.4})$$

The numerator of this expression can be computed from (4.22), yielding

$$\begin{aligned} \tau_0^3 \mathcal{S}(\varphi_3) &= \Sigma^{(3)} - \frac{X}{2\pi i \tau_0} \Sigma^{(4)} + \dots = \Sigma^{(3)} \Big|_0 + \epsilon_1 \epsilon_2 \left(\Sigma^{(3)} \Big|_1 - \frac{X_1}{2\pi i \tau_0} \Sigma^{(4)} \Big|_0 \right) + \dots \\ &= (2\pi i \tau_0)^3 \frac{\varphi_3}{(2\pi i \tau)^3} + \epsilon_1 \epsilon_2 (2\pi i \tau_0)^3 \left(\frac{1}{2} \frac{\varphi_5}{(2\pi i \tau)^4} + \frac{13}{4} \frac{\varphi_3 \varphi_4}{(2\pi i \tau)^5} + \frac{13}{4} \frac{\varphi_3^3}{(2\pi i \tau)^6} \right) + \dots \end{aligned} \quad (\text{D.5})$$

where the last step follows from the results given in appendix C. For the denominator we take into account that

$$\mathcal{S}(\tau) = \mathcal{S}(\tau_0) - \frac{\mathcal{S}(\varphi_2)}{2\pi i} = -\frac{1}{\tau_0} - \frac{\mathcal{S}(\varphi_2)}{2\pi i}. \quad (\text{D.6})$$

Resorting again to (4.22) to evaluate $\tau_0^2 \mathcal{S}(\varphi_2)$, we obtain

$$\begin{aligned} \mathcal{S}(\tau) &= -\frac{1}{\tau_0} - \frac{1}{2\pi i \tau_0^2} \left[\Sigma^{(2)} \Big|_0 + \epsilon_1 \epsilon_2 \left(\Sigma^{(2)} \Big|_1 - \frac{X_1}{2\pi i \tau_0} \Sigma^{(3)} \Big|_0 \right) + \dots \right] \\ &= -\frac{1}{\tau} - \epsilon_1 \epsilon_2 \left(\frac{1}{2} \frac{\varphi_4}{(2\pi i)^2 \tau^3} + \frac{3}{4} \frac{\varphi_3^2}{(2\pi i)^3 \tau^4} \right) + \dots \end{aligned} \quad (\text{D.7})$$

Inserting (D.5) and (D.7) into (D.4) and extracting the term of order $\epsilon_1 \epsilon_2$, we find

$$\tau_0 \mathcal{S}(X_1) \Big|_1 = -(2\pi i \tau_0) \left(\frac{1}{8} \frac{\varphi_5}{(2\pi i \tau)^3} + \frac{11}{16} \frac{\varphi_3 \varphi_4}{(2\pi i \tau)^4} + \frac{5}{8} \frac{\varphi_3^3}{(2\pi i \tau)^5} \right). \quad (\text{D.8})$$

Using this result and (D.3) into (D.1), we finally obtain the following constraint on X_2 :

$$\tau_0 \mathcal{S}(X_2) \Big|_0 - 2\pi i \tau_0 \left(\frac{X_2}{2\pi i \tau} \right) = (2\pi i \tau_0) \left(\frac{7}{48} \frac{\varphi_3 \varphi_4}{(2\pi i \tau)^4} + \frac{7}{32} \frac{\varphi_3^3}{(2\pi i \tau)^5} \right). \quad (\text{D.9})$$

Considering the structures involved in this relation, we can try an *Ansatz* such that X_2 is written as a total derivative:

$$\begin{aligned} X_2 &= \partial_a \left(-\lambda_1 \frac{\varphi_4}{(2\pi i\tau)^2} - \lambda_2 \frac{\varphi_3^2}{(2\pi i\tau)^3} \right) \\ &= \lambda_1 \frac{\varphi_5}{(2\pi i\tau)^2} + 2(\lambda_1 + \lambda_2) \frac{\varphi_3\varphi_4}{(2\pi i\tau)^3} + 3\lambda_2 \frac{\varphi_3^3}{(2\pi i\tau)^4}. \end{aligned} \tag{D.10}$$

With this position we find straightforwardly, using the formulæ of appendix C, that

$$\tau_0 \mathcal{S}(X_2) \Big|_0 - 2\pi i\tau_0 \left(\frac{X_2}{2\pi i\tau} \right) = 2\pi i\tau_0 \left((6\lambda_1 - 4\lambda_2) \frac{\varphi_3\varphi_4}{(2\pi i\tau)^4} + (9\lambda_1 - 6\lambda_2) \frac{\varphi_3^3}{(2\pi i\tau)^5} \right). \tag{D.11}$$

Comparing this result to the constraint (D.9), we obtain a single independent equation:

$$6\lambda_1 - 4\lambda_2 = \frac{7}{48}. \tag{D.12}$$

Our *Ansatz* thus satisfies the requirement (D.9) with still one free parameter. In section 5 we chose to fix this arbitrariness by requiring that the modified prepotential \widehat{F} introduced in (5.2) differs from F by a series of corrections in powers of $\epsilon_1\epsilon_2$ whose coefficients only involve φ_3 and the modified effective coupling $\widehat{\tau}$ defined in (5.5). It is easy to see that, starting with X_2 given in (D.10), the \widehat{F} term of order $(\epsilon_1\epsilon_2)^2$ in (5.7) would contain a part proportional to φ_4 unless λ_1 assumes a specific value which, by virtue of (D.12), fixes also λ_2 . These values are:

$$\lambda_1 = \frac{1}{16}, \quad \lambda_2 = \frac{11}{192}, \tag{D.13}$$

which are those used in (5.1).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] N. Seiberg and E. Witten, *Electric-magnetic duality, monopole condensation and confinement in $N = 2$ supersymmetric Yang-Mills theory*, *Nucl. Phys. B* **426** (1994) 19 [Erratum *ibid.* **B 430** (1994) 485] [[hep-th/9407087](#)] [[INSPIRE](#)].
- [2] N. Seiberg and E. Witten, *Monopoles, duality and chiral symmetry breaking in $N = 2$ supersymmetric QCD*, *Nucl. Phys. B* **431** (1994) 484 [[hep-th/9408099](#)] [[INSPIRE](#)].
- [3] E. D'Hoker and D. Phong, *Lectures on supersymmetric Yang-Mills theory and integrable systems*, [[hep-th/9912271](#)] [[INSPIRE](#)].
- [4] J. Minahan, D. Nemeschansky and N. Warner, *Instanton expansions for mass deformed $N = 4$ super Yang-Mills theories*, *Nucl. Phys. B* **528** (1998) 109 [[hep-th/9710146](#)] [[INSPIRE](#)].
- [5] M. Billó, M. Frau, L. Gallot and A. Lerda, *The exact 8d chiral ring from 4d recursion relations*, *JHEP* **11** (2011) 077 [[arXiv:1107.3691](#)] [[INSPIRE](#)].
- [6] N.A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, *Adv. Theor. Math. Phys.* **7** (2004) 831 [[hep-th/0206161](#)] [[INSPIRE](#)].

- [7] N. Nekrasov and A. Okounkov, *Seiberg-Witten theory and random partitions*, [hep-th/0306238](#) [INSPIRE].
- [8] A. Losev, N. Nekrasov and S.L. Shatashvili, *Testing Seiberg-Witten solution*, [hep-th/9801061](#) [INSPIRE].
- [9] G.W. Moore, N. Nekrasov and S. Shatashvili, *D particle bound states and generalized instantons*, *Commun. Math. Phys.* **209** (2000) 77 [[hep-th/9803265](#)] [INSPIRE].
- [10] R. Flume and R. Poghossian, *An algorithm for the microscopic evaluation of the coefficients of the Seiberg-Witten prepotential*, *Int. J. Mod. Phys. A* **18** (2003) 2541 [[hep-th/0208176](#)] [INSPIRE].
- [11] U. Bruzzo, F. Fucito, J.F. Morales and A. Tanzini, *Multiinstanton calculus and equivariant cohomology*, *JHEP* **05** (2003) 054 [[hep-th/0211108](#)] [INSPIRE].
- [12] R. Flume, F. Fucito, J.F. Morales and R. Poghossian, *Matone's relation in the presence of gravitational couplings*, *JHEP* **04** (2004) 008 [[hep-th/0403057](#)] [INSPIRE].
- [13] N. Nekrasov and S. Shadchin, *ABCD of instantons*, *Commun. Math. Phys.* **252** (2004) 359 [[hep-th/0404225](#)] [INSPIRE].
- [14] M. Mariño and N. Wyllard, *A note on instanton counting for $N = 2$ gauge theories with classical gauge groups*, *JHEP* **05** (2004) 021 [[hep-th/0404125](#)] [INSPIRE].
- [15] M. Billó et al., *Exotic instanton counting and heterotic/type-I-prime duality*, *JHEP* **07** (2009) 092 [[arXiv:0905.4586](#)] [INSPIRE].
- [16] F. Fucito, J.F. Morales and R. Poghossian, *Exotic prepotentials from $D(-1)D7$ dynamics*, *JHEP* **10** (2009) 041 [[arXiv:0906.3802](#)] [INSPIRE].
- [17] M. Billó et al., *Stringy instanton corrections to $N = 2$ gauge couplings*, *JHEP* **05** (2010) 107 [[arXiv:1002.4322](#)] [INSPIRE].
- [18] H. Nakajima and K. Yoshioka, *Lectures on instanton counting*, [math.AG/0311058](#) [INSPIRE].
- [19] M. Billó, M. Frau, F. Fucito and A. Lerda, *Instanton calculus in RR background and the topological string*, *JHEP* **11** (2006) 012 [[hep-th/0606013](#)] [INSPIRE].
- [20] K. Ito, H. Nakajima, T. Saka and S. Sasaki, *$N = 2$ instanton effective action in Ω -background and $D3/D(-1)$ -brane system in RR background*, *JHEP* **11** (2010) 093 [[arXiv:1009.1212](#)] [INSPIRE].
- [21] I. Antoniadis, E. Gava, K. Narain and T. Taylor, *Topological amplitudes in string theory*, *Nucl. Phys. B* **413** (1994) 162 [[hep-th/9307158](#)] [INSPIRE].
- [22] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Holomorphic anomalies in topological field theories*, *Nucl. Phys. B* **405** (1993) 279 [[hep-th/9302103](#)] [INSPIRE].
- [23] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes*, *Commun. Math. Phys.* **165** (1994) 311 [[hep-th/9309140](#)] [INSPIRE].
- [24] A. Klemm, M. Mariño and S. Theisen, *Gravitational corrections in supersymmetric gauge theory and matrix models*, *JHEP* **03** (2003) 051 [[hep-th/0211216](#)] [INSPIRE].
- [25] M.-X. Huang and A. Klemm, *Holomorphicity and modularity in Seiberg-Witten theories with matter*, *JHEP* **07** (2010) 083 [[arXiv:0902.1325](#)] [INSPIRE].

- [26] I. Antoniadis, S. Hohenegger, K. Narain and T. Taylor, *Deformed topological partition function and Nekrasov backgrounds*, *Nucl. Phys. B* **838** (2010) 253 [[arXiv:1003.2832](#)] [[INSPIRE](#)].
- [27] D. Krefl and J. Walcher, *Extended holomorphic anomaly in gauge theory*, *Lett. Math. Phys.* **95** (2011) 67 [[arXiv:1007.0263](#)] [[INSPIRE](#)].
- [28] M.-X. Huang and A. Klemm, *Direct integration for general Ω backgrounds*, *Adv. Theor. Math. Phys.* **16** (2012), no. 3 805–849 [[arXiv:1009.1126](#)] [[INSPIRE](#)].
- [29] N.A. Nekrasov and S.L. Shatashvili, *Quantization of integrable systems and four dimensional gauge theories*, [arXiv:0908.4052](#) [[INSPIRE](#)].
- [30] R. Poghossian, *Deforming SW curve*, *JHEP* **04** (2011) 033 [[arXiv:1006.4822](#)] [[INSPIRE](#)].
- [31] F. Fucito, J. Morales, D.R. Pacifici and R. Poghossian, *Gauge theories on Ω -backgrounds from non commutative Seiberg-Witten curves*, *JHEP* **05** (2011) 098 [[arXiv:1103.4495](#)] [[INSPIRE](#)].
- [32] L.F. Alday, D. Gaiotto and Y. Tachikawa, *Liouville correlation functions from four-dimensional gauge theories*, *Lett. Math. Phys.* **91** (2010) 167 [[arXiv:0906.3219](#)] [[INSPIRE](#)].
- [33] D. Gaiotto, *$N = 2$ dualities*, *JHEP* **08** (2012) 034 [[arXiv:0904.2715](#)] [[INSPIRE](#)].
- [34] A. Mironov, A. Morozov and S. Shakirov, *Matrix model conjecture for exact BS periods and Nekrasov functions*, *JHEP* **02** (2010) 030 [[arXiv:0911.5721](#)] [[INSPIRE](#)].
- [35] M.-X. Huang, A.-K. Kashani-Poor and A. Klemm, *The Ω deformed B-model for rigid $N = 2$ theories*, *Annales Henri Poincaré* **14** (2013) 425 [[arXiv:1109.5728](#)] [[INSPIRE](#)].
- [36] M.-X. Huang, *On gauge theory and topological string in Nekrasov-Shatashvili limit*, *JHEP* **06** (2012) 152 [[arXiv:1205.3652](#)] [[INSPIRE](#)].
- [37] A.-K. Kashani-Poor and J. Troost, *The toroidal block and the genus expansion*, *JHEP* **03** (2013) 133 [[arXiv:1212.0722](#)] [[INSPIRE](#)].
- [38] D. Galakhov, A. Mironov and A. Morozov, *S-duality as a beta-deformed Fourier transform*, *JHEP* **08** (2012) 067 [[arXiv:1205.4998](#)] [[INSPIRE](#)].
- [39] M. Billó et al., *Non-perturbative gauge/gravity correspondence in $N = 2$ theories*, *JHEP* **08** (2012) 166 [[arXiv:1206.3914](#)] [[INSPIRE](#)].
- [40] T. Okuda and V. Pestun, *On the instantons and the hypermultiplet mass of $N = 2^*$ super Yang-Mills on S^4* , *JHEP* **03** (2012) 017 [[arXiv:1004.1222](#)] [[INSPIRE](#)].
- [41] F. Fucito, J.F. Morales and R. Poghossian, *Multi instanton calculus on ALE spaces*, *Nucl. Phys. B* **703** (2004) 518 [[hep-th/0406243](#)] [[INSPIRE](#)].
- [42] J. Minahan, D. Nemeschansky and N. Warner, *Partition functions for BPS states of the noncritical E_8 string*, *Adv. Theor. Math. Phys.* **1** (1998) 167 [[hep-th/9707149](#)] [[INSPIRE](#)].
- [43] T.W. Grimm, A. Klemm, M. Mariño and M. Weiss, *Direct integration of the topological string*, *JHEP* **08** (2007) 058 [[hep-th/0702187](#)] [[INSPIRE](#)].
- [44] M. Billó, L. Gallot, A. Lerda and I. Pesando, *F-theoretic versus microscopic description of a conformal $N = 2$ SYM theory*, *JHEP* **11** (2010) 041 [[arXiv:1008.5240](#)] [[INSPIRE](#)].
- [45] T. Dimofte and S. Gukov, *Chern-Simons theory and S-duality*, [arXiv:1106.4550](#) [[INSPIRE](#)].
- [46] E. Witten, *Quantum background independence in string theory*, [hep-th/9306122](#) [[INSPIRE](#)].
- [47] M. Billó, M. Frau, L. Gallot, A. Lerda and I. Pesando, work in progress.