



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in [Neuroscience, 253 (2013) 67–77, DOI: 10.1016/j.neuroscience.2013.08.036].

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+ *Digital Object Identifier link to the published journal article on Elsevier's ScienceDirect® platform*]

1
2 **ANDROGEN RECEPTOR DEFICIENCY ALTERS THE ARGININE-**
3 **VASOPRESSIN SEXUALLY DIMORPHIC SYSTEM IN TFM RATS.**
4
5
6

7 **Allieri F.^a, Spigolon G.^{a,d}, Melcangi R.C.^c, Collado P.^b, Guillamón A.^b, Gotti S.**
8 **^{a,d}, Panzica G.C.^{a,d}**
9

10
11
12 ^a Dept. Neuroscience, University of Torino, Torino, Italy.

13 ^b Dept. Psychobiology, Universidad Nacional de Educación a Distancia, C/ Juan del
14 Rosal, 10, 28040-Madrid, Spain.
15
16

17 ^c Dept. of Pharmacological and Biomedical Sciences, Section of Biomedicine and
18 Endocrinology, University of Milano, Milano, Italy.
19

20 ^d Neuroscience Institute Cavalieri Ottolenghi (NICO), Orbassano (Torino), Italy
21
22
23
24
25
26

27 * Author for correspondence
28
29
30

31 Dr. Stefano Gotti

32 Neuroscience Institute Cavalieri Ottolenghi (NICO)

33 Laboratory of Neuroendocrinology,
34

35 Regione Gonzole, 10
36

37 10043 – Orbassano (Torino, Italy)
38

39 Phone +39 011 6706610
40

41 Fax +39 011 2367054
42

43 e-mail: stefano.gotti@unito.it
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

LIST OF ABBREVIATIONS

1		
2		
3		
4	AIS	androgen insensitivity syndrome
5	AR	androgen receptor
6		
7	AVP	arginine vasopressin
8		
9	AVP-ir	arginine vasopressin immunoreactive
10		
11	BST	bed nucleus of the stria terminalis
12		
13	DHT	dihydrotestosterone
14		
15	E₂	estradiol
16		
17	ER	estrogen receptor
18		
19	LS	lateral septum
20		
21	MeA	medial amygdaloid nucleus
22		
23	MPOA	medial preoptic area
24		
25	PVN	paraventricular nucleus
26		
27	SON	supraoptic nucleus
28		
29	T	testosterone
30		
31	Tfm	testicular feminilization mutation
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		
61		
62		
63		
64		
65		

1
2 **ABSTRACT**
3

4 In rodents as well as in many other mammalian and non-mammalian species, the
5 arginine-vasopressin (AVP) system includes a parvocellular sexually dimorphic
6 portion located within the bed nucleus of the stria terminalis (BST), the medial
7 amygdaloid nucleus (MeA) and the lateral septum. In this system, males have more
8 cells and denser projections than females, neurons show androgen and estrogen
9 receptors, and gonadal hormones are required for the activation. However, the role of
10 these hormones for the differentiation of the system is not clear. Previous studies
11 performed on aromatase knock out mice suggested that estradiol is not necessary for
12 the differentiation of the system, but it is important for its activation in adulthood.
13
14

15 To elucidate the role of androgens on differentiation and functioning of AVP
16 parvocellular system, we compared male and female rats with a non-functional
17 mutation of androgen receptor (*Tfm*, testicular feminization mutation) to their control
18 littermates. Our data show that the lack of a functional androgen receptor significantly
19 decreases the expression of AVP immunoreactivity within the BST and MeA of male
20 *Tfm*. Thus supporting the hypothesis that androgens, through the action of their
21 receptor, should have a relevant role in the organization and modulation of the AVP
22 parvocellular sexually dimorphic system.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

1
2
3
4 Gonadal steroids, through their interaction with intracellular receptors, induce several
5 effects in target cells; these receptors are members of ligand-activated transcription
6 factors family that bind responsive elements located on the genome and therefore are
7 able to modulate the transcription of a wide variety of genes (McCarthy, 2008).
8 Specific nuclear receptors for estrogens and androgens are present in the developing
9 brain of rodents (DonCarlos and Handa, 1994, DonCarlos, 1996, McAbee and
10 DonCarlos, 1998). Therefore, gonadal steroids are likely to play a crucial role in brain
11 plasticity and development; in particular, they may induce the differentiation of
12 sexually dimorphic structures and behaviors (Panzica et al., 1995, McCarthy and
13 Konkle, 2005, McCarthy and Arnold, 2011, Lenz et al., 2012).

14
15
16 Sex difference in testosterone (T) levels during development has been recognized as
17 the driving force of sexual differentiation of the mammalian central nervous system.
18 In male rats, T secretion reaches the peak from prenatal days 18-20 (Perakis and
19 Stylianopoulou, 1986). This early elevation in T levels coincides with the period
20 (*critical perinatal period*) when T can permanently influence the development of
21 behavior and neural structures (Cooke et al., 1998). Within the brain, T is metabolized
22 into estradiol (E₂) by aromatase or into 5 α -dihydrotestosterone (5 α -DHT) by 5 α -
23 reductase. These two steroids may bind their receptors inducing the differentiation of
24 neural circuits. In some cases both androgen (AR) and estrogen receptors (ER) may
25 contribute to structural plasticity of nervous circuits, as for the posterior part of the
26 amygdala (MeP) (Cooke et al., 2003).

27
28
29 The AR mediates age- and tissue-specific actions of androgens and is responsible for
30 the development, maintenance and regulation of the male reproductive system
31 (Raskin et al., 2009, Juntti et al., 2010). AR is encoded by a X-linked gene, which
32 consists of 8 exons, and it is essential for androgenic action, of both T and 5 α -DHT
33 (Gobinet et al., 2002).

34
35
36 In humans, mutations of the AR gene represent the molecular basis of androgen
37 insensitivity syndrome (AIS), a rare X-linked disorder characterized by a defective
38 virilization in genetic males ranging from partial forms, in which 46, XY phenotypic
39 males are infertile, to the complete syndrome showing a female phenotype
40 (Brinkmann, 2001, McPhaul, 2002, Hughes et al., 2012)

1 In mammals, as well as in most vertebrates, part of the arginine-vasopressin (AVP)
2 neuronal system is sexually dimorphic and gonadal steroids sensitive in adulthood
3 (for a review see De Vries and Panzica, 2006). The bed nucleus of the stria terminalis
4 (BST) and the amygdala contain more AVP-immunoreactive (AVP-ir) cells in male
5 than in female and projections from these nuclei to the lateral septum (LS) are denser
6 in male than in female; in addition, several other brain regions show sexually
7 dimorphic AVP innervation (De Vries et al., 1984b, Rood et al., 2013).

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Histological investigations demonstrated several sexually dimorphic features (volume, cell number and cell size) in BST and amygdala. The BST shows a consistent pattern of sex differences in several mammalian species [guinea pig (Hines et al., 1992); rat (Del Abril et al., 1987); rabbit (Segovia et al., 2006)] and even in humans (Zhou et al., 1995). Both nuclei have a crucial role in the regulation of rodent male and female sexual and aggressive behaviors (Veening and Coolen, 1998, Veening et al., 2005). In particular, the rodent parvocellular AVP system is involved in the control of social behaviors, such as pair bonding, social recognition, aggressivity, depression and anxiety (see Compaan et al., 1993, Veenema and Neumann, 2008, Choleris et al., 2009, Rotzinger et al., 2010).

Several studies have investigated the development of sexually dimorphic AVP system in mammals and other vertebrates and the role played by gonadal hormones for its regulation (see for a review De Vries and Panzica, 2006).

Gonadectomy and early T replacement therapy affect the organization of limbic AVP system and AR expression, suggesting that both T and AR are involved in the differentiation of this system (Bingham and Viau, 2008). Also aromatase (the enzyme converting T into E₂) expression is reduced by gonadectomy and restored by T replacement (Roselli et al., 1987). However, whereas the lack of a functional aromatase gene (ArKO) induces an apparent reduction of AVP cell number and fiber innervation in BST, amygdala and LS (Plumari et al., 2002), adult administration of E₂ in the same mice induced a full recovery of male AVP system (Pierman et al., 2008). Therefore, it seems that E₂ is not necessary for the differentiation of the sexually dimorphic AVP system, while it plays a major role in adulthood to stimulate the same system.

The *Tfm* rats (characterized by a testicular feminilization due to a spontaneous mutation of the AR gene) may help in elucidate the role of androgens and ARs in the activation/organization of the sexually dimorphic AVP system. These animals show a

1 low level of the aromatase protein itself, even in T-treated gonadectomized *Tfm* male
2 rats (Roselli et al., 1987). In addition, *Tfm* rodents show alterations in some behaviors
3 [i.e. adult playful attack and defense (Field et al., 2006), anabolic androgenic steroid-
4 induced aggressive behavior (Wu et al., 2009, Robinson et al., 2012), anxiety-related
5 behaviors (Zuloaga et al., 2008, Zuloaga et al., 2011) or social investigation (Tejada
6 and Rissman, 2012)] that are partially dependent by the AVP system (Veenema and
7 Neumann, 2008, Veenema et al., 2010). Also, some neuronal circuits are altered in
8 these animals, as the nitric oxide producing system of BST and other hypothalamic
9 nuclei (Martini et al., 2008).

10 As *Tfm* rodents have decreased AR binding but apparently normal ER binding
11 (Attardi et al., 1976), these animals offer a way to examine the contribution of the AR
12 to the sexual differentiation of rodent brain in gonadally intact individuals.
13
14
15

16 **MATERIALS AND METHODS**

17 Animals

18 Female *Tfm* rats from the colony of the Department of Psychobiology of UNED,
19 Madrid, Spain, (for the origin of the strain see Garcia-Falgueras et al., 2005) were
20 analyzed and identified by polymerase chain reaction (PCR, Fernandez et al., 2003) as
21 females heterozygous for the *Tfm* mutation (X^{Tfm}/X). These rats were mated with
22 Wistar male rats (Charles River; Criffa, Barcelona) to obtain the animals used in this
23 experiment (N = 20). The animals obtained in the matings have the same genetic
24 background (X^{Tfm}/X x Wistar male, X/Y) and therefore are comparable to each other.
25 The experimental groups, classified after PCR analyses, were as follows: (a) *Tfm*
26 males (X^{Tfm}/Y , N = 5); (b) *Tfm* carrier females (X^{Tfm}/X , N = 5); (c) control littermate
27 males (X/Y, N = 5), and (d) control littermate females (X/X, N = 5).

28 At the age of 21 days, the animals were housed in monosexual groups in standard
29 cages and maintained in 12:12 light/dark cycle (lights off at 8:00 PM). Food and
30 water were provided *ad libitum*. The animal room was kept at 22 ± 2 °C. Animal care
31 and handling throughout the experimental procedures were in accordance with the
32 European Union Council Directive of 24 November 1986 (86/609/EEC) and the
33 experiment was approved by the ethical committee of the UNED. The genotyping of
34 the animals was performed at the Department of Pharmacological and Biomedical
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Sciences (Milano, Italy) by using 200 μ l of peripheral blood according to the protocol
2 previously developed (Fernandez et al., 2003, Garcia-Falgueras et al., 2005).
3
4

5 Colchicine injection and sacrifice 6

7 At the age of 60 days, all the animals were subjected to an intracerebroventricular
8 injection of colchicine to increase the AVP immunostaining (Compaan et al., 1993,
9 Plumari et al., 2002, Pierman et al., 2008). Females were inspected for the stage of the
10 cycle (examination of vaginal smears) and injected only in estrus. Briefly, the animals
11 were deeply anaesthetized with a mixture of ketamine (80mg/Kg of body weight),
12 valium (10mg/Kg of body weight), and atropine (0,05mg/Kg of body weight) and
13 positioned in a stereotaxic apparatus (Kopf, Tujunga, CA, USA). They were then
14 stereotaxically injected into the left lateral ventricle with 10 μ l of solution
15 corresponding to 250 μ g/Kg of body weight of colchicine dissolved in distilled
16 water. Accuracy of the position of the injection cannula was controlled by observing
17 the increase in pressure in the cannula when it entered the ventricle. Later inspection
18 of brain sections revealed no trace of damaged tissue around the lateral ventricle
19 except for the penetration path of the needle. 48 hours after the colchicine injection,
20 rats were deeply anesthetized with the same anesthetic mixture (ketamine-valium-
21 atropine) and perfused through the heart with saline solution (0.9%) until vessels were
22 completely blood-free, and then with fixative (4% paraformaldehyde in 0,1 M
23 phosphate buffer, pH 7,3). Brains were dissected out of the skull, post-fixed for 2
24 hours in the same fixative and rinsed in 0.01 M phosphate buffer saline (PBS). They
25 were then placed overnight in a 30% sucrose solution in PBS, frozen in precooled
26 liquid isopentane at -35°C, and stored in a deep freezer at -80°C until sectioning.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 Immunohistochemistry 46

47 Brains were serially cut in the coronal plane at 25 μ m thickness using a cryostat and
48 collected in a cryoprotectant solution (Watson et al., 1986) at -20°C. Every forth
49 section one was stained for AVP by immunohistochemistry using the free floating
50 technique. Sections were immunostained using a commercial AVP antibody (ICN
51 Pharmaceuticals, CA; USA) diluted 1:20,000. The same antibody has been previously
52 used to stain the AVP cells and fibers in rodents species (Ferris et al., 1995, Villalba
53 et al., 1999, Plumari et al., 2002, Pierman et al., 2008). [The specificity of this](#)
54
55
56
57
58
59
60
61
62
63
64
65

1 antibody to vasopressin (and the absence of any cross-reactivity with oxytocin) has
2 been validated previously (Ferris et al., 1995). In the present study, we have
3 performed the following controls: a) the primary antibody was omitted or replaced
4 with an equivalent concentration of normal serum (negative controls); b) the
5 secondary antibody was omitted. In these conditions, cells and fibers were totally
6 unstained.
7
8
9

10 Briefly, the free-floating sections collected in the cryoprotectant solution were washed
11 overnight in PBS at pH 7.3. The following day, sections were first washed in PBS
12 containing 0.2 % Triton X-100 for 30 min and then treated to inhibit endogenous
13 peroxidase activity with a solution of PBS containing methanol/hydrogen peroxide for
14 20 min (Streefkerk, 1972). Sections were incubated for 30 min with normal goat
15 serum (Vector Laboratories, Burlingame, CA, USA) and incubated overnight at room
16 temperature with anti-AVP antibody diluted 1:20,000 in PBS, pH 7.3-7.4, containing
17 0.2% Triton X-100. A biotinilated goat anti-rabbit secondary antibody (Vector
18 Laboratories, Burlingame, CA, USA) was then employed at a dilution of 1:200 for 60
19 min at room temperature. The antigen-antibody reaction was revealed by 60 min
20 incubation with avidin-peroxidase complex (Vectastain ABC Kit Elite, Vector
21 Laboratories, Burlingame, CA, USA). The peroxidase activity was visualized with a
22 solution containing 0.400 mg/ml 3,3-diamino-benzidine (Sigma, Milan, Italy) and
23 0.004% hydrogen peroxide in 0.05M Tris-HCl buffer pH 7.6. Sections were mounted
24 on chromallum coated slides, air dried, cleared in xylene and cover slipped with
25 Entellan (Merck, Milan, Italy). Sections were collected on chromalum-coated slides,
26 air-dried, washed in xylene, and cover slipped with Entellan (Merck, Milano, Italy).
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 An alternate set of sections was stained with toluidine blue to set up a series of Nissl-
43 stained section that were used for anatomical orientation. Identification of all
44 structures was based on the rat brain atlas of Paxinos and Watson (1998).
45
46
47
48

49 Quantitative analysis

50 For quantitative analysis, all slides were first coded so that the observer was blind to
51 the experimental treatments. Sections were chosen to match similar levels of the
52 regions to be investigated. Selected fields were acquired through a Zeiss Axioplan I
53 microscope equipped with a Leica DFC 320 digital telecamera connected to an Apple
54 G4 Macintosh. Digital images were later analyzed with Image J program (version
55
56
57
58
59
60
61
62
63
64
65

1
2 1.46, a public domain program written by W. Rasband at the U.S. National Institutes
3 of Health, Bethesda, MD, USA).

4
5 The cells count and the fractional area, covered by AVP-ir structures, were evaluated
6 according to the following protocols in the regions schematically reported in Fig. 1:

7
8 - in the BST and amygdala we counted the number of AVP-ir cells (identified for the
9 presence of a clearly labeled cell body) using the cell counter plug-in of ImageJ. We
10 counted all the positive neurons within a rectangle of fixed size ($500,000 \mu\text{m}^2$) placed
11 over the nuclei (Fig. 1). The measures were performed in two separate rostro-caudal
12 levels of the bed nucleus of the stria terminalis (BST: -0.40 and -0.80 relative to
13 Bregma, (Paxinos and Watson, 1998), and medial amygdaloid nucleus (MeA: -2.12
14 and -2.56 relative to Bregma, (Paxinos and Watson, 1998) to control for a possible
15 heterogeneity of AVP distribution in these structures.
16
17

18
19 - In the LS, as well as in the BST and amygdala, we measured the density of AVP-ir
20 structures by calculating, in binary transformations of the images (threshold function
21 of Image J), the fractional area (percentages of pixels) covered by positive structures
22 in predetermined regions with a fixed extension ($500,000 \mu\text{m}^2$) as described in our
23 previous studies in mice and quails (Viglietti-Panzica et al., 2001, Plumari et al.,
24 2002, Viglietti-Panzica et al., 2007, Pierman et al., 2008). In particular we considered
25 for BST, and MeA the same levels described above and for LS two additional rostro-
26 caudal levels (coordinates 0.48 and -0.26 relative to Bregma in the rat brain atlas, see
27 Fig. 1).
28
29

30 Statistical analysis

31
32 Statistical analyses were performed using the software Statview 5.0 (Abacus
33 Concepts, Berkely, CA, USA).

34
35 The average from 2 coronal sections was used for data analysis, since a preliminary
36 analyses revealed no significant difference between the anterior-posterior levels for
37 each considered nucleus (BST, MeA, and LS). We have therefore performed a two-
38 way analysis of variance (ANOVA, genotype and sex as independent factors) on the
39 average cells count (BST and MeA) or on the average fractional area value (BST,
40 MeA, and LS). When appropriate, we applied a post-hoc comparison test (Fisher
41 PLSD). Differences were considered statistically significant for $p \leq 0.05$.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

RESULTS

The qualitative observation of the immunostained sections obtained from colchicine-injected WT male rats (XY) showed a distribution of AVP-ir structures in line with what has been previously described in rodents (De Vries et al., 1985, Plumari et al., 2002, Grassi et al., 2010, Rood and De Vries, 2011). In particular, we observed a dense population of AVP-ir parvocellular neurons and fibers in the BST (Fig.2), and the MeA (Fig.3), whereas the presence of positive fibers within the LS (Fig.4) was rather weak. The comparison with WT females evidenced an obvious sex difference (females have less immunoreactivity than males) for BST, MeA, and LS.

When comparing AVP immunostained sections of XY and *Tfm* (X^{tfm}/Y) male rats we observed a sharp decrease of AVP-ir elements in the BST and in the MeA of *Tfm* males (Figs.2-3). A slight decrease of the vasopressinergic innervation was detectable in the LS (Fig.4). These qualitative observations were confirmed by quantitative analysis of AVP immunoreactivity, whose results are summarized in Fig.5.

Regarding the average fractional area value, for BST, and MeA the two-way ANOVA reported significant effects for all the considered parameters (sex, genotype, and interaction sex-genotype). For the BST: sex ($F_{1,1}=32.76$, $p<0.0001$), genotype ($F_{1,1}=15.03$, $p=0.001$), interaction between sex and genotype ($F_{1,1,1}=10.32$, $p=0.005$). For the MeA: sex ($F_{1,1}=17.88$, $p=0.0006$), genotype ($F_{1,1}=9.61$, $p=0.007$), interaction between sex and genotype ($F_{1,1,1}=8.70$, $p=0.009$). The post-hoc Fisher PLSD test reported highly significant differences ($p<0.0001$ for BST, and $p<0.001$ for MeA) between male XY and male $X^{tfm}Y$, male XY and female XX, and male XY and female $X^{tfm}X$.

For the LS, the two-way ANOVA reported a significant effect only for sex ($F_{1,1}=10.32$, $p=0.003$). The post-hoc Fisher PLSD test reported significant differences ($p=0.01$) for comparisons of male XY against female XX or female $X^{tfm}X$.

For the average cell counts value in the BST and MeA the two-way ANOVA reported significant effects for the considered parameters (sex, genotype, and interaction sex-genotype). For the BST: sex ($F_{1,1}=161.63$, $p<0.0001$), genotype ($F_{1,1}=19.34$, $p=0.0004$), interaction between sex and genotype ($F_{1,1,1}=17.00$, $p=0.0008$). For the MeA: sex ($F_{1,1}=6.40$, $p=0.0223$), genotype ($F_{1,1}=4.83$, $p=0.0430$), interaction between

1 sex and genotype ($F_{1,1,1}=4.547$, $p=0.0488$). The post-hoc Fisher PLSD test reported
2 highly significant differences ($p<0.0001$ for BST, and $p<0.01$ for MeA) between male
3 XY and male $X^{tfm}Y$, male XY and female XX, and male XY and female $X^{tfm}X$.
4
5
6

7 **DISCUSSION**

8
9

10 Present results confirm the existence of significant sexual differences in the
11 parvocellular part of the AVP system of rat brain (De Vries and Panzica, 2006). In
12 particular, we found the presence of highly significant differences in the density of
13 AVP-ir elements (higher in males than in females) in BST, MeA, and LS nuclei. In
14 addition, we show that the absence of a fully functional AR, causing the testicular
15 feminization syndrome, induces significant decrease of the expression of AVP in
16 these nuclei. This decrease is so marked that in *Tfm* rats the sex dimorphism of AVP
17 is no longer observed. In fact, male $X^{tfm}Y$ shows a significant decrease of AVP-ir in
18 the BST, and the MeA nuclei in comparison to male XY, whereas the observed
19 decrease of the immunoreactivity in the LS is not significant. The lack of significant
20 effect of AR mutation on the AVP innervation of LS is intriguing, because these
21 fibers derive from the BST/medial amygdala system (De Vries and Buijs, 1983, Caffé
22 et al., 1987). However, the presence of a decrease not reaching the significance could
23 be due to the large variation of experimental measurements (see the extent of standard
24 error in the bar graphs), as well as to the use of colchicine injection to increase the
25 immunoreactivity in cell bodies of BST and MeA. This treatment is, in fact,
26 interfering with the axonal flux (Brat and Brimijoin, 1992) and we observed an
27 overall decrease of the AVP immunoreactivity in extra-hypothalamic fibers' systems
28 of our animals.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 The masculinization of the rodent hypothalamus (i.e. volume of the structure, number
46 of neurons, expression of neurotransmitters, and neuropeptides) is widely assumed to
47 be due to estrogenic metabolites of testosterone interacting with ERs during perinatal
48 critical period (aromatization hypothesis), while the α -fetoprotein (AFP) protects the
49 developing female brain from masculinization and defeminization by estrogens (for a
50 review see McCarthy et al., 2009). However, this is not the only mechanisms present
51 in the vertebrate brain, for example, in the avian hypothalamus estradiol has exactly
52 the opposite effect, inducing the demasculinization of those avian behaviors and
53
54
55
56
57
58
59
60
61
62
63
64
65

1 circuits that are sexually differentiated (for a review see Balthazart and Adkins-
2 Regan, 2002), while, in non human primates masculinization of brain and behavior is
3 an androgen dependent process (for a review see Wallen, 2005). Finally, recent
4 studies hypothesized an important role of epigenetic changes to induce brain sexual
5 differentiation (for a review see Lenz et al., 2012).
6
7

8
9 In adult rats, AVP projections from the BST and MeA are dependent by the levels of
10 circulating gonadal hormones: gonadectomy eliminates AVP expression and
11 replacement of hormones reinstates it (De Vries et al., 1984a). Many AVP-ir neurons
12 of BST and medial amygdala express estrogen and androgen receptors (Axelson and
13 Van Leeuwen, 1990, Zhou et al., 1994). However, the activational effect of gonadal
14 hormones cannot explain all differences in AVP expression because males and
15 females exposed to similar steroid levels still differ (De Vries and Al Shamma, 1990,
16 Wang et al., 1993), so the residual differences should be due to organizational effects
17 of hormones. Similar effects have been described for almost all vertebrates (Goodson
18 and Bass, 2001, Panzica et al., 2001, De Vries and Panzica, 2006).
19
20
21
22
23
24
25
26

27 In rodents, the mechanism by which gonadal hormones may influence the
28 differentiation of parvocellular AVP system has been studied in a variety of models.
29 Neonatal treatment of male and female rats gonadectomized at postnatal day 0 (PND-
30 0) with either T, E₂ or E₂-DHT increased the number of vasopressinergic cells in the
31 BST over that of DHT or oil treatment. However, treatment with DHT also
32 significantly increased the number of vasopressinergic cells over that of oil treatment.
33 Hence, these data confirm the role of E₂ in causing sexual differentiation of the brain;
34 in addition they provide a demonstration of a masculinizing effect of a
35 nonaromatizable androgen on a sexually dimorphic neuropeptide system (Han and De
36 Vries, 2003).
37
38
39
40
41
42
43
44

45 Scordalakes and Rissman (2004) performed a study of AVP expression in *Tfm*, ArKO
46 and *ArKO-Tfm* (DKO) mice. They looked the LS, the BNST and the MePD and
47 found a clear sex difference only in the LS. Moreover, they found also that the *Tfm*
48 background had no effect on AVP expression (WT and *Tfm* males had the greatest
49 amount of AVP-ir in the LS, and displayed the most aggression in resident–intruder
50 tests). These data seem in contradiction with our present results, but the study was
51 performed in mice and without colchicine treatment, so there are both methodological
52 and species differences.
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In the same experiment Scordalakes and Rissman (2004) found in the DKO mice less AVP immunoreactivity as compared to the mutants lacking either ERalpha or AR, suggesting again a contribution of both ERalpha and AR for the differentiation of this system. In partial contrast with these studies, in our previous experiments using aromatase knock out mice (ArKO) we demonstrated that early exposure to estrogens is not necessary for the full differentiation of the parvocellular AVP system of BST and MeA. In fact, ArKO adult males do not show AVP immunoreactivity in these regions (Plumari et al., 2002), but, if they are exposed in adulthood to E₂, AVP immunoreactivity is restored to levels observed in intact wild type males (Pierman et al., 2008). This suggests that the decrease in AVP expression in gonadally intact male ArKO mice was largely due to the absence of activational rather than to organizational effects of E₂ in these mice. These results are in line with previous studies on alpha-fetoprotein knockout mice (AFP-KO). Indeed, AFP-KO females, not protected by alpha-fetoprotein, are exposed to a large amount of maternal estrogens during development, but they did not show a male-typical AVP system suggesting that, [in these conditions](#), estrogens have probably no masculinizing effect on this sexually dimorphic neuropeptide system (Bakker et al., 2006). Finally, XY, XY^{-Sry}, XX^{Sry}, that are all male mice (in which Y⁻ symbolize the deletion of the gene Sry and Sry stand for the insertion of a Sry transgene) are more masculine than XX and XY⁻ female mice with regard to the density of AVP-expressing fibers in the lateral septum, suggesting a direct contribution of sex chromosome for the development of AVP sexual dimorphism (De Vries et al., 2002).

Recent studies, performed in mice, demonstrate that ARs have a role in masculinization of some behaviors (Raskin et al., 2009, Juntti et al., 2010). Our data on *Tfm* rats suggest that androgens, mediated by the AR, may have an important role, also in the differentiation of some circuits of the rodent brain, in particular in the differentiation of AVP-ir elements in the parvocellular sexually dimorphic system located in rat BST and MeA. This effects is sex-specific, in fact, the *Tfm* mutation apparently had no effect in heterozygous carrier female, that express a wild type copy and a mutated copy of the AR gene and, thus, can compensate the effect with the wild type AR. Based on similar studies on *Tfm* mutants, an important role for AR to induce sex differences has been proposed, in rodents, also for other sexually dimorphic neural circuits, such as spine formation in the prefrontal cortex (Hajszan et al., 2007), the size of neuronal somata in the rat MeP (Morris et al., 2005) and VMHvl (Dugger

1 et al., 2007), the calbindin immunoreactivity in the mouse medial preoptic area
2 (Edelmann et al., 2007), the expression of neuronal nitric oxide synthase (nNOS) in
3 rat BST and medial preoptic area (Martini et al., 2008), or the astrocyte number in the
4 rat medial amygdala (Johnson et al., 2008).
5

6
7 The expression of mRNA for AR in the rat BST and other forebrain regions has been
8 investigated by in situ hybridization, demonstrating its presence from embryonic day
9 20 (ED-20). In addition, AR mRNA levels increase throughout the perinatal period.
10 The developmental profile of AR mRNA in the forebrain is consistent with a
11 hypothetical role for androgens acting via the AR in the organization of male typical
12 neural circuits (McAbee and DonCarlos, 1998).
13

14 Although males and females both show increases in relative AR mRNA expression
15 between ED-20 and PND-10, the progression of AR mRNA levels is sex and region
16 specific. On ED-20, PND-0, and PND-4, the relative expression of AR mRNA is not
17 different in males and females in any of forebrain areas analyzed. Between PND-4
18 and PND-10 (i.e. during the second half of the critical period for sexual
19 differentiation), sex differences in AR mRNA expression arise in the BST and the
20 MPOA, but not in the other analyzed regions (lateral septum, ventromedial
21 hypothalamus, and arcuate nucleus). No data were provided for MeA in this study
22 (McAbee and DonCarlos, 1998). These region-specific patterns in the profile of AR
23 mRNA expression suggest a functional difference in the sensitivity of individual areas
24 to the potential developmental actions of androgens. Experimental studies involving
25 gonadectomy at PND-0 demonstrated that the onset of this sexual difference in the
26 AR mRNA expression at PND-10 is due to circulating levels of T (McAbee and
27 DonCarlos, 1999b), but males gonadectomized on PND-0 and treated daily with
28 DHT, a nonaromatizable androgen, had low levels of AR mRNA that were not
29 significantly different from AR mRNA levels in intact females. In contrast, males
30 gonadectomized on PND-0 and treated daily with diethylstilbestrol (DES), a synthetic
31 estrogen, maintained male-typical levels of AR mRNA in the BST and the MPOA.
32 Treatment with the androgen receptor antagonist, flutamide had no effect on AR
33 mRNA expression. The conclusion is that AR mRNA is up-regulated by estrogen but
34 is not regulated by androgen during the early postnatal period (McAbee and
35 DonCarlos, 1999a).
36

37 In summary, available data demonstrate that AR and ER are largely present in the
38 AVP system of BST/MeA, that the expression of these receptors in BST is sexually
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 differentiated from early stage of postnatal development, and that the development of
2 AR mRNA differences is due to the presence of estrogens. Therefore we can
3 hypothesize that both ER and AR are necessary to develop the sexual dimorphism of
4 AVP parvocellular system of BST and MeA, and that their action is sequential during
5 the perinatal critical period. The lack of effect of aromatase deletion (Pierman et al.,
6 2008) could be due to a compensatory effect of the DHT metabolite 3beta-diol that
7 can stimulate the AVP promoter through ERbeta (Pak et al., 2007, 2009).

8
9
10
11
12 The present analysis cannot determine whether the morphological effects of a
13 defective AR are due to the protein dysfunction during the perinatal, pubertal, or adult
14 period because *Tfm* rats have defective AR throughout development. Whether AR
15 stimulation acts on the developing and/or the adult parvocellular AVP system must
16 await further study. The mechanism could also involve epigenetic changes. In fact, a
17 recent study demonstrated that a transient reduction in the amygdala of the expression
18 of methyl-CpG-binding protein 2 (MeCP2, a protein binding to methylated DNA)
19 decreased AR mRNA in two-week old males (but not in the adult), whereas it
20 permanently eliminated the sex difference in the AVP system (Forbes-Lorman et al.,
21 2012). Thus, the reduction in AR expression during early postnatal life may play a
22 role in the decrease in AVP expression in MeCP2-siRNA treated males.

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
In conclusion, present data indicate that androgens, through the action of AR and in
cooperation with estrogens, may have a relevant role in the organization and
modulation of gender-specific components of the neural circuits that mediates male
social behaviors, in particular the AVP producing population of BST and MeA.

Acknowledgements: This work was supported by Fondazione Cavalieri Ottolenghi
and by grants of the University of Torino (GCP), University of Milan (RCM),
Fondazione San Paolo (Progetto Neuroscienze, PF-2009.1180, GCP & RCM),
SEJ2007-65686 (AG) and SEJ2007-62942 (PC)

Legends to the figures

1
2
3
4 Fig.1. Schematic diagrams showing the levels and the regions (grey squares) where
5 the counts were performed. Letters are referred to the different level of Bregma: a)
6 0.48 for lateral septum; b) -0.26 for lateral septum; c) -0.40 for bed nucleus of the
7 stria terminalis; d) -0.80 for bed nucleus of the stria terminalis; e) -2.12 for medial
8 amygdaloid nucleus; f) -2.56 for medial amygdaloid nucleus. Drawings from the atlas
9 of Paxinos and Watson (1998).
10
11
12
13
14
15

16 Fig.2. Representative photomicrographs showing AVP-immunoreactive staining in
17 the bed nucleus of the stria terminalis (BST) of male (XY) and female (XX) wild type
18 rats as well as in *Tfm* rats of both sexes ($X^{tfm}Y$ and $X^{tfm}X$). f – fornix; opt – optic tract.
19 The scale bar represents 150 μ m.
20
21
22
23
24

25 Fig.3. Representative photomicrographs showing AVP-immunoreactive staining in
26 medial amygdaloid nucleus (MeA) of male (XY) and female (XX) wild type rats as
27 well as in *Tfm* rats of both sexes ($X^{tfm}Y$ and $X^{tfm}X$). f – fornix; opt – optic tract. The
28 scale bar represents 150 μ m.
29
30
31
32
33

34 Fig.4. Representative photomicrographs showing AVP-immunoreactive staining in
35 the lateral septum (LS) of male (XY) and female (XX) wild type rats as well as in *Tfm*
36 rats of both sexes ($X^{tfm}Y$ and $X^{tfm}X$). * : third ventricle. The scale bar represents 100
37 μ m.
38
39
40
41
42

43 Fig.5. Bar graphs illustrating the fractional area covered by AVP-ir elements
44 (means \pm S.E.M.; top panel) and the cells mean (means \pm S.E.M.; bottom panel) in
45 the bed nucleus of the stria terminalis (BST), medial amygdaloid nucleus (MeA), and
46 lateral septum (LS), of male (XY) and female (XX) wild type rats as well as in *Tfm*
47 rats of both sexes ($X^{tfm}Y$ and $X^{tfm}X$). *** p <0.001, ** p <0.01, * p <0.05 different from
48 XY males;
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES

- Attardi B, Geller LN, Ohno S (1976) Androgen and estrogen receptors in brain cytosol from male, female, and testicular feminized (tfm/y hermaphrodite) mice. *Endocrinology* 98:864-874.
- Axelsson JF, Van Leeuwen FW (1990) Differential localization of estrogen receptors in various vasopressin synthesizing nuclei of the rat brain. *J Neuroendocrinol* 2:209-216.
- Bakker J, De Mees C, Douhard Q, Balthazart J, Gabant P, Szpirer J, Szpirer C (2006) Alpha-fetoprotein protects the developing female mouse brain from masculinization and defeminization by estrogens. *Nat Neurosci* 9:220-226.
- Balthazart J, Adkins-Regan E (2002) Sexual differentiation of brain and behavior in birds. In: *Hormones, Brain and Behavior*, vol. 4 (Pfaff, D. W. et al., eds), pp 223-301 New York: Academic Press.
- Bingham B, Viau V (2008) Neonatal gonadectomy and adult testosterone replacement suggest an involvement of limbic arginine vasopressin and androgen receptors in the organization of the hypothalamic-pituitary-adrenal axis. *Endocrinology* 149:3581-3591.
- Brat DJ, Brimijoin S (1992) A paradigm for examining toxicant effects on viability, structure, and axonal transport of neurons in culture. *Mol Neurobiol* 6:125-135.
- Brinkmann AO (2001) Molecular basis of androgen insensitivity. *Mol Cell Endocrinol* 179:105-109.
- Caffé AR, Van Leeuwen FW, Luiten PGM (1987) Vasopressin cells in the medial amygdala of the rat project to the lateral septum and ventral hippocampus. *J Comp Neurol* 261:237-252.
- Choleris E, Clipperton-Allen AE, Phan A, Kavaliers M (2009) Neuroendocrinology of social information processing in rats and mice. *Front Neuroendocrinol* 30:442-459.
- Compaan JC, Buijs RM, Pool CW, De Ruiter AJH, Koolhaas JM (1993) Differential lateral septal vasopressin innervation in aggressive and nonaggressive male mice. *Brain Res Bull* 30:1-6.
- Cooke B, Hegstrom CD, Villeneuve LS, Breedlove SM (1998) Sexual differentiation of the vertebrate brain: principles and mechanisms. *Front Neuroendocrinol* 19:323-362.
- Cooke BM, Breedlove SM, Jordan CL (2003) Both estrogen receptors and androgen receptors contribute to testosterone-induced changes in the morphology of the medial amygdala and sexual arousal in male rats. *Horm Behav* 43:336-346.
- De Vries GJ, Al Shamma HA (1990) Sex differences in hormonal responses of vasopressin pathways in the rat brain. *J Neurobiol* 21:686-693.
- De Vries GJ, Buijs RM (1983) The origin of the vasopressinergic and oxytocinergic innervation of the rat brain with special reference to the lateral septum. *Brain Res* 273:307-317.
- De Vries GJ, Buijs RM, Sluiter AA (1984a) Gonadal hormone actions on the morphology of the vasopressinergic innervation of the adult rat brain. *Brain Res* 298:141-145.

- 1 De Vries GJ, Buijs RM, Van Leeuwen FW (1984b) Sex differences in vasopressin
2 and other neurotransmitter systems in the brain. *Prog Brain Research* 61:185-
3 203.
- 4 De Vries GJ, Buijs RM, Van Leeuwen FW, Caffé AR, Swaab DF (1985) The
5 vasopressinergic innervation of the brain in normal and castrated rats. *J Comp*
6 *Neurol* 233:236-254.
- 7 De Vries GJ, Panzica GC (2006) Sexual differentiation of central vasopressin and
8 vasotocin systems in vertebrates: different mechanisms, similar endpoints.
9 *Neuroscience* 138:947-955.
- 10 De Vries GJ, Rissman EF, Simerly RB, Yang LY, Scordalakes EM, Auger CJ, Swain
11 A, Lovell-Badge R, Burgoyne PS, Arnold AP (2002) A model system for
12 study of sex chromosome effects on sexually dimorphic neural and behavioral
13 traits. *J Neurosci* 22:9005-9014.
- 14 Del Abril A, Segovia S, Guillamon A (1987) The bed nucleus of the stria terminalis in
15 the rat: regional sex differences controlled by gonadal steroids early after
16 birth. *Dev Brain Res* 32:295-300.
- 17 DonCarlos LL (1996) Developmental Profile and Regulation of Estrogen Receptor
18 (Er) mRNA Expression in the Preoptic Area of Prenatal Rats. *Dev Brain Res*
19 94:224-233.
- 20 DonCarlos LL, Handa RJ (1994) Developmental profile of estrogen receptor mRNA
21 in the preoptic area of male and female neonatal rats. *Dev Brain Res* 79:283-
22 289.
- 23 Dugger BN, Morris JA, Jordan CL, Breedlove SM (2007) Androgen receptors are
24 required for full masculinization of the ventromedial hypothalamus (VMH) in
25 rats. *Horm Behav* 51:195-201.
- 26 Edelman M, Wolfe C, Scordalakes EM, Rissman EF, Tobet S (2007) Neuronal nitric
27 oxide synthase and calbindin delineate sex differences in the developing
28 hypothalamus and preoptic area. *Dev Neurobiol* 67:1371-1381.
- 29 Fernandez R, Collado P, Garcia Doval S, Garcia-Falgueras A, Guillamon A, Pasaro E
30 (2003) A molecular method for classifying the genotypes obtained in a
31 breeding colony from testicular feminized (Tfm) rats. *Horm Metab Res*
32 35:197-200.
- 33 Ferris CF, Delville Y, Miller MA, Dorsa DM, De Vries GJ (1995) Distribution of
34 small vasopressinergic neurons in golden hamsters. *J Comp Neurol* 360:589-
35 598.
- 36 Field EF, Whishaw IQ, Pellis SM, Watson NV (2006) Play fighting in androgen-
37 insensitive tfm rats: evidence that androgen receptors are necessary for the
38 development of adult playful attack and defense. *Dev Psychobiol* 48:111-120.
- 39 Forbes-Lorman RM, Rautio JJ, Kurian JR, Auger AP, Auger CJ (2012) Neonatal
40 MeCP2 is important for the organization of sex differences in vasopressin
41 expression. *Epigenetics* 7:230-238.
- 42 Garcia-Falgueras A, Pinos H, Collado P, Pasaro E, Fernandez R, Jordan CL, Segovia
43 S, Guillamon A (2005) The role of the androgen receptor in CNS
44 masculinization. *Brain Res* 1035:13-23.
- 45 Gobinet J, Poujol N, Sultan C (2002) Molecular action of androgens. *Mol Cell*
46 *Endocrinol* 198:15-24.
- 47 Goodson JL, Bass AH (2001) Social behavior functions and related anatomical
48 characteristics of vasotocin/vasopressin systems in vertebrates. *Brain Res Rev*
49 35:246-265.

- 1 Grassi D, Amorim MA, Garcia-Segura LM, Panzica G (2010) Estrogen receptor alpha
2 is involved in the estrogenic regulation of arginine vasopressin
3 immunoreactivity in the supraoptic and paraventricular nuclei of
4 ovariectomized rats. *Neurosci Lett* 474:135-139.
- 5 Hajszan T, Maclusky NJ, Johansen JA, Jordan CL, Leranath C (2007) Effects of
6 androgens and estradiol on spine synapse formation in the prefrontal cortex of
7 normal and testicular feminization mutant male rats. *Endocrinology* 148:1963-
8 1967.
- 9 Han TM, De Vries GJ (2003) Organizational Effects of Testosterone, Estradiol, and
10 Dihydrotestosterone on Vasopressin mRNA Expression in the Bed Nucleus of
11 the Stria Terminalis. *J Neurobiol* 54:502-510.
- 12 Hines M, Allen LS, Gorski RA (1992) Sex Differences in Subregions of the Medial
13 Nucleus of the Amygdala and the Bed Nucleus of the Stria Terminalis of the
14 Rat. *Brain Res* 579:321-326.
- 15 Hughes IA, Davies JD, Bunch TI, Pasterski V, Mastroyannopoulou K, MacDougall J
16 (2012) Androgen insensitivity syndrome. *Lancet* 380:1419-1428.
- 17 Johnson RT, Breedlove SM, Jordan CL (2008) Sex differences and laterality in
18 astrocyte number and complexity in the adult rat medial amygdala. *J Comp*
19 *Neurol* 511:599-609.
- 20 Juntti SA, Tollkuhn J, Wu MV, Fraser EJ, Soderborg T, Tan S, Honda S, Harada N,
21 Shah NM (2010) The androgen receptor governs the execution, but not
22 programming, of male sexual and territorial behaviors. *Neuron* 66:260-272.
- 23 Lenz KM, Nugent BM, McCarthy MM (2012) Sexual differentiation of the rodent
24 brain: dogma and beyond. *Front Neurosci* 6:26.
- 25 Martini M, Di Sante G, Collado P, Pinos H, Guillamon A, Panzica GC (2008)
26 Androgen receptors are required for full masculinization of nitric oxide
27 synthase system in rat limbic-hypothalamic region. *Horm Behav* 54:557-564.
- 28 McAbee MD, DonCarlos LL (1998) Ontogeny of region-specific sex differences in
29 androgen receptor messenger ribonucleic acid expression in the rat forebrain.
30 *Endocrinology* 139:1738-1745.
- 31 McAbee MD, DonCarlos LL (1999a) Estrogen, but not androgens, regulates androgen
32 receptor messenger ribonucleic acid expression in the developing male rat
33 forebrain. *Endocrinology* 140:3674-3681.
- 34 McAbee MD, DonCarlos LL (1999b) Regulation of Androgen Receptor Messenger
35 Ribonucleic Acid Expression in the Developing Rat Forebrain. *Endocrinology*
36 140:1807-1814.
- 37 McCarthy MM (2008) Estradiol and the developing brain. *Physiol Rev* 88:91-134.
- 38 McCarthy MM, Arnold AP (2011) Reframing sexual differentiation of the brain. *Nat*
39 *Neurosci* 14:677-683.
- 40 McCarthy MM, Konkle ATM (2005) When is a sex difference not a sex difference?
41 *Front Neuroendocrinol* 26:85-102.
- 42 McCarthy MM, Wright CL, Schwarz JM (2009) New tricks by an old dogma:
43 mechanisms of the Organizational/Activational Hypothesis of steroid-
44 mediated sexual differentiation of brain and behavior. *Horm Behav* 55:655-
45 665.
- 46 McPhaul MJ (2002) Androgen receptor mutations and androgen insensitivity. *Mol*
47 *Cell Endocrinol* 198:61-67.
- 48 Morris JA, Jordan CL, Dugger BN, Breedlove SM (2005) Partial demasculinization of
49 several brain regions in adult male (XY) rats with a dysfunctional androgen
50 receptor gene. *J Comp Neurol* 487:217-226.
- 51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 Pak TR, Chung WC, Hinds LR, Handa RJ (2007) Estrogen receptor-beta mediates
2 dihydrotestosterone-induced stimulation of the arginine vasopressin promoter
3 in neuronal cells. *Endocrinology* 148:3371-3382.
- 4 Pak TR, Chung WC, Hinds LR, Handa RJ (2009) Arginine vasopressin regulation in
5 pre- and postpubertal male rats by the androgen metabolite 3beta-diol. *Am J*
6 *Physiol Endocrinol Metab* 296:E1409-1413.
- 7 Panzica GC, Aste N, Castagna C, Viglietti-Panzica C, Balthazart J (2001) Steroid-
8 induced plasticity in the sexually dimorphic vasotocinergic innervation of the
9 avian brain: behavioral implications. *Brain Res Rev* 37:178-200.
- 10 Panzica GC, Aste N, Viglietti-Panzica C, Ottinger MA (1995) Structural sex
11 differences in the brain: influence of gonadal steroids and behavioral
12 correlates. *J Endocrinol Invest* 18:232-252.
- 13 Paxinos G, Watson C (1998) The rat brain in stereotaxic coordinates. Fourth Edition.
14 San Diego: Academic Press.
- 15 Perakis A, Stylianopoulou F (1986) Effects of a prenatal androgen peak on rat brain
16 sexual differentiation. *J Endocrinol* 108:281-285.
- 17 Pierman S, Sica M, Allieri F, Viglietti-Panzica C, Panzica GC, Bakker J (2008)
18 Activational effects of estradiol and dihydrotestosterone on social recognition
19 and the arginine-vasopressin immunoreactive system in male mice lacking a
20 functional aromatase gene. *Horm Behav* 54:98-106.
- 21 Plumari L, Viglietti Panzica C, Allieri F, Honda S, Harada N, Absil P, Balthazar J,
22 Panzica GC (2002) Changes in the Arginine-Vasopressin Immunoreactive
23 Systems in Male Mice Lacking a Functional Aromatase Gene. *J*
24 *Neuroendocrinol* 14:971-978.
- 25 Raskin K, de Gendt K, Duittoz A, Liere P, Verhoeven G, Tronche F, Mhaouty-Kodja
26 S (2009) Conditional inactivation of androgen receptor gene in the nervous
27 system: effects on male behavioral and neuroendocrine responses. *J Neurosci*
28 29:4461-4470.
- 29 Robinson S, Penatti CA, Clark AS (2012) The role of the androgen receptor in
30 anabolic androgenic steroid-induced aggressive behavior in C57BL/6J and
31 Tfm mice. *Horm Behav* 61:67-75.
- 32 Rood BD, De Vries GJ (2011) Vasopressin innervation of the mouse (*Mus musculus*)
33 brain and spinal cord. *J Comp Neurol* 519:2434-2474.
- 34 Rood BD, Stott RT, You S, Smith CJ, Woodbury ME, De Vries GJ (2013) Site of
35 origin of and sex differences in the vasopressin innervation of the mouse (*Mus*
36 *musculus*) brain. *J Comp Neurol* 521:2321-2358.
- 37 Roselli CE, Salisbury RL, Resko JA (1987) Genetic evidence for androgen-dependent
38 and independent control of aromatase activity in the rat brain. *Endocrinology*
39 121:2205-2210.
- 40 Rotzinger S, Lovejoy DA, Tan LA (2010) Behavioral effects of neuropeptides in
41 rodent models of depression and anxiety. *Peptides* 31:736-756.
- 42 Scordalakes EM, Rissman EF (2004) Aggression and arginine vasopressin
43 immunoreactivity regulation by androgen receptor and estrogen receptor
44 alpha. *Genes Brain Behav* 3:20-26.
- 45 Segovia S, Garcia-Falgueras A, Carrillo B, Collado P, Pinos H, Perez-Laso C,
46 Vinader-Caerols C, Beyer C, Guillamon A (2006) Sexual dimorphism in the
47 vomeronasal system of the rabbit. *Brain Res* 1102:52-62.
- 48 Streefkerk JG (1972) Inhibition of erythrocyte peroxidase activity by treatment with
49 hydrogen peroxide following methanol. *J Histochem Cytochem* 20:829-831.
- 50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 Tejada LD, Rissman EF (2012) Sex differences in social investigation: effects of
2 androgen receptors, hormones and test partner. *J Neuroendocrinol* 24:1144-
3 1153.
- 4 Veenema AH, Beiderbeck DI, Lukas M, Neumann ID (2010) Distinct correlations of
5 vasopressin release within the lateral septum and the bed nucleus of the stria
6 terminalis with the display of intermale aggression. *Horm Behav* 58:273-281.
- 7 Veenema AH, Neumann ID (2008) Central vasopressin and oxytocin release:
8 regulation of complex social behaviours. *Prog Brain Res* 170:261-276.
- 9 Veening JG, Coolen LM (1998) Neural activation following sexual behavior in the
10 male and female rat brain. *Behav Brain Res* 92:181-193.
- 11 Veening JG, Coolen LM, de Jong TR, Joosten HW, de Boer SF, Koolhaas JM, Olivier
12 B (2005) Do similar neural systems subservise aggressive and sexual behaviour
13 in male rats? Insights from c-Fos and pharmacological studies. *Eur J*
14 *Pharmacol* 526:226-239.
- 15 Viglietti-Panzica C, Balthazart J, Fratesi S, Plumari L, Absil P, Panzica GC (2001)
16 Estradiol mediates effects of testosterone on vasotocin-immunoreactivity in
17 the adult quail brain. *Horm Behav* 40:445-461.
- 18 Viglietti-Panzica C, Mura E, Panzica GC (2007) Effects of early embryonic exposure
19 to genistein on male copulatory behavior and vasotocin system of Japanese
20 quail. *Horm Behav* 51:355-363.
- 21 Villalba C, Auger CJ, de Vries GJ (1999) Androstenedione effects on the vasopressin
22 innervation of the rat brain. *Endocrinology* 140:3383-3386.
- 23 Wallen K (2005) Hormonal influences on sexually differentiated behavior in
24 nonhuman primates. *Front Neuroendocrinol* 26:7-26.
- 25 Wang Z, Bullock NA, De Vries GJ (1993) Sexual differentiation of vasopressin
26 projections of the bed nucleus of the stria terminals and medial amygdaloid
27 nucleus in rats. *Endocrinology* 132:2299-2306.
- 28 Watson RE, Wiegand SJ, Clough RW, Hoffman GE (1986) Use of cryoprotectant to
29 maintain long-term peptide immunoreactivity and tissue morphology. *Peptides*
30 7:155-159.
- 31 Wu MV, Manoli DS, Fraser EJ, Coats JK, Tollkuhn J, Honda S, Harada N, Shah NM
32 (2009) Estrogen masculinizes neural pathways and sex-specific behaviors.
33 *Cell* 139:61-72.
- 34 Zhou JN, Hofman MA, Gooren LJG, Swaab DF (1995) A sex difference in the human
35 brain and its relation to transsexuality. *Nature* 378:68-70.
- 36 Zhou L, Blaustein JD, De Vries GJ (1994) Distribution of androgen receptor
37 immunoreactivity in vasopressin- and oxytocin-immunoreactive neurons in the
38 male rat brain. *Endocrinology* 134:2622-2627.
- 39 Zuloaga DG, Morris JA, Jordan CL, Breedlove SM (2008) Mice with the testicular
40 feminization mutation demonstrate a role for androgen receptors in the
41 regulation of anxiety-related behaviors and the hypothalamic-pituitary-adrenal
42 axis. *Horm Behav* 54:758-766.
- 43 Zuloaga DG, Poort JE, Jordan CL, Breedlove SM (2011) Male rats with the testicular
44 feminization mutation of the androgen receptor display elevated anxiety-
45 related behavior and corticosterone response to mild stress. *Horm Behav*
46 60:380-388.
- 47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)




Figure 5
[Click here to download high resolution image](#)

