

IRIS A_{per}TO



UNIVERSITÀ
DEGLI STUDI
DI TORINO

This is the author's final version of the contribution published as:

A. ZIPPO; A. DE ROBERTIS; R. SERAFINI; S. OLIVIERO.
PIM1-dependent phosphorylation of Histone H3 at Serine 10 is required for
MYC-dependent transcriptional activation and oncogenic transformation..
NATURE CELL BIOLOGY. 9 pp: 932-944.

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/141074>

This full text was downloaded from iris - AperTO: <https://iris.unito.it/>

iris - AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

PIM1-dependent phosphorylation of Histone H3 at Serine 10 is required for MYC-dependent transcriptional activation and oncogenic transformation.

Alessio Zippo, Alessandra De Robertis, Riccardo Serafini, and Salvatore Oliviero

Dipartimento di Biologia Molecolare Universita' di Siena

Via Fiorentina 1, 53100 Siena, Italy.

Running title: PIM1 phosphorylates H3S10

Key words: transcriptional regulation, MYC, PIM1, chromatin, nucleosome phosphorylation.

Corresponding Author :

Salvatore Oliviero

Dipartimento di Biologia Molecolare Universita' di Siena

Via Fiorentina 1, 53100 Siena, Italy.

e mail oliviero@unisi.it

The serine/threonine kinase PIM1, cooperates with MYC in cell cycle progression and tumorigenesis. However, the nature of this cooperation remains elusive. Here we show that following growth factor stimulation, PIM1 forms a complex with the MYC/MAX dimer via the MYC BoxII domain (MBII). MYC recruits PIM1 to the E boxes of the MYC-target genes FOSL1 (FRA1) and ID2 and PIM1 phosphorylates Serine 10 of histone H3 (H3S10) on the nucleosome at the MYC-binding sites contributing to their transcriptional activation. MYC and PIM1 co-localize at sites of active transcription and expression profile analysis revealed that PIM1 contributes to the regulation of 20% of the MYC-regulated genes. Moreover, PIM1-dependent H3S10 phosphorylation contributes to MYC transforming capacity. These results establish a new function for PIM1 as a MYC cofactor that phosphorylates the chromatin at MYC-target loci and suggest that nucleosome phosphorylation, at E boxes, contributes to MYC-dependent transcriptional activation and cellular transformation.

Chromatin structure plays a crucial role in eukaryotic gene transcription. Several transcription activators have been shown to recruit chromatin-modifying activities, which either mediate histone covalent modifications or ATP-dependent nucleosome mobilization.

Covalent histone modifications including acetylation, methylation, phosphorylation and ubiquitinylation have been associated with gene regulation^{1,2}. Phosphorylation of H3S10 has been connected with transcriptional activation in different organisms and with chromosome condensation during mitosis^{3,4}. In mammalian cells, growth factor stimulation induces rapid phosphorylation of histone H3 at Serine 10 (H3S10) at c-Jun and c-Fos promoters mediated by MSK1, MSK2, and RSK2 kinases⁴⁻⁷. In response to inflammatory cytokines, IKK- γ phosphorylates H3S10 at NF-kB-responsive promoters^{8,9}.

In this study, we analyzed the function of PIM1, a constitutively active serine/threonine kinase. Pim1 was first identified as a common insertion site in T cell lymphomas induced by Molony murine leukemia virus (Mo-MuLV)¹⁰. RNA interference (RNAi) experiments demonstrated that human Pim1 (hereinafter referred to as PIM1) is essential for cell cycle progression of human umbilical vein endothelial cells (HUVEC) and for differentiation of endothelial precursors *in vitro*¹¹. Myc and Pim1 have been shown to co-operate in cell growth and transformation. Pim1 over-expression in transgenic mice demonstrated its cooperation with c-Myc in cell transformation¹²⁻¹⁵ and microarray expression profiling showed that Pim1 is over-expressed in Myc-driven prostate tumours¹⁶. However, the nature of c-Myc and Pim1 cooperation in cell transformation remains elusive.

Human c-Myc (hereinafter referred as MYC) codes for a basic helix-loop-helix leucine zipper (B-HLH) transcription factor that binds to the E box (CACGTG) when dimerized with MAX and regulates the transcription of distinct genes involved in cell cycle progression, apoptosis, cell growth, and differentiation¹⁷⁻²³. The DNA binding domain of MYC is at the C-terminus while the N-terminal domain through the MYC boxII (MBII), essential for MYC-dependent cell transformation, has been shown to recruit multiprotein enzymatic complexes to the MYC-activated genes²⁴.

To understand the function of PIM1, we investigated the molecular connection between PIM1 and MYC. We propose that the recruitment of PIM1 to the chromatin by MYC contributes to transcriptional activation of MYC-target genes by phosphorylating H3S10 at the E box element and suggest that this cooperation is relevant for MYC-dependent tumour formation.

RESULTS

PIM1 phosphorylates the nucleosome at H3S10.

PIM1 mRNA is induced with fast kinetics in HUVEC treated with VEGF-A¹¹. Upon VEGF-A treatment PIM1 is induced and localizes in HUVEC nuclei within 40 minutes (see Supplementary Information Fig. S1). As PIM1 is a serine/threonine kinase with putative consensus sites present on histones, its nuclear localization could account for the increase of H3 phosphorylation observed in these cells between 60 and 90 minutes after VEGF-A

treatment (see Supplementary Information Fig. S1). We found that PIM1 could directly phosphorylate nucleosomes *in vitro* by incubating the recombinant GST-PIM1 protein with chromatin fractions obtained from 293 cells in the presence of [γ - 32 P] ATP. This analysis revealed that a single protein, with molecular weight corresponding to H3, was phosphorylated by wild type GST-PIM1, but not by the kinase inactive mutant GST-PIM1-K67M (Fig. 1a, lanes 1-3). Specific inhibition of nucleosome phosphorylation was observed in the presence of a peptide derived from the H3 N-terminus but not with a peptide derived from H2B N-terminus or with an unrelated peptide (Fig. 1a, lanes 4-12). PIM1 could phosphorylate reconstituted nucleosomes containing either recombinant wild type H3 or H3S28A, but not nucleosomes assembled with the H3S10A mutant or the double mutant (Fig. 1b) demonstrating that PIM1 phosphorylates H3 at S10. Histone H3 co-immunoprecipitated with transiently expressed PIM1 or the kinase inactive mutant suggesting that PIM1 associates *in vivo* with the nucleosome (Fig. 1c). Incubation of the immunoprecipitate in the presence of [γ - 32 P] ATP showed that the wild type PIM1, but not the kinase inactive mutant PIM1-K67M phosphorylated H3. We conclude that PIM1 associates *in vitro* and *in vivo* with H3 and directly phosphorylates H3S10 on the nucleosome.

PIM1 forms a complex with MYC and MAX.

VEGF-A treatment induces MYC and PIM1 co-localization in the cell nuclei at 60 minutes, decreasing thereafter (Fig. 2a). We tested whether PIM1 coimmunoprecipitates with MYC.

Time course analysis in HUVEC treated with VEGF-A showed that upon induction, both MYC and PIM1 were induced with fast kinetics. Immunoprecipitations with anti-MAX antibodies at various time points showed the formation of the MAX/MYC complex from 60 minutes after growth factor stimulation thus corresponding to the appearance of MYC in the nucleus (Fig. 2b, right panel). Importantly, PIM1 co-immunoprecipitated with MYC and MAX suggesting that a complex containing PIM1 is formed. Endogenous PIM1 was also found to co-immunoprecipitate with endogenous MYC and MAX from 293 cells after serum treatment for 120 minutes (Fig. 2c), thus demonstrating that a MAX/MYC/PIM1 endogenous complex is also formed in these cells.

As MYC-dependent cell transformation requires MYC boxII (MBII) domain to bind cofactors either in a TRRAP-dependent or -independent manner²⁴, we tested whether PIM1 interacts with MYC at MBII via TRRAP. Wild type FLAG-MYC or MYC deleted in the MBII domain (FLAG-MYC^ΔMBII) were co-expressed with PIM1 in 293 cells and immunoprecipitated either with FLAG or PIM1 antibodies. Western blot analysis showed reciprocal immunoprecipitation of FLAG-MYC with PIM1, while co-immunoprecipitation was strongly reduced between FLAG-MYC^ΔMBII and PIM1 suggesting that MBII domain of MYC is required for the formation of a complex with PIM1 (Fig. 2d). Endogenous TRRAP associated with MYC but not with the FLAG-MYC^ΔMBII as previously described²⁵. In contrast, PIM1 did not co-immunoprecipitate the endogenous TRRAP (lanes 5 and 10) suggesting that MYC forms independent complexes with PIM1 or TRRAP via the MBII domain.

PIM1 co-localizes with MYC on the chromatin at actively transcribing loci.

To measure PIM1 and MYC association at MYC-target genes, we performed a nuclear distribution analysis of PIM1 with respect to MYC, nascent transcripts, and chromatin markers. Quantitative analysis of double staining with anti-PIM1 and antibodies against different nuclear markers by sequential laser scanning (Fig. 3) showed an elevated degree of PIM1 co-localization with MYC with a Pearson's correlation coefficient (Rr) of 0.82 ± 0.06 . PIM1 also showed an elevated overlap with H3S10ph (Rr 0.85 ± 0.07) and H3S10phK14ac while little co-localization could be detected between PIM1 and H3K9me2 (Rr 0.37 ± 0.03), a marker of silent heterochromatin. Furthermore, we analyzed the nuclear distribution of PIM1 with respect to active transcription sites by in situ transcriptional run-on experiments. Double labelling revealed a high degree of overlap between PIM1 and sites of active transcription (Rr 0.77 ± 0.05) with a pattern resembling that of transcription factories²⁶. A similar pattern of nuclear localization with respect to nascent transcripts was observed for MYC (Rr 0.81 ± 0.09). Taken together these results suggest that following growth factor treatment PIM1 kinase is mostly localized with the transcription factor MYC at actively transcribing sites and with phosphorylated histones.

H3S10 phosphorylation at the FOSL1 enhancer correlates with PIM1 association.

Next we analysed PIM1 association and H3S10 phosphorylation at FOSL1 (also known as FRA1), a well characterized MYC-target gene^{27,28}. FOSL1 contains an upstream

functional SRE, which is not sufficient to confer promoter activation, but requires the activity of a downstream enhancer that is activated by MYC and AP-1²⁷⁻²⁹. Upon VEGF-A treatment of HUVEC FOSL1 expression is induced with a peak at 60 minutes (Fig. 4a). To demonstrate a direct binding of MYC and PIM1 to the FOSL1 regulatory sequences ChIP followed by real-time quantitative PCR analysis was performed at various timepoints after VEGF-A treatment. As shown in Figure 4b, we generated primers to analyze four FOSL1 regions: a FOSL1 upstream region containing a functional SRE and a non canonical E box element^{30,31} (probe A), the downstream enhancer containing a canonical E box (probe B), a downstream region within the first intron (probe C), and a FOSL1 region within the second intron (probe D). Quantitative ChIP analysis with probe B revealed the transient recruitment to the FOSL1 enhancer of MYC and PIM1 from 60 minutes concomitant with a marked increase of H3S10ph signal (Fig. 4c). ChIP with probe B also showed a significant increase of H3S10phK9ac and H3S10phK14ac while H3K9acK14ac showed elevated levels before induction suggesting that, at the FOSL1 enhancer, H3 acetylation precedes MYC binding and H3S10 phosphorylation takes place at the acetylated H3.

Analysis of the upstream element (probe A) revealed MYC association with this DNA fragment. PIM1 did not associate with this element suggesting that the FOSL1 promoter is bound by an alternative MYC complex. This FOSL1 upstream element showed an increase of H3S10 phosphorylation at 15 minutes after VEGF-A treatment, which was therefore not dependent on PIM1. As the FOSL1 SRE element is constitutively occupied by the ternary

complex factor (TCF), which is activated by the RAS/ERK pathway³⁰, we further analyzed FOSL1 transcription and H3S10 phosphorylation in the presence of H89, a specific MSK1/MSK2 inhibitor⁵. Cell treatment with H89 affected FOSL1 transcription although a VEGF-A-dependent induction was still observed (Fig. 4d). The analysis of H3S10ph signal showed specific inhibition of H3S10 phosphorylation at the FOSL1 upstream element but not at the enhancer (Fig. 4e) demonstrating that phosphorylation at the upstream element and at the downstream enhancer are independent from each other and mediated by different kinases. The acetylation pattern at the FOSL1 upstream element showed a significant increase of H3 phosphorylation and acetylation with a peak between 15 and 30 minutes suggesting that, at this site, H3S10 phosphorylation precedes or is concomitant with H3 acetylation at K9 and K14 as previously described for IE promoters^{3,4}.

The downstream FOSL1 fragments (analyzed with probes C and D) did not reveal MYC or PIM1 binding, nor the increase of phosphorylated H3S10, while presented elevated levels of acetylated H3. These results demonstrate specific recruitment of PIM1 to the FOSL1 enhancer concomitant with an increase of H3S10 phosphorylation at this site and with transcriptional activation of the gene.

H3S10 phosphorylation at FOSL1 enhancer is mediated by PIM1.

PIM1 depletion using small hairpin RNA (shRNA) was performed to test whether PIM1

association with the FOSL1 enhancer was required for H3S10 phosphorylation. In 293 cells treated with serum MYC, PIM1, and FOSL1 showed a slower kinetics of induction with respect to HUVEC treated with VEGF-A possibly due to the different cell types (cell line versus primary cells). PIM1 silencing strongly inhibited FOSL1 mRNA expression (Fig. 7b and see Supplementary Information Fig. S3). Quantitative ChIP analysis (Fig. 5b and see Supplementary Information Fig. S2) demonstrated a peak of H3S10 phosphorylation at the FOSL1 enhancer (probe B as described in Fig. 4b) at 120 minutes that was significantly reduced by PIM1 silencing, while ChIP analysis at the FOSL1 upstream element (probe A) showed a faster kinetics of H3S10 phosphorylation that was not affected by PIM1 knockdown. No H3S10 phosphorylation was observed at the downstream fragments (probes C and D). We concluded that H3S10 phosphorylation at the FOSL1 enhancer is PIM1-dependent.

PIM1 and MYC are found to associate at the FOSL1 enhancer with a peak between 90 and 120 minutes. PIM1 silencing significantly reduced PIM1 association at the enhancer without affecting MYC association. The overall pattern of H3 phospho-acetylation at K9 and K14 in 293 cells showed elevated levels of H3 acetylation already present before H3 phosphorylation at the FOSL1 enhancer (probe B) and along the FOSL1 gene (probe C and D) which was not affected by PIM1 silencing.

Moreover, expressing the kinase inactive mutant PIM1-K67M, significantly reduced H3S10ph levels at the FOSL1 enhancer (see Supplementary Information Fig. S3) demonstrating that the kinase activity of PIM1 is required for H3S10 phosphorylation at

this site.

PIM1 is recruited by MYC at the FOSL1 and ID2 E box elements.

MYC silencing was performed to analyze whether its expression is required for PIM1 recruitment to the FOSL1 enhancer (Fig. 6a and see Supplementary Information Fig. S4). We also analyzed MYC-dependent recruitment of PIM1 at the ID2 E box. ID2 is a well characterized MYC-regulated gene that, differently from FOSL1, contains its functional MYC-binding sites upstream the promoter^{28,32}. As expected, ChIP analysis on control cells showed MYC association at the FOSL1 enhancer and the ID2 E box at 120 minutes after serum treatment, while MYC silencing significantly reduced MYC association at these sites (Fig. 6b and see Supplementary Information Fig. S4). ChIP analysis revealed PIM1 association with both genes in control cells while MYC knockdown significantly reduced PIM1 association at these E boxes (Fig. 6c and see Supplementary Information Fig. S4). Moreover, H3S10 phosphorylation showed significant reduction in MYC silenced cells at these sites (Fig. 6d and see Supplementary Information Fig. S4). In contrast, H3K9acK14ac was unaffected by MYC silencing (Fig. 6e and see Supplementary Information Fig. S4) confirming that H3 acetylation is not dependent on MYC binding. Importantly, the reintroduction of wild type FLAG-MYC, but not the FLAG-MYC^{ΔMBII} in MYC-silenced cells restored PIM1 recruitment and H3S10 phosphorylation at both sites (Fig. 6 f, g, h and see Supplementary Information Fig. S4). These results establish that MYC, via its MBII domain, recruits PIM1 to the E box elements of these MYC-regulated

genes and its recruitment is required for phosphorylating H3S10 at these sites.

PIM1 is required for transcriptional activation of FOSL1 and ID2 genes.

After serum treatment of 293 cells, both FOSL1 and ID2 mRNA were induced with similar kinetics (Fig. 7b, e) corresponding to the peak of PIM1 and MYC association and H3S10 phosphorylation in these cells. The same results were obtained by measuring FOSL1 and ID2 unprocessed heterogeneous nuclear RNA (hnRNA) indicating that increased mRNA level depend on transcriptional activation. PIM1 silencing, as well as the expression of PIM1-K67M, inhibited induction of both FOSL1 and ID2 transcripts (Fig 7b, e and see Supplementary Information Fig. S3) demonstrating that PIM1 kinase activity is required for full transcriptional activation of both genes.

Quantitative ChIP analysis of RNA Polymerase II (Pol II) and as well as Pol II phosphorylated in Ser2 (Ser2P) at the C-terminal domain (CTD) of control cells revealed a significant increase in signal at FOSL1 and ID2 promoters and along the genes in response to serum treatment (Fig. 7c, f and see Supplementary Information Fig. S3) mirroring of the increase in transcript levels. PIM1 silencing led to a specific reduction of Pol II signal at the promoters and within the coding regions. This reduction, however, could not account for the inhibition of the transcriptional activity mediated by the PIM1 silencing. By contrast, the reduction of the elongating Pol II (Pol II-Ser2P) along the gene more closely correlated with transcript levels of both FOSL1 and ID2 suggesting that PIM1 silencing mostly affects the presence of Pol II engaged in transcript elongation.

PIM1 contributes to the regulation of a subgroup of MYC-regulated genes.

To understand the extent of PIM1 and MYC co-operation in gene transcription we performed expression profile analysis of 293 cells silenced either for MYC or PIM1 at 120 minutes after serum treatment. MYC silencing affected the expression of 1026 genes of which 818 were up-regulated and 208 were down-regulated (see Supplementary Information Table SI) in line with previous analysis of MYC-regulated genes³³. Comparison of genes regulated by MYC with those regulated by PIM1 showed that PIM1 contributes to the regulation of 207 genes out of the 1026 MYC-regulated genes. Thus, 20% of MYC-regulated genes, are also regulated by PIM1. Functional characterization of these genes, based on Gene Ontology categorization through the Panther database (<http://www.pantherdb.org>), revealed that nucleic acid metabolism, transcriptional regulation, RNA processing, and oncogenesis are statistically overrepresented ($P < 0.05$) (see Supplementary Information Table SII, SIII). Based on this analysis we conclude that PIM1 act as a cofactor for transcriptional regulation in a significant subset of MYC-dependent genes.

PIM1 recruitment to the chromatin by MYC contributes to cell transformation.

Next we investigated the biological relevance of MYC - PIM1 co-operation. First we verified whether PIM1 contributes to MYC-dependent regulation of cell cycle progression by silencing PIM1 in P493-6 cells, a cell line that depends on MYC expression for its

proliferation^{27,34,35}. Incorporation of [³H] thymidine was significantly decreased in PIM1 silenced cells and analysis of the cell cycle by FACS showed that PIM1 depletion determines a delay in S phase progression of these cells (see Supplementary Information Fig. S5).

To test whether PIM1 cooperation with MYC in cell transformation and tumour growth is due to PIM1's role in transcriptional activation of MYC-target genes, we performed transformation experiments in Rat-1 fibroblasts. Over-expression of MYC transforms these cells morphologically and induces them to grow in soft agar³⁶. PIM1 silencing strongly inhibited the formation of soft agar foci, induced by MYC over-expression, and MYC silencing affected the number of foci induced by PIM1 over-expression to a similar extent, (Fig. 8a) thus, demonstrating that PIM1 is indeed required for MYC-dependent cell transformation. Moreover, the over-expression of MYC and PIM1 generates a significantly higher number of soft agar colonies than cells over-expressing either MYC or PIM1 (Fig. 8a, b).

To understand whether PIM1 acts as a MYC cofactor in cell transformation we generated chimeric constructs in which PIM1 was directly fused to the N terminal domain of MYC (PIM1-MYC). We found that the PIM1-MYC fusion molecule induced a comparable number of soft agar colonies relative to the co-expression of MYC and PIM1 in the same cells (Fig. 8b) suggesting that PIM1 recruitment to the chromatin by MYC is required for cooperation between MYC and PIM1 in cell transformation. Since formation of the MYC complex with PIM1 and its recruitment to MYC binding sites requires the MYC MBII

domain (Fig. 2, 6), which is essential for MYC oncogenic activity, PIM1 was also fused to the transformation-defective MYC deleted in the MBII domain. Interestingly, we found that the fusion PIM1-MYC[?]MBII, but not co-transfection of MYC[?]MBII with PIM1, was able to rescue, at least in part, the defective MYC-transformation (Fig. 8b). Importantly, this fusion rescues both transcriptional activity and ability to phosphorylate H3S10 at its recruiting sites (Fig. 8d, f). In contrast, a chimeric molecule in which PIM1 was fused to a MYC[?]C, a mutant unable to bind to the DNA, was defective for transforming activity and for phosphorylation of H3S10 at the target genes analysed (see Supplementary Information Figure S6). Interestingly, the transformed foci induced by the PIM1-MYC fusion grew vigorously in soft agar forming bigger colonies as compared with PIM1KD-MYC (see Supplementary Information Fig. S6).

Finally, we tested whether H3 phosphorylation at S10 is required for MYC-dependent transformation by analysing MYC transforming potential in the presence of either yellow fluorescent protein (YFP) fused to H3 (YFP-H3) or to H3 mutated in S10 (YFP-H3S10A). Both fusion proteins co-localised in the nucleus during interphase and mitosis together with the endogenous H3S10ph (see Supplementary Information Fig. S7) and they did not interfere with cell cycle progression (see Supplementary Information Fig. S7) as previously reported for other histone fusion proteins³⁷. Then we performed transformation assays by either expressing MYC, PIM1 or PIM1-MYC fusion proteins in these cells (see Supplementary Information Fig. S8). We observed that, while YFP-H3 did not affect the number of foci, YFP-H3S10A mutant significantly reduced the number of soft agar

colonies, FOSL1 and ID2 transcription as well as H3 phosphorylation at their MYC binding sites (see Supplementary Information Fig. S8).

Taken together these results show that H3 phosphorylation at MYC-target genes by PIM1 is required for the cooperative effects of PIM1 and MYC in cell transformation.

DISCUSSION

This study defines a new role for the serine/threonine kinase PIM1 as a MYC-dependent modifier of the chromatin. Based on biochemical and functional evidence, we show that following growth factor stimulation a MYC/MAX/PIM1 complex, formed in the nucleus, recruits PIM1 to the chromatin where it phosphorylates H3 at S10. This PIM1-dependent nucleosome phosphorylation is required for the transcriptional activation of a subset of MYC-target genes and MYC-dependent cell transformation.

Immunoprecipitation experiments showed that following growth factor stimulation MYC, in conjunction with MAX, forms a complex with PIM1 in the nucleus. Silencing experiments demonstrated that MYC is required for PIM1 recruitment to the chromatin while PIM1 is necessary to phosphorylate the H3 at S10 on the nucleosome at E boxes to activate the transcription. According to the current view phosphorylation at the N-terminal domain of H3 could either be required to loosen the interaction between DNA and nucleosome and/or to generate a platform to recruit additional regulatory factors as described in the “histone code” hypothesis¹. PIM1-dependent phosphorylation at the

FOSL1 and ID2 genes is specific for its recruitment sites and does not spread over the genes, suggesting that this phosphorylation is necessary to trigger the enhancer to its active state.

It was previously shown that upon growth factors treatment H3 is phosphorylated at S10 with fast kinetics by MSK1/MSK2⁴⁻⁷. Interestingly, FOSL1 shows two regions with different phosphorylation kinetics of H3S10: a phosphorylation site at the upstream SRE element, occurring within 15-30 minutes due to MSK1/MSK2, and a phosphorylation at the enhancer that was coincident with the peak of transcriptional activation of the gene mediated by PIM1. These two regions behaved differently with respect to acetylation of H3 at K9 and K14. While at the FOSL1 promoter H3 phosphorylation and acetylation are tightly coupled as previously shown for other IE promoters^{3,38-40}, at the FOSL1 enhancer, phosphorylation of H3S10 takes place in a region highly acetylated at H3 prior to MYC and PIM1 binding in agreement with the finding that for MYC recognition its target sites must be highly acetylated in H3 as well as methylated in H3K4/K79⁴¹. On the FOSL1 enhancer phosphorylation and acetylation of H3 might have distinct functional roles as PIM1-dependent H3S10 phosphorylation is transient and limited to the E box element while acetylation at K9 and K14 is more stable and distributed along the gene suggesting that phosphorylation at these elements represents a critical step for the transcriptional activation. The fast turnover of H3 phosphorylation observed in our experiments and its temporal coincidence with transcriptional activation might be a general phenomenon as H3

phosphorylation was previously detected with the transcriptional activation of heat shock genes in *Drosophila* and with the RAR[?]2 gene induction by retinoic agonists in murine P19 embryonal cells while acetylation was constitutive at these loci^{42,43}.

It has been estimated that about 11% of cellular genes present a functional E box to which MYC can associate on the genome²⁸. MYC silencing demonstrated that MYC recruits PIM1 at specific MYC binding sites and confocal microscopy showed an elevated degree of nuclear co-localization of PIM1 with nascent transcripts and with MYC following growth factor treatment, suggesting that, at this stage of the cell cycle, PIM1 is recruited by MYC to a large number of sites. Gene expression profile analysis of MYC and PIM1 regulated genes in cells treated with serum at 120 minutes, revealed that 207 transcripts, corresponding to 20% of MYC regulated genes, also require PIM1 for their regulation. The co-regulation includes genes involved in cell metabolism, protein synthesis, cycle progression, and oncogenesis. Interestingly, a large number of genes are transcriptional factors, which suggests that PIM1 participates in MYC-dependent regulatory networks. In agreement with these data cell cycle analysis, in PIM1 knockdown cells, revealed reduced entry in the S phase. Transformation analysis showed a more dramatic PIM1 effect as PIM1 silencing strongly inhibited the formation of MYC-dependent transformed soft agar colonies. Although our experiments do not exclude that specific PIM1-regulated genes cooperate with MYC in cell transformation, our data strongly suggest that PIM1-dependent phosphorylation of H3 at MYC-target genes is necessary to regulate key genes required for MYC-dependent oncogenic transformation. In fact, PIM1 fusion to the transformation

defective MYC^ΔMBII could significantly rescue its transforming potential as well as H3 phosphorylation at MYC binding sites while PIM1 co-expression with MYC^ΔMBII did not rescue.

In conclusion, this study provides a molecular mechanism for MYC and PIM1 cooperation in gene regulation and oncogenic transformation. As mutations that alter MYC expression are among the most common found in human and animal cancers⁴⁴, it is conceivable that inhibiting MYC association with PIM1 and/or inhibiting PIM1 kinase activity by specific drugs might represent a method for the treatment of cancers in humans.

METHODS

DNA constructs. Human PIM1 and MYC cDNA were cloned into pcDNA5/FRT (Invitrogen) vector under the control of the CMV/TetO2 tetracycline inducible promoter. The PIM1-K67M construct was generated by introducing a site-specific mutation in the DNA sequence corresponding to the lysine 67 to mutate it into a methionine (K67M). The MYC^ΔMBII construct was obtained by deleting the DNA sequence corresponding to the amino acids 119-152. The MYC^ΔC construct was obtained by deleting the DNA sequence corresponding to the amino acids 353-439. The PIM1-MYC fusion constructs were obtained by fusing the cDNA corresponding to PIM1 or PIM1-K67M via a 10 glycine encoding linker sequence to the cDNA corresponding either to MYC, MYC^ΔMBII or MYC^ΔC.

shRNA constructs were obtained by cloning a double-stranded DNA cassettes into the

pENTR/H1/TO vector, which allows the expression of shRNA under the control of the tetracycline inducible H1 promoter/TetO2.

To generate shRNA cassettes the following oligonucleotides were designed following the manufacturer's protocol. The 21-nucleotides positive strands are:

PIM1 shRNA#1 (shPIM1#1): 5'-GTCTCTTCAGAGTGTCAGCAC-3'

PIM1 shRNA scrambled control#1 (shsP#1): 5'-GACTCATCAGTATGTTAGCAT-3'

PIM1 shRNA#2 (shPIM1#2): 5'-GAGGAAGAGAGTATCTATGGG

PIM1 shRNA scramble control#2 (shsP#2): 5'-GAAGCCGAGAGTATCTATGTT -3'

PIM1 shRNA#3 (shPIM1#3): 5'-CGAAGAAATCCAGGACCATCC -3'

PIM1 shRNA scrambled control#3 (shsP#3): 5'-AAGTCATCCGAAGACCACAAC-3'

PIM1 shRNA#4 (shPIM1#4): 5'-ACGTGGAGAAGGACCGGATTT-3'

PIM1 shRNA scrambled control #4 (shsP#4): 5'-GATTTCGAGGACATGGACTGAG-3'

c-MYC shRNA#1 (shMYC#1): 5'-GCAATCACCTATGAACTTGTT-3'

c-MYC shRNA scramble control#1(shsM#1): 5'-GCATTCACCTAAGAAGTTCTT-3'

c-MYC shRNA#2 (shMYC#2): 5'- GACGACGAGACCTTCATCAAA -3'

c-MYC shRNA2 scramble control #2 (shMYC#2): 5'-AGATCCAACCAGTCAGGACAT-
3'

c-MYC shRNA#3 (shMYC#3): 5'-CCTGAGACAGATCAGCAACAA-3'

c-MYC shRNA scramble control#3 (shsM#3): 5'-CAACATGGTAACGAACAGCAA -3'

c-MYC shRNA#4 (shMYC#4): 5'-CAGTTGAAACACAAACTTGAA-3'

c-MYC shRNA scramble control#4(shsM#4): 5'-AACTAAGGTA ACTATGCACAA-3'

c-MYC shRNA#5 (shMYC#5): 5'-CCATAATGTAAACTGCCTCAA-3'

c-MYC shRNA scramble control #5(shsM#5): 5'-GCAATGACCACCTATTCAATA-3'

The recombinant GST-PIM1 construct was obtained by cloning the PIM1 WT and PIM1-K67M cDNA into the bacterial expression vector pGEX-4T and the fused protein was expressed and purified using the standard protocol.

The vector coding for *Xenopus leavis* histones were expressed and purified as previously described⁴⁵. Mutant recombinant histones H3 (H3S10A, H3S28A, H3S10A/S28A) were obtained by site directed mutagenesis as previously described⁴⁶. YFP-H3 was kindly provided by Marco Bianchi and mutagenized as previously described⁴⁶.

Cell culture condition. We generated tetracycline inducible 293 cell line stably expressing PIM1 WT or PIM1-K67M by using the Invitrogen's Flp-T-Rex System.

pCDNA5/FRT/TO expression vector containing the gene of interest was inserted into the cells via Flp recombinase-mediated DNA recombination at the FRT site. The transfected cells were selected on the basis of hygromycin B (Sigma) resistance and pool clones were isolated.

Selected clones were cultivated in Dulbecco modified essential medium (D-MEM) containing 10% fetal bovine serum (FBS; Sigma) and 150µg/ml hygromycin for 2/3 weeks. To perform time course experiments 293 cells were synchronized and then treated with 50% serum shock as described⁴⁷. Primary HUVEC cells were grown in M199 medium

supplemented with 15% fetal bovine serum, brain protein extracts and heparin (50 μ g/ml). The cells were then starved for 24 hours in M199 medium containing 1% FBS and 0.5% BSA and induced with 50 ng/ml of VEGF-A. The cell line P493-6 (kindly provided by Bruno Amati) were grown in RPMI 1640 containing 10% fetal calf serum (FCS) 100 U/ml penicillin, 100 μ g/ml streptomycin 2mM L-glutamine. MYC repression was obtained with 0.1 μ g/ml tetracycline. MYC induction was obtained by washing the cells three times with tetracycline free phosphate buffered saline containing 10% FCS as previously described²⁷.

Immunostaining and co-localization analysis. Co-localization experiments were performed on fixed cells as previously described¹¹ using a Leica TCS SP2 confocal microscopy. To label nascent RNA, anti-BrUTP (Sigma cat. B2531) has been utilized in permeabilized cells by immunodetection with small modifications from previously described protocol⁴⁸. The degree of double staining was measured by quantitative analysis of the overlapping pixel using the Image J 1.32j (NIH, USA) software. All pixels having the same position in both images are considered coincident and the brightness level of each pixel is plotted as x and y axis of the scatter diagram. Two completely overlapping images will result in a straight diagonal line in scatter diagram.

Protein immunoprecipitation, interaction, kinase assays, chromatin

immunoprecipitation. Proteins were extracted using F-buffer (10mM TRIS pH7.0, 50mM NaCl, 30mM Na₂HPO₄, 50mM NaF, 5μM ZnCl₂, 1% Triton-X 100) and treated with 1U/ml Micrococcal nuclease (Sigma) at 4⁰C for 1hr to obtain the chromatin proteins (Chromatin fraction).

To identify interacting protein, 0.5 mg of total proteins was incubated with 3 μg of specific antibody overnight at 4° C. Immunocomplexes were incubated with Protein A- or G-conjugated magnetic beads (DYNAL) for 30 minutes at 4°C. Samples were washed four times with F buffer and 2 times with 0,15M NaCl F-buffer for 20 minutes at 4°C. Proteins were eluded by boiling the samples and analysed by western blotting.

GST-interaction assay was performed by incubating 1mg of protein extracts (derived from chromatin fractions) with recombinant proteins GST, GST-PIM1WT and GST-PIM1-K67M bound to Glutathione-sepharose resin (Amersham). To allow specific interactions, proteins were incubated in Interaction buffer (TRIS 20mM; KCl 200mM; DTT 10mM; EGTA 0,1mM; Glycerol 20%; Inhibitor protease cocktail) for 2 hours at 4°C in presence of 21-aa peptides. Unbound proteins were eliminated by 3 washes in the same buffer followed by protein equilibration in the Kinase buffer (TRIS 30mM; DTT 10mM; EGTA 0,1mM MnCl 2mM; Glycerol 10%). Kinase assay was performed by incubating samples in the same buffer in presence of 4 pmol of [³²P] ATP for 2 minutes at 30°C. Reaction was stopped by adding 10mM EDTA and samples were washed 6 times with Kinase buffer containing 10mM EDTA and 200mM KCl. Phosphorylated proteins were separated by SDS-PAGE

and visualized by autoradiography.

Nucleosome core particles (NCPs) reconstitution was carry out as described⁴⁹.

Chromatin Immunoprecipitation (ChIP) assays were performed as described by Upstate protocols (www.upstate.com). DNA was analyzed by quantitative real-time PCR by using Light Cycler DNA Amplification Kit SYBR Green I (Roche, Mannheim, Germany). All experiment values were normalized to those obtained with a non-immune serum and divided by input, using the procedure previously described⁵⁰. Experiments of MSK1/MSK2 inhibition were performed by incubating cells with H89 (Calbiochem) as previously described⁵. Oligonucleotide sequences will be provided upon request.

Quantitative RT-PCR analysis was performed as previously described¹¹.

Cell transformation and colony formation in soft agarose

The analysis of cell transformation and anchorage independent growth were performed as previously described³⁶. Cells transfected with the constructs indicated were

suspended at approximately 1×10^4 cells in 2 ml of 0.35% low-melting agarose onto 35-mm culture dish containing 0.7% agarose base. Triplicates were prepared for each tested construct and inspected by eye after crystal violet staining for colony formation after incubation at 37⁰C for 10-14 days. Efficiency of transfection was measured by reporter

GFP gene and the levels of proteins were compared by Western blotting.

Antibodies

The antibodies used for in this work are purchased by Santa Cruz Biotechnology (anti-PIM1 SC-7856 for ChIP and immunofluorescence experiments; anti-PIM1 SC-13513 for Western blot analysis; anti-Myc SC-764 for ChIP and immunofluorescence experiments and anti-Myc SC-40 for Western blot analysis; anti-MAX SC-197 and SC-765; anti-SP1 SC-59; anti-Pol II SC-899), by Covance anti Pol II Ser2 H5, by Upstate (anti-Histone H3 06-755; anti-acetyl Histone H3 06-599; anti-phospho (Ser10)-acetyl (Lys14)-Histone H3 07-081; anti-phospho (Ser10)- Histone H3 05-817), and by Abcam anti phospho (Ser10)-acetyl (Lys9)-Histone H3 Ab 12181-50.

ACKNOWLEDGEMENTS

This work was supported by Associazione Italiana Ricerca sul cancro (AIRC), Ministero Italiano Università e Ricerca (MIUR), and Fondazione Monte dei Paschi di Siena. We are indebted to all members of the laboratory for reagents, helpful suggestions, and encouragement; Marina Rocchigiani for VEGF-A preparations and Beatrice Grandi for technical support. Marco Bianchi for YFP-H3 construct, Roberto Mantovani for *Xenopus leavis* histone constructs, Bruno Amati for Rat-1 and P493-6 cells. Isabel Delany, Marco Bianchi, and Ernesto Guccione for the critical reading of the manuscript.

AUTHOR CONTRIBUTION

A.Z. planned and performed the experiments and analysed the data. A.D. generated stable clones and performed immunoprecipitation experiments. R.S. performed the transformation assays. S.O. planned the

experimental design, analysed the data and wrote the manuscript.

COMPETING FINANCIAL INTEREST

The authors declare that they have no competing financial interests.

REFERENCES

1. Strahl, B.D. & Allis, C.D. The language of covalent histone modifications. *Nature* **403**, 41-5 (2000).
2. Berger, S.L. Histone modifications in transcriptional regulation. *Curr Opin Genet Dev* **12**, 142-8 (2002).
3. Cheung, P., Allis, C.D. & Sassone-Corsi, P. Signaling to chromatin through histone modifications. *Cell* **103**, 263-71 (2000).
4. Nowak, S.J. & Corces, V.G. Phosphorylation of histone H3: a balancing act between chromosome condensation and transcriptional activation. *Trends Genet* **20**, 214-20 (2004).
5. Thomson, S., Mahadevan, L.C. & Clayton, A.L. MAP kinase-mediated signalling to nucleosomes and immediate-early gene induction. *Semin Cell Dev Biol* **10**, 205-14 (1999).
6. Sassone-Corsi, P. et al. Requirement of Rsk-2 for epidermal growth factor-activated phosphorylation of histone H3. *Science* **285**, 886-91 (1999).
7. Soloaga, A. et al. MSK2 and MSK1 mediate the mitogen- and stress-induced

- phosphorylation of histone H3 and HMG-14. *Embo J* **22**, 2788-97 (2003).
8. Yamamoto, Y., Verma, U.N., Prajapati, S., Kwak, Y.T. & Gaynor, R.B. Histone H3 phosphorylation by IKK-alpha is critical for cytokine-induced gene expression. *Nature* **423**, 655-9 (2003).
 9. Anest, V. et al. A nucleosomal function for IkappaB kinase-alpha in NF-kappaB-dependent gene expression. *Nature* **423**, 659-63 (2003).
 10. Cuypers, H.T. et al. Murine leukemia virus-induced T-cell lymphomagenesis: integration of proviruses in a distinct chromosomal region. *Cell* **37**, 141-50 (1984).
 11. Zippo, A., De Robertis, A., Bardelli, M., Galvagni, F. & Oliviero, S. Identification of Flk-1-target genes in vasculogenesis: Pim-1 is required for endothelial and mural cell differentiation in vitro. *Blood* **103**, 4536-44 (2004).
 12. van Lohuizen, M. et al. Predisposition to lymphomagenesis in pim-1 transgenic mice: cooperation with c-myc and N-myc in murine leukemia virus-induced tumors. *Cell* **56**, 673-82 (1989).
 13. van Lohuizen, M. et al. Identification of cooperating oncogenes in E mu-myc transgenic mice by provirus tagging. *Cell* **65**, 737-52 (1991).
 14. Verbeek, S. et al. Mice bearing the E mu-myc and E mu-pim-1 transgenes develop pre-B-cell leukemia prenatally. *Mol Cell Biol* **11**, 1176-9 (1991).
 15. Wang, Z. et al. Pim-1: a serine/threonine kinase with a role in cell survival, proliferation, differentiation and tumorigenesis. *J Vet Sci* **2**, 167-79 (2001).

16. Ellwood-Yen, K. et al. Myc-driven murine prostate cancer shares molecular features with human prostate tumors. *Cancer Cell* **4**, 223-38 (2003).
17. Amati, B., Alevizopoulos, K. & Vlach, J. Myc and the cell cycle. *Front Biosci* **3**, d250-68 (1998).
18. Grandori, C., Cowley, S.M., James, L.P. & Eisenman, R.N. The Myc/Max/Mad network and the transcriptional control of cell behavior. *Annu Rev Cell Dev Biol* **16**, 653-99 (2000).
19. Levens, D.L. Reconstructing MYC. *Genes Dev* **17**, 1071-7 (2003).
20. Murphy, M.J., Wilson, A. & Trumpp, A. More than just proliferation: Myc function in stem cells. *Trends Cell Biol* **15**, 128-37 (2005).
21. Grandori, C. et al. c-Myc binds to human ribosomal DNA and stimulates transcription of rRNA genes by RNA polymerase I. *Nat Cell Biol* **7**, 311-8 (2005).
22. Arabi, A. et al. c-Myc associates with ribosomal DNA and activates RNA polymerase I transcription. *Nat Cell Biol* **7**, 303-10 (2005).
23. Trumpp, A. et al. c-Myc regulates mammalian body size by controlling cell number but not cell size. *Nature* **414**, 768-73 (2001).
24. Adhikary, S. & Eilers, M. Transcriptional regulation and transformation by Myc proteins. *Nat Rev Mol Cell Biol* **6**, 635-45 (2005).
25. McMahon, S.B., Van Buskirk, H.A., Dugan, K.A., Copeland, T.D. & Cole, M.D. The novel ATM-related protein TRRAP is an essential cofactor for the c-Myc

- and E2F oncoproteins. *Cell* **94**, 363-74 (1998).
26. Osborne, C.S. et al. Active genes dynamically colocalize to shared sites of ongoing transcription. *Nat Genet* **36**, 1065-71 (2004).
 27. Schuhmacher, M. et al. The transcriptional program of a human B cell line in response to Myc. *Nucleic Acids Res* **29**, 397-406 (2001).
 28. Fernandez, P.C. et al. Genomic targets of the human c-Myc protein. *Genes Dev* **17**, 1115-29 (2003).
 29. Bergers, G., Graninger, P., Braselmann, S., Wrighton, C. & Busslinger, M. Transcriptional Activation of the Fra-1 gene by AP-1 is mediated by regulatory sequences in the first intron. *Mol. Cell. Biol.* **15**, 3748-3758 (1995).
 30. Adisheshaiah, P., Peddakama, S., Zhang, Q., Kalvakolanu, D.V. & Reddy, S.P. Mitogen regulated induction of FRA-1 proto-oncogene is controlled by the transcription factors binding to both serum and TPA response elements. *Oncogene* **24**, 4193-205 (2005).
 31. Blackwell, T.K. et al. Binding of myc proteins to canonical and noncanonical DNA sequences. *Mol Cell Biol* **13**, 5216-24 (1993).
 32. Lasorella, A., Nosedà, M., Beyna, M., Yokota, Y. & Iavarone, A. Id2 is a retinoblastoma protein target and mediates signalling by Myc oncoproteins. *Nature* **407**, 592-8 (2000).
 33. Zeller, K.I. et al. Global mapping of c-Myc binding sites and target gene networks in human B cells. *Proc Natl Acad Sci U S A* **103**, 17834-9 (2006).

34. Schuhmacher, M. et al. Control of cell growth by c-Myc in the absence of cell division. *Curr Biol* **9**, 1255-8 (1999).
35. Pajic, A. et al. Cell cycle activation by c-myc in a burkitt lymphoma model cell line. *Int J Cancer* **87**, 787-93 (2000).
36. Eilers, M., Picard, D., Yamamoto, K.R. & Bishop, J.M. Chimaeras of myc oncoprotein and steroid receptors cause hormone-dependent transformation of cells. *Nature* **340**, 66-8 (1989).
37. Kanno, T. et al. Selective recognition of acetylated histones by bromodomain proteins visualized in living cells. *Mol Cell* **13**, 33-43 (2004).
38. Barratt, M.J., Hazzalin, C.A., Cano, E. & Mahadevan, L.C. Mitogen-stimulated phosphorylation of histone H3 is targeted to a small hyperacetylation-sensitive fraction. *Proc Natl Acad Sci U S A* **91**, 4781-5 (1994).
39. Cheung, P. et al. Synergistic coupling of histone H3 phosphorylation and acetylation in response to epidermal growth factor stimulation. *Mol Cell* **5**, 905-15 (2000).
40. Clayton, A.L., Rose, S., Barratt, M.J. & Mahadevan, L.C. Phosphoacetylation of histone H3 on c-fos- and c-jun-associated nucleosomes upon gene activation. *Embo J* **19**, 3714-26 (2000).
41. Guccione, E. et al. Myc-binding-site recognition in the human genome is determined by chromatin context. *Nat Cell Biol* (2006).
42. Nowak, S.J. & Corces, V.G. Phosphorylation of histone H3 correlates with

- transcriptionally active loci. *Genes Dev* **14**, 3003-13 (2000).
43. Lefebvre, B., Ozato, K. & Lefebvre, P. Phosphorylation of histone H3 is functionally linked to retinoic acid receptor beta promoter activation. *EMBO Rep* **3**, 335-40 (2002).
 44. Cole, M.D. & McMahon, S.B. The Myc oncoprotein: a critical evaluation of transactivation and target gene regulation. *Oncogene* **18**, 2916-24 (1999).
 45. Caretti, G., Motta, M.C. & Mantovani, R. NF-Y associates with H3-H4 tetramers and octamers by multiple mechanisms. *Mol Cell Biol* **19**, 8591-603 (1999).
 46. Salameh, A., Galvagni, F., Bardelli, M., Bussolino, F. & Oliviero, S. Direct recruitment of CRK and GRB2 to VEGFR-3 induces proliferation, migration, and survival of endothelial cells through the activation of ERK, AKT, and JNK pathways. *Blood* **106**, 3423-31 (2005).
 47. Balsalobre, A., Damiola, F. & Schibler, U. A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell* **93**, 929-37 (1998).
 48. Zhang, S., Herrmann, C. & Grosse, F. Nucleolar localization of murine nuclear DNA helicase II (RNA helicase A). *J Cell Sci* **112 (Pt 16)**, 2693-703 (1999).
 49. Cirillo, L.A. & Zaret, K.S. Preparation of defined mononucleosomes, dinucleosomes, and nucleosome arrays in vitro and analysis of transcription factor binding. *Methods Enzymol* **375**, 131-58 (2004).
 50. Kouskouti, A. & Talianidis, I. Histone modifications defining active genes

persist after transcriptional and mitotic inactivation. *Embo J* **24**, 347-57 (2005).

FIGURE LEGENDS

Figure 1. PIM1 phosphorylates H3S10 on the nucleosome.

(a) PIM1 phosphorylates histone H3 on nucleosomes from chromatin fraction. GST-recombinant proteins, as indicated, were incubated with purified nucleosomes from 293 cells and the kinase reactions were carried out in the presence of [γ - 32 P] ATP. GST-PIM1 interactions with nucleosomes were performed in the absence (-) or presence of 20 fold molar excess of unrelated (unr), H3 or H2B peptides as indicated. GST recombinant proteins and nucleosomes were visualized by Coomassie staining. Histone phosphorylation was revealed by autoradiography. H3 but not H2B N-terminal domain competes for PIM1 association to the nucleosome inhibiting the kinase reaction. (Full scans are shown in Supplementary Information Figure S9).

(b) PIM1 phosphorylates histone H3 at serine 10. Recombinant GST-PIM1 or the PIM1 kinase inactive mutant were incubated with nucleosomes reconstituted *in vitro* from recombinant histones. Nucleosomes were assembled containing respectively the histone H3 wild type (lanes 1, 2), the mutant H3S10A (lanes 3,4), the mutant H3S28A (lanes 5,6) or the double mutant H3S10A/S28A (lanes 7,8). Histone H3 autoradiography resulting from [γ - 32 P] ATP incorporation (first panel)

is shown. The input kinases and nucleosomes were visualized by Coomassies staining (lower panels).

(c) PIM1 interacts with histone H3 *in vivo*. PIM1 was immunoprecipitated from 293 stable clones expressing the inducible PIM1 (WT) or the kinase inactive mutant PIM1-K67M (K67M) as indicated. PIM1 and H3 were revealed by immunoblotting (IB) with specific antibodies. A kinase reaction (KA) was carried out incubating the PIM1/H3 co-immunoprecipitate in the presence of [γ - 32 P] ATP. Histone phosphorylation was visualized by autoradiography. 2% of the total immunoprecipitated protein samples were loaded as inputs (lower panel). (Full scans are shown in Supplementary Information Figure S9).

Figure 2. PIM1 forms a complex with MYC.

(a) PIM1 co-localizes with MYC in HUVEC upon VEGF-A treatment. Indirect immunofluorescence analysis of HUVEC treated with VEGF was performed by using antibodies against PIM1 (red) and MYC (green) and double immunostaining are shown (Scale bar = 70 μ m).

(b) VEGF-A treatment induced interaction of endogenous PIM1 with the MYC/MAX dimer. Cell proteins extracted from VEGF-A treated HUVEC were either immunostained (INPUT) or subjected to immunoprecipitation (IP) with anti-IgG antibodies (-), or anti-MAX antibodies as indicated (+). Immunostaining (IB) analysis was performed by using anti-MAX, -MYC and -PIM1 antibodies as

indicated. 10% of the total immunoprecipitated protein samples were loaded as inputs. (Full scans are shown in Supplementary Information Figure S9).

(c) Endogenous PIM1 forms a complex with MYC/MAX in serum-stimulated 293. Western Blot analysis of input extracts, IgG, MAX, MYC or PIM1 immunoprecipitates from serum induced or not induced as indicated. Interacting proteins were revealed by using anti-MAX, -MYC, or -PIM1 antibodies respectively.

(d) MYC forms a complex with PIM1 through its MBII domain. 293 were transiently co-transfected with FLAG-MYC constructs as indicated and PIM1. Cell lysates were immunoprecipitated with anti-IgG, anti-FLAG, or anti-PIM1 antibodies followed by Western blot analysis to detect MAX, MYC, PIM1 and TRRAP proteins as indicated. Asterisk indicated an unspecific band.

Figure 3. PIM1 co-localizes with H3S10ph, active chromatin, and MYC.

(a) Immunofluorescence analysis of PIM1 nuclear sub-localization in HUVEC treated with VEGF for 60 minutes. Fixed cells were double-stained for PIM1 (red) and one of the indicated markers (green) (Scale bar = 8 μ m). The corresponding pictures were merged and all pixels having the same position in both images are considered coincident and were represented as white spots (co-localization points). The brightness level of each pixel is plotted as x (PIM1) and y (marker) axis of the scatter diagram of the correlation plot. The pixels co-localization results of 7 individual images represented in the correlation plot were quantified as

Pearson's correlation coefficient (R_r) as mean \pm SD with P values determined using Student's *t*-test.

(b) Immunofluorescence analysis of MYC localization at transcriptional active sites. Cells were labeled with BrUTP and detected with anti-bromodeoxyuridine antibody (green). Double labeling was revealed at transcriptional active sites by staining cells with an anti-MYC antibody (red). The pixels co-localization were performed as in (a).

Figure 4. PIM1 associates to chromatin at the FOSL1 enhancer.

(a) FOSL1 is transiently induced by VEGF-A treatment. Serum-starved HUVEC were treated with VEGF-A and total RNAs were collected at different time points. FOSL1 transcript levels were quantified by quantitative RT-PCR and relative values normalized on GAPDH expression level. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$). **(b)** Schematic representation of the FOSL1 gene region including the first two exons (not in scale). Exons are represented as black squares within the gene and the E box is figured as a black spot. The probes used in ChIP analysis are indicated. PCR probes intervals are numbered relatively to the first nucleotide of exon 1 (+1). **(c)** VEGF-A induces PIM1 association with chromatin at the FOSL1 enhancer. Protein interactions on FOSL1 chromatin were measured by quantitative ChIP assay on VEGF-A-treated HUVEC. Chromatin extracts were immunoprecipitated using antibodies as indicated. Real-time quantitative PCR measurements of the

immunoprecipitated DNA of the corresponding gene regions identified as probes A, B, C or D at different time points are expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

(d) FOSL1 expression in the presence of MSK1/MSK2 inhibitor H89. Serum-starved HUVEC were treated with VEGF-A in the presence or absence of 10 μ M H89 and total RNAs were collected at different time points. FOSL1 transcript levels were quantified by real-time quantitative RT-PCR and relative values normalized on GAPDH expression level. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

(e) Phosphorylation of H3S10 at the FOSL1 upstream element is dependent from MSK1/MSK2 kinase activities. H3S10 phosphorylation was measured by quantitative ChIP assay on VEGF-A-treated HUVEC. Chromatin extracts were immunoprecipitated using antibodies as indicated. Real-time quantitative PCR measurements of the immunoprecipitated DNA of the corresponding gene regions identified as probes A, B, C or D at different time points are expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

Figure 5. PIM1 mediates H3S10 phosphorylation and contributes to FOSL1 transcription.

(a) Inducible shRNA expression targeting PIM1. Western blot analysis of 293

stable clones expressing a control scrambled shRNA (shsP#1) or PIM1 shRNA (shPIM1#1) inducible by tetracycline. Immunostaining with antibodies as indicated. Protein extracts were obtained from not treated or serum-treated cells as indicated, in presence (Tet +) or absence (Tet -).

(b) PIM1 knockdown alters H3S10 phosphorylation at the FOSL1 enhancer. Probes A, B, C and D are schematized as in Figure 4. Time course analysis by ChIP assay was performed by using antibodies as specified from 293 cells expressing either a control scrambled shRNA or PIM1 shRNA. Real-time quantitative PCR measurements of the immunoprecipitated DNA are expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

Figure 6. PIM1 phosphorylates histone H3 in a MYC-dependent manner.

(a) Inducible MYC gene silencing. 293 were transiently transfected with either a control scrambled shRNA (shsM#1) or MYC shRNA (shMYC#1) as indicated by tetracycline treatment. Western blot analysis on protein extracts was performed by immunoblotting, using antibodies recognising MYC, MAX, or SP1 as indicated.

(b-e) MYC recruits PIM1 on chromatin. Chromatin samples were obtained from serum-induced 293 cells expressing the control scrambled shRNA or MYC shRNA as indicated. ChIP assays were performed using antibodies as indicated. Immunoprecipitated DNA was analyzed by quantitative PCR with primers to amplify the FOSL1 enhancer and ID2 upstream MYC-binding sites (-1705/-1460).

Real-time quantitative PCR measurements of the immunoprecipitated DNA are expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

(f) Western blot analysis of 293 transiently transfected with control scrambled shRNA (shsM#1), MYC shRNA (shMYC#1), or the shRNA together with plasmids expressing either FLAG-MYC or FLAG-MYC[?]MBII as indicated. shMYC#1 recognizes MYC 3'-untranslated region which is not present in the FLAG constructs co-transfected. Protein expression was performed by immunoblotting, using antibodies as indicated.

(g-h) MYC but not FLAG-MYC[?]MBII rescues the recruitment of PIM1 to the chromatin. Chromatin samples were obtained from 293 cells expressing the control scrambled shRNA, MYC shRNA, or the shRNA together with FLAG-MYC or FLAG-MYC[?]MBII (F-MYC[?]MBII) as indicated. CHIP assays were performed using antibodies as indicated. Immunoprecipitated DNA was analyzed by quantitative PCR with primers to amplify the FOSL1 enhancer and ID2 upstream MYC-binding sites (-1705/-1460). Real-time quantitative PCR measurements of the immunoprecipitated DNA and expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

Figure 7. PIM1 is required for the transcriptional activation of FOSL1 and ID2.

(a, d) Schematic representation of the FOSL1 and ID2 gene regions (not in scale).

Exons are represented as black squares E boxes are figured as a black spot. The probes used for ChIP analysis are indicated. PCR probes are numbered relatively to the first nucleotide of exon 1 (+1).

(b) Time-course analysis by quantitative real time RT-PCR of FOSL1 mRNA and unprocessed heterogeneous nuclear FOSL1 RNA (hnRNA) in 293 cells expressing either a control scrambled shRNA or PIM1 shRNA as indicated. Gene specific RNA levels were measured and normalized on GAPDH expression and represented as fold induction (vertical bars). Real-time quantitative RT-PCR measurements. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

(c) Time course analysis by ChIP assay was performed by using antibodies recognising RNA Polymerase II (Pol II) or RNA Polymerase II phosphorylated in its C terminal domain at Serine 2 (Ser2P) from 293 cells expressing either a control scrambled shRNA or PIM1 shRNA at the FOSL1 promoter or coding region as specified in panel A. Real-time quantitative PCR measurements of the immunoprecipitated DNA and expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

(e) Time-course analysis by quantitative real time RT-PCR of ID2 mRNA and ID2 hnRNA in 293 cells expressing either a control scrambled shRNA (shsP#1) or PIM1 shRNA (shPIM1#1). RNA levels were measured and normalized on GAPDH expression and represented as fold induction (vertical bars) by Real-time

quantitative RT-PCR measurements. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

(f) Time course analysis by ChIP assay was performed by using antibodies recognising RNA Polimerase II (Pol II) or RNA Polimerase II phosphorylated in its C terminal domain at Serine 2 (Ser2P) from 293 cells expressing either a control scrambled shRNA or PIM1 shRNA at the ID2 promoter or coding region as specified in panel D.

Real-time quantitative PCR measurements of the immunoprecipitated DNA are expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

Figure 8. PIM1 contributes to MYC-dependent transformation. (a b) Soft agar colony formation of Rat-1 cell lines. Rat-1 fibroblasts were transfected with the constructs indicated. Transforming potential of the transfected constructs was determined by examining the soft agar colonies formed within two weeks post-transfection. Transforming potential was estimated by the average number of visible soft agar colonies observed in three plates. A representative experiment is shown.

(c-d) Quantitative real time RT-PCR of FOSL1 and ID2 mRNA in Rat-1 cells expressing the constructs indicated. Gene specific RNA levels were measured and normalized on GAPDH expression and represented as fold induction (vertical bars)

by real-time quantitative RT-PCR measurements. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).

(e-f) ChIP assay performed in Rat-1 cells by using antibodies as indicated. The primers used for the FOSL1 gene correspond to probe B shown in Figure 4 and for the ID2 MYC binding sites as in Figure 7. Real-time quantitative PCR measurements of the immunoprecipitated DNA are expressed as 1/1000 (‰) of the DNA inputs. The values represent the mean s. d. of three replicate samples from one representative experiment ($n = 3$).