

Metadata of the chapter that will be visualized in SpringerLink

Book Title	Engineering Geology for Society and Territory - Volume 1	
Series Title		
Chapter Title	Metamorphic CO ₂ Degassing in the Active Himalayan Orogen: Exploring the Influence of Orogenic Activity on the Long-Term Global Climate Changes	
Copyright Year	2014	
Copyright HolderName	Springer International Publishing Switzerland	
Corresponding Author	Family Name	Franco
	Particle	
	Given Name	Rolfo
	Prefix	
	Suffix	
	Division	Department of Earth Sciences
	Organization	University of Torino
	Address	10125, Turin, Italy
	Division	
	Organization	IGG – CNR
	Address	10125, Turin, Italy
	Email	franco.rolfo@unito.it
Author	Family Name	Chiara
	Particle	
	Given Name	Groppo
	Prefix	
	Suffix	
	Division	Department of Earth Sciences
	Organization	University of Torino
	Address	10125, Turin, Italy
	Email	
Author	Family Name	Pietro
	Particle	
	Given Name	Mosca
	Prefix	
	Suffix	
	Division	
	Organization	IGG – CNR
	Address	10125, Turin, Italy
	Email	
Author	Family Name	Simona
	Particle	
	Given Name	Ferrando
	Prefix	

Suffix
Division Department of Earth Sciences
Organization University of Torino
Address 10125, Turin, Italy
Email

Author Family Name **Emanuele**
Particle
Given Name **Costa**
Prefix
Suffix
Division Department of Earth Sciences
Organization University of Torino
Address 10125, Turin, Italy
Email

Author Family Name **Kaphle**
Particle
Given Name **Krishna P.**
Prefix
Suffix
Division
Organization Nepal Geological Society
Address Kathmandu, Nepal
Email

Abstract A number of studies suggest that mountain ranges have strong impact on the global carbon cycle; metamorphic degassing from active collisional orogens supplies a significant fraction of the global solid-Earth derived CO₂ to the atmosphere, thus playing a fundamental role even in today's Earth carbon cycle. The Himalayan belt, a major collisional orogen still active today, is a likely candidate for the production of a large amount of metamorphic CO₂ that may have caused changes in long-term climate of the past, present and near future. Large metamorphic CO₂ fluxes are facilitated by rapid prograde metamorphism of big volumes of impure carbonate rocks coupled with facile escape of CO₂ to the Earth's surface. So far, the incomplete knowledge of the nature, magnitude and distribution of the CO₂-producing processes hampered a reliable quantitative modeling of metamorphic CO₂ fluxes from the Himalayan belt. This study, integrated in the framework of the Ev-K2-CNR SHARE (Stations at High Altitude for Research on the Environment) Project, focuses on the metamorphic decarbonation processes occurring during the Himalayan collision. We hereby present preliminary results focusing on the distribution of different types of metacarbonate rocks in the Eastern Himalaya, their petrographic description and the first reported petrological data about the nature of the CO₂-producing reactions in garnet-bearing calc-silicate rocks. These results represent a contribution toward a better understanding of the influence exerted by orogenic processes on climatic changes at global scale.

Keywords (separated by '-') Orogenic CO₂ - Climate changes - Decarbonation processes - Himalaya - Metacarbonate rocks



Metamorphic CO₂ Degassing in the Active Himalayan Orogen: Exploring the Influence of Orogenic Activity on the Long-Term Global Climate Changes

Rolfo Franco, Groppo Chiara, Mosca Pietro, Ferrando Simona, Costa Emanuele, and Krishna P. Kaphle

Abstract

A number of studies suggest that mountain ranges have strong impact on the global carbon cycle; metamorphic degassing from active collisional orogens supplies a significant fraction of the global solid-Earth derived CO₂ to the atmosphere, thus playing a fundamental role even in today's Earth carbon cycle. The Himalayan belt, a major collisional orogen still active today, is a likely candidate for the production of a large amount of metamorphic CO₂ that may have caused changes in long-term climate of the past, present and near future. Large metamorphic CO₂ fluxes are facilitated by rapid prograde metamorphism of big volumes of impure carbonate rocks coupled with facile escape of CO₂ to the Earth's surface. So far, the incomplete knowledge of the nature, magnitude and distribution of the CO₂-producing processes hampered a reliable quantitative modeling of metamorphic CO₂ fluxes from the Himalayan belt. This study, integrated in the framework of the Ev-K2-CNR SHARE (Stations at High Altitude for Research on the Environment) Project, focuses on the metamorphic decarbonation processes occurring during the Himalayan collision. We hereby present preliminary results focusing on the distribution of different types of metacarbonate rocks in the Eastern Himalaya, their petrographic description and the first reported petrological data about the nature of the CO₂-producing reactions in garnet-bearing calc-silicate rocks. These results represent a contribution toward a better understanding of the influence exerted by orogenic processes on climatic changes at global scale.

Keywords

Orogenic CO₂ • Climate changes • Decarbonation processes • Himalaya • Metacarbonate rocks

5.1 Introduction and Aim of the Study

Metamorphic degassing from active collisional orogens supplies a significant amount of CO₂ to the atmosphere, playing a fundamental role in the long-term (>1 Ma) global carbon cycle (e.g. Kerrick and Caldeira 1993; Selverstone and Gutzler 1993; Bickle 1996; Berner 1999; Gaillardet and Galy 2008; Evans 2011). The Himalaya is the most prominent collisional orogen on Earth, where tectonic and erosional processes are still active today. Therefore, it is a likely candidate for the production of a large amount of metamorphic CO₂ that may have caused changes in long-term climate of the past and that may still influence the

R. Franco (✉) · G. Chiara · F. Simona · C. Emanuele
Department of Earth Sciences, University of Torino, 10125,
Turin, Italy
e-mail: franco.rolfo@unito.it

R. Franco · M. Pietro
IGG – CNR, 10125, Turin, Italy

K. P. Kaphle
Nepal Geological Society, Kathmandu, Nepal

atmospheric composition in the near future (Gaillardet and Galy 2008). Large metamorphic CO₂ fluxes should have been (and should be) facilitated by rapid metamorphism of large volumes of metacarbonate rocks, coupled with facile escape of CO₂ to the Earth's surface. The nature and magnitude of the metamorphic CO₂ cycle in Himalaya, however, is still poorly known. This study, integrated since 2012 in the framework of the Ev-K2-CNR SHARE (Stations at High Altitude for Research on the Environment) Project, focuses on the metamorphic decarbonation processes occurring during the Himalayan collision. Fieldwork activity is combined with petrographic, petrologic, structural, geochronological geochemical and fluid inclusion studies with the aims of clarifying: (i) abundance and types of CO₂-source rocks, (ii) nature and rate of CO₂-producing reactions, (iii) nature and composition of the released CO₂-rich fluids, (iv) nature and distribution of the CO₂ escape-paths toward the Earth's surface, and (v) chronology of metamorphic CO₂-producing reactions occurred in the Himalayas at different structural levels and at different times. We hereby present a preliminary sketch map reporting the distribution of the different types of metacarbonate rocks in the Eastern Himalaya. The first petrological results on the nature of the CO₂-producing reactions in garnet-bearing calc-silicate rocks are summarized and discussed in the global perspective of the orogenic CO₂-cycle. These results represent a contribution toward a better understanding of the influence exerted by the orogenic processes on climatic changes at global scale.

5.2 Metacarbonate Rocks in Eastern Himalaya

5.2.1 Field Occurrences

In the eastern Himalaya, calc-silicate rocks are widespread in the lower and upper structural levels of the Greater Himalayan Sequence (GHS) (e.g. Goscombe et al. 2006), but they have received so far little notice. Field data acquired in more than 10 years in central-eastern Nepal and Sikkim allowed us to distinguish two different modes of occurrence (Fig. 5.1): (i) in the lower portion of the GHS (GHS-L), calc-silicate rocks generally occur as decimetre to metre-thick levels or boudins (Fig. 5.2a, b) within medium- to high-grade, locally anatectic, staurolite- and/or kyanite-bearing metapelites (e.g. Groppo et al. 2009; Mosca et al. 2012); (ii) structurally upward (GHS-U), calc-silicate rocks are hosted in anatectic kyanite-sillimanite-bearing gneisses (i.e. Barun Gneiss, see Groppo et al. 2012) and often occur as tens to hundreds of meter thick, folded or boudinated, levels occasionally associated to layers of impure marbles

(Fig. 5.2c, d). The transition between the hosting paragneiss and the calc-silicate granofels is generally gradual and is characterized by the progressive disappearance of biotite, the appearance of clinopyroxene and the modal increase of plagioclase. A banded structure is locally observed in the calc-silicate rocks, defined by the different modal proportion of the rock-forming minerals in adjacent layers. This suggests that calc-silicate rocks derive from former marly intercalations within a thick sedimentary sequence.

5.2.2 Petrography

The studied samples have granofelsic structure, and sometimes show evidence of a brittle- to ductile deformation resulting in a local grain size reduction. Mineral assemblages are systematically different in the GHS-L and GHS-U, respectively.

5.2.2.1 Garnet-Bearing Assemblages (GHS-L)

The equilibrium assemblage consists of plagioclase + clinopyroxene + quartz + garnet ± zoisite. Garnet is locally very abundant and it is often intergrown with quartz (Fig. 5.3a, b). Microstructural evidence suggest that garnet grew at the expense of zoisite, clinopyroxene and calcite, the latter being only locally observed as inclusion in garnet. Coarse-grained graphite is locally very abundant and it has been interpreted as precipitated from a H₂O-CO₂ fluid released through decarbonation reactions during prograde and/or early retrograde metamorphic evolution (Groppo et al. 2013). Titanite and apatite are ubiquitous. Thin layers of phlogopite + white mica impure marbles are only occasionally associated to these calc-silicate rocks.

5.2.2.2 K-Feldspar + Scapolite Assemblages (GHS-U)

Calc-silicate rocks from the GHS-U consist of K-feldspar + clinopyroxene + quartz ± scapolite ± calcite, and later plagioclase, epidote, green amphibole and interstitial carbonates (Fig. 5.3c, d). Relict biotite often occurs within clinopyroxene and/or is replaced by K-feldspar. Clinopyroxene is often partially replaced by later green Ca-amphibole ± epidote ± calcite, whereas scapolite is locally partially replaced by fine-grained dusty aggregates of plagioclase + calcite and/or it is overgrown by coarse-grained epidote. In addition to the ubiquitous titanite, a strongly pleochroic allanite and a bluish to colorless tourmaline locally occur, whereas graphite is always absent. The decimetric to metric levels of marbles often intercalated with these calc-silicate granofels are characterized by the same mineral assemblage, but in different modal proportions.




Fig. 5.1 Simplified geological map of the central-eastern sector of the Himalayan belt (modified from Goscombe et al. 2006; Dasgupta et al. 2004; Mosca et al. 2012, 2013) showing sample locations (*squares*: Grt + Zo calc-silicate assemblages and Phl marbles of the GHS-L; *circles*: Kfs + Di + Scp calc-silicate rocks and marbles of the GHS-U). *White lines* are the geotraverses investigated since 2004. The *double-dashed grey line* is the approximate political boundary between

Nepal to the south, west, China (Tibet) to the north, India (Sikkim) to the east. MCT: Main Central Thrust; STDS: South Tibetan Detachment System; E: Everest, K: Kangchenjunga, M: Makalu. Inset shows the location of the study area (*black rectangle*) in the framework of the Himalayan chain. The *grey shaded belt* approximates the location of the Higher Himalayan Crystallines. MFT: Main Frontal Thrust; MBT: Main Boundary Thrust




Fig. 5.2 Field occurrence of calc-silicate rocks from the GHS-L (a, b) and GHS-U (c, d). The *arrows* in a and b indicate the calc-silicate boudins

5.3 CO₂-Producing Processes in Garnet-Bearing Assemblages

The nature of the CO₂-producing reactions in the garnet-bearing calc-silicate rocks from the GHS-L has been petrologically investigated in the CFAS-CO₂-H₂O system using activity-corrected P-T phase diagrams at fixed fluid composition, isobaric T-X(CO₂) phase diagram sections,

and phase diagram projections in which fluid composition is not explicitly constrained (Groppo et al. 2013).

The petrological results allowed to define the P-T-X_{fluid} regime during the metamorphic evolution of the studied calc-silicate rocks. A prograde heating up to peak-T of ca. 800° C (at about 10–11 kbar) involved the growth of grossular-rich garnet (Grs_{67–81}) in equilibrium with quartz at the expenses of zoisite, clinopyroxene and calcite; this

Author Proof

 143
 144
 145
 146
 147
 148
 149
 150

 151
 152
 153
 154
 155
 156
 157
 158




Fig. 5.3 Representative microstructures of calc-silicate granulites from the GHS-L (a, b) and GHS-U (c, d). *Top*: plane polarized light. *Bottom*: crossed polarized light

159 reaction released a CO₂-rich fluid (XCO₂ = 0.20–0.44). A
160 grossular-rich garnet (Grs_{67–76}) additionally grew in equi-
161 librium with plagioclase and quartz during either prograde
162 heating or early decompression. Also this second reaction
163 released a CO₂-rich fluid, with approximately the same
164 composition (XCO₂ = 0.30–0.44) as that released through
165 the first reaction. It follows that the amount of CO₂ released
166 during garnet growth increases with the increase in gros-
167 sular component of garnet (Groppo et al. 2013). The study
168 of primary and secondary carbonic and aqueo-carbonic fluid
169 inclusions occurring in quartz, plagioclase, clinopyroxene
170 and, possibly, garnet and calcite will provide direct con-
171 straints on nature and composition of the involved CO₂-rich
172 fluids.

173 Our results also demonstrate that the studied calc-silicate
174 rocks behaved as a closed-system during their prograde and
175 early retrograde evolution, although their volume abun-
176 dance within the hosting paragneiss is low. In such a closed
177 system, the CO₂-rich fluid released during prograde and
178 early retrograde evolution induced the hydration of the
179 adjacent silicates (mainly plagioclase and clinopyroxene);
180 these hydration reactions, depleting the fluid in H₂O (i.e.
181 enriching the fluid in carbon) triggered the precipitation of
182 graphite.

183 5.4 Conclusions

184

185 In the eastern Himalaya, calc-silicate rocks are widespread
186 in the lower and upper structural levels of the GHS. From
187 field data and petrographic observations, two main calc-
188 silicate assemblages have been recognized, reflecting dif-
189 ferences in the protolith composition: garnet + zoisite
190 assemblages in the GHS-L, and K-feldspar + scapolite

191 assemblages in the GHS-U. Our preliminary petrologic
192 study (Groppo et al. 2013) demonstrates that calc-silicate
193 rocks of appropriate composition may act as CO₂-source
194 during prograde heating and/or early decompression,
195 releasing internal-derived CO₂-rich fluids through garnet-
196 forming reactions. However, if the system remains closed,
197 fluid-rock interactions may induce hydration of the calc-
198 silicate assemblages and the in situ graphite precipitation,
199 thereby removing carbon from the fluid. The interplay
200 between these two contrasting processes—i.e. production of
201 metamorphic CO₂-rich fluids versus carbon sequestration
202 through graphite precipitation—must be taken in account
203 when dealing with a global estimate of the role exerted by
204 decarbonation processes on the orogenic CO₂-cycle.

205 References

206

- 207 Berner RA (1999) A new look at the long-term carbon cycle. *GSA*
208 *Today* 9(11):1–6
- 209 Bickle MJ (1996) Metamorphic decarbonation, silicate weathering and
210 the long-term carbon cycle. *Terra Nova* 8:270–276
- 211 Dasgupta S, Ganguly J, Neogi S (2004) Inverted metamorphic
212 sequence in the Sikkim Himalayas: crystallization history, P–T
213 gradient and implications. *J metamorphic Geol* 22:395–412
- 214 Evans KA (2011) Metamorphic carbon fluxes: how much and how
215 fast? *Geology* 39:95–96
- 216 Gaillardet J, Galy A (2008) Himalaya-carbon sink or source? *Science*
217 320:1727–1728
- 218 Goscombe B, Gray D, Hand M (2006) Crustal architecture of the
219 Himalayan metamorphic front in eastern Nepal. *Gondwana Res*
220 10:232–255
- 221 Groppo C, Rolfo F, Lombardo B (2009) P–T evolution across the Main
222 Central Thrust Zone (Eastern Nepal): hidden discontinuities
223 revealed by petrology. *J Petrol* 50:1149–1180
- 224 Groppo C, Rolfo F, Indares A (2012) Partial melting in the Higher
225 Himalayan Crystallines of Eastern Nepal: the effect of



- 226 decompression and implications for the “channel flow” model. 235
227 J Petrol 53:1057–1088 236
- 228 Groppo C, Rolfo F, Castelli D, Connolly JAD (2013) Metamorphic 237
229 CO₂ production from calc-silicate rocks via garnet-forming 238
230 reactions in the CFAS-H₂O-CO₂ system. Contrib Mineral Petrol 239
231 166:1655–1675 240
- 232 Kerrick DM, Caldeira K (1993) Paleoatmospheric consequences of 241
233 CO₂ released during early Cenozoic regional metamorphism in the 242
234 Tethyan orogen. Chem Geol 108:201–230 243
- Mosca P, Groppo C, Rolfo F (2012) Structural and metamorphic features 235
of the Main Central Thrust Zone and its contiguous domains in the 236
eastern Nepalese Himalaya. J Virtual Expl Electronic Edition, 41:2 237
- Mosca P, Groppo C, Rolfo F (2013) Main geological features of the 238
Rolwaling-Khumbu Himal between the Khimti Khola and Dudh 239
Kholi valleys (eastern-central Nepal Himalaya). Rend Online Soc 240
Geol It 29:112–115 241
- Selverstone J, Gutzler DS (1993) Post-125 Ma carbon storage 242
associated with continent-continent collision. Geology 21:885–888 243

Author Proof

UNCORRECTED PROOF

Author Query Form

244

245

248

246

249

247

250

Book ID : **326568_1_En**
Chapter No.: **5**



251

252

253

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

254

255

Dear Author

256

257

258

During the process of typesetting your chapter, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

259

264

263

266

Query Refs.	Details Required	Author's Response
AQ1	No Queries	

Author Proof

UNCORRECTED PROOF

MARKED PROOF

Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	∧	New matter followed by ∧ or ∧ [Ⓢ]
Delete	/ through single character, rule or underline or ┌───┐ through all characters to be deleted	Ⓞ or Ⓞ [Ⓢ]
Substitute character or substitute part of one or more word(s)	/ through letter or ┌───┐ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↙
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⊕
Change bold to non-bold type	(As above)	⊖
Insert 'superior' character	/ through character or ∧ where required	Υ or Υ under character e.g. Υ or Υ
Insert 'inferior' character	(As above)	∧ over character e.g. ∧
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	ʹ or ʸ and/or ʹ or ʸ
Insert double quotation marks	(As above)	“ or ” and/or “ or ”
Insert hyphen	(As above)	⊞
Start new paragraph	┌	┌
No new paragraph	┐	┐
Transpose	┌┐	┌┐
Close up	linking ○ characters	○
Insert or substitute space between characters or words	/ through character or ∧ where required	Υ
Reduce space between characters or words		↑